



# LUND UNIVERSITY

## Osteoarthritis: Does post-injury ACL reconstruction prevent future OA?

Wen, Chunyi; Lohmander, Stefan

*Published in:*  
Nature Reviews Rheumatology

*DOI:*  
[10.1038/nrrheum.2014.120](https://doi.org/10.1038/nrrheum.2014.120)

2014

[Link to publication](#)

*Citation for published version (APA):*  
Wen, C., & Lohmander, S. (2014). Osteoarthritis: Does post-injury ACL reconstruction prevent future OA? *Nature Reviews Rheumatology*, 10(10), 577-578. <https://doi.org/10.1038/nrrheum.2014.120>

*Total number of authors:*  
2

### General rights

Unless other specific re-use rights are stated the following general rights apply:  
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00



## OSTEOARTHRITIS

Does post-injury ACL reconstruction prevent later OA?

Chunyi Wen and L. Stefan Lohmander

*Young adults with an acute rupture of the anterior cruciate ligament of the knee are faced with the decision of whether or not to undergo early reconstructive surgery. However, a lack of high-quality evidence means questions remain about whether this surgical strategy protects against later development of osteoarthritis.*

Wen, C. & Lohmander, L. S. *Nat. Rev. Rheumatol.* advance online publication XX Month 2014; doi:10.1038/

Return to an active lifestyle and risk of future osteoarthritis (OA) are two, sometimes conflicting, concerns of the young adult with an acute rupture of the anterior cruciate ligament (ACL) of the knee, which could influence the decision of whether or not to undergo early surgery to reconstruct the torn ligament. Brophy and colleagues recently reported the results of a retrospective study into the prevalence of previous knee surgery, such as anterior cruciate ligament (ACL) reconstruction and meniscectomy, in an arthroplasty registry of 1,286 patients with a diagnosis of late-stage OA or post-traumatic arthritis<sup>1</sup>. Strikingly, they found that patients with a history of ACL reconstruction received their knee replacement at ~50 years of age, compared with age ~67 years for those without a history of knee surgery. With knee replacement at a young age markedly increasing the risk of later revision surgery, the results of Brophy *et al.*'s study highlight the problem of OA resulting from knee injury, and raise the question of whether we can prevent this serious late sequel.

The rate of radiographic OA after ACL rupture and reconstruction varies widely in different reports, with a crude estimate of 50% at about 15 years after injury<sup>2</sup>. This high rate of radiographic signs of OA has remained unchanged despite refinements to surgical reconstruction techniques<sup>2</sup>.

The patient-reported outcome of an ACL rupture and reconstruction is influenced by patient-related factors such as sex, BMI, smoking status, pre-surgery activity level, whether the patient returns to sports and the patient's expectations (Figure 1). Trauma-related factors, such as concomitant injuries to the meniscus or joint cartilage, are highly relevant as well<sup>3</sup>. The only recent randomized controlled trial (RCT) to compare early ACL reconstruction plus structured rehabilitation with rehabilitation alone failed to show a difference between patient-reported outcomes of the two strategies at 2 or 5 years<sup>4,5</sup>, suggesting that many patients do as well for at least 5 years without undergoing early surgical reconstruction. Furthermore, no high-level evidence exists to support a protective effect of ACL reconstruction against later development of OA. On the contrary, an RCT comparing early reconstruction and

structured rehabilitation found no difference in the rate of radiographic or clinical signs of OA 5 years after the injury<sup>5</sup>.

The development of OA following an ACL rupture and reconstructive surgery remains an unsolved problem. To better understand the role of patient-related and injury-related factors in the choice of treatment and in the outcome, we need large and long-term prospective cohort studies that include those treated with and without ACL reconstruction, to complement additional RCTs comparing the efficacy of different interventions.

We also need further basic research to better understand the role of the immediate joint trauma at the time of the ACL rupture in the development of OA, as well as the relative contribution of long-term chronic derangement of joint loading. Chondral injury and bone contusion is present in essentially all patients with acute traumatic ACL rupture<sup>6</sup>. This immediate mechanical insult activates inflammatory cytokine and protease cascades in cartilage, synovial and bone cells, and triggers apoptosis and catabolic responses in the articular cartilage that degrade the cartilage matrix<sup>7</sup>. These processes release matrix molecule fragments that represent damage associated molecular patterns (DAMPs), which activate Toll-like receptors, to potentially prolong the inflammatory response. The possibility needs to be considered that surgery in this acute phase adds an additional trauma that might enhance the early-phase pathological processes and extend the joint damage: to replace a torn ACL, bone tunnels are drilled for the tendon graft, resulting in stress deprivation and substantial bone loss<sup>8</sup>. This intervention might compromise not only graft fixation but also the long-term outcome.

The loading patterns of the knee with a ruptured ACL, whether reconstructed or not, are not normal, with increased mechanical load on the cartilage and altered location of the load on the joint surfaces<sup>9</sup>. A damaged meniscus will further enhance this abnormal loading. Joint cartilage with an impaired matrix, such as occurs soon after injury, is most sensitive to high loading rates<sup>10</sup>. Recognizing this consequence of injury is important for planning the rehabilitation and activity counseling for patients with these injuries.

These early consequences of ACL rupture suggest that, in order to rectify the continued high rate of OA following ACL rupture, we need to direct our attention to the acute phase after injury. We need to explore if early interventions to decrease cartilage cell death, harness inflammatory cascades, prevent activation of Toll-like receptors or slow the breakdown of cartilage matrix could prevent or decrease some of the downstream, late consequences of these common injuries. To save the acutely injured joint, we might, in the future, need the same attitudes and urgent actions as now exercised when trying to save myocardium or brain cells in patients with acute infarction or stroke, respectively.

The clinical management of the young active person with OA from a previous knee injury remains a challenge. A structured, personalized exercise program should be the basic and primary approach, together with advice and support to maintain a normal body weight. Patients should be encouraged to stay physically active, but to avoid high-impact, pivoting activities. Intermittent use of analgesics or a brace might be helpful for some. Knee replacement can be effective for those with severe symptoms of OA, and should be considered before they have lost too much function and become deconditioned. Long-term risk of need for implant revision surgery remains a concern for those who undergo knee replacement at a young age.

#### **Authors' affiliations:**

Department of Orthopaedics & Traumatology, Li Ka Shing Faculty of Medicine, The University of Hong Kong, 21 Sassoon Road, Pokfulam, Hong Kong (C.W.).  
Department of Orthopaedics, Clinical Sciences Lund, Lund University, SE-22185 Lund, Sweden (L.S.L.)

#### **Correspondence to:**

C.W., [paulwen@hku.hk](mailto:paulwen@hku.hk)

#### **Competing interests:**

The authors declare no competing interests.

#### **About the authors:**

Chunyi Wen is a postdoctoral fellow and honorary Assistant Professor in the Department of Orthopaedics & Traumatology at Li Ka Shing Faculty of Medicine, The University of Hong Kong.

L. Stefan Lohmander MD, PhD is Senior Professor in the Department of Orthopaedics at Clinical Sciences Lund, Lund University, Sweden, and Professor in the Research Unit for Musculoskeletal Function & Physiotherapy and the Department of Orthopaedics & Traumatology, University of Southern Denmark, Denmark.

#### **References**

1. Brophy, R.H., Gray, B.L., Nunley, R.M., Barrack, R.L. & Clohisy, J.C. Total knee arthroplasty after previous knee surgery: expected interval and the effect on patient age. *J Bone Joint Surg Am* **96**, 801-5 (2014).
2. Lohmander, L.S., Englund, M., Dahl, L.L., Roos, E.M. The long-term consequence of anterior cruciate ligament and meniscus injuries: Osteoarthritis. *Am J Sports Med* **35**:1756-69 (2007).
3. Cox, C.L., et al. Are articular cartilage lesions and meniscus tears predictive of IKDC, KOOS, and Marx activity level outcomes after anterior cruciate ligament reconstruction? A 6-year multicenter cohort study. *Am J Sports Med* **42**:1058-67 (2014).

- 132 4. Frobell, R.B., Roos, E.M., Roos, H.P., Ranstam, J., Lohmander, L.S. A  
133 randomized trial of treatment for acute anterior cruciate ligament tear. *New*  
134 *Engl J Med* **363**:331-342 (2010).
- 135 5. Frobell, R.B., et al. Treatment for acute anterior cruciate ligament tear: five  
136 year outcome of randomised trial. *BMJ* **346**, f232 (2013).
- 137 6. Potter, H.G., et al. Cartilage injury after acute, isolated anterior cruciate  
138 ligament tear: immediate and longitudinal effect with clinical/MRI follow-up.  
139 *Am J Sports Med* **40**, 276-85 (2012).
- 140 7. Anderson, D.D., et al. Post-traumatic osteoarthritis: improved understanding  
141 and opportunities for early intervention. *J Orthop Res* **29**: 802–9 (2011).
- 142 8. Wen, C.Y., Qin, L., Lee, K.M., Wong, M.W. & Chan, K.M. Influence of bone  
143 adaptation on tendon-to-bone healing in bone tunnel after anterior cruciate  
144 ligament reconstruction in a rabbit model. *J Orthop Res* **27**, 1447-56 (2009).
- 145 9. Hosseini, A., Van de Velde, S., Gill, T.J. & Li, G. Tibiofemoral cartilage  
146 contact biomechanics in patients after reconstruction of a ruptured anterior  
147 cruciate ligament. *J Orthop Res* **30**, 1781-8 (2012).
- 148 10. Nia et al. High-Bandwidth AFM-Based rheology reveals that cartilage is most  
149 sensitive to high loading rates at early stages of impairment. *Biophysical J*  
150 **104**:1529–37 (2013).

151  
152

**Figure 1.** Development of OA after acute rupture of the ACL of the knee. OA following ACL rupture is the consequence of the interaction between many risk factors, some associated with the person, such as heritability, and others with the environment, such as the severity of the trauma. Evidence that current interventions are able to alter the course from ACL rupture to OA is lacking. (Modified from Lohmander et al. AJSM 2007).

Abbreviations: ACL, anterior cruciate ligament; OA, osteoarthritis.

