



# LUND UNIVERSITY

## Disordered eating and emotion dysregulation among adolescents and their parents

Hansson, Erika; Daukantaitė, Daiva; Johnsson, Per

*Published in:*  
BMC psychology

*DOI:*  
[10.1186/s40359-017-0180-5](https://doi.org/10.1186/s40359-017-0180-5)

2017

*Document Version:*  
Publisher's PDF, also known as Version of record

[Link to publication](#)

*Citation for published version (APA):*

Hansson, E., Daukantaitė, D., & Johnsson, P. (2017). Disordered eating and emotion dysregulation among adolescents and their parents. *BMC psychology*, 5(12), Article 12. <https://doi.org/10.1186/s40359-017-0180-5>

*Total number of authors:*  
3

### General rights

Unless other specific re-use rights are stated the following general rights apply:  
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00

RESEARCH ARTICLE

Open Access



# Disordered eating and emotion dysregulation among adolescents and their parents

Erika Hansson<sup>1\*</sup>, Daiva Daukantaitė<sup>2</sup> and Per Johnsson<sup>2</sup>

## Abstract

**Background:** Research on the relationships between adolescent and parental disordered eating (DE) and emotion dysregulation is scarce. Thus, the aim of this study was to explore whether mothers' and fathers' own DE, as measured by SCOFF questionnaire, and emotion dysregulation, as measured by the difficulties in emotion regulation scale (DERS), were associated with their daughters' or sons' DE and emotion dysregulation. Furthermore, the importance of shared family meals and possible parent-related predictors of adolescent DE were explored.

**Method:** The total sample comprised 1,265 adolescents ( $M_{age} = 16.19$ ,  $SD = 1.21$ ; age range 13.5–19 years, 54.5% female) whose parents had received a self-report questionnaire via mail. Of these, 235 adolescents (18.6% of the total sample) whose parents completed the questionnaire were used in the analyses. Parents' responses were matched and compared with those of their child.

**Results:** Adolescent girls showed greater levels of DE overall than did their parents. Furthermore, DE was associated with emotion dysregulation among both adolescents and parents. Adolescent and parental emotion dysregulation was associated, although there were gender differences in the specifics of this relationship. The frequency of shared dinner meals was the only variable that was associated to DE and emotion dysregulation among adolescents, while parental eating disorder was the only variable that enhanced the probability of adolescent DE.

**Conclusion:** The present study contributes to the literature by demonstrating that there are significant associations between parents and their adolescent children in terms of DE, emotion dysregulation, and shared family meals. Future studies should break down these relationships among mothers, fathers, girls, and boys to further clarify the specific associational, and possibly predictive, directions.

**Keywords:** Disordered eating, Emotion dysregulation, Adolescents, Parents, Shared meals

## Background

Evidence for the parental influence on adolescents' eating disorders and disordered eating (DE)-namely, the behaviors and attitudes toward body perception, eating habits, weight regulation, and self-evaluation [46]-is inconclusive at present. Research exploring the specific pathways through which parental behaviors operate in this capacity is needed to clarify why some adolescents develop DE while others do not [22, 47]. Traditionally, responsibility has been attributed to mothers for the

occurrence of eating disorders in children (primarily daughters) [11], which has led most studies to explore different aspects of DE among mothers and daughters, such as mothers' talk about their own and their daughters' weight with their daughters [4], or the similarity of coping styles between mothers and daughters with eating disorders [25]. Despite this focus on mother-daughter relations, associations between parenting practices and eating behaviors have also been found among adolescent boys [28], making it clear that further efforts to understand the parental influences on boys' and girls' DE are needed. This is especially important since the prevalence of DE, which may lead to a clinical eating disorder [35] that renders high morbidity [19, 42] is rather

\* Correspondence: Erika.Hansson@hkr.se

<sup>1</sup>Kristianstad University, Kristianstad, Sweden

Full list of author information is available at the end of the article



high, varying from 15 to 24% in boys and to 25 to 52% in girls [16–18]. According to Wertheim [47], mothers' and fathers' food abstaining behaviors have predicted the food abstaining behaviors of their adolescent daughters but there is insufficient research on adolescents of both genders [28], and thus, the present study sought to fill this gap by exploring in what ways mothers' and fathers' DE is associated with that of their adolescent children.

Parental comments on a child's weight do appear to be the most consistent correlate of child weight and shape concerns and behaviors; however, one study found that children's modeling of parental weight concerns through observation of parents' own dieting behavior, weight-related concerns, and weight loss attempts gave rise to DE among children in fourth and fifth grade [43]. Such indirect parental influences suggests that children are influenced by their parents through modelling certain behaviors and the receiving of covert messages that communicate parental beliefs and expectations, as explained by social learning theory [2, 48]. Furthermore, such indirect parental influences, might have a stronger effect in adolescence due to their unobtrusive nature, which is especially relevant when considering adolescents' increased independence from parental control with age [37, 49].

DE has also been found to relate to emotion dysregulation among adults [50] and adolescents of both genders [30]. Children of both genders have been shown to express their emotions in different ways—girls are more likely to exhibit more positive and internalizing emotions, whereas boys are more likely to exhibit externalizing emotions [8]. However, as with DE, most research on emotion dysregulation has focused on mothers and daughters, finding evidence that children of depressed mothers are more likely to engage in maladaptive emotion regulation strategies than are children of non-depressed mothers [14, 40]. Furthermore, emotion regulation during middle childhood and adolescence appears to be more closely related to the emotion regulation ability of the mother than of the father [3]. There is also evidence that mothers and fathers give unique contributions to their child's development of emotion regulatory skills by responding differently to their children's sadness behavior [7].

Overall, research on the relationships between parent and adolescent emotion regulation and DE is scarce [3, 21, 32]. As such, the central question examined in this paper was whether parents' own DE and emotion dysregulation are implicitly associated with their adolescents'. One opportunity for children to observe and model healthy eating behaviors (provided that the eating behaviors of the parents are indeed healthy) is during shared family meals. The number of shared meals has

been found to be inversely associated with DE among adolescents [26, 36]; notably, this relationship is most commonly noted among females [41]. It is debated what specific problematic behaviors are reduced by regular shared meals, as some studies have suggested that shared meals serve a protective function against bulimia or extreme weight control behaviors [41]. Evidence does suggest that shared meals serve a protective function for both girls and boys, although in different ways. For example, it appears as if shared meals are associated with a decreased risk of engaging in risk behaviours among girls but not among boys [41].

In summary, to extend the knowledge of factors associated with adolescent DE, we explored the relations between parental and adolescent DE, emotion dysregulation, and shared family meals. We hypothesized that there would be associations as well as differences between parents and adolescents in regards to DE and emotion dysregulation, and that high levels of emotion dysregulation among parents would be associated with high levels of DE among the adolescents. Furthermore, we hypothesized that associations would be found between shared meals, emotion dysregulation, and DE among both adolescents and their parents, and that parental DE, parental eating disorders, parental emotion dysregulation, and shared family meals would affect the probability of DE among the adolescents. The analyses were stratified by gender in order to examine possible gender differences in these relationships.

## Method

### Participants

#### Adolescents

The study was conducted in a municipality in southern Sweden between January and March 2014. The initial sample comprised 1,265 adolescents ( $M_{\text{age}} = 16.19$ ,  $SD = 1.21$ ; age range 13.5 – 19 years, 54.5% female); in this study, however, we only analyzed a subsample of 235 adolescents (18.6% of the total sample) whose parents had completed the necessary questionnaires.

#### Parents

The parents of the students who attended the participating schools received a letter with a questionnaire that they were asked to complete and return in a pre-paid envelope. Two hundred ninety mainly Swedish (95.1%) parents (83.8% mothers) responded. Of those, 235 could be matched to their children using the personal identity number of the child. The remaining 55 responses could not be matched because of various reasons. An analysis of differences between children whose parents participated and those whose parents did not is presented in the Result section.

## Measures

### Disordered eating

**SCOFF** The SCOFF questionnaire [31] comprises five questions concerning eating habits and attitudes toward weight and body shape, as follows (with the SCOFF acronym letters printed in bold): “Do you make yourself sick (vomit) because you feel uncomfortably full?”; “Do you worry that you have lost control over how much you eat?”; “Have you recently lost more than one stone (15 pounds or about 6.8 kg) in a 3-month period?”; “Do you believe yourself to be fat when others say you are thin?”; and “Would you say that food dominates your life?” A threshold of two positive answers was proposed to indicate the probability of an existing eating disorder [29, 31], and this threshold was used in the present study, although it has been challenged with the idea that only one positive answer would be necessary to raise suspicion of DE, at least among adolescents [16]. The SCOFF was translated into Swedish [16] by one of the authors by using the back translation procedure with the help of another Swede competent in English. The reliability of the SCOFF-scale, assessed with Kuder and Richardson’s formula 20 (KR20), was .42 for the adolescents, and .29 for the parents.

**Eating disorder** Parents were asked to answer the following question: “Do you suffer from, or have you earlier in life suffered from, a clinical eating disorder?” The response options were “yes,” “no,” or “I don’t want to answer.”

### Emotion regulatory behaviors

**Emotion regulation** The Difficulties in Emotion Regulation Scale (DERS) [15] is a 36-item self-report questionnaire that assesses the six dimensions of emotion regulation: (a) lack of emotional *awareness* (e.g., “I am attentive to my feelings” (reversed)); (b) lack of emotional *clarity* (e.g., “I have no idea how I am feeling”); (c) impulse *control* difficulties (e.g., “When I’m upset I feel out of control”); (d) difficulties in engaging in *goal* directed behaviors (e.g., “When I’m upset I have difficulties getting work done”); (e) *non-acceptance* of emotional responses (e.g., “When I’m upset, I feel guilty for feeling that way”); and (f) limited access to emotion regulation *strategies* (e.g., “When I am upset, my emotions feel overwhelming”). The items are rated in terms of how frequently each statement applies to them on a five-point Likert scale ranging from 1 (“almost never”) to 5 (“almost always”). The total score for the DERS is calculated by adding all answers and the subscale scores are also added to a total number. The DERS has demonstrated high internal consistency [15], with a Cronbach’s  $\alpha$  of .93. In the current study, the Cronbach’s alpha was .94 for the total scale for adolescents (.77 for the

parents), .76 (.34) for awareness, .81(.64) for clarity, .85 (.58) for impulsivity, .86 (.60) for goals, .87 (.51) for non-acceptance, and .86 (.65) for strategies.

### Shared Meals

The adolescents answered three questions on the frequency of shared meals: “How often do the majority of your family members share breakfast/lunch/dinner with you?” Each question was rated on a 7-point Likert-type scale ranging from 1 (“1 day per week”) to 7 (“7 days per week”). The answers were added into a total score of a maximum of 21 meals per week.

## Procedure

### Adolescents

Both the legal guardians of the students below age 15 and the adolescents themselves, irrespective of age, received written information on the study aims, procedures, and the right to decline participation, and the same information was given orally to the adolescents on the day of data collection as well. The adolescents were also assured of confidentiality. Parents provided their passive consent—that is, they had to sign and return a form if they refused to allow their child to participate in the study. The adolescents consented to participation by completing the questionnaire, which took approximately one hour.

### Parents

A questionnaire, together with a letter explaining the survey and the right to decline participation, was sent out to all parents. By filling in the questionnaire and sending it back, the respondents gave their informed consent to participate. The parent questionnaire took approximately 20 min to complete.

## Statistical analysis

All analyses were calculated by use of the IBM SPSS Statistics program, and Pearson’s ( $r$ ), as well as Point-Biserial ( $r_{pb}$ ) correlation coefficients were used for all correlations. The analyses involving independent group comparisons were calculated with independent t-tests and the Mann-Whitney U-test, while paired-samples t-tests, as well as the McNemar’s test were used to explore the differences between adolescents and their parents. A logistic regression analysis was used to explore which variables among the parents possibly affect adolescent DE. All  $p$ -values were two-tailed.

## Results

### Preliminary analysis

Overall, 235 parents (86.4% mothers) with matched adolescents ( $M_{age} = 16.18$ ,  $SD = 1.20$ ; 56.6% female) participated in this study. Preliminary analyses were conducted

to examine whether there were any significant differences in the study variables between the 235 adolescents whose parents had completed the questionnaire and those whose parents had not, and as can be seen in Table 1, no significant differences were found among the two samples.

No significant differences were found between mothers and fathers regarding DE, as measured by the SCOFF, or in emotion dysregulation, as measured by the DERS. Therefore, mothers and fathers were not separated in the following analyses.

**Parental and adolescent DE**

Parental and adolescent DE was associated among the girls ( $r_{pb} = .19, p \leq .05$ ), but not among the boys. As Table 2 shows, about 90% of adolescents and 95% of parents reported fewer than two “yes” answers on the SCOFF and none of the participants answered “yes” to more than three SCOFF questions in total.

To further illustrate the presence of DE among adolescents and their parents, a McNemar’s chi-square test (Table 3) was conducted, and the results indicated significantly less DE among the parents than among the adolescents ( $p \leq .05$ ).

Five mothers and one father answered “yes” to the question of whether they suffered from, or had earlier in life suffered from, a clinical eating disorder.

**Parental and adolescent emotion dysregulation**

An independent *t*-test showed that girls ( $M = 79.49, SD = 20.76$ ) tended to have higher total scores on the DERS than did boys ( $M = 65.16, SD = 19.60, t(171) = 4.63, p \leq .001, Cohen’s d = 0.72$ ), which indicates that boys tend to have less emotion dysregulatory problems than girls. Furthermore, there was a significant positive correlation between adolescents’ total DERS scores and their parents’ ( $r = .18, p \leq .01$ ). Table 4 shows the means and standard deviations of the girls, boys, and parents for the total and subscale scores of the DERS.

**Girls**

Paired samples *t*-tests showed that girls ( $M = 79.49, SD = 20.76$ ) significantly differed in DERS total score

from their parents ( $M = 58.53, SD = 13.52, t(89) = 7.19, p \leq .001, Cohen’s d = 0.99$ ), indicating that girls experienced greater emotion dysregulation problems than did parents. Parents’ *strategies* subscale score correlated with girls’ scores on the *strategies* ( $r = .25, p \leq .01$ ), *impulse-control* ( $r = .19, p \leq .05$ ), and *emotional goals* ( $r = .20, p \leq .05$ ) subscales of the DERS. The parents’ *impulse control* subscale score also correlated with that of girls ( $r = .22, p \leq .05$ ).

**Boys**

Boys ( $M = 65.16, SD = 19.60$ ) also showed significantly higher DERS total scores than did their parents ( $M = 58.53, SD = 13.52, t(67) = 3.15, p \leq .002, Cohen’s d = 0.50$ ). Regarding the correlations, parents’ scores on the *clarity* subscale correlated with boys’ scores on the *clarity* ( $r = .27, p \leq .01$ ), *strategies* ( $r = .22, p \leq .05$ ), and *impulse control* ( $r = .30, p \leq .01$ ) subscales. Furthermore, vparents’ scores on the *strategies* subscale significantly correlated with boys’ scores on the *strategies* ( $r = .35, p \leq .01$ ) subscale, while parental *impulse control* scores correlated with boys’ *non-acceptance* ( $r = .24, p \leq .05$ ) and *strategies* ( $r = .27, p \leq .05$ ) scores. Finally, parents’ *non-acceptance* scores correlated with boys’ *non-acceptance* ( $r = .32, p \leq .01$ ) and *strategies* ( $r = .28, p \leq .01$ ) scores.

**The relation between DE and emotion dysregulation among adolescents and parents**

DE and emotion dysregulation were significantly correlated in adolescent girls ( $r_{pb} = .30, p \leq .01$ ) and in parents ( $r_{pb} = .27, p \leq .01$ ), but not in adolescent boys. No significant correlation was found between adolescent DE and parental emotion regulation.

**The relation between family meals, DE, and emotion dysregulation**

In regards to shared family meals, no significant correlation was found regarding the total number of meals shared and DE among the adolescents, and neither between the total number of meals shared and emotion dysregulation. However, when only family *dinner* meals were included in the analysis, and the breakfast and lunch meals omitted, weak but significant negative

**Table 1** The children whose parents participated and children whose parents did not were compared to determine structural differences among the groups

	Parent participated <i>M (SD) (Mdn for SCOFF)</i>	Parent did not participate <i>M (SD) (Mdn for SCOFF)</i>	<i>t</i> (u for SCOFF)
SCOFF	0.00	0.00	$u = 99716.5, p = .07$
DERS	73.11 (21.42)	75.19 (23.70)	$t(295.59) = 1.11, p = .30$
<i>M<sub>age</sub></i>	16.18 (1.20)	16.19 (1.22)	$t(1262) = .084, p = .93$

Note: For each analysis, we excluded the data of adolescents with missing values in a pairwise fashion. Samples whose parents did not participate and whose parents did participate ranged from 642 to 1029 adolescents and 173 to 235 adolescents, respectively. As can be seen from the *p*-values no such significant differences were found

**Table 2** Number of “yes” answers on the SCOFF among adolescents and parents

“Yes” answers	Adolescents (n)	Adolescent Girls (n)	Adolescent Boys (n)	Parents (n)	Mothers (n)	Fathers (n)
0	(164) 73.9%	(78) 63.4%	(86) 86.9%	(190) 81.9%	(163) 81.5%	(27) 84.5%
1	(35) 15.8%	(23) 18.7%	(12) 12.1%	(32) 13.8%	(29) 14.5%	(3) 9.5%
2	(19) 8.6%	(18) 14.7%	(1) 1.0%	(9) 3.9%	(8) 4%	(1) 3%
3	(4) 1.7%	(4) 3.2%	(0)	(1) 0.4%	(0) 0%	(1) 3%

More than two “yes” answers are indicative of an eating disorder

correlations with DE ( $r_{pb} = -.16, p \leq .01$ ) as well as with emotion dysregulation ( $r = -.16, p \leq .05$ ) were found. When exploring girls and boys separately, no significant correlations were observed.

**The effect of parental eating disorders**

A logistic regression analysis was performed to assess the impact of a number of factors, including parents’ DE and earlier diagnosed eating disorder, emotion dysregulation, and frequency of shared meals, on the odds that adolescents would report having DE. However, only the results of the girls are presented, as only one boy gave two “yes” answers on SCOFF. The model contained 12 independent variables: eating breakfast, lunch, or dinner together with parent; parental awareness; clarity; impulse control; goals; non-acceptance of emotions; parental emotional strategies; parental SCOFF score; and parental self-reported eating disorder. The full model containing all predictors was significant (for girls)  $\chi^2 (1, N = 133) = 6.31, p = .012$ , indicating that the model could distinguish the girls who reported DE and girls who did not. The model as a whole explained between 5.7% (Cox and Snell R-square) and 9.3% (Nagelkerke’s R-squared) of the variance in DE and correctly classified 83.2% of the cases. Surprisingly, only one of the independent variables made a significant contribution to the model: whether the parent reported suffering from an eating disorder. The odds ratio was .07, indicating that there is slightly higher odds that girls of parents suffering, or having suffered from an eating disorder would report DE.

**Discussion**

In this study the relationships between DE, emotion dysregulation, frequency of shared family meals among adolescents and their parents, as well as possible parental

predictors of adolescent DE were explored. Associations between girls’ DE and their parents’ DE were found, and DE was also associated with difficulties with emotion dysregulation among adolescents as well as parents. Furthermore, associations were found among adolescent and parental emotion dysregulation, although the specifics of this relationship showed gender differences such as the parents’ emotion dysregulatory strategies were associated with impulse control and emotional goals for girls but not for boys. Regarding the boys, the parents’ scores on the emotional clarity subscale were associated with emotional clarity, strategies and impulse control. These associations were not found among girls. Parents’ non-acceptance scores were also positively associated with the non-acceptance scores of the boys’, but not the girls’. Only the shared family dinner meal was associated with less problems with DE and emotion dysregulation among adolescents, and finally, only parental eating disorder enhanced the probability of adolescent DE.

One boy and 22 girls reported 2 or more “yes” answers on the SCOFF, which is indicative of an eating disorder. However, only one answer on SCOFF might, in some cases, be indicative of DE [16] and about 13.1% of the boys and 35.2% of the girls reported 1 or more “yes” answers on SCOFF. These numbers are roughly comparable to studies where the prevalence of DE varied from 15 to 17% among boys and to 33 to 52% among girls in German and Finnish adolescent samples [17, 18].

Adolescents were more likely to experience DE than were their parents, which is in line with research saying that more than 50% of teenage girls and 33% of teenage boys are using restrictive measures to lose weight at any given time [33] while “only” 13 and 11% of women and men, respectively, scored in the DE range [6]. A possible explanation as to why DE is more common among

**Table 3** Observed frequencies of DE among adolescents and parents (based on two “yes” answers on SCOFF)

	Adolescents				Total
	Girls		Boys		
	DE	No DE	DE	No DE	
Parents with DE	4 (18%)	5 (5%)	0 (0%)	1 (1%)	10 (4.5%)
Parents without DE	18 (82%)	93 (95%)	1 (100%)	97 (99%)	209 (95.5%)
Total	22 (100%)	98 (100%)	1 (100%)	98 (100%)	219 (100%)

Note: The boys’ association was not significant according to McNemar’s test

**Table 4** Girls', boys' and parents' means and standard deviations on the total and subscale scores of the DERS

DERS	Girls		Boys		Parents	
	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>
Total	79.49	20.76	65.16	19.60	58.53	13.52
Awareness	16.27	5.05	15.68	5.37	12.40	4.11
Clarity	11.82	3.98	8.68	3.62	6.86	2.06
Impulse control	11.23	5.25	9.17	4.28	8.24	2.48
Goals	13.13	4.95	10.91	4.51	9.63	3.42
Non-acceptance	12.72	5.40	9.22	3.50	10.04	3.98
Strategies	16.39	6.76	12.72	5.10	11.32	3.28

younger people is that emotion regulation skills develop well into adulthood, and therefore might not be as advanced in adolescence as in later life [38]. Thus, if emotion dysregulation is indeed associated with DE, greater emotion regulation skills might be one contributor to healthier eating behaviors as a person gets older.

Girls' DE was associated with their parents DE, which is in line with earlier research [47], and on a speculative note, considering the majority of the parents were mothers, a gender link between daughters' and mothers' DE could be possible, as well as a parallel relationship between boys' and fathers' DE. Furthermore, the girls showed greater problems with DE than did their parents, but boys did not. The number of boys who reported DE was small, and it is possible that the SCOFF is interpreted in different ways by girls and boys [16].

Girls in the study showed higher emotion dysregulation than boys, and adolescents in general had higher emotion dysregulation scores than did their parents. As mentioned earlier, the difference between adolescents and parents could be explained by prior findings saying that emotion regulation develops well into adulthood [38] but this does not explain why girls in this study experience more emotion dysregulation than boys. On the contrary, girls, in general, tend to mature earlier than boys do [27], which ought to give them an advantage in terms of emotion regulation skills. On the other hand, adolescent girls, compared to adolescent boys, are more likely to suffer from low self-esteem and depression [39], which are correlated with DE [10, 24]; these might in some way inhibit the development of emotion regulation skills among girls.

Both girls' and boys' emotion regulation skills were related to their parents', which is in line with the modeling hypothesis proposed earlier [2]. According to this hypothesis, children imitate their parents' manner of regulating their emotions implicitly. Although such modeling may explain the relations of overall DE and emotion dysregulation among adolescents and parents, it might not explain why the specific parental emotion

dysregulation *strategies* were associated with adolescents' emotion dysregulation. More specifically, this association suggests that parents have an explicit *strategy* to teach the child his or her emotion regulation skills [32].

In line with previous studies [9, 23, 30], an association between DE and emotion dysregulation was found in adolescents as well as their parents, however, parental emotion dysregulation was not associated with their children's DE. Furthermore, there was no association between frequency of shared family meals and DE among adolescents [26]. We did observe, however, that frequency of shared family *dinners* was negatively associated with adolescent DE and emotion dysregulation, although this association was very weak. This could be because dinners are more relevant than are other meals, and might be considered more of a "family affair" in Sweden, where lunch is provided at school (and, incidentally, skipped by about 50% of adolescents) [20] and breakfast, if not completely disregarded [20], often is a stressful affair for most families with children in school and two parents working full time, although frequency of breakfast meals was earlier found to be associated with DE [1]. Perhaps "family" time is more important for reducing DE as opposed to mere "food intake," which occurs during breakfast and lunch as well, and this might also be true for the development of emotion regulation. Furthermore, it should be noted that we only measured adolescents' reports of how many times per week the family came together for different meals. It is possible that they have different views on what should constitute a "shared family meal" than their parents would have. It is also important to notice that not all shared meals would be considered joyful occasions, and we did not consider the wider family constellation, including siblings. A somewhat surprising result was that parental eating disorder was the only variable that raised the odds of adolescent DE. However, this result might further strengthen the importance of modeling among adolescents and their parents [2].

#### Strengths and limitations

Although parents' response rate was poor and there was a possible volunteer bias, the knowledge of which parent (mother or father) answered the questions and our matching of the parent to his or her child can be considered strengths of this study. The data were collected in common public schools in a municipality in southern Sweden and the students came from both urban and rural settings, making our sample similar to the composition in other Swedish municipalities. The final sample comprised 235 adolescent-parent pairs, which is a rather large sample overall. All data were obtained via self-report and no diagnostic interviews were used to confirm answers to the questionnaires which can be

considered a limitation of the study. On the other hand, it has been argued that the anonymous self-report format might yield more valid data than interviews because they are less personally intrusive [12]. Another limitation is that, although the reliability of the DERS scale seemed appropriate for adolescents (based on the Cronbach's alpha), it was much lower for parents. If the "awareness" subscale was removed, the Cronbach's alpha for parents would become equivalent to that of adolescents which should be considered in further studies. The initial psychometric testing of the DERS scale, which yielded high internal consistency, was made on undergraduate students [15], and when using adult samples, the awareness subscale has been found to fail to correlate with the four of the five other subscales, while the remaining five all correlate significantly with one another [44, 45]. The fact that the DERS did not work sufficiently for the adults was a limitation, but there was an obvious advantage of using the exact same scale for the adolescents and parents. As a final limitation, we did not measure adolescents' possible eating disorder and our question on shared meals did not contain a "never" answer.

## Conclusions and further directions

Although evidence for parental influence on DE and emotion dysregulation is inconsistent [22, 47], likely due to methodological differences [34], the present study contributes to the research by showing significant associations between parents and their adolescent children with regard to DE and emotion dysregulation as well as shared meals. Future studies should specifically ask for answers from both parents separately to further clarify the specific influences mothers and fathers have on their children. Studies should also continually include adolescents of both genders, as already suggested by Berge et al. [5]. On this note, it would also be of interest to focus on boys with DE to further explore how boys' DE relates to their parents' DE and emotion dysregulation. Another avenue to explore is fathers' possible influence on their sons' and daughters' eating, which would offer a contrast to studies focusing solely on mothers and their responsibility for DE [11, 13], which seems to be quite outdated in contemporary society.

## Abbreviation

DE: Disordered eating; DERS: Difficulties in emotion regulation scale

## Funding

The collection of data from the parents was made possible by funding from Academia Gastronomica Scaniensis.

## Availability of data and materials

The data are, in accordance to the ruling of the ethics committee, not to be shared.

## Authors' contributions

EH designed the study, carried out the data collection, performed the statistical analysis, and wrote the paper. DD designed the study, performed the statistical analysis, and gave conceptual advice. PJ designed the study and gave conceptual advice. All authors discussed the results and implications and approved the final manuscript.

## Competing interests

The authors declare that they have no competing interests.

## Consent for publication

Not applicable.

## Ethics approval and consent to participate

The study was approved by the Central Ethical Review Board in Lund, Sweden. Parents and adolescents consented to participation by filling in the questionnaires and returning them to the researchers. Parents consented to the children's participation by an "opt-out" procedure.

Trial registration number provided by the Central Ethical Review Board, Lund Sweden: 2012/499 (October 17, 2012) for the adolescents, and 2013/642 (October 15, 2013) for the parents.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## Author details

<sup>1</sup>Kristianstad University, Kristianstad, Sweden. <sup>2</sup>Lund University, Lund, Sweden.

Received: 29 November 2016 Accepted: 24 March 2017

Published online: 04 April 2017

## References

- Ackard D, Neumark-Sztainer DM. Family mealtime while growing up: Associations with symptoms of bulimia nervosa. *Eat Disord.* 2001;9(3):239–49.
- Bandura A. *Social learning theory.* Englewood Cliffs: Prentice-Hall; 1977.
- Bariola E, Hughes EK, Gullone E. Relationships Between Parent and Child Emotion Regulation Strategy Use: A Brief Report. *J Child Fam Stud.* 2012; 21(3):443–8. doi:10.1007/s10826-011-9497-5.
- Bauer KW, Bucchianeri MM, Neumark-Sztainer D. Mother-reported parental weight talk and adolescent girls' emotional health, weight control attempts, and disordered eating behaviors. *J Eat Disord.* 2013;1:45. doi:10.1186/2050-2974-1-45.
- Berge JM, Wall M, Larson N, Eisenberg ME, Loth KA, Neumark-Sztainer D. The unique and additive associations of family functioning and parenting practices with disordered eating behaviors in diverse adolescents. *J Behav Med.* 2014;37(2):205–17. doi:10.1007/s10865-012-9478-1.
- Büyükgöze-Kavas A. Eating attitudes and depression in a Turkish sample. *Eur Eat Disord Rev.* 2007;15(4):305–10.
- Cassano M, Perry-Parrish C, Zeman J. Influence of gender on parental socialization of children's sadness regulation. *Soc Dev.* 2007;16(2):210–31. doi:10.1111/j.1467-9507.2007.00381.x.
- Chaplin TM, Aldao A. Gender differences in emotion expression in children: a meta-analytic review. *Psychol Bull.* 2013;139(4):735–65. doi:10.1037/a0030737.
- Davis-Becker K, Peterson CM, Fischer S. The relationship of trait negative urgency and negative affect to disordered eating in men and women. *Personal Individ Differ.* 2014;56:9–14. doi:10.1016/j.paid.2013.08.010.
- Dennard EE, Richards CS. Depression and coping in subthreshold eating disorders. *Eat Behav.* 2013;14(3):325–9. doi:10.1016/j.eatbeh.2013.05.011.
- Eliot AO, Baker CW. Maternal stressors and eating-disordered adolescent girls. *Fam Ther.* 2000;27(3):165.
- Fairburn CG, Beglin SJ. Assessment of eating disorders: interview or self-report questionnaire? *Int J Eat Disord.* 1994;16(4):363–70.
- Francis LA, Birch LL. Maternal influences on daughters' restrained eating behavior. *Health Psychol.* 2005;24(6):548–54. doi:10.1037/0278-6133.24.6.548.
- Garber J, Braafladt N, Zeman J. The Regulation of Sad Affect - an Information-Processing Perspective. *Development of Emotion Regulation and Dysregulation,* 208–240. 1991. doi:10.1017/Cbo9780511663963.011.
- Gratz KL, Roemer L. Multidimensional assessment of emotion regulation and dysregulation: Development, factor structure, and initial validation of the difficulties in emotion regulation scale. *J Psychopathol Behav Assess.* 2004; 26(1):41–54. doi:10.1023/B:Joba.0000007455.08539.94.



16. Hansson E, Daukantaite D, Johnsson P. SCOFF in a general Swedish adolescent population. *J Eat Disord.* 2015;3:48. doi:10.1186/s40337-015-0087-6.
17. Hautala L, Helenius H, Karukivi M, Maunula AM, Nieminen J, Aromaa M, Saarjarvi S. The role of gender, affectivity and parenting in the course of disordered eating: a 4-year prospective case-control study among adolescents. *Int J Nurs Stud.* 2011;48(8):959–72. doi:10.1016/j.ijnurstu.2011.01.014.
18. Herpertz-Dahlmann B, Wille N, Holling H, Vloet TD, Ravens-Sieberer U. Disordered eating behaviour and attitudes, associated psychopathology and health-related quality of life: results of the BELLA study. *Eur Child Adolesc Psychiatry.* 2008;1(51):82–91. doi: 10.1007/s00787-008-1009-9.
19. Hill LS, Reid F, Morgan JF, Lacey JH. SCOFF, the development of an eating disorder screening questionnaire. *Int J Eat Disord.* 2010;43(4):344–51. doi:10.1002/eat.20679.
20. Höglund D, Samuelson G, Mark A. Food habits in Swedish adolescents in relation to socioeconomic conditions. *Eur J Clin Nutr.* 1998;52(11):784–9. doi:10.1038/sj.ejcn.1600644.
21. John OP, Gross JJ. Healthy and unhealthy emotion regulation: personality processes, individual differences, and life span development. *J Pers.* 2004; 72(6):1301–33. doi:10.1111/j.1467-6494.2004.00298.x.
22. Kluck AS. Family influence on disordered eating: the role of body image dissatisfaction. *Body Image.* 2010;7(1):8–14. doi:10.1016/j.bodyim.2009.09.009.
23. Lafrance Robinson A, Kosmerly S, Mansfield-Green S, Lafrance G. Disordered eating behaviours in an undergraduate sample: Associations among gender, body mass index, and difficulties in emotion regulation. *Can J Behav Sci.* 2014;46(3):320–6. doi:10.1037/a0031123.
24. Lampard AM, Byrne SM, McLean N. Does Self-Esteem Mediate the Relationship between Interpersonal Problems and Symptoms of Disordered Eating? *Eur Eat Disord Rev.* 2011;19(5). doi: 10.1002/erv.1120.
25. Lantzouni E, Cox MH, Salvator A, Crosby RD. Mother-daughter coping and disordered eating. *Eur Eat Disord Rev.* 2015;23(2):126–32. doi:10.1002/erv.2343.
26. Larson NI, Neumark-Sztainer D, Hannan PJ, Story M. Family meals during adolescence are associated with higher diet quality and healthful meal patterns during young adulthood. *J Am Diet Assoc.* 2007;107(9):1502–10. doi:10.1016/j.jada.2007.06.012.
27. Lim S, Han CE, Uhlhaas PJ, Kaiser M. Preferential detachment during human brain development: age- and sex-specific structural connectivity in diffusion tensor imaging (DTI) data. *Cereb Cortex.* 2015;25(6):1477–89.
28. Loth KA, MacLehose RF, Fulkerson JA, Crow S, Neumark-Sztainer D. Are food restriction and pressure-to-eat parenting practices associated with adolescent disordered eating behaviors? *Int J Eat Disord.* 2014;47(3):310–4. doi:10.1002/eat.22189.
29. Luck AJ, Morgan JF, Reid F, O'Brien A, Brunton J, Price C, Lacey JH. The SCOFF questionnaire and clinical interview for eating disorders in general practice: comparative study. *BMJ.* 2002;325(7367):755–6.
30. Mills P, Newman EF, Cossar J, Murray G. Emotional maltreatment and disordered eating in adolescents: testing the mediating role of emotion regulation. *Child Abuse Negl.* 2015;39:156–66. doi:10.1016/j.chiabu.2014.05.011.
31. Morgan JF, Reid F, Lacey JH. The SCOFF questionnaire: assessment of a new screening tool for eating disorders. *BMJ.* 1999;319(7223):1467–8.
32. Morris AS, Silk JS, Steinberg L, Myers SS, Robinson LR. The Role of the Family Context in the Development of Emotion Regulation. *Soc Dev.* 2007;16(2): 361–88. doi:10.1111/j.1467-9507.2007.00389.x.
33. Neumark-Sztainer D. I'm Like, SO Fat!: Helping Your Teen Make Healthy Choices about Eating and Exercise in a Weight-obsessed World: Guilford Press. 2005.
34. Neumark-Sztainer D, Bauer KW, Friend S, Hannan PJ, Story M, Berge JM. Family weight talk and dieting: how much do they matter for body dissatisfaction and disordered eating behaviors in adolescent girls? *J Adolesc Health.* 2010;47(3):270–6. doi:10.1016/j.jadohealth.2010.02.001.
35. Neumark-Sztainer D, Wall M, Guo J, Story M, Haines J, Eisenberg M. Obesity, disordered eating, and eating disorders in a longitudinal study of adolescents: how do dieters fare 5 years later? *J Am Diet Assoc.* 2006;106: 559–68. doi:10.1016/j.jada.2006.01.003.
36. Neumark-Sztainer D, Wall M, Story M, Fulkerson JA. Are family meal patterns associated with disordered eating behaviors among adolescents? *J Adolesc Health.* 2004;35(5):350–9. doi:10.1016/j.jadohealth.2004.01.004.
37. Riediger M, Schmiedek F, Wagner GG, Lindenberger U. Seeking pleasure and seeking pain: differences in prohedonic and contra-hedonic motivation from adolescence to old age. *Psychol Sci.* 2009;20(12):1529–35. doi:10.1111/j.1467-9280.2009.02473.x.
38. Rutherford HJ, Wallace NS, Laurent HK, Mayes LC. Emotion Regulation in Parenthood. *Dev Rev.* 2015;36:1–14. doi:10.1016/j.dr.2014.12.008.
39. Santrock JW. *Adolescence.* New York: McGraw-Hill; 2016.
40. Silk JS, Shaw DS, Skuban EM, Oland AA, Kovacs M. Emotion regulation strategies in offspring of childhood-onset depressed mothers. *J Child Psychol Psychiatry.* 2006;47(1):69–78. doi:10.1111/j.1469-7610.2005.01440.x.
41. Skeer MR, Ballard EL. Are family meals as good for youth as we think they are? A review of the literature on family meals as they pertain to adolescent risk prevention. *J Youth Adolesc.* 2013;42(7):943–63. doi:10.1007/s10964-013-9963-z.
42. Smink FR, van Hoeken D, Hoek HW. Epidemiology of eating disorders: incidence, prevalence and mortality rates. *Curr Psychiatry Rep.* 2012;14(4): 406–14. doi:10.1007/s11920-012-0282-y.
43. Smolak L, Levine MP, Schermer F. Parental input and weight concerns among elementary school children. *Int J Eat Disord.* 1999;25(3):263–71. doi: 10.1002/(Sici)1098-108x(199904)25:3<263:Aid-Eat3>3.0.Co;2-V.
44. Tull MT, Barrett HM, McMillan ES, Roemer L. A preliminary investigation of the relationship between emotion regulation difficulties and posttraumatic stress symptoms. *Behav Ther.* 2007;38(3):303–13. doi:10.1016/j.beth.2006.10.001.
45. Tull MT, Gratz KL, Litzman RD, Kimbrel NA, Lejuez CW. Reinforcement Sensitivity Theory and emotion regulation difficulties: A multimodal investigation. *Personal Individ Differ.* 2010;49(8):989–94. doi:10.1016/j.paid.2010.08.010.
46. Waaddegaard M, Thoning H, Petersson B. Validation of a screening instrument for identifying risk behaviour related to eating disorders. *Eur Eat Disord Rev.* 2003;11(6):433–55. doi:10.1002/erv.537.
47. Wertheim EH. Relationships among adolescent girls' eating behaviors and their parents' weight-related attitudes and behaviors. *Sex Roles.* 1999;41(3–4):169–87. doi:10.1023/A:1018850111450.
48. White HR, Johnson V, Buyske S. Parental modeling and parenting behavior effects on offspring alcohol and cigarette use - A growth curve analysis. *J Subst Abuse.* 2000;12(3):287–310. doi:10.1016/S0899-3289(00)00056-0.
49. Zimmer-Gembeck MJ, Collins A. Autonomy Development during Adolescence. In: Adams GR, Berzonsky MD, editors. *Blackwell Handbook of Adolescence.* Hoboken: Blackwell Publishing; 2005.
50. Zysberg L, Rubanov A. Emotional intelligence and emotional eating patterns: a new insight into the antecedents of eating disorders? *J Nutr Educ Behav.* 2010;42(5):345–8.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

