



LUND UNIVERSITY

Phosphorylation of the activation loop tyrosine 823 in c-Kit is crucial for cell survival and proliferation.

Agarwal, Shruti; Kazi, Julhash U.; Rönnstrand, Lars

Published in:
Journal of Biological Chemistry

DOI:
[10.1074/jbc.M113.474072](https://doi.org/10.1074/jbc.M113.474072)

2013

[Link to publication](#)

Citation for published version (APA):
Agarwal, S., Kazi, J. U., & Rönnstrand, L. (2013). Phosphorylation of the activation loop tyrosine 823 in c-Kit is crucial for cell survival and proliferation. *Journal of Biological Chemistry*, 288(31), 22460-22468.
<https://doi.org/10.1074/jbc.M113.474072>

Total number of authors:
3

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Phosphorylation of the activation loop tyrosine 823 in c-Kit is crucial for cell survival and proliferation*

Shruti Agarwal, Julhash U. Kazi and Lars Rönnstrand¹

Experimental Clinical Chemistry, Wallenberg Laboratory, Department of Laboratory Medicine, Lund University, Skåne University Hospital, 20502 Malmö, Sweden,

*Running title: Phosphorylation of Y823 is crucial for c-Kit signaling

¹Corresponding author: Lars Rönnstrand, Experimental Clinical Chemistry, Wallenberg Laboratory, Department of Laboratory Medicine, Lund University, Skåne University Hospital, 20502 Malmö, Sweden, Tel.: +46 40 33 72 22, Fax: +46 40 33 11 04, E-mail: Lars.Ronnstrand@med.lu.se

Keywords: c-Kit; Activation loop; Y823, tyrosine phosphorylation, signaling

Background: Phosphorylation of Y823 in the A-loop of c-Kit is not required for kinase activity but might have other functions.

Results: Mutation of Y823 in c-Kit causes alterations in downstream signaling and a reduction in ligand-dependent cell survival and proliferation.

Conclusion: Phosphorylation of Y823 regulates c-Kit stability.

Significance: This provides a function for the activation loop tyrosine of c-Kit.

SUMMARY

The receptor tyrosine kinase c-Kit, also known as the stem cell factor receptor, plays a key role in several developmental processes. Activating mutations in c-Kit lead to alteration of these cellular processes and have been implicated in many human cancers such as gastrointestinal stromal tumors (GISTs), acute myeloid leukemia (AML), testicular seminomas and mastocytosis. Regulation of the catalytic activity of several kinases is known to be governed by phosphorylation of tyrosine residues in the activation loop of the kinase domain. However, in the case of c-Kit phosphorylation of Y823 has been demonstrated to be a late event that is not required for kinase activation. However, since phosphorylation of Y823 is a ligand-activated event, we sought to investigate the functional consequences of Y823 phosphorylation. By using a tyrosine to phenylalanine mutant of tyrosine 823 we investigated the impact of Y823 on c-Kit signaling. We here demonstrate that Y823 is crucial for cell survival and proliferation and mutation of Y823 to phenylalanine leads to decreased sustained phosphorylation and

ubiquitination of c-Kit as compared to the wild-type receptor. Furthermore, the mutated receptor was upon ligand-stimulation quickly internalized and degraded. Phosphorylation of the E3 ubiquitin ligase, Cbl was transient followed by a substantial reduction in phosphorylation of downstream signaling molecules such as Akt, Erk, p38, Shc and Gab2. Thus, we propose that activation loop tyrosine 823 is crucial for activation of both the MAPK and PI3K pathways and that its disruption leads to a destabilization of the c-Kit receptor and decreased survival of cells.

c-Kit belongs to the family of type III receptor tyrosine kinases (RTKs)² and is known for its critical role in hematopoiesis, pigmentation and reproduction. It is also important for survival, proliferation and differentiation of hematopoietic progenitor cells (1,2). Several activating mutations in c-Kit lead to a deregulation of the signaling cascades causing malignancies such as acute myeloid leukemia, gastrointestinal stromal tumors, testicular seminomas (3-6). Architecturally, c-Kit is comprised of five extracellular Immunoglobulin-like domains, a transmembrane domain, a juxtamembrane region, a kinase domain split into two parts by a kinase insert and a carboxyterminal tail. The activation loop is present in the C-terminal lobe of the kinase domain. Activation of the receptor is initiated by binding of its ligand, stem cell factor (SCF) which leads to receptor dimerization and transphosphorylation on specific tyrosine residues. These phosphorylated residues serve as docking sites for signal transduction molecules containing SH2 (Src homology 2) domains such

as Cbl, Gab2, Shc, SHP2. They get activated upon phosphorylation and/or binding to the receptor and mediate signaling downstream of the receptor that eventually leads to various cellular responses. Many receptor tyrosine kinases undergo monoubiquitination or polyubiquitination following ligand stimulation, which targets them for degradation in the lysosomes or proteasomes, respectively. Members of Cbl family of ubiquitin E3 ligases play a crucial role in ubiquitination of c-Kit (1,7,8).

In many tyrosine kinases phosphorylation of tyrosine residue(s) in the activation loop is a crucial early event that leads to activation of the kinase. Examples of this type of receptors include the insulin receptor and the FGFR1 (9,10). However, in the case of c-Kit the phosphorylation of tyrosine residues in the juxtamembrane region is the most important activating event. In the absence of phosphorylation of the juxtamembrane region, it is inserted into a cleft between the N-terminal and C-terminal lobes of the kinase domain. Thereby the so-called C-helix is disrupted and the DFG motif is prevented from establishing an active conformation. Upon phosphorylation of the tyrosine residues in the juxtamembrane region, it is released together with the activation loop from the active site, the C-helix can move into position in the active site and thereby correctly orient the DFG residues that are important for catalytic activity (11,12).

Kinetic studies on recombinant c-Kit in vitro using mass spectrometry revealed that phosphorylation of Y823 is a late event that was observed when 90% of the kinase was already phosphorylated (13). Mutation of Y823 to a phenylalanine residue did not impair kinase activity. Further, Mol et al. showed that in a crystal structure of unactivated enzyme Y823 is bound to the catalytic base D792 blocking the access of substrates to the catalytic site (11). Therefore, phosphorylation of Y823 may disengage the activation loop from its inhibitory state. Another possibility is that phosphorylation and dephosphorylation of Y823 stabilizes the receptor structure and downstream signaling without being directly involved in kinase activity. However, the role of Y823 and its effects on c-Kit signaling has not been studied until now.

In this study we have investigated the cellular and biochemical effects of mutating Y823 to phenylalanine (Y823F). We show that

Y823 is crucial for cell survival and proliferation. Cells expressing the Y823F mutant of c-Kit showed much lower proliferation and survival as compared to cells expressing the wild-type receptor despite the fact that the kinase activity was intact. Furthermore, the Y823F mutant receptor was internalized and degraded much faster than the wild-type receptor. A reduction in phosphorylation of the adaptor proteins Cbl, Shc, and Gab2 was also seen. The PI3-kinase/Akt, Ras/Erk and p38 pathways were also affected, in that the phosphorylation of Akt, Erk and p38 was very transient and not sustained as in wild-type c-Kit. Taken together, this study adds a novel perspective towards understanding the role of the activation loop tyrosine in c-Kit that is related to downstream signaling rather than kinase activity.

EXPERIMENTAL PROCEDURES

Reagents and antibodies-Transfection reagent Lipofectamine 2000 was from Invitrogen and jetPEI was from Polyplus. Cycloheximide was purchased from Sigma. Human recombinant SCF and murine recombinant IL-3 (Interleukin-3) were obtained from ProSpec Tany Technogene (Rehovot, Israel). Rabbit polyclonal anti-c-Kit serum was raised against a synthetic peptide corresponding to the carboxyterminus of c-Kit and purified as described (14). The anti-Cbl antibodies have been described elsewhere (15). The phospho-tyrosine antibody 4G10 was purchased from Millipore and ubiquitin antibody from Covance Research Products. Antibodies against phospho-p38, p38 and Shc were from BD Transduction Laboratories. Phycoerythrin(PE)-labeled c-Kit antibody was from BD Biosciences. Anti-phospho-Akt antibody was from Epitomics. Polyclonal anti-Gab2, anti-Akt, anti-phospho-Erk, anti-Erk and horseradish peroxidase-coupled secondary anti-goat antibodies were purchased from Santa Cruz Biotechnology. Secondary Horseradish peroxidase-coupled anti-mouse and anti-rabbit antibodies were from Invitrogen.

Cell culture-Ba/F3 cells and M07e cells were cultured in RPMI 1640 medium containing 10% heat-inactivated fetal bovine serum (FBS), 100 µg/ml streptomycin, 100 units/ml penicillin and 10 ng/ml recombinant murine interleukin-3 (IL-3) or 10 ng/ml recombinant human IL3, respectively. Kasumi-1 cells were cultivated in above medium lacking IL-3. Dulbecco's modified Eagle's medium supplemented with 10% FBS, 100 µg/ml streptomycin and 100

units/ml penicillin was used to culture COS-1 and EcoPack cells.

Expression constructs-pcDNA3-c-Kit-WT, pMSCV-c-Kit-WT constructs were described previously (1,8). pcDNA3-c-Kit Y823F and pMSCV-c-Kit-Y823F constructs were generated by site-directed mutagenesis using QuikChange mutagenesis XL kit (Stratagene). All plasmids were verified by sequencing.

Transient and stable transfection-Transient transfection of COS1 cells was performed using transfection reagent, JetPEI according to the manufacturer's instructions. Transfected cells were incubated for about 30 h before they were serum-starved overnight. Cell stimulation, lysis and immunoprecipitation was performed essentially according to (16). Stable transfections were performed essentially as described (17). Cells expressing wild-type c-Kit or c-Kit Y823F were confirmed by flow cytometry.

Immunoprecipitation and Western blotting-Cell lysis, immunoprecipitation and Western blotting was performed as described (16). Immunodetection was performed by enhanced chemiluminescence using horseradish peroxidase substrate (Millipore Corporation, Billerica, MA, USA) and the signals were detected by a CCD camera (LAS-3000, Fujifilm, Tokyo, Japan). Signal intensities were further quantified by Multi-Gauge software (Fujifilm).

In vitro kinase activity assays-COS-1 cells transfected with pcDNA3-cKit-WT and pcDNA3-c-Kit-Y823F plasmids were starved of serum overnight. Cells were stimulated with 100 ng/ml SCF ligand and cell lysates were prepared. c-Kit receptor was immunoprecipitated and processed for in vitro kinase reaction essentially as described (18) except that 5 μ g of myelin basic protein (MBP) was used as exogenous kinase substrate. The reaction mixture was incubated at room temperature for variable periods of time and the reaction was stopped with 2x sample buffer. Samples were heated at 95°C for 5 min and separated by SDS-PAGE and transferred on Immobilon-P membrane. Phosphorylation signals were detected using a Phosphorimager (FLA-3000, Fujifilm, Tokyo, Japan). Equal loading was verified by Western blotting with a c-Kit antibody. To eliminate background phosphorylation on phosphoserine, alkali treatment of filters was performed essentially according to (19)

Cell proliferation and survival assay-Ba/F3 cells were washed three times with RPMI-1640 medium and seeded in 24-well plates (70,000 cells/well). Cells were then incubated either with 100 ng/ml SCF, without cytokine or with 10 ng/ml IL-3 for 48 hours. Viable cells were counted using trypan blue exclusion method. Alternatively, cells were stained by Click-iT EdU Alexa 647 (Life Technologies) cell proliferation kit employing the manufacturer's protocol. Stained cells were then analyzed by flow cytometry (BD FACSCalibur). Apoptosis was measured using an Annexin-V, 7-Amino-actinomycin D (7-AAD) kit (BD Biosciences Pharmingen), according to the manufacturer's instructions; double negative (Annexin-V)/(7-AAD) cells represent viable cells.

Internalization experiment-For internalization assay, Ba/F3 cells were incubated with 100 μ g/ml of cycloheximide and starved for 4 hours at 37°C in RPMI-1640 medium lacking serum and cytokines. Cells were stimulated with 100 ng/ml SCF for the indicated periods of time. In order to assess internalization, cells were stained with a PE-labeled anti-c-Kit antibody and surface expression of c-Kit receptor was determined by flow cytometry. Alternatively, receptor internalization was verified by Western blotting. Freshly prepared solution of 0.2 mg/ml EZ-Link Sulfo-NHS-Biotin (Thermo Scientific) dissolved in PBS was added to cells. The cells were then incubated on ice for 40 minutes to allow biotinylation of cell surface proteins. After incubation, excessive biotin was removed by a washing once with cold PBS. In addition, cold 50 mM Tris was added to the cells to block excessive reactive biotin. After 5 min. of blocking, cells were lysed and processed for pulldown with immobilized Avidin agarose (Thermo Scientific). The supernatant obtained after centrifugation was subjected to immunoprecipitation with anti-c-Kit antibody. c-Kit from both fractions was detected by Western blotting.

Degradation experiments-Ba/F3 cells were incubated with 100 μ g/ml of cycloheximide, starved followed by stimulation with SCF for the indicated periods of time. For protein degradation experiments, ligand stimulation was followed by lysis and immunoprecipitation with a c-Kit antibody. Cell lysates obtained after stimulation were subjected to immunoprecipitation followed by detection of c-Kit by Western blotting using an antibody against c-Kit.

RESULTS

The activation loop Y823F mutation accelerates receptor phosphorylation. The kinase domain of c-Kit is known to interact with juxtamembrane (JM) domain to maintain the kinase in an inhibitory state. When four tyrosine residues of JM domain at positions 547, 553, 568 and 570 are phosphorylated upon ligand stimulation, the JM domain is released from the kinase domain and the substrate can interact with the catalytic site. In addition, the activation loop transits from the DFG-out to the DFG-in state in transition from the unactivated state to the activated state. Together these two changes make the kinase catalytically active (13). We wanted to assess the changes in kinase activity of the receptor if the activation loop tyrosine, Y823, was mutated. Therefore, we generated the Y823F mutant of c-Kit and stably transfected it into Ba/F3 cells. Ba/F3 is an immortalized murine bone marrow-derived pro-B-cell line which is dependent on IL-3 for its growth(20). We have used Ba/F3 cells in the current study because this cell line does not express c-Kit endogenously and therefore can be used to overexpress both wild-type and mutant forms of human c-Kit. We also performed experiments with wild-type c-Kit and Y823F c-Kit by transient transfections in Cos1 cells. Both cell lines were tested for cell surface expression of c-Kit by flow cytometry (Fig. 1A). In order to verify, that overexpression of c-Kit wt and c-Kit Y823F was comparable to endogenous expression of c-Kit, we used two leukemic cell lines, Mo7e and Kasumi as controls (Fig. 1B). We further observed that both the wild-type c-Kit as well as Y823F c-Kit are rapidly phosphorylated after stimulation with SCF. However, the onset of phosphorylation in the mutant was faster and stronger than in wild-type c-Kit, it rapidly declined in the Y823F mutant whereas wild-type c-Kit showed a more sustained response (Fig. 1 C). This suggests that there is either a conformational change in the Y823F mutant or that the mutant c-Kit stability is altered.

Ubiquitination and degradation are more rapid in the Y823F mutant than in wild-type c-Kit. It has been previously shown that RTKs upon ligand stimulation undergo mono-ubiquitination or poly-ubiquitination that targets them to be internalized and degraded in the lysosomes or the proteasomes, respectively (1). Previous studies demonstrate that Cbl which belongs to a family of ubiquitin E3 ligases is

crucial for ubiquitination of c-Kit (1,7,8). Furthermore, Cbl can either directly interact with c-Kit at Y568 and Y936 (1) or indirectly through the adapter protein Grb2 (8). This leads to mono-ubiquitination of c-Kit and targets it for degradation in lysosomes. We wanted to investigate whether mutation of Y823 in the activation loop of c-Kit affects the phosphorylation of Cbl which could result in changes in ubiquitination which in turn could affect receptor degradation. Cbl was immunoprecipitated from both wild-type and Y823F mutant c-Kit following ligand stimulation and extent of phosphorylation of Cbl was analyzed. The results clearly demonstrate that Cbl is phosphorylated already after 2 min. of SCF stimulation in both c-Kit wt and c-Kit-Y823F but the mutant is unable to sustain the phosphorylation (Fig. 2A). In order to assess the degree of ubiquitination, c-Kit was immunoprecipitated and Western blotting was performed using an antibody against ubiquitin. In agreement with the Cbl phosphorylation data, ubiquitination of c-Kit Y823F is very strong after 2 min. of SCF stimulation but then rapidly declines. In contrast, ubiquitination of wild-type c-Kit is weaker but increases over a longer period of time (Fig. 2B). Degradation of c-Kit was analyzed by starving Ba/F3 cells in presence of cycloheximide, an inhibitor of protein synthesis, for 4h followed by immunoprecipitation and detection using anti-c-Kit antibody. Degradation of c-Kit receptor was analysed for up to 30 min. of SCF stimulation. We observed that in wild-type c-Kit expressing cells, c-Kit is detectable for up to 30 min. whereas in cells expressing the Y823F mutant, c-Kit is degraded before 15 min. of SCF stimulation (Fig., 2 C and D).

A change in activation loop Y823 to Y823F causes the receptor to internalize faster. It is known that together with JM domain, activation loop also plays a role in maintaining the receptor in an unactivated state. Phosphorylation of Y823 and of tyrosine residues in JM domain is crucial to maintain the receptor in an active state. Moreover, previous reports show that Cbl in addition to regulating degradation of RTKs also regulates their internalization. Thus, we analyzed the pattern of receptor internalization in response to ligand activation. Ba/F3 cells were serum-starved in presence of cycloheximide for 4h and stimulated with SCF for indicated periods of time. Cells were stained with a PE-labeled anti-c-Kit

antibody and surface expression was determined by flow cytometry (Fig 3A). As an alternative method we verified the rate of internalization of cell surface c-Kit compared with the total c-Kit by Western blotting (Fig. 3B and C). The data clearly demonstrate that the Y823F mutant of c-Kit is more rapidly internalized than wild-type c-Kit.

Mutation of activation loop tyrosine 823 has no effect on kinase activity. Activation of c-Kit by its ligand triggers activation of numerous downstream signal transduction molecules. Some of these molecules are tyrosine kinases by themselves, e. g. Src family kinases such as Lyn and the Fes kinase (21,22) and have been demonstrated to be able to phosphorylate c-Kit (23). Thus, the phosphorylation status of c-Kit in living cells could be influenced by other factors than the intrinsic kinase activity of c-Kit itself. Therefore we wanted to evaluate the kinase activity of wild-type c-Kit and the Y823F c-Kit mutant in an in vitro assay. To test whether activation loop tyrosine has an effect on kinase activity, both wildtype and Y823F mutant c-Kit were transiently transfected into Cos1 cells and stimulated with SCF. Immunoprecipitates of c-Kit from these cells were incubated with myelin basic protein (MBP) as an exogenous substrate and an in vitro kinase reaction was performed. No change in kinase activity was observed between the wild-type and the Y823F mutated c-Kit receptor (Fig. 4 A and B) which is in agreement with previous reports (13).

Mutation of Y823 negatively affects Akt and Erk downstream signaling pathways. Activation of c-Kit upon ligand stimulation initiates sequential recruitment of several signaling molecules which form a network between the receptor and the cell nucleus. This networking is governed by several regulatory signal pathways including the Ras/Erk, p38, and PI3K/Akt pathways (24). In order to investigate how mutation of Y823 affects signal transduction, the phosphorylation of Akt, Erk, and p38 was examined by Western blotting using respective phospho-specific antibodies. Whereas Ba/F3 cells expressing wild-type c-Kit responded to SCF stimulation with phosphorylation of Akt that persisted for longer time, activation of Ba/F3 cells expressing the Y823F mutant of c-Kit showed strong but transient phosphorylation of Akt (Fig., 5A and B). A similar trend was observed with Erk and p38 phosphorylation (Fig., 5A and B).

Effect of Y823F mutation on phosphorylation of adaptor proteins. Adaptor proteins such as Cbl, Shc, SHP2 and Gab2 are known to regulate the signal transduction through c-Kit either directly or through other adaptor proteins which in turn affect the downstream signaling pathways (24). We further analyzed the up-stream adaptor proteins that affect cell proliferation and cell survival. SCF stimulated Ba/F3 cells were subjected to immunoprecipitation using specific antibodies and phosphorylation of each adaptor protein was detected using phospho-specific antibodies. We observed that phosphorylation of Cbl, Shc and Gab2 was reduced in cells expressing Y823F c-Kit, compared to cells expressing wild-type c-Kit, while SHP2 phosphorylation was not significantly affected by Y823F mutation (Fig., 6A and B).

Mutation of Y823 leads to a significant decrease in both cell survival and proliferation. We wanted to ascertain if the mutation of activation loop tyrosine in c-Kit had an effect on survival and proliferation of cells. The cells lacking Y823 showed a significant reduction in proliferation as confirmed by EdU incorporation in proliferating cells and trypan blue exclusion method (Fig., 7A and B). The effect of the Y823F mutation of c-Kit on cell survival was also examined by staining the cells with Annexin V and 7-Amino Actinomycin D and analyzing by flow cytometry. Ba/F3 cell expressing the Y823F mutant of c-Kit showed almost 60% reduction in cell survival compared to cells expressing wild-type c-Kit (Fig. 7C).

DISCUSSION

The activation loop of c-Kit spans about 20-25 amino acids in the C-lobe of kinase domain and together with the juxtamembrane domain maintains the kinase in an auto-inhibitory state. Studies indicate that two key processes are required for activation of kinase, one the release of JM domain that exposes the catalytic site to the substrate and second, the activation loop coming to the DFG-in state. We have observed that c-Kit phosphorylation is not hampered by the Y823F mutation. Y823 is the only tyrosine phosphorylation site in the activation loop and yet the previous structural studies point to its irrelevance in the kinase activation process. So what is Y823 doing in the activation loop? Does it have any role in downstream signaling through c-Kit receptor?

To date very little is known about the role of the activation loop tyrosines in other receptors other than in the regulation of kinase activity. In non-receptor tyrosine kinase, Syk phosphorylation of activation loop tyrosines is described crucial for propagation of signaling through the immunoreceptor but does not affect the kinase activity of Syk (25). In an early report from Server and co-workers (26), Y821 in murine c-Kit (homologous to Y823 in human c-Kit) was shown to be of importance for proliferation and survival in murine bone marrow-derived mast cells without affecting PI3-kinase, p21ras or Erk activation. However, the discrepancy with our data on Erk activation could be explained by the use of different cell systems. The EGFR Y845, analogous to Y823 in c-Kit, was demonstrated to be required in mitogenic pathways and as a mediator for an association between CoxII and EGFR crucial for cell survival (27,28). In both c-Kit Y821 and EGFR Y845, a phenylalanine mutant showed reduced cell survival. In PDGF receptor, activation loop Y857 was shown to be crucial for *in-vitro* kinase activity and for cell proliferation but did not affect internalization (29). In our study we show that Y823F affects downstream signaling pathways of c-Kit, it is fully dispensable for kinase activity and that the mutated receptor is internalized and degraded at a much accelerated rate compared to the wild-type receptor. Mutation of Y823 to Y823F further significantly reduced cell proliferation and survival. Thus, our study adds new perspectives to existing knowledge about the role of activation loop tyrosine (Y823) in c-Kit signaling.

Previous studies on the kinetics of phosphorylation of c-Kit have provided valuable information on the role of Y823 in c-Kit activation (13). However, those studies were performed on a recombinant intracellular fragment of c-Kit, and thus lacking the ligand-binding domain. Furthermore, in a cell free system the other tyrosine kinases and molecules that affect c-Kit signaling (such as Src and Fes) are lacking and their contribution to phosphorylation of the receptor is not seen. Phosphorylation, ubiquitination, and degradation experiments on ligand-induced c-Kit activation demonstrate that the Y823F mutant of c-Kit is able to transduce a phosphorylation signal but at a much accelerated rate and considerably more transient in its nature compared to wild-type c-Kit. Cbl, a ubiquitin E3 ligase is known to

regulate ubiquitination and degradation of receptor tyrosine kinases. We observed that phosphorylation of Cbl increases when we increase the ligand stimulation time in wild type receptor whereas it decreases in Y823F. The effect on short lived ubiquitination and faster degradation could also be due to instant dephosphorylation after a brief phosphorylation of Cbl. The mutant receptor is also internalized much faster than the wild type receptor. This could again be explained by the phosphorylation status of Cbl which is known to regulate internalization (1). Thus, the effects on ubiquitination, internalization and degradation could be the consequence of Cbl inactivation which can interact with c-Kit either directly or through activation of Src kinases. Further, phosphorylation of adaptor proteins such as Gab2 and Shc is also reduced in Y823F as compared to the wild type receptor. These molecules are downstream to SFK (Src Family Kinases) which directly interact with c-Kit (30,31). Gab2 and Shc are connected to Akt and Erk pathways downstream (21,24). Therefore a reduction in Akt and Erk1/2 phosphorylation could be explained by altered Shc and Gab2 signaling molecules. Reduced Akt and Erk1/2 phosphorylation reduce cell survival and cell proliferation which are linked to Akt and Erk pathways. Similar observations have been made in PDGF β receptor where the activation loop Y857F mutation hampers complete activation of Akt and Erk (29). On the contrary, Y857F also led to reduced SHP2 activation which was not the case with Y823F in c-Kit. A reduction in SHP2 phosphorylation has been proposed to lead to reduction in Erk phosphorylation. Since SHP2 is not affected in our case, there are likely to be an alternate pathway affecting Erk signaling (29,32). One likely candidate for this is the adapter protein Shc, since we have previously shown that Src-phosphorylation of Shc is essential for the ability of c-Kit to mediate activation of Erk (21). Similarly, transient phosphorylation of p38 could also be caused by a reduction in Src-mediated Shc phosphorylation (33,34). Its role in cell migration has also been previously described (35). Thus, mutation in activation loop tyrosine, Y823 affects multiple signaling pathways as also described for Syk kinase where activation loop tyrosines are crucial for sustained downstream signaling without being involved in catalysis (36,37). Based on previous studies and our own current data, a perspective towards function of activation loop

tyrosine is that when the c-Kit receptor dimerizes and autophosphorylates, the potential tyrosine residues on JM domain are phosphorylated and release the auto-inhibitory state. Next is the phosphorylation of activation loop tyrosine that upon phosphorylation makes the loop in DFG-in state and locks the kinase from an unactivated state into an activated state. But when Y823 is mutated to F823, there is a conformational change and a de-stabilization of the kinase such that the activated state is no longer maintained. It could be due to this reason that signaling through the mutated receptor initiates normally but is not sustained due to a destabilized receptor which leads to a faster internalization, degradation and a marked reduction in cell viability and cell proliferation capacity. However, a change in conformation is suspected but it excludes the domain binding to kinase inhibitor, sunitinib. As proposed by DiNitto et al, the Y823F mutation could render the enzyme more flexible such that after activation it comes back to DFG-out unactivated state preferred by most kinase inhibitors.

It turns out that the effect of mutating Y823 is very much dependent on which amino acid replaces the tyrosine at position 823 is critical. DiNitto and coworkers demonstrated in their in vitro cell-free system that the Y823A and Y823D were devoid of kinase activity while the Y823F mutant has full activity (13). It is interesting that the Y823D mutation of c-Kit has been described in human testicular seminomas (38), malignant melanomas (39), gastrointestinal stromal tumors (40) and pediatric core-factor binding acute myeloid leukemia (41). The Y823D mutant was found to be constitutively active in living cells (38). This apparent

discrepancy in results might be attributable to the fact that in the living cell c-Kit has access to accessory molecules, that is missing in a cell free system, that are necessary for its signaling. Furthermore, the Y823C (38) and Y823N (42) mutants have also been identified in tumors, but no data are available as to whether these mutations also are activating. Thus, the amino acid that substitutes the tyrosine residues is critical for signaling outcome. In one case it might lead to constitutive activation, in another case to modulation of signaling and in a third case to inactivation of the kinase.

In view of our results, Y823 appears to be a good target for cancer therapy. The Y823F makes the receptor not only more sensitive to therapeutic targets such as Sunitinib and Imatinib but also de-stabilizes the receptor such that activation of Akt, Erk1/2 and p38 is reduced. This in turn leads to a significant reduction in cell proliferation and cell survival. Thus, a therapy targeting Y823 such that its phosphorylation is prevented could, in combination with chemotherapy, provide an improved treatment option for tumors caused by c-Kit mutations. This study exemplifies that even when sequence of kinase domains is highly conserved across members of the RTK family, point mutations of various receptors play divergent roles in cellular outcomes and in the mechanism of signal transduction. This underlines the importance of investigating further point mutations in c-Kit and the mechanisms by which they influence cell signaling, to improve our current understanding of the association of receptor mutations with cancer prognosis.

REFERENCES

1. Masson, K., Heiss, E., Band, H., and Rönnstrand, L. (2006) Direct binding of Cbl to Tyr568 and Tyr936 of the stem cell factor receptor/c-Kit is required for ligand-induced ubiquitination, internalization and degradation. *Biochem J* **399**, 59-67
2. Lennartsson, J., and Rönnstrand, L. (2012) Stem cell factor receptor/c-Kit: from basic science to clinical implications. *Physiol Rev* **92**, 1619-1649
3. Hirota, S., Isozaki, K., Moriyama, Y., Hashimoto, K., Nishida, T., Ishiguro, S., Kawano, K., Hanada, M., Kurata, A., Takeda, M., Muhammad Tunio, G., Matsuzawa, Y., Kanakura, Y., Shinomura, Y., and Kitamura, Y. (1998) Gain-of-function mutations of c-kit in human gastrointestinal stromal tumors. *Science* **279**, 577-580
4. Nishida, T., Hirota, S., Taniguchi, M., Hashimoto, K., Isozaki, K., Nakamura, H., Kanakura, Y., Tanaka, T., Takabayashi, A., Matsuda, H., and Kitamura, Y. (1998) Familial gastrointestinal stromal tumours with germline mutation of the KIT gene. *Nat Genet* **19**, 323-324

5. Kitamura, Y., Hirota, S., and Nishida, T. (2001) A loss-of-function mutation of c-kit results in depletion of mast cells and interstitial cells of Cajal, while its gain-of-function mutation results in their oncogenesis. *Mutat Res* **477**, 165-171
6. Lennartsson, J., Jelacic, T., Linnekin, D., and Shivakrupa, R. (2005) Normal and oncogenic forms of the receptor tyrosine kinase kit. *Stem Cells* **23**, 16-43
7. Zeng, S., Xu, Z., Lipkowitz, S., and Longley, J. B. (2005) Regulation of stem cell factor receptor signaling by Cbl family proteins (Cbl-b/c-Cbl). *Blood* **105**, 226-232
8. Sun, J., Pedersen, M., Bengtsson, S., and Rönnstrand, L. (2007) Grb2 mediates negative regulation of stem cell factor receptor/c-Kit signaling by recruitment of Cbl. *Exp Cell Res* **313**, 3935-3942
9. Furdui, C. M., Lew, E. D., Schlessinger, J., and Anderson, K. S. (2006) Autophosphorylation of FGFR1 kinase is mediated by a sequential and precisely ordered reaction. *Mol Cell* **21**, 711-717
10. Hubbard, S. R. (1997) Crystal structure of the activated insulin receptor tyrosine kinase in complex with peptide substrate and ATP analog. *EMBO J* **16**, 5572-5581
11. Mol, C. D., Dougan, D. R., Schneider, T. R., Skene, R. J., Kraus, M. L., Scheibe, D. N., Snell, G. P., Zou, H., Sang, B. C., and Wilson, K. P. (2004) Structural basis for the autoinhibition and STI-571 inhibition of c-Kit tyrosine kinase. *J Biol Chem* **279**, 31655-31663
12. Mol, C. D., Lim, K. B., Sridhar, V., Zou, H., Chien, E. Y., Sang, B. C., Nowakowski, J., Kassel, D. B., Cronin, C. N., and McRee, D. E. (2003) Structure of a c-kit product complex reveals the basis for kinase transactivation. *J Biol Chem* **278**, 31461-31464
13. DiNitto, J. P., Deshmukh, G. D., Zhang, Y., Jacques, S. L., Coli, R., Worrall, J. W., Diehl, W., English, J. M., and Wu, J. C. (2010) Function of activation loop tyrosine phosphorylation in the mechanism of c-Kit auto-activation and its implication in sunitinib resistance. *J Biochem* **147**, 601-609
14. Blume-Jensen, P., Siegbahn, A., Stabel, S., Heldin, C. H., and Rönnstrand, L. (1993) Increased Kit/SCF receptor induced mitogenicity but abolished cell motility after inhibition of protein kinase C. *Embo J* **12**, 4199-4209
15. Lupher, M. L., Jr., Reedquist, K. A., Miyake, S., Langdon, W. Y., and Band, H. (1996) A novel phosphotyrosine-binding domain in the N-terminal transforming region of Cbl interacts directly and selectively with ZAP-70 in T cells. *J Biol Chem* **271**, 24063-24068
16. Sun, J., Pedersen, M., and Rönnstrand, L. (2008) Gab2 is involved in differential phosphoinositide 3-kinase signaling by two splice forms of c-Kit. *J Biol Chem* **283**, 27444-27451
17. Sun, J., Pedersen, M., and Rönnstrand, L. (2009) The D816V mutation of c-Kit circumvents a requirement for Src family kinases in c-Kit signal transduction. *J Biol Chem* **284**, 11039-11047
18. Hansen, K., Johnell, M., Siegbahn, A., Rorsman, C., Engstrom, U., Wernstedt, C., Heldin, C. H., and Ronnstrand, L. (1996) Mutation of a Src phosphorylation site in the PDGF beta-receptor leads to increased PDGF-stimulated chemotaxis but decreased mitogenesis. *EMBO J* **15**, 5299-5313
19. Kamps, M. P., and Sefton, B. M. (1989) Acid and base hydrolysis of phosphoproteins bound to immobilon facilitates analysis of phosphoamino acids in gel-fractionated proteins. *Anal Biochem* **176**, 22-27
20. Kazi, J. U., Sun, J., and Rönnstrand, L. (2013) The presence or absence of IL-3 during long-term culture of Flt3-ITD and c-Kit-D816V expressing Ba/F3 cells influences signaling outcome. *Exp Hematol*

21. Lennartsson, J., Blume-Jensen, P., Hermanson, M., Pontén, E., Carlberg, M., and Rönstrand, L. (1999) Phosphorylation of Shc by Src family kinases is necessary for stem cell factor receptor/c-kit mediated activation of the Ras/MAP kinase pathway and c-fos induction. *Oncogene* **18**, 5546-5553
22. Smith, J. A., Samayawardhena, L. A., and Craig, A. W. (2010) Fps/Fes protein-tyrosine kinase regulates mast cell adhesion and migration downstream of Kit and beta1 integrin receptors. *Cell Signal* **22**, 427-436
23. Lennartsson, J., Wernstedt, C., Engström, U., Hellman, U., and Rönstrand, L. (2003) Identification of Tyr900 in the kinase domain of c-Kit as a Src-dependent phosphorylation site mediating interaction with c-Crk. *Exp Cell Res* **288**, 110-118
24. Masson, K., and Rönstrand, L. (2009) Oncogenic signaling from the hematopoietic growth factor receptors c-Kit and Flt3. *Cell Signal* **21**, 1717-1726
25. Zhang, J., Billingsley, M. L., Kincaid, R. L., and Siraganian, R. P. (2000) Phosphorylation of Syk activation loop tyrosines is essential for Syk function. An in vivo study using a specific anti-Syk activation loop phosphotyrosine antibody. *J Biol Chem* **275**, 35442-35447
26. Serve, H., Yee, N. S., Stella, G., Sepp-Lorenzino, L., Tan, J. C., and Besmer, P. (1995) Differential roles of PI3-kinase and Kit tyrosine 821 in Kit receptor-mediated proliferation, survival and cell adhesion in mast cells. *EMBO J* **14**, 473-483
27. Boerner, J. L., Demory, M. L., Silva, C., and Parsons, S. J. (2004) Phosphorylation of Y845 on the epidermal growth factor receptor mediates binding to the mitochondrial protein cytochrome c oxidase subunit II. *Mol Cell Biol* **24**, 7059-7071
28. Boerner, J. L., Biscardi, J. S., Silva, C. M., and Parsons, S. J. (2005) Transactivating agonists of the EGF receptor require Tyr 845 phosphorylation for induction of DNA synthesis. *Mol Carcinog* **44**, 262-273
29. Wardega, P., Heldin, C. H., and Lennartsson, J. (2010) Mutation of tyrosine residue 857 in the PDGF beta-receptor affects cell proliferation but not migration. *Cell Signal* **22**, 1363-1368
30. Daub, H., Weiss, F. U., Wallasch, C., and Ullrich, A. (1996) Role of transactivation of the EGF receptor in signalling by G-protein-coupled receptors. *Nature* **379**, 557-560
31. Luttrell, L. M., Daaka, Y., Della Rocca, G. J., and Lefkowitz, R. J. (1997) G protein-coupled receptors mediate two functionally distinct pathways of tyrosine phosphorylation in rat 1a fibroblasts. Shc phosphorylation and receptor endocytosis correlate with activation of Erk kinases. *J Biol Chem* **272**, 31648-31656
32. Rönstrand, L., Arvidsson, A. K., Kallin, A., Rorsman, C., Hellman, U., Engström, U., Wernstedt, C., and Heldin, C. H. (1999) SHP-2 binds to Tyr763 and Tyr1009 in the PDGF beta-receptor and mediates PDGF-induced activation of the Ras/MAP kinase pathway and chemotaxis. *Oncogene* **18**, 3696-3702
33. Ueda, S., Mizuki, M., Ikeda, H., Tsujimura, T., Matsumura, I., Nakano, K., Daino, H., Honda, Z., Sonoyama, J., Shibayama, H., Sugahara, H., Machii, T., and Kanakura, Y. (2002) Critical roles of c-Kit tyrosine residues 567 and 719 in stem cell factor-induced chemotaxis: contribution of src family kinase and PI3-kinase on calcium mobilization and cell migration. *Blood* **99**, 3342-3349
34. Samayawardhena, L. A., Hu, J., Stein, P. L., and Craig, A. W. (2006) Fyn kinase acts upstream of Shp2 and p38 mitogen-activated protein kinase to promote chemotaxis of mast cells towards stem cell factor. *Cell Signal* **18**, 1447-1454
35. Sundström, M., Alfredsson, J., Olsson, N., and Nilsson, G. (2001) Stem cell factor-induced migration of mast cells requires p38 mitogen-activated protein kinase activity. *Exp Cell Res* **267**, 144-151

36. Carsetti, L., Laurenti, L., Gobessi, S., Longo, P. G., Leone, G., and Efremov, D. G. (2009) Phosphorylation of the activation loop tyrosines is required for sustained Syk signaling and growth factor-independent B-cell proliferation. *Cell Signal* **21**, 1187-1194
37. Kulathu, Y., Grothe, G., and Reth, M. (2009) Autoinhibition and adapter function of Syk. *Immunol Rev* **232**, 286-299
38. Kemmer, K., Corless, C. L., Fletcher, J. A., McGreevey, L., Haley, A., Griffith, D., Cummings, O. W., Wait, C., Town, A., and Heinrich, M. C. (2004) KIT mutations are common in testicular seminomas. *Am J Pathol* **164**, 305-313
39. Beadling, C., Jacobson-Dunlop, E., Hodi, F. S., Le, C., Warrick, A., Patterson, J., Town, A., Harlow, A., Cruz, F., 3rd, Azar, S., Rubin, B. P., Muller, S., West, R., Heinrich, M. C., and Corless, C. L. (2008) KIT gene mutations and copy number in melanoma subtypes. *Clin Cancer Res* **14**, 6821-6828
40. Calabuig-Farinas, S., Lopez-Guerrero, J. A., Navarro, S., Machado, I., Poveda, A., Pellin, A., and Llombart-Bosch, A. (2011) Evaluation of prognostic factors and their capacity to predict biological behavior in gastrointestinal stromal tumors. *Int J Surg Pathol* **19**, 448-461
41. Goemans, B. F., Zwaan, C. M., Miller, M., Zimmermann, M., Harlow, A., Meshinchi, S., Loonen, A. H., Hahlen, K., Reinhardt, D., Creutzig, U., Kaspers, G. J., and Heinrich, M. C. (2005) Mutations in KIT and RAS are frequent events in pediatric core-binding factor acute myeloid leukemia. *Leukemia* **19**, 1536-1542
42. Willmore-Payne, C., Holden, J. A., Chadwick, B. E., and Layfield, L. J. (2006) Detection of c-kit exons 11- and 17-activating mutations in testicular seminomas by high-resolution melting amplicon analysis. *Mod Pathol* **19**, 1164-1169

ACKNOWLEDGEMENTS

We would like to thank Susanne Bengtsson for kind technical assistance. We thank Jianmin Sun, Elena Razumovskaya and Vaibhav Agarwal for help with usage of various software.

FOOTNOTES

* This work was supported by grants from the Swedish Research Council, the Swedish Cancer Foundation, Gunnar Nilssons Cancer foundation.

¹To whom correspondence may be addressed: Experimental Clinical Chemistry, Wallenberg Laboratory, Department of Laboratory Medicine, Lund University, Skåne University Hospital, 20502 Malmö, Sweden, Tel.: +46 40 33 72 22, Fax: +46 40 33 11 04, E-mail: Lars.Ronnstrand@med.lu.se

²The abbreviations used are: RTK, receptor tyrosine kinase; Mitogen-activated protein kinase; PI3K, Phosphoinositide 3-kinase; SCF, stem cell factor; SH2, Src-homology2; EGFR, Epidermal growth factor receptor; IL-3, Interleukin-3; 7-AAD, 7-Amino Actinomycin D; JM, Juxtamembrane; PE, Phycoerythrin.

FIGURE LEGENDS

FIGURE 1: Y823F mutation leads to enhanced and faster phosphorylation of c-Kit receptor.

A, stably transfected Ba/F3 cells were labeled with phycoerythrin-conjugated anti-c-Kit antibodies or an isotype control and analyzed by flow cytometry. The *black peak* indicates cells labeled with the isotype control, and the *gray peak* corresponds to the cells labeled with anti-c-Kit antibody. B, Mo7e, and Kasumi cell lines expressing endogenous c-Kit and Ba/F3 cells expressing c-Kit and c-Kit/Y823F were lysed and subjected to immunoprecipitation with anti-c-Kit antibody. C-Kit expression levels were detected by Western blotting. C Ba/F3 cells were stably transfected with Ba/F3-c-Kit and Ba/F3-c-Kit/Y823F constructs. Cells were serum-starved overnight at 37°C and stimulated by SCF for indicated time periods. Cell lysates were prepared, immunoprecipitated (IP) with anti-c-Kit antibody

and analyzed by Western blotting. Total receptor phosphorylation was detected using phosphotyrosine (pY) antibody and c-Kit was used as a loading control.

FIGURE 2. The Y823F mutant of c-Kit mediates increased phosphorylation of Cbl concomitant with increased ubiquitination and degradation of c-Kit. *A*, Ba/F3-c-Kit and Ba/F3-c-Kit/Y823F cells were serum-starved and stimulated with 100 ng/ml SCF for the indicated times. Cells were lysed, and lysates were immunoprecipitated (*IP*) with anti-Cbl antibody, followed by Western blotting with phosphotyrosine antibodies and with a c-Kit antibody. *IB*, immunoblot. *B*, Lysates from serum-starved Ba/F3-c-Kit and Ba/F3-c-Kit/Y823F cells were immunoprecipitated (*IP*) with anti-c-Kit antibody. Ubiquitination of receptor was detected using anti-Ub antibody. *C*, Receptor degradation was assessed by serum starvation of Ba/F3-c-Kit and Ba/F3-c-Kit/Y823F cells for 4h at 37°C in presence of 100µg/ml of cycloheximide. Cells were then stimulated with 100 ng/ml SCF for indicated periods of time and cells were immediately placed on ice. Cell lysates were prepared and were subjected to immunoprecipitation (*IP*) with c-Kit antibodies followed by immunodetection using c-Kit antibody. *D*, Signal intensities from two independent experiments were quantified using Multi-Gauge software to calculate the percentage of receptor degradation.

FIGURE 3. The Y823F mutation of the activation loop of c-Kit enhances receptor internalization. Ba/F3-c-Kit and Ba/F3-c-Kit/Y823F cells were serum-starved for 4h at 37°C in presence of cycloheximide and stimulated with 100 ng/ml SCF for the indicated times. *A*, Cells were transferred to ice followed by incubation with phycoerythrin-conjugated anti-c-Kit antibody. The c-Kit surface expression level was analyzed by flow cytometry. Internalization of c-Kit was quantified at various time points compared with unstimulated cells. To determine mean fluorescence intensities (MFI) of wild-type receptor with mutated receptor, FloJo software was used. *B*, Cells were labeled with Sulfo-NHS-Biotin and incubated on ice for 40 minutes to allow biotinylation of cell surface proteins. Cells were lysed and processed for pulldown with immobilized Avidin. The supernatant obtained after centrifugation was subjected to immunoprecipitation with anti-c-Kit antibody. c-Kit from both fractions was detected by Western blot. *C*, Internalization of c-Kit was quantified at various time points and statistically analyzed using GraphPad Prism, ***, $p < 0.001$.

FIGURE 4. Y823F mutation in activation loop does not affect receptor kinase activity. Cos1-c-Kit-WT and Cos1-c-Kit/Y823F cells were serum-starved overnight and stimulated with 100 ng/ml SCF. Cell lysates were prepared and c-Kit was immunoprecipitated. *A*, Cell lysates were incubated with protein G beads, washed and subjected to in vitro kinase assay with $\gamma^{32}\text{P}$ -ATP and MBP as exogenous substrate. Phosphorylation signal was detected using L process software. c-Kit was detected by Western blotting. *B*, Calculation of relative MBP phosphorylation and statistical analysis was performed using GraphPad Prism, ns, not significant.

FIGURE 5. The Y823F mutation in c-Kit alters down-stream signaling. Ba/F3-c-Kit WT and Ba/F3-c-Kit/Y823F cells were serum-starved and treated with or without 100 ng/ml SCF for indicated periods of time. *A*, Total cell lysates were separated by SDS-PAGE, electrotransferred to Immobilon P membrane and probed with phospho-Akt antibody, phospho-Erk1/2 and phosphor p38 antibodies. Antibodies against Akt, Erk and p38 were used as loading controls. *B*, Signal intensities from three independent experiments were quantified using Multi-Gauge software to calculate the reduction and GraphPad Prism was used to calculate significance. ns, not significant, **, $p < 0.01$.

FIGURE 6. Phosphorylation mutant Y823F negatively regulates activation of adaptor proteins. *A*, Ba/F3-c-Kit-WT and Ba/F3-c-Kit/Y823F cells were serum-starved and incubated in the presence or absence of SCF for indicated period of time. Cell lysates were subjected to immunoprecipitation with antibodies against Gab2, Shc, Cbl and SHP2 respectively. Proteins were separated by SDS-PAGE, electrotransferred to Immobilon P membranes and probed either with phosphospecific antibodies or general phosphotyrosine antibodies (pY). *B*, Signal intensities from three independent experiments were quantified with Multi-Gauge software and statistical analysis was done using GraphPad Prism.

FIGURE 7. Cells expressing the Y823F mutant of c-Kit display reduced cell proliferation and cell survival in response to SCF stimulation. Ba/F3-c-Kit WT and Ba/F3-c-Kit/Y823F cells were grown for 48 h in the presence or absence of SCF and IL-3. *A*, cells were incubated with EdU for 2 h, fixed,

labeled with Alexa Fluor 647 and analyzed by flow cytometry. *B*, Viable cells were counted by trypan blue exclusion method *C*, cells were labeled with annexin V and 7-aminoactinomycin D, and living cells were measured by flow cytometry. IL-3 was used as a positive control. Quantification of labeled cells was obtained with FloJo software and results from three independent experiments were statistically analysed using GraphPad Prism, *ns*, not significant, $**p < 0.01$, $***, p < 0.001$.

Figure 1




Figure 2






Figure 3

A.



B.



C.





Figure 4

A.



B.





Figure 5

A.



B.





Figure 6

A.



B.




Figure 7

