

LUND UNIVERSITY

Can the APO method be used for measuring soft data?: A pilot study.

Strandberg, Eva-Lena; Ovhed, Ingvar; Håkansson, Anders; Troein, Margareta

Published in: Scandinavian Journal of Primary Health Care

DOI: 10.3109/02813432.2011.628233

2011

Link to publication

Citation for published version (APA): Strandberg, E.-L., Ovhed, I., Håkansson, A., & Troein, M. (2011). Can the APO method be used for measuring soft data?: A pilot study. Scandinavian Journal of Primary Health Care, 29(4), 241-246. https://doi.org/10.3109/02813432.2011.628233

Total number of authors: 4

General rights

Unless other specific re-use rights are stated the following general rights apply: Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

· Users may download and print one copy of any publication from the public portal for the purpose of private study

or research.
You may not further distribute the material or use it for any profit-making activity or commercial gain

· You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: https://creativecommons.org/licenses/

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117 221 00 Lund +46 46-222 00 00

Can the APO method be used for measuring soft data?

A pilot study.

Eva Lena Strandberg^{1,2 §}, Ingvar Ovhed², Anders Håkansson (deceased)¹, Margareta Troein¹

¹Lund University, Department of Clinical Sciences in Malmö, General

Practice/Family Medicine, Malmö, Sweden

²Blekinge County Council, Karlskrona, Sweden

[§]Corresponding author

Correspondence:

Eva Lena Strandberg, PhD, Researcher

Lund University, Department of Clinical Sciences, Malmö, General Practice/Family

Medicine, Skåne University Hospital,

SE-205 02 Malmö. Sweden

Email addresses:

ELS: <u>eva-lena.strandberg@med.lu.se</u>, <u>eva-lena.strandberg@ltblekinge.se</u>

IO: ingvar.ovhed@ltblekinge.se

MT: margareta.troein@med.lu.se

No. of words: 2000

Abstract: 250 words

Running title: Can the APO method measure soft data?

Key Points

The APO process is a pronounced bottom-up method for quality development and for influencing medical decision making in general practice.

- It was possible to use the APO method for measuring soft data, using knowledge to exemplify one aspect of holistic view.
- Our results indicate that the APO method could be an alternative to timeconsuming methods for studying the consultation.

Abstract

Objectives

The aims were to develop auditing according to the APO (Audit Project Odense) method for measuring soft data, exemplified by a holistic view, and to test the instrument.

Design

A descriptive study of the development of an APO chart and a test registration.

Setting

Primary health care, Blekinge County, Sweden.

Subjects

Ten general practitioners (GPs) were invited to transform categories of the concept of holistic view obtained in an earlier study, into 30 variables on an APO registration chart. The participants chose to study different kinds of knowledge as aspects of holistic care.

Main outcome measure

An APO registration chart and test of the instrument.

Results

After three meetings the group had drawn up an APO registration chart supplemented with Likert scales. A pilot audit was performed. Eight doctors registered 255 consultations. In assessment of the patients' problems, factual medical knowledge was important in 83% of the cases, familiarity in 53%, and a capacity for judgement in 36%. In decision making factual medical knowledge was used in 88% and capacity for judgement in 58%. A holistic view was necessary for the outcome in 43% and valuable in 25%. The GPs used the Likert scales in a majority of the cases. *Conclusions*

In this first step in developing an instrument, the results indicate that the APO method could be an alternative for studying what happens in the consultation, and the occurrence of an abstract phenomenon such as the use of different kinds of knowledge as part of a holistic view.

Keywords: general practice, consultation, development of measurement instruments, quality development, the APO method

Introduction

The National Board of Health and Welfare was commissioned by the Swedish government in 2002–2005 to follow up the national action plan for the development of medical care in Sweden. This included studying the watchwords of primary care: continuity, accessibility, cooperation, and a holistic view. A group of researchers was commissioned to perform the study. A qualitative focus group study of the meaning of the concept of holistic view was performed with general practitioners (GPs) and district nurses [1,2]. According to the participants the term holistic view could be divided into three main categories: attitude, knowledge, and circumstances (Table I).

The National Board also raised the question whether it was possible to measure the prevalence of consultations in which a holistic view was used in the decision making (Table II). We wanted to test the APO (Audit Project Odense) method, a wellestablished tool for quality development among Nordic GPs [3-7]. An APO audit is a simple and feasible method for common medical problems, which to our knowledge has never been used for soft data. We chose this opportunity to examine the possibilities and limitations of the APO tool. This participatory bottom-up process is based on a registration of consultations, where each physician registers a considerable amount of consultations during a limited period on a simple registration chart [3]. The participants decide about the variables, test them in a pilot audit, and adjust them for the real registration. Next, each participant receives his/her personal result, for comparison with colleagues in follow-up discussions between professionals.

The aims of this pilot study were to develop an audit instrument to measure soft data exemplified by the concept of holistic view, and to test the instrument.

- 5 -

Material and Methods

Previously, 22 GPs defined criteria for a holistic view. In this pilot study, all ten members of a CME (Continuing Medical Education) group were invited to develop an APO audit. Eight of them participated in the first study [1,2]. Their task was to transform concepts based on the earlier results into 30 measurable variables, and to test the registration chart.

All participants were specialists in family medicine with long experience, four women and six men. Eight doctors worked in villages, two in urban settings. All of them worked in clinics with several GPs, each with their own patient list, and they helped each other with patients.

Participation in an APO audit is voluntary and participant-centred. Every process starts with the participants defining central concepts of the topic in order to operationalize them on a registration chart [3]. This process started with a summary of the meaning of the categories from the previous study to reach common ground. The group met three times, led by a person with long experience of constructing APO registration charts.

The pilot audit was conducted in March 2007. The participants were asked to register all patients. The results were compiled in a report presenting the results of the entire group. The report and the final registration chart can be ordered from the corresponding author.

As described in the Results, Likert scales were introduced. The Likert scale assessments were analysed by dividing the ten-grade scale into three main groups (low value 1–3, neutral value 4–7, and high value 8–10).

- 6 -

Results

The registration chart takes form

The *first meeting* began with a run-through of the findings from the interview study about the concept of holistic view. The participants were encouraged to discuss the concept and how it could be measured. Each person stated his/her opinion about the different categories and how their presence in the consultation could be captured, in accordance with the APO method. The group chose to focus on one aspect of holistic view, the *concept of knowledge*, and on the presence and significance of knowledge for medical decision making.

The participants discussed the need for detailed information about the patient. At first, a traditional APO audit was regarded as too blunt to capture the presence of abstract phenomena in the decision making. Therefore they decided to add three tenstep Likert scales to capture more detailed information: Familiarity with the patient, Symptomatology, and Agreement/Discrepancy concerning the patient's stated reason for the consultation (Figure 1).

The scales were regarded as important cornerstones in the practical application of the concept of holistic view. The third scale was envisaged as a way for doctors to capture the very core of holistic thinking, i.e. to reveal a hidden agenda (if any); in other words, what is initially concealed to both the doctor and the patient.

At the *second meeting* further definitions of the concept of "knowledge" were added to the concepts in the previous study. The group introduced a subordinate level concerning familiarity. The group discussed the importance of knowledge about patients' networks, knowledge from the primary care team, written and oral knowledge, and the doctor's own "capacity for judgement" inspired by Pörn [8], concerning both well-founded factual knowledge and what the GP has learned from practice related to the particular individual in the actual consultation. The capacity for judgement is thus not transferable from one GP to another. Regarding factual knowledge, the group stressed that in medical contexts this is an essential type of knowledge. Sometimes no factual knowledge is available "here and now", and the group considered it important to reveal such situations, and situations where factual knowledge had not been used.

At the *third meeting* the variables were adjusted and instructions for registration were completed (Tables IIIa, IIIb).

The pilot audit

Eight of ten doctors handed in completed charts after two reminders. The GPs had registered a total of 255 consultations with an average of 32 per doctor. The distribution was between 25 and 52 (median=30) consultations per doctor. Of the 255 consultations, 60% were with a patient from the doctor's list.

In the *assessment* of the patient's problem (criteria 6–9), the doctors regarded special factual medical knowledge as important in 83% of the cases, familiarity with the patient in 53%, and the capacity for judgement in 36%.

In the *decision making* (criteria 10–27), factual knowledge was considered in 88% of the cases and the capacity for judgement in 58%. In contrast, knowledge obtained from the primary care team, orally from colleagues or others, and from

written sources had been considered in the decision making only to a limited extent, from 11% to 28%.

Evaluating the patient's problem, the doctors assessed a holistic view (criteria 28–31) necessary for the outcome in 43% of the consultations and valuable in 25%. In 31% of the cases a holistic view was not regarded as necessary. A holistic view was never considered negative.

In 88% of the consultations the doctors had filled in the Likert scale about *familiarity* with the patient, in 85% the scale about *symptomatology*, and in 85% the scale about the stated and the ultimate *reason for visit*.

Of all the patients, 47% were well known to the doctors. Fifteen percent had a complex symptomatology, and in 3% of the consultations the GP changed opinion about the main problem during the consultation. In a subset of the consultations when GPs met patients on their own lists only and completed the Likert scales, 72% of the patients were well known, 18% had a complex symptomatology, and in 2% of the consultations the doctor changed view on the reason for the visit.

Discussion

This pilot study concerns the possibilities and limitations of the APO method. In this first step in developing an instrument, we wanted to examine whether it is possible to measure abstract phenomena in GPs' clinical work with an APO audit. The collegial brainstorming that starts every APO process also worked in this context [3]. The registration chart was supplemented with three Likert scales, used in a majority of the consultations, giving more detailed information about the consultation. It was possible to *construct* an APO chart with almost exclusively "soft" variables. It was also

- 9 -

possible to *measure* the presence of one part of holistic view if one shares these doctors' idea of what a holistic view is.

A full APO audit includes construction of the registration chart, a pilot registration, a meeting for adjustments of variables and registration chart and then the intended registration, followed by a meeting to compare results. In this study the process stopped after the pilot registration.

The significance of our findings

An APO audit reflects everyday reality. The accuracy depends on how well the variables reflect important actions. It is a snapshot, which is used for discussions among colleagues about important values [3,4,9,10]. This pilot study shows alternative uses for the APO method, and introducing Likert scales led to a development of the instrument. Although this is a crude way to study a GP's everyday work, the advantage is the speed of data collection and processing/analysis compared with recordings, interviews, etc. An APO audit captures subjective experiences which describe and influence reality [9,11-14].

The formation of variables is strongly participant-centred. We lack an evaluation of the participants' views on the audit. In a full-scale audit concerning the use of different kinds of knowledge and a holistic view in GPs' decision making, these experiences should be utilized in the design of a final registration chart.

An APO audit, with its quick prospective and consecutive collection of data, is a method to reflect clinical work and decision making. The significance of factual knowledge was particularly discussed during the construction process of the registration chart as the basic form of knowledge for a GP, as stressed by Pendleton and Wulff & Götzsche [15,16].

Learning and development do not follow a set pattern and are not the same for everyone. In the personal learning process it is therefore essential to proceed from how "I" act and think in given situations, in order to improve. An APO audit, if honestly completed, exposes defects and demolishes myths. Behaviour is influenced best if discussions with colleagues are combined with feedback of personal data [10,17,18].

The pilot audit

The pilot audit shows that it is possible to use the APO method to measure soft data in decision making.

The Likert scales gave more detailed knowledge about the situations in which the doctors use their capacity for judgement, alone and in different combinations. Pendleton and Malterud stress the importance of the right questioning technique to arrive at difficult-to-obtain knowledge about the patients and their symptoms [16,19-21]. The APO instrument can serve to reflect the extent to which the doctor uses such models to discover what patients keep concealed.

The Likert scale concerning familiarity has a direct link to continuity, which is one of the watchwords of primary care also defined by WONCA Europe [15,22-27]. Continuity and the relationship to the patient are also an important part of treatment [28]. The concept of cooperation is reflected through the variables concerning consultation with colleagues and others, as well as the Likert scales concerning symptomatology and reason for visit.

Conclusions

We have not tried to answer the question whether the APO method could be used to measure holistic view in full. We have only focussed on the aspect of what kind of *knowledge* is used in the decision making. We know that an APO audit influences doctors' medical decision making in well-demarcated and commonly occurring cases such as respiratory tract infections [5,9,11,13]. We have tested the instrument in a new context and with a partly new methodology. To gain some idea about the prevalence of other aspects of a holistic view in the everyday work of GPs, other methods could be useful.

Soft data, such as a holistic view, can be studied with the APO method, provided that the definition of *knowledge* is accepted as an aspect of holistic view. The use of Likert scales made it possible to reach a deeper understanding of the consultation. The results indicate that the APO method could be an alternative to more time-consuming methods for studying the consultation.

Competing interests

The authors declare that they have no competing interests.

Acknowledgements

We thank the participating GPs for their important contribution. This study was supported by Blekinge County Council.

References

- Strandberg EL, Wilhelmsson S. "Utan helhetssyn ingen primärvård".
 (Without a holistic view no primary care. In Swedish). Karlshamn: Blekinge FoU-enhet, Rapport 2004:1, Sweden.
- Strandberg EL, Ovhed I, Borgquist L, Wilhelmsson S. The perceived meaning of a (w)holistic view among general practitioners and district nurses in Swedish primary care: a qualitative study. *BMC Fam Pract* 2007;8:8.
- Bentzen N. Medical audit The APO method in general practice. Scand J Prim Health Care 1993;11(Suppl 1):13-18.
- Munck AP, Hansen DG, Lindman A, Ovhed I, Førre S, Thorsteinsson JB. A Nordic collaboration on medical audit – The APO method for quality development and continuous medical education (CME) in primary health care. *Scand J Prim Health Care* 1998;16:2-6.
- Søndergaard J, Hansen DG, Aarslev P and Munck AP. A multifaceted intervention according to the Audit Project Odense method improved secondary prevention of ischemic heart disease: a randomised controlled trial. *Family Practice* 2006; 23: 198–202.
- Sheldon MG. *Medical Audit in General Practice*. Occasional Paper 20. London: The Royal College of General Practitioners, 1982.
- Crombie D, Fleming D. *Practice Activity Analysis*. Occasional Paper 41. London: The Royal College of General Practitioners, 1988.
- Pörn I. *Filosofi visdom för vår tid?* (In Swedish; Philosophy wisdom for our time?) In Venkula J. (ed.) Ajatuksen Voima. Jyväskylä: Edistyksellinen tiedeliitto, 1990.

- Hansen CN, Hansen DG, Kragstrup J, Busch O, Munck A. The role of the APO method in improving diabetes care in general practice: the results of a Danish prospective multipractice audit circle. *Qual Prim Care* 2003;1:225-232.
- Strandberg EL, Ovhed I, Håkansson A, Troein M. The meaning of quality work from the general practitioner's perspective: an interview study. *BMC Fam Pract* 2006;7:60.
- Melander E, Björgell A, Björgell P, Ovhed I, Mölstad S. Medical audit changes physicians' prescribing of antibiotics for respiratory tract infections. *Scand J Prim Health Care* 1999;17:180-184.
- Munck A, Damsgaard J, Hansen DG, Bjerrum L, Søndergaard J. The Nordic method for quality improvement in general practice. *Qual Prim Care* 2003;1:73-78.
- Munck AP, Gahrn-Hansen B, Søgaard P, Søgaard J. Long-lasting improvement in general practitioners' prescribing of antibiotics by means of medical audit. *Scand J Prim Health Care* 1999;17:185-190.
- 14. Søndergaard J, Hansen DG, Aarslev P, Munck AP. A multifaceted intervention according to the Audit Project Odense method improved secondary prevention of ischemic heart disease: a randomised controlled trial. *Fam Pract* 2006;23:1998-202.
- 15. Pendleton D, Schofield T, Tate P, Havelock P. Konsultationen kommunikation mellan läkare och patient. (The consultation: an approach to learning and teaching.) Lund: Studentlitteratur, 1994.
- 16. Wulff HR, Götzsche PC. *Den kliniska beslutsprocessen* (The clinical decisionmaking process. In Swedish). Lund: Studentlitteratur, 2000.

- Armson H, Kinzie S, Hawes D, Roder S, Wakefield J, Elmslie T. Translating learning into practice: Lessons from the practice-based small group learning program. *Canadian Family Physician* 2007;53:1477-1485.
- Eliasson G, Mattsson B. Experiences of small CME group work in general practice in Sweden. *Scand J Prim Health Care* 1999;17:196-200.
- Malterud K. Key questions a strategy for modifying clinical communication. Transforming tacit skills into a clinical method. *Scand J Prim Health Care*. 1994 Jun;12(2):121-7.
- Malterud K, Hollnagel H. Key questions in consultation a clinical communication method. (In Danish). Ugeskr Læger. 2004 May 17;166(21):2034-5.
- 21. Undeland M, Malterud K. Diagnostic interaction: The patient as a source of knowledge? A qualitative case study from general practice. *Scand J Prim Health Care*, 2008; 26:222-227.
- Bøgelund Frederiksen H, Elverdam B: FynPEP Praktiserende læger og deres patienters kvalitetskriterier i hverdagen og deres erfaringer med at deltage i DanPEP. (In Danish). Kvalitetsudviklingsvalget, Fyns Amt, April 2004.
- 23. Mercer SW, Cawston PG, Bikker AP. Quality in general practice consultations; a qualitative study of the views of patients living in an area of high socio-economic deprivation in Scotland. *BMC Fam Pract* 2007;8:22.
- 24. Guthrie B, Wyke S. Personal continuity and access in UK general practice: a qualitative study of general practitioners' and patients' perceptions of when and how they matter. *BMC Fam Pract* 2006;7:11.

- 25. Ridd M, Shaw A, Salisbury C. 'Two sides of the coin' the value of personal continuity to GPs: a qualitative interview study. *Fam Pract* 2006;23:461-468.
- 26. Balint M. The doctor, his patient and the illness. London: Pitman, 1957.
- 27. WONCA Europe. The European definition of general practice/family medicine. 2005. http://www.woncaeurope.org/Definition%20GP-FM.htm
- Johansen M-L, Holtedahl KA, Rudebeck CE. A doctor close at hand: How GPs view their role in cancer care. *Scand J Prim Health Care*. 2010; 28:249-255.

Tables

Table I – Categories and subcategories from the content analysis of the concept of "holistic view"

Table II – A holistic view in primary care – two linked projects

Table III a – Main groups, groups and variables in the APO registration chart concerning holistic view in the general practitioner's medical decision making

Table III b – Definitions: The participants' explanations of the meaning of the different types of knowledge assessed in the pilot audit

Figures

Figure 1 – The Likert Scales

 Table I. Categories and subcategories from the content analysis of the concept of

 "holistic view" [2]

Category	Subcategory
Attitude	Professional attitude
	Political/administrative attitude
Knowledge	Factual knowledge
	Tacit knowledge
Circumstances	Motivating factor
	Organisation
	Sphere of activity
	Tool

Period	Earlier study	
	(Strandberg EL, Ovhed I, Borgquist L, Wilhelmsson S. The	
	perceived meaning of a (w)holistic view among general	
	practitioners and district nurses in Swedish primary care: a	
	qualitative study. BMC Fam Pract 2007;8:8)	
January 2002	Invitation and start of planning the study	
November 2002 – March 2003	7 focus group interviews with 22 GPs and 20 nurses	
August – December 2003	Analysis	
August 2007	Publication	
	Current study	
Spring 2006	Planning of the current study	
September 2006 – February 2007	The initial APO audit brainstorming process	
2 weeks in March 2007	Audit registration	

Table II. A holistic view in primary care – two linked projects

Table III a. The main groups, groups and variables of the audit registration chart

concerning holistic view in the general practitioner's medical decision making.

	Groups	Variables (in tilted boxes)	#
Patient characteristics	Patient	Own patient	1
		Other doctor's patient	2
		No one's/everyone's patient	3
	Problem	Well-defined problem	4
		Difficult to define	5
Assessment	Important type of knowledge	Purely factual knowledge	6
		Familiarity	7
		Knowledge from the patient's network	8
		The capacity for judgement	9
Considered for medical	Factual knowledge	Considered	10
decision		Not considered	11
		Not obtainable	12
	Team knowledge	Considered	13
		Not considered	14
		Lacking	15
	District knowledge	Considered	16
		Not considered	17
		Not obtainable	18
	Oral knowledge	Considered	19
		Not considered	20
		Not obtainable	21
	Written knowledge	Considered	22
		Not considered	23
		Not obtainable	24
	Capacity for judgement	Considered	25
		Not considered	26
		Not obtainable	27
Evaluation	Significance of holistic view	Necessary	28
		Not necessary	29
		Valuable	30
		Negative/added nothing	31

Table III b. Definitions: The participants' explanations of the meaning of the different types of knowledge assessed in the pilot audit

• Purely factual knowledge:

The patient's problem requires factual medical knowledge.

• *Familiarity*:

The overview and interpretation of a situation is not always fully underpinned by medical knowledge. <u>The patient's problem requires</u> familiarity (sometimes expressed as tacit and implicit knowledge), which is acquired through extensive experience.

• Knowledge from the patient's network :

<u>The patient's problem requires</u> knowledge about and from the patient's network, family, and social surroundings (may have reached the doctor different ways, but) is (regarded as) essential for helping the patient.

• The capacity for judgement:

<u>The patient's problem requires</u> that you use your personal judgement, sometimes in contradiction to EBM, accepted practice or your own former experience, since you believe that this is best for your patient.

Familiarity with the patient



Figure 1. The three Likert Scales for capturing a holistic consultation technique