



LUND UNIVERSITY

Errata: G. Kristensson, Scattering of Electromagnetic Waves by Obstacles, SCITECH, 2016

Kristensson, Gerhard

2025

[Link to publication](#)

Citation for published version (APA):

Kristensson, G. (2025, Feb 8). Errata: G. Kristensson, Scattering of Electromagnetic Waves by Obstacles, SCITECH, 2016.

Total number of authors:

1

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

**Errata: G. Kristensson,
Scattering of Electromagnetic Waves by Obstacles,
SCITECH, 2016**

Page	Line	Read	Should read
xix	10	stuck	struck
47	10	$\chi = 0$	$\chi = 0$ $\chi \neq 0$
67	2 ⁻	dispite	despite
74	4–5	to solve the Maxwell equations in the homogeneous isotropic material, see Problem 1.16.	to solve the Maxwell equations in the homogeneous isotropic material asymptotically in the limit as $r \rightarrow \infty$, see Problem 1.16.
98	6–7	Show that the fields in (1.73) solve the Maxwell equations in a homogeneous isotropic material at all points $r > 0$.	Show that the fields in (1.73) solve the Maxwell equations in a homogeneous isotropic material in the limit as $r \rightarrow \infty$.
102	8	chance	change
136	1 ⁻	$C_+ = \{(\mathbf{r}, t) : t \geq 0, \mathbf{r} - \mathbf{r}_0 = ct\}$	$C_+ = \{(\mathbf{r}, t) : t \geq 0, \mathbf{r} - \mathbf{r}_1 = ct\}$
137	Figure 2.8	$B(r_1, ct)$	$B(\mathbf{r}_1, ct)$
137	Figure 2.8	$(r_1, 0)$	$(\mathbf{r}_1, 0)$
137	1	$C_0 +$	C_+
173	2 ⁻	Section 3.7	Section 3.6
183	6, 10		Change the order of temporal and spatial integration
202	(4.22)	$ \mathbf{E}_0 ^2$	$ \mathbf{E}_0 ^4$
208	5 ⁻	perpendicular	parallel
230	16 ⁻	$\hat{\mathbf{p}}_i = \mathbf{E}_0 / \mathbf{E}_0 $	$\hat{\mathbf{p}}_e = \mathbf{E}_0 / \mathbf{E}_0 $
266	Figure 6.1	GTD, FO, GO	GTD, PO, GO

272	1 ⁻	$\mathbf{E}_0$	$\mathbf{E}_0$
275		$\hat{\nu} \cdot \epsilon_1(\mathbf{r}, 0) \cdot \nabla \phi(\mathbf{r}) _- = \epsilon \hat{\nu} \cdot \nabla \phi(\mathbf{r}) _+$	$\hat{\nu} \cdot \epsilon_1(\mathbf{r}, 0) \cdot \nabla \phi(\mathbf{r}) _- = \epsilon \hat{\nu} \cdot \nabla \phi(\mathbf{r}) _+$
289	12	the analysis in not	the analysis is not
344	9 ⁻	the electric field polarized parallel to the side a	the electric field polarized is parallel to the side a
345	11	$\theta \neq 0, \pi$ or $\phi = \pi/2$	$\theta \neq 0, \pi$ or $\phi \neq \pi/2$
345	6 ⁻	$\theta \neq 0, \pi$ or $\phi = \pi/2$	$\theta \neq 0, \pi$ or $\phi \neq \pi/2$
350	15	[4,195]	[4,196]
351	8 ⁻	$\psi \times$	$\nabla \psi \times$
351	7 ⁻	The first term disappears	The first and last terms disappear
351	4 ⁻	(23)	(24)
372	11	$\nabla \times \mathbf{v}_{\tau n}(k\mathbf{r}') = k\mathbf{v}_{\bar{\tau}n}(k\mathbf{r}')$	$\nabla \times \mathbf{v}_{\tau n}(k\mathbf{r}) = k\mathbf{v}_{\bar{\tau}n}(k\mathbf{r})$
377	1	(4.18)	(4.19)
404	1 ⁻	$\tau_l(x)$	$\tau_l(\cos \theta)$
461	Figure 8.28	The curve should be constant 1 for $s/a > 1$	
463	Figure 8.29	The curve should be constant 1 for $s/a > 1$	
471	2	$\chi = \kappa = 0.1$	$\chi/\sqrt{\epsilon\mu} = \kappa/\sqrt{\epsilon\mu} = 0.1$
471	9	$\chi = \kappa = 0.1$	$\chi/\sqrt{\epsilon\mu} = \kappa/\sqrt{\epsilon\mu} = 0.1$
473	7	$t_{121} = \frac{2k^3a^3(i\chi - \kappa)}{(\epsilon_1 + 2\epsilon)(\mu_1 + 2\mu) - \chi^2 - \kappa^2}$	$t_{121} = \frac{2k^3a^3\epsilon\eta(i\chi - \kappa)}{(\epsilon_1 + 2\epsilon)(\mu_1 + 2\mu) - \chi^2 - \kappa^2}$
473	8	$t_{211} = \frac{2k^3a^3(i\chi + \kappa)}{(\epsilon_1 + 2\epsilon)(\mu_1 + 2\mu) - \chi^2 - \kappa^2}$	$t_{211} = \frac{2k^3a^3\epsilon\eta(i\chi + \kappa)}{(\epsilon_1 + 2\epsilon)(\mu_1 + 2\mu) - \chi^2 - \kappa^2}$
473	10	$6i\mathbf{J}\chi$	$6i\mathbf{J}\epsilon\eta\chi$
474	6	$6\Pi\chi$	$6\Pi\epsilon\eta\chi$

474	12, 15	J	$\epsilon\eta\mathbf{J}$
474	1 ⁻	II	$\epsilon\eta\mathbf{II}$
475	3, 7	II	$\epsilon\eta\mathbf{II}$
475	7	$(\epsilon_1 + 2)$ (numerator and denominator)	$(\epsilon_1 + 2\epsilon)$
475	7	$(\mu_1 + 2)$	$(\mu_1 + 2\mu)$
477	5	$t_{121} = \frac{2k^3a^3(\mathrm{i}\chi - \kappa)}{(\epsilon_1 + 2\epsilon)(\mu_1 + 2\mu) - \chi^2 - \kappa^2}$	$t_{121} = \frac{2k^3a^3\epsilon\eta(\mathrm{i}\chi - \kappa)}{(\epsilon_1 + 2\epsilon)(\mu_1 + 2\mu) - \chi^2 - \kappa^2}$
477	6	$t_{211} = \frac{2k^3a^3(\mathrm{i}\chi + \kappa)}{(\epsilon_1 + 2\epsilon)(\mu_1 + 2\mu) - \chi^2 - \kappa^2}$	$t_{211} = \frac{2k^3a^3\epsilon\eta(\mathrm{i}\chi + \kappa)}{(\epsilon_1 + 2\epsilon)(\mu_1 + 2\mu) - \chi^2 - \kappa^2}$
483	1 ⁻	$f_n = -\sum_{n'} T_{nn'} a_{n'}$	$f_n = \sum_{n'} T_{nn'} a_{n'}$
486	5	$\frac{1}{r^2(\theta)} = r_0 + r_1 P_1(\cos \theta) + r_2 P_2(\cos \theta)$	$\frac{1}{r^2(\theta)} = r_0 + r_2 P_2(\cos \theta)$
486	6	$i = 0, 1, 2$	$i = 0, 2$
486	1 ⁻	$\frac{1}{r^2(\hat{\mathbf{r}})} = \sum_{l=0}^2 \sum_{m=0}^l \sum_{\sigma=\mathrm{e},\mathrm{o}} r_{\sigma ml} Y_{\sigma ml}(\hat{\mathbf{r}})$	$\frac{1}{r^2(\hat{\mathbf{r}})} = +r_0 Y_{\mathrm{e}00}(\hat{\mathbf{r}}) + r_1 Y_{\mathrm{e}02}(\hat{\mathbf{r}}) + r_2 Y_{\mathrm{e}22}(\hat{\mathbf{r}})$
487	1 ⁻	where $\hat{\mathbf{p}}_{\mathrm{e}\parallel} = \hat{\mathbf{e}}_{\mathrm{i}\parallel} \cdot \mathbf{E}_0 / \mathbf{E}_0 $ and $\hat{\mathbf{p}}_{\mathrm{e}\perp} = \hat{\mathbf{e}}_{\mathrm{i}\perp} \cdot \mathbf{E}_0 / \mathbf{E}_0 $.	where $\hat{\mathbf{p}}_{\mathrm{e}\parallel} = \hat{\boldsymbol{\theta}}$ and $\hat{\mathbf{p}}_{\mathrm{e}\perp} = \hat{\boldsymbol{\phi}}$.
492	5	homogenous	homogeneous
493	2	$\sum_{n'} \iint_{S_s}$	$\iint_{S_s}$
493	4	$= 0$	$= \mathbf{0}$
494	4	$\frac{\mu}{\mu_1} B_{\mathcal{W}'} + \frac{k}{k_1} C_{\mathcal{W}'}$	$\frac{\mu}{\mu_1} B_{\mathcal{W}'} - \frac{k}{k_1} C_{\mathcal{W}'}$
494	5	$\frac{k}{k_1} B_{\mathcal{W}'} + \frac{\mu}{\mu_1} C_{\mathcal{W}'}$	$-\frac{k}{k_1} B_{\mathcal{W}'} + \frac{\mu}{\mu_1} C_{\mathcal{W}'}$
494	6	$-\frac{\mu}{\mu_1} B_{\mathcal{W}'} - \frac{k}{k_1} C_{\mathcal{W}'}$	$-\frac{\mu}{\mu_1} B_{\mathcal{W}'} + \frac{k}{k_1} C_{\mathcal{W}'}$

494	7	$-\frac{k}{k_1}B_{ll'} - \frac{\mu}{\mu_1}C_{ll'}$	$\frac{k}{k_1}B_{ll'} - \frac{\mu}{\mu_1}C_{ll'}$
496	8 ⁻	Reference [264, Chapter 6]	Reference 264
496	6 ⁻	Reference [141] and references given therein	Reference 141 and references given therein
505	13	not that $\mathcal{R}_{n'n}(k\mathbf{r}_p)$ is real valued	note that $\mathcal{R}_{n'n}(k\mathbf{r}_p)$ is real-valued
507	5 ⁻	[134]	[135]
509	9	$\pi_l^m(t)$	$\Pi_l^m(t)$
517	6	$\frac{1}{r^2(\theta)} = r_0 + r_1P_1(\cos\theta) + r_2P_2(\cos\theta)$	$\frac{1}{r^2(\theta)} = r_0 + r_2P_2(\cos\theta)$
528	7 ⁻	The relation	The relations
581	2 ⁻	$\mathbf{r} = -\mathbf{T}_{22}^{-1} \cdot \mathbf{T}_{21} = (\mathbf{P}_{11} + \mathbf{W} \cdot \mathbf{P}_{21} \dots$	$\mathbf{r} = -\mathbf{T}_{22}^{-1} \cdot \mathbf{T}_{21} = -(\mathbf{P}_{11} + \mathbf{W} \cdot \mathbf{P}_{21} \dots$
608	10	$y_n(z) = -\frac{1}{z^{n+1}} \sum_{k=0}^{\infty} \frac{(2n-2k-1)!!}{k!} \left(\frac{z^2}{2}\right)^k$	$y_n(z) = -\frac{1}{z^{n+1}} \sum_{k=0}^n \frac{(2n-2k-1)!!}{k!} \left(\frac{z^2}{2}\right)^k$ $+ \frac{(-1)^{n+1}}{z^{n+1}} \sum_{k=n+1}^{\infty} \frac{1}{k!(2k-2n-1)!!} \left(-\frac{z^2}{2}\right)^k$
619	2 ⁻	$G_n(z) = -\frac{n}{z} - \frac{1}{n/z - G_{n-1}(z)}$	$G_n(z) = -\frac{n}{z} + \frac{1}{n/z - G_{n-1}(z)}$
628	10	$Y_{\sigma ml} = \frac{C_{lm}}{2C'_{lm}} \begin{Bmatrix} Y_{lm} + Y_{l-m} \\ -iY_{lm} + iY_{l-m} \end{Bmatrix}$	$Y_{\sigma ml} = \frac{C_{lm}}{2C'_{lm}} \begin{Bmatrix} Y_{lm} + Y_{lm}^* \\ -iY_{lm} + iY_{lm}^* \end{Bmatrix}$
634	13	[38, page 170]	[40, page 216]
641	8	$f(n)$	$f(-n)$
641	10	$f(n)$	$f(-n)$
643	3	inverse	
643	4	$f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \hat{f}(\xi) e^{-ix\xi} d\xi = 0, \quad x < 0$	$\hat{f}(\xi) = \int_{-\infty}^{\infty} f(x) e^{-ix\xi} dx = 0, \quad \xi < 0$

675	Five times	$C_{ml,m'l'}(d, \eta)$ or $C_{ml,-m'l'}(d, \eta)$	$C_{ml,m'l'}(kd, \eta)$ or $C_{ml,-m'l'}(kd, \eta)$
675-6	Five times	$D_{ml,m'l'}(d, \eta)$ or $D_{ml,-m'l'}(d, \eta)$	$D_{ml,m'l'}(kd, \eta)$ or $D_{ml,-m'l'}(kd, \eta)$
677	4^-	$= \begin{pmatrix} j_1 & j_2 & j_3 - 1 \\ 0 & 0 & 0 \end{pmatrix}$	$= - \begin{pmatrix} j_1 & j_2 & j_3 - 1 \\ 0 & 0 & 0 \end{pmatrix}$
720	3^-	$\mathbf{K}(\hat{\mathbf{r}}) = \hat{\phi} \frac{ika^2\eta_0}{2a} J_1(ka \sin \theta)$	$\mathbf{K}(\hat{\mathbf{r}}) = \hat{\phi} \frac{Ika\eta_0}{2} J_1(ka \sin \theta)$