



LUND UNIVERSITY

Use of steroids to reduce high fever in head injury-is it acceptable?

Grände, Per-Olof

Published in:
Intensive Care Medicine

DOI:
[10.1007/s00134-006-0396-y](https://doi.org/10.1007/s00134-006-0396-y)

2006

[Link to publication](#)

Citation for published version (APA):

Grände, P.-O. (2006). Use of steroids to reduce high fever in head injury-is it acceptable? *Intensive Care Medicine*, 32(12), 2079-2079. <https://doi.org/10.1007/s00134-006-0396-y>

Total number of authors:

1

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

This is an author produced version of a paper published in Intensive Care Medicine. This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Citation for the published paper:

Grande, Per-Olof.

"Use of steroids to reduce high fever in head injury-is it acceptable?"

Intensive Care Medicine, 2006, Issue: Oct 13.

<http://dx.doi.org/10.1007/s00134-006-0395-z>

Access to the published version may require journal subscription.

Published with permission from: Springer

Steroids to reduce high fever in head injury – is it acceptable?

Per-Olof Grände

Department of Anaesthesiology and Intensive Care
Lund University and Lund University Hospital, Sweden

Correspondence:

Per-Olof Grände, Dep of Anaesthesiology and Intensive Care, Lund University Hospital, SE-221 85, Lund, Sweden

Tel: +46 46 2227753

Fax. +46 46 2224546

E-mail: per-olof.grande@med.lu.se

Steroids to reduce high fever in head injury – is it acceptable?

Dear Sir,

I will thank Dr Llompарт- Pou and co-authors for the valuable and adequate comments on the use of steroids in head injured patients. In principle I share your view. Steroids should not be used as a general therapy to head injured patients due to its potential side effects, as supported by the large clinical MRC CRASH study referred by you. As also commented by you, steroid treatment will induce deviation from normal physiology, and therefore is against the very essence of the Lund therapy. This does not mean, however, that the use of steroids is prohibited under all circumstances. I will explain why steroid treatment still was accepted under specific conditions in the Lund therapy (1).

Mainly due to less lung complications, the number of patients with high fever has been reduced with the Lund therapy (1), but patients with extravascular blood or large posttraumatic cerebral inflammation or direct hypothalamic damage, may still suffer from high fever. As the effect of paracetamol is small, active cooling and steroids are the only known ways to effectively reduce a high temperature. This gives us 3 choices 1) accept the high fever, 2) use active cooling or 3) treat fever with a bolus dose of steroids. All 3 choices comprise side effects, and we are on the horns of a dilemma, as we do not know the dignity and the importance of these effects for outcome of the 3 choices.

The current literature strongly supports the view that high fever is deleterious and lowering of a high fever will improve outcome for the head injured patient (2). If accepting this view, we have to choose between active cooling and steroids. By creating a difference between the set

thermostat temperature and body temperature, active cooling reduces temperature in an unphysiological way, activating endocrine, autonomic and motor systems - also the deeply sedated patients will be exposed to such adaptations and may be extremely stressed. This situation must be evaluated against a condition where the temperature has been lowered in a more physiological way, by resetting the thermostat temperature level towards more normal values.

This is the reason for why steroid therapy is used in the Lund concept instead of active cooling. After all, the adverse effects of one bolus dose of a steroid (1) must be small compared with the more long term steroid treatment used in clinical studies (3), and most likely also smaller than those induced by long term active cooling. The temperature-reducing effect of one bolus dose is marked and often remaining. The patient becomes more stable hemodynamically and a raised ICP is often reduced. The induced hyperglycemia can be treated by insulin and a transient adrenal suppression may be compensated by a small dose of hydrocortisone. The number of patients suffering from high fever (>38.5 C) when treated strictly according to the Lund Concept is relatively few (1), and number of candidates to be treated with the recommended single bolus dose of Solu-Medrol therefore will be limited.

Sincerely

Per-Olof Grande

Department of Anaesthesiology and Intensive Care,

University and University Hospital of Lund, Sweden

per-olof.grande@med.lu.se

References

1. Grände PO (2006) The “Lund Concept” for treatment of severe head trauma – physiological principles and clinical application. Intensive Care Med. In press
2. Thompson HJ, Tkacs NC, Saatman KE, Raghupathi R, McIntosh TK. (2003) Hyperthermia following traumatic brain injury: a critical evaluation. Neurobiol of disease 12:163-73
3. Alderson P, Roberts I (2005) Corticosteroids for acute traumatic brain injury. Cochrane Database of systemic Reviews, Issue 1. Art. No.: CD000196. DOI: 10.1002/14651858.CD000196.pub2.