



# LUND UNIVERSITY

## Reduced oxidative pentose phosphate pathway flux in recombinant xylose-utilizing *Saccharomyces cerevisiae* strains improves the ethanol yield from xylose.

Jeppsson, Marie; Johansson, Björn; Hahn-Hägerdal, Bärbel; Gorwa-Grauslund, Marie-Francoise

*Published in:*  
Applied and Environmental Microbiology

*DOI:*  
[10.1128/AEM.68.4.1604-1609.2002](https://doi.org/10.1128/AEM.68.4.1604-1609.2002)

2002

[Link to publication](#)

### *Citation for published version (APA):*

Jeppsson, M., Johansson, B., Hahn-Hägerdal, B., & Gorwa-Grauslund, M.-F. (2002). Reduced oxidative pentose phosphate pathway flux in recombinant xylose-utilizing *Saccharomyces cerevisiae* strains improves the ethanol yield from xylose. *Applied and Environmental Microbiology*, 68(4), 1604-1609.  
<https://doi.org/10.1128/AEM.68.4.1604-1609.2002>

*Total number of authors:*  
4

### General rights

Unless other specific re-use rights are stated the following general rights apply:  
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00

## Reduced Oxidative Pentose Phosphate Pathway Flux in Recombinant Xylose-Utilizing *Saccharomyces cerevisiae* Strains Improves the Ethanol Yield from Xylose

Marie Jeppsson, Björn Johansson, Bärbel Hahn-Hägerdal, and Marie F. Gorwa-Grauslund\*

Department of Applied Microbiology, Lund University, 221 00 Lund, Sweden

Received 18 September 2001/Accepted 7 January 2002

**In recombinant, xylose-fermenting *Saccharomyces cerevisiae*, about 30% of the consumed xylose is converted to xylitol. Xylitol production results from a cofactor imbalance, since xylose reductase uses both NADPH and NADH, while xylitol dehydrogenase uses only NAD<sup>+</sup>. In this study we increased the ethanol yield and decreased the xylitol yield by lowering the flux through the NADPH-producing pentose phosphate pathway. The pentose phosphate pathway was blocked either by disruption of the *GND1* gene, one of the isogenes of 6-phosphogluconate dehydrogenase, or by disruption of the *ZWF1* gene, which encodes glucose 6-phosphate dehydrogenase. Decreasing the phosphoglucose isomerase activity by 90% also lowered the pentose phosphate pathway flux. These modifications all resulted in lower xylitol yield and higher ethanol yield than in the control strains. TMB3255, carrying a disruption of *ZWF1*, gave the highest ethanol yield (0.41 g g<sup>-1</sup>) and the lowest xylitol yield (0.05 g g<sup>-1</sup>) reported for a xylose-fermenting recombinant *S. cerevisiae* strain, but also an 84% lower xylose consumption rate. The low xylose fermentation rate is probably due to limited NADPH-mediated xylose reduction. Metabolic flux modeling of TMB3255 confirmed that the NADPH-producing pentose phosphate pathway was blocked and that xylose reduction was mediated only by NADH, leading to a lower rate of xylose consumption. These results indicate that xylitol production is strongly connected to the flux through the oxidative part of the pentose phosphate pathway.**

Fuel ethanol produced from fermentation of lignocellulosic hydrolysates is an attractive replacement for liquid fossil fuels because its production is renewable and it does not generate net carbon dioxide. Hydrolysis of lignocellulose generates mostly hexose but also some pentose sugars. The pentose sugars cannot be metabolized by *Saccharomyces cerevisiae*, the preferred ethanol-producing microorganism. In hydrolysate made from hardwood, xylose must be fermented to ethanol for the process to be economically feasible (39). The yeast *Pichia stipitis* metabolizes xylose through expression of the *XYL1* gene, encoding xylose reductase (XR), and the *XYL2* gene, encoding xylitol dehydrogenase (XDH). XR catalyzes the reduction of xylose to xylitol by using NADH or NADPH (30), whereas XDH oxidizes xylitol to xylulose exclusively with NAD<sup>+</sup> (31). Unfortunately, *P. stipitis* is sensitive to ethanol (10) and requires low and carefully controlled oxygenation (34), which prevents its use for industrial ethanol production.

Recombinant *S. cerevisiae* strains expressing the *XYL1* and *XYL2* genes from *P. stipitis* have been constructed and can ferment xylose (26); however, most of the consumed xylose is secreted as xylitol (20, 26, 36, 42). Xylitol production can be lowered by overexpression of the *XKS1* gene, which encodes the native xylulokinase (XK) (23), but still about one third of the consumed xylose is converted to xylitol under anaerobic conditions (13). Xylitol formation may result from the cofactor imbalance between the NADPH-consuming XR and NADH-producing XDH reactions (26).

Eliasson et al. (12) reported improved ethanol yield from xylulose in *S. cerevisiae* strains that have low phosphoglucose isomerase activity (PGI) or a deletion in the *GND1* gene, which encodes 6-phosphogluconate dehydrogenase (6-PGDH). Glucose 6-phosphate, a branch point metabolite between glycolysis and the pentose phosphate pathway (PPP), is reversibly converted to fructose-6-phosphate using PGI, while 6-PGDH converts 6-phosphogluconate to ribulose-5-phosphate in the PPP (35). This difference suggests that the increased ethanol yield from xylulose observed for these strains could be related to altered flux through the PPP.

We tested this hypothesis by lowering the PPP flux by genetic engineering. We evaluated the effects of these changes on xylose utilization and product formation in *S. cerevisiae* strains expressing XR and XDH from *P. stipitis* and overproducing native XK. The PPP flux was modified by (i) lowering PGI activity, (ii) deleting the *GND1* gene, and (iii) deleting the *ZWF1* gene, which encodes glucose-6-phosphate dehydrogenase (G6PDH). We used a flux model (40) to compare internal fluxes in the  $\Delta zwf1$  strain and a control strain.

### MATERIALS AND METHODS

**Strains.** We used seven different *S. cerevisiae* strains in this study (Table 1). *Escherichia coli* DH5 $\alpha$  (Life Technologies, Rockville, Md.) was used for subcloning. All strains were stored in 20% glycerol at –80°C. Yeast cells from freshly streaked YPD (4) plates were used for inoculation.

**Nucleic acid manipulation.** Plasmid DNA was prepared with the Qia miniprep kit (Qiagen, Valencia, Calif.) or Bio-Rad plasmid miniprep kit (Hercules, Calif.). Restriction and modification enzymes were obtained from Roche (Roche Diagnostics AB, Bromma, Sweden) and from Life Technologies (Rockville, Md.), respectively. DNA extractions from agarose gel were made by using a Qiagen gel extraction kit.

**Transformation.** Competent cells of *E. coli* DH5 $\alpha$  were prepared and transformed by the method of Inoue et al. (22), and yeast transformations were made

\* Corresponding author. Mailing address: Department of Applied Microbiology, Lund University, P.O. Box 124, 221 00 Lund, Sweden. Phone: 46 46 222 0619. Fax: 46 46 222 4203. E-mail: Marie-Francoise.Gorwa@tmb.lth.se.

TABLE 1. *S. cerevisiae* strains used in this study

Strain	Relevant genotype	Relevant phenotype	Reference
ENY.WA-1A	<i>MAT<math>\alpha</math> ura3-52 leu2-3,112 his3-<math>\Delta</math>1 trp1-289 MAL2-8c MAL3 SUC3</i>	1,000–2,500 mU of PGI mg of protein <sup>-1</sup>	24
RBV6-1 <sup>a</sup>	ENY.WA-1A <i>pgi1-1<math>\Delta</math>::URA3 PGI1::LEU2</i>	190–270 mU of PGI mg of protein <sup>-1</sup>	7
TMB3250 <sup>a</sup>	ENY.WA-1A <i>his3::YIpXR/XDH/XK</i>	Expresses XR, XDH, and XK	This work
TMB3251 <sup>a</sup>	RBV6-1 <i>his3::YIpXR/XDH/XK</i>	Expresses XR, XDH, and XK; 10-fold-decreased PGI activity	This work
TMB3001 <sup>b</sup>	CEN.PK 113-7A ( <i>MAT<math>\alpha</math> his3-<math>\Delta</math>1 MAL2-8c SUC2</i> ) <i>his3::YIpXR/XDH/XK</i>	Expresses XR, XDH, and XK	13
TMB3255 <sup>b</sup>	TMB3001 <i>zwf1::KanMX</i>	Expresses XR, XDH, and XK; no G6PDH activity	This work
TMB3008 <sup>c</sup>	CEN.HJ5-1B ( <i>MAT<math>\alpha</math> leu2-3,112 his3-<math>\Delta</math>1 ura3-52 trp1-289 gnd1::HIS3 MAL2-8c SUC2</i> ) <i>his3::YIploxZEO</i>	Expresses XR, XDH, and XK; no G6PDH activity	This work

<sup>a</sup> Parental strain ENY.WA-1A.<sup>b</sup> Parental strain CEN.PK 113-7A.<sup>c</sup> Parental strain CEN.PK2-1C.

by a modified lithium acetate method (15). *E. coli* transformants were selected on Luria-Bertani (LB) medium (4) plates with kanamycin (30  $\mu$ g ml<sup>-1</sup>) (ICN Biochemical Inc., Aurora, Ohio) and/or ampicillin (100  $\mu$ g ml<sup>-1</sup>) (IBI Shelton Scientific Inc., Shelton, Conn.). *S. cerevisiae* transformants were selected on YPD plates with zeocin (100  $\mu$ g ml<sup>-1</sup>) (Invitrogen, Groningen, The Netherlands) or geneticin (200  $\mu$ g ml<sup>-1</sup>) (Life Technologies, Rockville, Md.) or on yeast nitrogen base without amino acids (Difco, Detroit, Mich.) supplemented for auxotrophic requirements.

**Construction of TMB3250 and TMB3251.** The YIpXR/XDH/XK vector (13) was digested with *Pst*I within the *HIS3* gene and used to transform strains ENY.WA-1A and RBV6-1, generating *S. cerevisiae* TMB3250 and TMB3251, respectively, after selection for histidine prototrophy.

**Construction of TMB3008.** Plasmid pUG6 (17) was digested with *Sac*I and *Xba*I, and the *KanMX* gene was replaced with the zeocin resistance gene from pTEF1/Zeo (Invitrogen, Groningen, The Netherlands) using the same restriction sites, resulting in pUG6 Zeo. The zeocin resistance gene, flanked by *loxP* sequences, was removed from pUG6 Zeo using *Pvu*II and *Spe*I and blunted with Klenow DNA polymerase. The YIpXR/XDH/XK plasmid (13) was digested with *Pvu*II and ligated to the zeocin resistance gene fragment. The resulting plasmid, YIpLoxZeo, was digested with *Nde*I within the *HIS3* gene and used to transform CEN.HJ5-1B, generating *S. cerevisiae* TMB3008 after selection for zeocin resistance.

**Construction of TMB3255.** The *ZWF1* gene was amplified from *S. cerevisiae* TMB3001 chromosomal DNA by PCR using oligonucleotides 5'-CGGGATCC AAAATGTCACCTGACCGCGGC-3', with a *Bam*HI restriction site added at the 3' end (bold), and 5'-GTTTCGGCTCGGCCGAGGAGG-3'. The *ZWF1* PCR product was inserted in the pUC19 vector (44) after restriction cleavage with *Eco*RI and *Bam*HI. The *KanMX* gene with *loxP* sequences was amplified from pUG6 by PCR using oligonucleotides 5'-TCCCCCGGGAGCTTCGTACGCTG CAG-3', with an added *Sma*I restriction site (bold), and 5'-GGGGTACCAT AGG GAGACCGGCAGATCC-3', with an added *Kpn*I restriction site (bold). The *KanMX* PCR product was inserted into the *ZWF1* gene using restriction sites *Msc*I and *Kpn*I within the *ZWF1* gene and *Sma*I and *Kpn*I flanking the *KanMX* gene. The plasmid was digested with *Bam*HI and *Eco*RI, and the linear product was used for transformation of TMB3001 (13), generating TMB3255.

**Small-flask cultivations.** All cultivations were performed using defined mineral medium (38). The medium (50 ml), supplemented with 40 g of glucose liter<sup>-1</sup> in a 250-ml baffled shake flask, was inoculated with approximately 10<sup>7</sup> cells liter<sup>-1</sup> and incubated overnight at 30°C on an orbital incubator at 130 rpm (INR-200; Gallenkamp, Leicester, United Kingdom). TMB3250 and TMB3251 were grown on 40 g of fructose liter<sup>-1</sup> instead of glucose because the low PGI activity causes a growth defect on glucose (7). These precultures were used to inoculate a second culture of 200 ml in a 1,000-ml baffled shake flask, which was incubated under the same conditions. The cells were harvested in exponential phase by centrifugation at 4,400  $\times$  g (10 min, 4°C) and washed twice with 0.9% NaCl.

Stirred 25-ml vials containing 20 ml of mineral medium with 50 g of xylose liter<sup>-1</sup> as the sole carbon source were inoculated with ca. 5 g of cells liter<sup>-1</sup>, corresponding to ca. 3  $\times$  10<sup>9</sup> cells liter<sup>-1</sup>. The medium was supplemented with 100 mM citrate buffer (pH 5.5) and the required amino or nucleic acids (50 mg of L-tryptophan liter<sup>-1</sup> and/or 250 mg of L-leucine liter<sup>-1</sup> and/or 50 mg of uracil liter<sup>-1</sup>). Ergosterol and Tween 80 were added to final concentrations of 10 and 400 mg liter<sup>-1</sup>, respectively (2, 3). Fermentation was conducted at 30°C in duplicate. Samples were withdrawn with a hypodermic needle and syringe.

**Continuous cultivations.** Cultivations were performed using defined mineral medium (38). Yeast cells were grown under oxygen-limited conditions in 200 ml of medium containing 20 g of xylose liter<sup>-1</sup>, 20 g of glucose liter<sup>-1</sup>, 10 mg of ergosterol liter<sup>-1</sup>, and 400 mg of Tween 80 liter<sup>-1</sup> in a 250-ml baffled shake flask. The culture was incubated overnight at 30°C. Cells were centrifuged at 4,400  $\times$  g for 5 min at 4°C and used to inoculate 1.5 liters of the same medium to ca. 5  $\times$  10<sup>7</sup> cells liter<sup>-1</sup> in a Bioflo III fermentor (New Brunswick Scientific, Edison, N.J.). Antifoam was added at 0.05% (vol/vol) (Dow Corning Antifoam RD Emulsion; BDH Laboratory Supplies, Poole, United Kingdom). Continuous cultivation was set up at dilution rates of 0.06 and 0.12 h<sup>-1</sup> at 30°C and pH 5.5, controlled by addition of 3 M NaOH, and a stirring speed of 200 rpm. The fermentor was sparged with 0.2 liter of nitrogen min<sup>-1</sup> (containing less than 5 ppm O<sub>2</sub>) as measured with a gas mass flowmeter (Bronkhorst, Ruurlo, The Netherlands).

**Analyses of substrates and products.** Glucose, xylose, xylitol, succinate, glycerol, acetate, and ethanol concentrations were determined by column liquid chromatography (CLC) using a Gilson CLC system (Gilson, Villiers-le-Bel, France). An Aminex HPX-87H column (Bio-Rad, Richmond, Calif.) and an RID-10A refractive index detector (Shimadzu, Kyoto, Japan) were used. The column temperature was 45°C, and 5 mM H<sub>2</sub>SO<sub>4</sub> was used as the mobile phase at a flow rate of 0.6 ml min<sup>-1</sup>.

During continuous cultivation, cells from the outlet were used to measure RNA (5), proteins, and polysaccharides (19). Samples for CLC and cell dry weight determination were withdrawn from the fermentor. Growth was followed by measuring the optical density at 620 nm (OD<sub>620</sub>). Cell dry weight was determined by filtering 1 volume of sample through a 0.45- $\mu$ m filter and washing with 3 volumes of water. The filter was dried in a microwave oven at 350 W for 8 min, cooled in a desiccator, and weighed. The composition of the outgoing gas was monitored with a carbon dioxide and oxygen monitor (type 1308; Brüel & Kjør, Copenhagen, Denmark).

**Enzymatic measurements.** Crude extracts for enzyme measurements were made using the Y-PER yeast protein extraction reagent (Pierce, Rockford, Ill.). Protein concentration was determined by the Coomassie protein assay reagent (Pierce), using bovine serum albumin as a standard. The phosphoglucose isomerase (PGI, EC 5.3.1.9) activity was measured as described previously (27). The glucose-6-phosphate dehydrogenase activity (G6PDH, EC 1.1.1.49) and 6-phosphogluconate dehydrogenase activity (6-PGDH, EC 1.1.1.44) were measured according to the method of Bergmeyer (6).

Intracellular NADPH and NADP<sup>+</sup> were measured enzymatically (6) on a TD-700 fluorometer (Turner Designs, Sunnyvale, Calif.) after extraction with boiling ethanol (16) from samples withdrawn from anaerobic growth on glucose.

## RESULTS

**Effect of decreased PGI activity on xylose fermentation.** The xylose pathway, consisting of XR and XDH enzymes from *P. stipitis* and overproduced XK from *S. cerevisiae*, was introduced via the YIpXR/XDH/XK vector (13) into ENY.WA-1A (control) and RBV6-1 strains (10-fold decrease in PGI activity [7]), generating *S. cerevisiae* TMB3250 and TMB3251, respectively.

TABLE 2. Specific xylose consumption and ethanol, xylitol, acetate, and glycerol yields after 70 h of 25-ml batch fermentations with xylose (50 g/liter) as the sole carbon source<sup>a</sup>

Strain	Relevant phenotype or genotype	Xylose consumed (g/g of biomass)	Yield (g/g of xylose consumed)			
			Ethanol	Xylitol	Acetate	Glycerol
TMB3250		7.2 ± 0.4	0.30 ± 0.01	0.30 ± 0.01	0.023 ± 0.001	0.047 ± 0.001
TMB3251	Low PGI	4.6 ± 0.3	0.34 ± 0.01	0.21 ± 0.01	0.034 ± 0.001	0.058 ± 0.002
TMB3001		9.2 ± 0.1	0.31 ± 0.00	0.29 ± 0.01	0.025 ± 0.001	0.052 ± 0.004
TMB3008	<i>Δgnd1</i>	5.6 ± 0.7	0.38 ± 0.01	0.13 ± 0.01	0.051 ± 0.005	0.052 ± 0.011
TMB3255	<i>Δzwf1</i>	1.5 ± 0.1	0.41 ± 0.02	0.05 ± 0.01	0.084 ± 0.005	0.054 ± 0.008

<sup>a</sup> Values are the averages of the results of two independent fermentations and deviations from the averages.

TMB3251 had about 10% of the PGI activity of its control strain, TMB3250 (data not shown).

After 70 h of xylose fermentation, TMB3250 and TMB3251 had consumed 7.2 and 4.6 g of xylose g of biomass<sup>-1</sup>, respectively (Table 2). Ethanol yields were 0.30 g of xylose g<sup>-1</sup> for TMB3250 and 0.34 g of xylose g<sup>-1</sup> for TMB3251. The 11% higher ethanol yield of TMB3251 was accompanied by a lower xylitol yield (0.21 g g<sup>-1</sup>) than that of TMB3250 (0.30 g g<sup>-1</sup>). The acetate and glycerol yields were slightly higher in TMB3251 (0.03 g g<sup>-1</sup> and 0.06 g g<sup>-1</sup>, respectively) than in TMB3250 (0.02 g g<sup>-1</sup> and 0.05 g g<sup>-1</sup>, respectively).

**Xylose fermentation by a *ΔGND1* strain.** *GND1* encodes one of the two NADP<sup>+</sup>-dependent isoenzymes of 6-phosphogluconate dehydrogenase that catalyze the conversion of 6-phosphogluconate to ribulose 5-phosphate in the PPP. The xylose pathway was introduced in the CEN.HJ5-1B strain, which has an inactive *GND1* gene (24), to study the effect of an altered PPP on xylose fermentation. *S. cerevisiae* strain TMB3008 was generated by integration of the YIpLoxZEO vector in a glucose-positive revertant of CEN.HJ5-1B. A glucose-positive revertant was used instead of the original strain to facilitate glucose metabolism (33).

TMB3008 and its control, TMB3001, were used for batch fermentation of xylose. TMB3001 and TMB3008 consumed 9.2 and 5.6 g of xylose g of biomass<sup>-1</sup>, respectively (Table 2). Ethanol yields were 0.31 g g<sup>-1</sup> for TMB3001 and 0.38 g g<sup>-1</sup> for TMB3008 (Table 2). TMB3008 showed 24% higher ethanol yield and a lower xylitol yield (0.13 g g<sup>-1</sup>) than TMB3001 (0.29 g g<sup>-1</sup>). The acetate yield was higher in TMB3008 (0.05 g g<sup>-1</sup>) than in TMB3001 (0.03 g g<sup>-1</sup>). The glycerol yields were similar in the two strains.

**Xylose fermentation by a *ΔZWFI* strain.** The strain with low PGI activity and the *ΔGND1* strain had phenotypes similar to those of their respective wild-type strains with respect to xylose fermentation (Table 2). Both of these genetic alterations decreased the flux through the PPP during xylose fermentation. Lowering the PPP flux could result from abolishing the 6-PGDH activity or from lower gluconeogenic flux as a consequence of low PGI activity. Therefore, the disruption of *ZWF1* also should result in lower xylitol and higher ethanol yields. *Δzwf1* mutations have no negative side effects with respect to glucose utilization (24), whereas both the *GND1* deletion and the reduction in PGI activity resulted in defective glucose metabolism (7, 33). The *ZWF1* gene was deleted in TMB3001, resulting in TMB3255, and the strains were compared for xylose fermentation.

TMB3255 had the highest ethanol yield (0.41 g g<sup>-1</sup>), the lowest xylitol yield (0.05 g g<sup>-1</sup>), and the highest acetate yield

(0.08 g g<sup>-1</sup>) of the three strains (Table 2). However, the specific xylose consumption after 70 h (1.5 g per g of biomass) was also the lowest of the three strains (Table 2). This qualitatively similar product pattern suggests that the phenotypes of *Δgnd1*, low-PGI, and *Δzwf1* strains are all related to lower PPP flux.

The *Δzwf1* genotype has been reported to cause a requirement for organic sulfur (37). We did not observe this requirement, and xylose uptake was not affected by the addition of methionine.

**Enzyme activities.** The *ΔZWFI* strain (TMB3255) showed a more pronounced effect on xylose fermentation than that observed in the *ΔGND1* strain (TMB3008) (Table 2). These results suggest that the oxidative part of the PPP might not be totally blocked in TMB3008, since we inactivated only one of the two 6-PGDH isoenzymes in *S. cerevisiae*.

We measured the enzyme activities for G6PDH and 6-PGDH of TMB3001, TMB3008, and TMB3255 in cultures grown on glucose and xylose (Table 3). The enzyme activities of G6PDH and 6-PGDH in TMB3001 (control strain) were comparable with previously published values for *S. cerevisiae* (27). The 6-PGDH enzyme activity increased in TMB3001 when cultivated on xylose instead of glucose. In TMB3008 (*Δgnd1*), 6-PGDH and G6PDH activities were undetectable. This might be an effect of the reversion to a glucose-positive phenotype of this strain, since *ΔGND1* mutants often lose G6PDH activity when exposed to high concentrations of glucose (33). As expected, G6PDH activity was not detectable in TMB3255 (*Δzwf1*). A fourfold decrease in 6-PGDH activity was measured in TMB3255 compared with the control strain (TMB3001). The hypothesis of a partly active PPP in TMB3008 was therefore not confirmed by higher activities of G6PDH or 6-PGDH in TMB3008 relative to those in TMB3255.

TABLE 3. Specific G6PDH and 6-PGDH activities measured after growth on glucose or incubation on xylose

Strain	Relevant genotype	Sp act <sup>a</sup> (U/mg of protein)			
		G6PDH		6-PGDH	
		Glucose	Xylose	Glucose	Xylose
TMB3001		0.90	0.95	0.54	0.83
TMB3008	<i>Δgnd1</i>	<0.01	<0.01	<0.01	<0.01
TMB3255	<i>Δzwf1</i>	<0.01	<0.01	0.21	0.21

<sup>a</sup> Displayed values are the averages of the results of duplicate experiments with less than a 10% difference.






FIG. 1. Internal metabolic fluxes of TMB3001 (*ZWF1*) and TMB3255 ( $\Delta zwf1$ ) (bold) in chemostat cultures at a dilution rate of  $0.06 \text{ h}^{-1}$  (upper value of each pair) and  $0.12 \text{ h}^{-1}$  (lower value) with a feed containing  $20 \text{ g}$  of xylose  $\text{liter}^{-1}$  and  $20 \text{ g}$  of glucose  $\text{liter}^{-1}$ . TMB3001 has a wild-type level of G6PDH. TMB3255 is deficient in G6PDH. All fluxes are normalized to a total specific sugar consumption of  $100 \text{ mmol per g}$  (dry weight) of cells per h. Substances shown inside boxes are substrates or products excreted into the medium. P, phosphate; DHAP, dihydroxyacetone phosphate; G-3-P, glyceraldehyde 3-phosphate.

**Continuous cultivations and flux analyses.** A flux analysis was performed to compare intracellular fluxes in TMB3001 and TMB3255 using a stoichiometric model (40). Chemostat cultivations on a mixture of  $20 \text{ g}$  of xylose  $\text{liter}^{-1}$  and  $20 \text{ g}$  of glucose  $\text{liter}^{-1}$  were performed at dilution rates of  $0.06$  and  $0.12 \text{ h}^{-1}$ . The flux values for TMB3001 and TMB3255 (Fig. 1) were normalized to  $100 \text{ mmol}$  of glucose and xylose consumed per g of biomass per h.

The xylose consumption rate and the xylitol production rate were lower in TMB3255 than in TMB3001 (Table 4), as expected from the batch fermentation. However, the difference in xylose consumption rate between the two strains was not as pronounced in chemostat cultivation with glucose and xylose as the substrate (Table 4) as it was in batch fermentations with xylose as the sole carbon source (Table 2). At the highest dilution rate,  $0.12 \text{ h}^{-1}$ , xylose consumption decreased and glycerol production increased, as found previously (40). The xylose consumption rate was  $37\%$  lower at  $0.12 \text{ h}^{-1}$  than at  $0.06 \text{ h}^{-1}$  for TMB3001. It was only  $12\%$  lower at  $0.12 \text{ h}^{-1}$  than at  $0.06 \text{ h}^{-1}$  for TMB3255.

The flux analyses confirmed that there was no flux through the oxidative part of the PPP in TMB3255, from which the *ZWF1* gene was deleted (Fig. 1). In contrast,  $14$  to  $18\%$  of the

consumed glucose was channeled through the PPP of the control strain TMB3001 (Fig. 1). The flux from ribulose 5-phosphate to xylulose 5-phosphate was reversed in TMB3255 compared to that in TMB3001. The model also predicted that XR used only NADH in TMB3255, whereas XR used  $57$  and  $53\%$  NADPH in TMB3001 at  $0.06 \text{ h}^{-1}$  and  $0.12 \text{ h}^{-1}$ , respectively. Intracellular concentrations of NADPH and  $\text{NADP}^+$  in TMB3001 and TMB3255 were analyzed using enzymatic assays with fluorimetric detection. Average values indicated that the  $\text{NADPH}/\text{NADP}^+$  ratio was three to five times lower in TMB3255 than in TMB3001 (data not shown).

## DISCUSSION

The design of an efficient recombinant *S. cerevisiae* strain for xylose fermentation has been a major challenge for many years. Xylitol is a major fermentation by-product following xylose catabolism by *S. cerevisiae* strains expressing *XYL1* and *XYL2* (26, 36, 42). One hypothesis is that xylitol formation results from the apparent cofactor imbalance between the XR and XDH enzymes (8). The XR enzyme has been engineered by site-directed mutagenesis for lower affinity for NADPH; however, lower enzymatic activity and substantially higher  $K_m$  for xylose were observed (25, 45). The expression in *S. cerevisiae* of several bacterial xylose isomerases, which convert xylose to xylulose without cofactors, has also failed (1, 18, 21, 28, 32), with the exception of the xylose isomerase from *Thermus thermophilus*, for which a very low activity was obtained (43).

Both the XK level and the XR/XDH ratio have an effect on xylitol formation in *S. cerevisiae*. Overexpression of *XKS1* increases the ethanol yield (20, 23). However, strains overexpressing *XKS1* still excrete about one-third of the consumed xylose as xylitol (40) under anaerobic conditions. Strains with a low XR/XDH ratio form less xylitol than strains with a high ratio (42). In the xylose-fermenting *P. stipitis*, the XR/XDH ratio is much higher than in *S. cerevisiae* (11, 34). Despite this difference, *P. stipitis* produces less xylitol (26). A mathematical core model of the ratio of the XR, XDH, and XK enzymes indicated that cofactor concentrations may influence the formation of xylitol more than the activity ratios of the enzymes do (14). These results suggest that the levels and ratios of

TABLE 4. Measured specific uptake rates (negative values) and production rates (positive values) of substrates and products at dilution rates of  $0.06$  and  $0.12 \text{ h}^{-1}$  for TMB3001 and TMB3255 ( $\Delta zwf1$ )<sup>a</sup>

Substrate or product	Uptake or production (mmol/g of biomass/h)			
	TMB3001		TMB3255	
	$D = 0.06 \text{ h}^{-1}$	$D = 0.12 \text{ h}^{-1}$	$D = 0.06 \text{ h}^{-1}$	$D = 0.12 \text{ h}^{-1}$
Xylose	-1.5	-1.9	-0.80	-1.3
Glucose	-3.4	-6.9	-4.0	-7.3
$\text{CO}_2$	6.6	11.7	7.6	11.9
Ethanol	5.4	10.3	7.1	12.0
Xylitol	0.64	0.71	0.10	0.17
Glycerol	0.81	1.6	0.75	1.8
Acetate	0.04	0.07	0.36	0.78
Succinate	0.02	0.04	0.00	0.00

<sup>a</sup> Displayed values are the averages of the results of duplicate experiments with less than a  $5\%$  difference.

NADPH and/or NADH are regulated differently in *P. stipitis* than in *S. cerevisiae*.

We constructed a recombinant xylose-fermenting *S. cerevisiae* strain with a low XR/XDH ratio, *XKS1* overexpression, and an inactivated oxidative PPP. To the best of our knowledge, TMB3255 has the highest ethanol yield ( $0.41 \text{ g g}^{-1}$ ) and the lowest xylitol yield ( $0.05 \text{ g g}^{-1}$ ) reported for any xylose-fermenting recombinant *S. cerevisiae*. PPP activity was lowered or inactivated by deleting either *GND1* or *ZWF1* or by lowering PGI activity. A low PGI activity decreases the flux from fructose 6-phosphate to glucose 6-phosphate, which is the substrate of G6PDH, the first enzyme of the PPP. Since PPP is the main source of NADPH (9), we suggest that the low xylitol yield is directly linked to depletion of NADPH in strains with defective PPP. The higher acetate yield is an indication of NADPH depletion, since oxidation of acetaldehyde to acetate requires an  $\text{NADP}^+$ -dependent acetaldehyde dehydrogenase (29, 41). Furthermore, the NADPH/ $\text{NADP}^+$  ratio was lower in TMB3255 than in TMB3001.

The XR enzyme uses both NADPH and NADH (30), whereas the XDH enzyme uses  $\text{NAD}^+$  exclusively. We therefore propose the following model. In strains with a low flux through the PPP, the level of NADPH is low and a greater fraction of xylose is reduced with NADH. Since NADH is consumed in the XR step and produced in the XDH step, the xylose conversion to xylulose is balanced with respect to cofactors, which would explain why only minor amounts of xylitol are formed. This model is also supported by chemostat results with the control strain TMB3001. At higher dilution rates, the faster anabolism leads to reduced NADPH availability and results in increased use of NADH versus NADPH by XR, concomitant with a decreased xylitol yield (40; this study).

Lowering the oxidative PPP activity resulted in a reduced rate of xylose consumption. One reason for this decrease is that the NADPH-dependent reduction of xylose to xylitol is reduced, resulting in a reduced overall xylose fermentation rate. This was observed for TMB3255 cultivated in a chemostat on a mixture of glucose and xylose. The NADPH formed, which occurred mainly via acetate formation, was sufficient only for biomass synthesis. A strictly NADH-dependent xylose reduction was observed, and the xylose consumption rate was reduced accordingly. In xylose fermentation by nongrowing cells, the rate of xylose consumption by TMB3255 was 84% lower than that by TMB3001, which could not be explained only by the lack of the  $\text{NADP}^+$ -dependent reaction. We suggest that the difference is connected to the nongrowing conditions and to the absence of glucose as a cosubstrate to ensure maintenance.

This work has demonstrated the connection between the level of the oxidative PPP and xylitol by-product formation in recombinant xylose-utilizing *S. cerevisiae*. By removing the NADPH-dependent xylose reduction, we obtained the highest ethanol yield from xylose ever reported for *S. cerevisiae*. However, the xylose consumption rate decreased accordingly, suggesting a limitation of XR activity. Strains with enhanced XR activity may overcome this limitation.

#### ACKNOWLEDGMENTS

We thank E. Boles, Düsseldorf University, Düsseldorf, Germany, for *S. cerevisiae* RBY6-1 and H. Juhnke, J. W. Goethe University, Frankfurt am Main, Germany, for *S. cerevisiae* CEN.HJ5-1B.

This work was financially supported by the Swedish National Energy Administration and the Nordic Energy Research Programme. M.F.G.-G. was supported by Marie Curie fellowship QLK3-CT-1999-51355 from the European Community.

M. Jeppsson and B. Johansson contributed equally to this work.

#### REFERENCES

- Amore, R., and C. P. Hollenberg. 1989. Xylose isomerase from *Actinoplanes missouriensis*: primary structure of the gene and the protein. *Nucleic Acids Res.* **17**:7515.
- Andreasen, A., and T. Stier. 1953. Anaerobic nutrition of *Saccharomyces cerevisiae*. I. Ergosterol requirement for growth in a defined medium. *J. Cell Comp. Physiol.* **41**:23–36.
- Andreasen, A., and T. Stier. 1954. Anaerobic nutrition of *Saccharomyces cerevisiae*. II. Unsaturated fatty acid requirement for growth in a defined medium. *J. Cell Comp. Physiol.* **44**:271–281.
- Ausubel, F. M., R. Brent, R. E. Kingston, D. D. Moore, J. G. Seidman, J. A. Smith, and K. Struhl (ed.). 1995. *Current protocols in molecular biology*. Wiley, New York, N.Y.
- Benthin, S., J. Nielsen, and J. Villadsen. 1991. A simple and reliable method for the determination of cellular RNA content. *Biotechnol. Tech.* **5**:39–42.
- Bergmeyer, H. 1974. *Methods in enzymatic analysis*, 2nd ed. Academic Press, New York, N.Y.
- Boles, E., and F. K. Zimmermann. 1994. Open reading frames in the antisense strands of genes coding for glycolytic enzymes in *Saccharomyces cerevisiae*. *Mol. Gen. Genet.* **243**:363–368.
- Bruinenberg, P. M., P. H. M. de Bot, J. P. van Dijken, and W. A. Scheffers. 1983. The role of redox balances in the anaerobic fermentation of xylose by yeasts. *Appl. Microbiol. Biotechnol.* **18**:287–292.
- Bruinenberg, P. M., J. P. van Dijken, and W. A. Scheffers. 1983. A theoretical analysis of NADPH production and consumption in yeasts. *J. Gen. Microbiol.* **129**:953–964.
- du Preez, J. C., M. Bosch, and B. A. Prior. 1987. Temperature profiles of growth and ethanol tolerance of the xylose fermenting yeasts *Candida shehatae* and *Pichia stipitis*. *Appl. Microbiol. Biotechnol.* **25**:521–525.
- du Preez, J. C., B. van Driessell, and B. A. Prior. 1989. Effect of aerobiosis on fermentation and key enzyme levels during growth of *Pichia stipitis*, *Candida shehatae* and *Candida tenuis* on D-xylose. *Arch. Microbiol.* **152**:143–147.
- Eliasson, A., E. Boles, B. Johansson, M. Österberg, J. M. Thevelein, I. Spencer-Martins, H. Juhnke, and B. Hahn-Hägerdal. 2000. Xylulose fermentation by mutant and wild-type strains of *Zygosaccharomyces* and *Saccharomyces cerevisiae*. *Appl. Microbiol. Biotechnol.* **53**:376–382.
- Eliasson, A., C. Christensson, C. F. Wahlbom, and B. Hahn-Hägerdal. 2000. Anaerobic xylose fermentation by recombinant *Saccharomyces cerevisiae* carrying *XYL1*, *XYL2*, and *XKS1* in mineral medium chemostat cultures. *Appl. Environ. Microbiol.* **66**:3381–3386.
- Eliasson, A., J. S. Hofmeyr, S. Pedler, and B. Hahn-Hägerdal. 2001. The xylose reductase/xylitol dehydrogenase/xylulokinase ratio affects product formation in recombinant xylose-utilising *Saccharomyces cerevisiae*. *Enzyme Microb. Technol.* **29**:288–297.
- Gietz, R. D., and R. H. Schiestl. 1995. Transforming yeast with DNA. *Methods Mol. Cell. Biol.* **5**:255–269.
- Gonzalez, B., J. Francois, and M. Renaud. 1997. A rapid and reliable method for metabolite extraction in yeast using boiling buffered ethanol. *Yeast* **13**:1347–1355.
- Guldener, U., S. Heck, T. Fielder, J. Beinhauer, and J. H. Hegemann. 1996. A new efficient gene disruption cassette for repeated use in budding yeast. *Nucleic Acids Res.* **24**:2519–2524.
- Hallborn, J. 1995. Ph.D. thesis. Lund University, Lund, Sweden.
- Herbert, D., P. Phipps, and R. Strange. 1971. Chemical analysis of microbial cells. *Methods Microbiol.* **5B**:209–344.
- Ho, N. W. Y., Z. Chen, and A. P. Brainard. 1998. Genetically engineered *Saccharomyces* yeast capable of effective cofermentation of glucose and xylose. *Appl. Environ. Microbiol.* **64**:1852–1859.
- Ho, N. W. Y., P. Stevis, J. Rosenfeld, J. Huang, and G. T. Tsao. 1983. Expression of the *Escherichia coli* xylose isomerase gene by a yeast promoter. *Biotechnol. Bioeng. Symp.* **13**:245–250.
- Inoue, H., H. Nojima, and H. Okayama. 1990. High efficiency transformation of *Escherichia coli* with plasmids. *Gene* **96**:23–28.
- Johansson, B., C. Christensson, T. Hobley, and B. Hahn-Hägerdal. 2001. Xylulokinase overexpression in two strains of *Saccharomyces cerevisiae* also expressing xylose reductase and xylitol dehydrogenase and its effect on fermentation of xylose and lignocellulosic hydrolysate. *Appl. Environ. Microbiol.* **67**:4249–4255.
- Juhnke, H., B. Krems, P. Kötter, and K. D. Entian. 1996. Mutants that show increased sensitivity to hydrogen peroxide reveal an important role for the pentose phosphate pathway in protection of yeast against oxidative stress. *Mol. Gen. Genet.* **252**:456–464.
- Kostrzynska, M., C. R. Sopher, and H. Lee. 1998. Mutational analysis of the role of the conserved lysine-270 in the *Pichia stipitis* xylose reductase. *FEMS Microbiol. Lett.* **159**:107–112.

26. Kötter, P., and M. Ciriacy. 1993. Xylose fermentation by *Saccharomyces cerevisiae*. Appl. Microbiol. Biotechnol. **38**:776–783.
27. Maitra, P. K., and Z. Lobo. 1971. A kinetic study of glycolytic enzyme synthesis in yeast. J. Biol. Chem. **246**:475–488.
28. Moes, C. J., I. S. Pretorius, and W. H. van Zyl. 1996. Cloning and expression of the *Clostridium thermosulfurogens* D-xylose isomerase gene (*xylA*) in *Saccharomyces cerevisiae*. Biotechnol. Lett. **18**:269–274.
29. Nissen, T. L., U. Schulze, J. Nielsen, and J. Villadsen. 1997. Flux distributions in anaerobic, glucose-limited continuous cultures of *Saccharomyces cerevisiae*. Microbiology **143**:203–218.
30. Rizzi, M., P. Erleman, N. A. Bui-Thanh, and H. Dellweg. 1988. Xylose fermentation by yeast. 4. Purification and kinetic studies of xylose reductase from *Pichia stipitis*. Appl. Microbiol. Biotechnol. **29**:148–154.
31. Rizzi, M., K. Harwart, P. Erlemann, N.-A. Bui-Thanh, and H. Dellweg. 1989. Purification and properties of the NAD<sup>+</sup>-xylitol-dehydrogenase from the yeast *Pichia stipitis*. J. Ferment. Bioeng. **67**:20–24.
32. Sarthy, A. V., B. L. McConaughy, Z. Lobo, J. A. Sundstrom, C. E. Furlong, and B. D. Hall. 1987. Expression of the *Escherichia coli* xylose isomerase gene in *Saccharomyces cerevisiae*. Appl. Environ. Microbiol. **53**:1996–2000.
33. Sinha, A., and P. K. Maitra. 1992. Induction of specific enzymes of the oxidative pentose phosphate pathway by glucono-delta-lactone in *Saccharomyces cerevisiae*. J. Gen. Microbiol. **138**:1865–1873.
34. Skoog, K., and B. Hahn-Hägerdal. 1990. Effect of oxygenation on xylose fermentation by *Pichia stipitis*. Appl. Environ. Microbiol. **56**:3389–3394.
35. Stryer, L. 1988. Biochemistry. W. H. Freeman & Co., New York, N.Y.
36. Tantirungkij, M., N. Nakashima, T. Seki, and T. Yoshida. 1993. Construction of xylose-assimilating *Saccharomyces cerevisiae*. J. Ferment. Bioeng. **75**:83–88.
37. Thomas, D., H. Cherest, and Y. Surdin-Kerjan. 1991. Identification of the structural gene for glucose-6-phosphate dehydrogenase in yeast. Inactivation leads to a nutritional requirement for organic sulfur. EMBO J. **10**:547–553.
38. Verduyn, C., E. Postma, W. A. Scheffers, and J. P. Van Dijken. 1992. Effect of benzoic acid on metabolic fluxes in yeasts: a continuous-culture study on the regulation of respiration and alcoholic fermentation. Yeast **8**:501–517.
39. von Sivers, M., and G. Zacchi. 1995. A techno-economical comparison of three processes for the production of ethanol from pine. Bioresour. Technol. **51**:43–52.
40. Wahlbom, C. F., A. Eliasson, and B. Hahn-Hägerdal. 2001. Intracellular fluxes in a recombinant xylose-utilizing *Saccharomyces cerevisiae* cultivated anaerobically at different dilution rates and feed concentrations. Biotechnol. Bioeng. **72**:289–296.
41. Wahlbom, C. F., and B. Hahn-Hägerdal. Furfural, 5-hydroxymethyl furfural and acetoin act as external electron acceptors during anaerobic fermentation of xylose in recombinant *Saccharomyces cerevisiae*. Biotechnol. Bioeng., in press.
42. Walfridsson, M., M. Anderlund, X. Bao, and B. Hahn-Hägerdal. 1997. Expression of different levels of enzymes from the *Pichia stipitis* *XYL1* and *XYL2* genes in *Saccharomyces cerevisiae* and its effects on product formation during xylose utilisation. Appl. Microbiol. Biotechnol. **48**:218–224.
43. Walfridsson, M., X. Bao, M. Anderlund, G. Lilius, L. Bülow, and B. Hahn-Hägerdal. 1996. Ethanolic fermentation of xylose with *Saccharomyces cerevisiae* harboring the *Thermus thermophilus* *xylA* gene, which expresses an active xylose (glucose) isomerase. Appl. Environ. Microbiol. **62**:4648–4651.
44. Yanisch-Perron, C., J. Vieira, and J. Messing. 1985. Improved M13 phage cloning vectors and host strains: nucleotide sequences of the M13mp18 and pUC19 vectors. Gene **33**:103–119.
45. Zhang, Y., and H. Lee. 1997. Site-directed mutagenesis of the cysteine residues in the *Pichia stipitis* xylose reductase. FEMS Microbiol. Lett. **147**: 227–232.