



LUND UNIVERSITY

The identity of some Hippolais specimens from Eritrea and the United Arab Emirates examined by mtDNA analysis: a record of Sykes's Warbler H-rama in Africa

Pearson, D J; Ash, J S; Bensch, Staffan

Published in:
Ibis

DOI:
[10.1111/j.1474-919x.2004.00305.x](https://doi.org/10.1111/j.1474-919x.2004.00305.x)

2004

[Link to publication](#)

Citation for published version (APA):

Pearson, D. J., Ash, J. S., & Bensch, S. (2004). The identity of some Hippolais specimens from Eritrea and the United Arab Emirates examined by mtDNA analysis: a record of Sykes's Warbler H-rama in Africa. *Ibis*, *146*(4), 683-684. <https://doi.org/10.1111/j.1474-919x.2004.00305.x>

Total number of authors:
3

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Short communication

The identity of some *Hippolais* specimens from Eritrea and the United Arab Emirates examined by mtDNA analysis: a record of Sykes's Warbler *H. rama* in Africa

DAVID J. PEARSON,^{1*} JOHN S. ASH² & STAFFAN BENSCH³

¹Lupin Close, Reydon, Southwold, Suffolk IP18 6NW, UK

²Godshill Wood, Fordingbridge, Hampshire SP6 2LR, UK

³Department of Animal Ecology, Ecology Building, Lund University, S-223 62 Lund, Sweden

A confusing group of *Hippolais* and *Acrocephalus* warblers is found in coastal habitat in northeast Africa and the nearby area of the Red Sea, Gulf of Aden and Persian Gulf. The group includes migrant and resident forms, some of which bear great similarity to other species in the same habitat. They are difficult to identify in the field, and some also pose problems in the museum.

We recently examined specimens of a small, apparently resident *Hippolais* warbler from northern Somalia (Ash & Pearson 2002). While comparing these with other material at the Natural History Museum, Tring, our attention was drawn to three very similar birds, two from the United Arab Emirates (UAE) and one from coastal Eritrea, whose measurements and details are given in Table 1.

The UAE specimens, collected by M.D. Gallagher, were both worn. One, collected from Khor Khalba on 24 March 1971, Reg. no. 1977.18.27, was labelled as a Sykes's Warbler *H. (caligata) rama*; the other, collected from Ras el Khaimah on 16 June 1972, Reg. no. 1972.6.5, was labelled as an Olivaceous Warbler *H. pallida elaeica*. Both birds were much shorter- and rounder-winged than *elaieca*. Their measurements, wing formula details and tail/wing ratios agreed well with those given for *rama* by Svensson (2001), and we judged that both were of this form.

The Eritrean specimen, Reg. no. 1952. 4.7, collected by K.D. Smith at Arafale on 5 November 1951, was in fresh plumage. It was labelled *H. p. elaeica*, but was certainly not this form. It had a much shorter wing and its measurements fitted well with *H. rama*. The pattern on the outer

tail-feathers, with whitish fringes near the tip merging into the dark centres, and the narrow, inconspicuous pale edges to the secondaries and tertiaries were also features consistent with *rama* rather than with *elaieca*. Wing formula details fitted with *rama*, but were not entirely typical of this species. The second primary (numbered ascendantly) was equal to the seventh (it is usually shorter in *rama*) and the sixth primary lacked emargination (it is emarginated in many *rama*). The small forms of *H. pallida* breeding in and south of the Sahara can show an identical wing structure. We were therefore unable at this stage to assign this specimen conclusively to *rama*. We have now been able to check the identity of all three specimens by reference to their mitochondrial DNA structure.

DNA was isolated from toe pads using standard methods, including proteinase K digestion and one step each of phenol and chloroform purification (Bensch & Pearson 2002). Because it is often difficult to amplify longer fragments from old specimens, presumably because of DNA degradation, we used the primers orinus1 and cyt 4 (Bensch & Pearson 2002). This protocol targets a 141-bp fragment located in the mitochondrial *cytochrome b* gene between the universal vertebrate primers L14841 and H15149 (Kocher *et al.* 1989). The PCR products were sequenced using the Big Dye terminator cyclic sequencing kit loaded on an ABI PRISM™ 310 (Perkin Elmer). The sequences were aligned using BioEdit (Hall 1999) and compared with published sequences retrieved from the GenBank International Nucleotide Sequence Database.

The mtDNA sequences of the three specimens were identical, and matched perfectly over the 141 bp investigated with the published sequence of *H. rama* (GenBank AJ004792, Krasnovodsk, Turkmenistan; Helbig & Seibold 1999). Comparison of this region with the published sequence for *H. caligata* (GenBank AJ004793, western Mongolia; Helbig & Seibold 1999) revealed 13 substitutions, with *H. pallida elaeica* (GenBank AJ004791, Tsavo, Kenya; Helbig & Seibold 1999) 11 substitutions, and with *H. p. pallida* (eight specimens, Egypt, S. Bensch unpubl. data) 12–13 substitutions. Hence, it is clear that these specimens are all representatives of *H. rama*.

The identity of the UAE specimens came as no surprise. However, confirmation that the Eritrean specimen was also *rama* was of considerable interest, as this appears to represent the first confirmed record for the species from Africa. There have been past claims of wintering *rama* in northeast Africa, and the species may well have been overlooked there, but none of these reports was accepted in reviews by Moreau (1972), Curry-Lindahl (1981) or Urban *et al.* (1997). Four specimens from Eritrea that Zedlitz thought were *rama* were later re-identified by him as *H. pallida* (Zedlitz 1910–1911, pp. 61 and 611), and we are informed by L. Svensson (*in litt.*) that these are in fact *H. p. elaeica*. Another specimen from Somalia, collected as *pallida* but re-identified by Meinertzhagen as *rama* (Archer & Godman 1961) appears to us to be *pallida*,

*Corresponding author.
Email: davidpearson@freeUK.com

Table 1. Details and measurements (in mm) of *Hippolais* specimens from Eritrea and the United Arab Emirates held by the Natural History Museum, Tring, UK. The primaries are numbered ascendantly.

Reg. no.	1977.18.27	1972.6.5	1952.4.7
Country	UAE	UAE	Eritrea
Locality	Khor Khalba (25°01'N, 56°22'E)	Ras el Khaimah (25°57'N, 56°03'E)	Arafale (15°05'N, 39°41'E)
Date	24.3.1971	16.6.1972	5.6.1951
Sex	M?	M	
Wing	59	61	59
Tail	52	52	49
Bill (to skull)	14.5	15.5	16
Tarsus	21.5	21	20
Longest primary	3–4	3–4	3–5
P2 equal to?	P8/9	P7/8	P7
P2 < wing tip	6.5	5.5	4.5
P6 < wing tip	1.5	2	2.5
P10 < wing tip	9	9.5	9
P1 > primary coverts	6.5	8	7
Emarginated primaries	3–6	3–6	3–5

probably of the race *elaieca*. Svensson (2001) has drawn attention to yet another specimen at Tring (Reg. no. 1934.1.1.5183), collected in southern Sudan on 7 April 1900, registered as *pallida* but later relabelled *rama*. We agree with him that this bird is difficult to assign with certainty. In conclusion, we would stress that particular care is required in the separation of *H. rama* from the races of *H. pallida* in Arabia and northeast Africa.

We thank Dr Robert Prŷs-Jones and Mark Adams for facilities and support at the Natural History Museum, Tring. Michael Gallagher kindly provided details of his collected specimens.

REFERENCES

- Archer, G. & Godman, E.M. 1961. *The Birds of British Somaliland and the Gulf of Aden*, Vol. IV. Edinburgh: Oliver & Boyd.
- Ash, J.S. & Pearson, D.J. 2002. *Hippolais* warblers apparently breeding on the north Somalia coast. *Bull. Brit. Orn. Club* **122**: 222–228.
- Bensch, S. & Pearson, D. 2002. The Large-billed Reed Warbler *Acrocephalus orinus* revisited. *Ibis* **144**: 259–267.
- Curry-Lindahl, K. 1981. *Bird Migration in Africa*. London: Academic Press.
- Hall, T.A. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* **41**: 95–98.
- Helbig, A.J. & Seibold, I. 1999. Molecular phylogeny of Palearctic–African *Acrocephalus* and *Hippolais* warblers (Aves: Sylviidae). *Mol. Phylogenetics Evol.* **11**: 246–260.
- Kocher, T.D., Thomas, W.K., Meyer, A., Edwards, S.V., Pääbo, S., Villablanca, F.X. & Wilson, A.C. 1989. Dynamics of mitochondrial DNA evolution in animals: amplification and sequencing with conserved primers. *Proc. Natl Acad. Sci. USA* **86**: 6196–6200.
- Moreau, R.E. 1972. *The Palaearctic–African Bird Migration Systems*. London: Academic Press.
- Svensson, L. 2001. Identification of Western and Eastern Olivaceous, Booted and Sykes's Warblers. *Birding World* **14**: 192–219.
- Urban, E.K., Fry, C.H. & Keith, G.S. (eds) 1997. *The Birds of Africa*, Vol. V. London: Academic Press.
- Zedlitz, O.G. 1910–11. Meine ornithologische Ausbeute in Nordost-Afrika. *J. Ornithol.* **58**: 290–394, 731–807; **59**: 1–92, 591–693.

Received 13 August 2003; revision accepted 15 March 2004.

First published online on 28 July 2004;

doi: 10.1111/j.1474-919x.2004.00305.x