LUND UNIVERSITY

Environmental monitoring using optical techniques

Svanberg, Sune

Published in:

Proceedings of SPIE - The International Society for Optical Engineering

DOI:

[10.1117/12.519493](https://doi.org/10.1117/12.519493)

2002

[Link to publication](#)

Citation for published version (APA):

Svanberg, S. (2002). Environmental monitoring using optical techniques. In *Proceedings of SPIE - The International Society for Optical Engineering* (Vol. 5226, pp. 234-237). SPIE. <https://doi.org/10.1117/12.519493>

Total number of authors:

1

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Environmental monitoring using optical techniques

Sune Svanberg

Department of Physics, Lund Institute of Technology, S-221 00 Lund, Sweden

and

Lund Laser Center, Lund University, S-221 00 Lund, Sweden

ABSTRACT

An overview of optical techniques for environmental monitoring is presented. Range-resolved measurements of atmospheric pollutants can be performed using the differential absorption lidar technique. Fluorescence lidar allows assessment of vegetation status and also the conditions of the facades of historical buildings. Diode lasers provide particularly realistic schemes for atmospheric gas analysis, where certain wavelength ranges, which are not easily directly assessed, can be reached by sum- and difference frequency generation. Finally, the gas correlation principle can be used for real-time imaging of hydrocarbons. Several types of such optical environmental monitoring are illustrated with examples from research at the Lund Institute of Technology, Sweden.

Keywords: Lidar, laser-induced fluorescence, diode laser spectroscopy, gas correlation, air pollution, hydrocarbons

1. INTRODUCTION

Optical spectroscopy, and in particular laser spectroscopy, provide many possibilities for real-world applications. Powerful techniques have been developed for chemical analysis, combustion diagnostics, environmental monitoring and biomedical diagnostics (see, e.g., [1,2]). A comprehensive treatment, where a vast number of applications of atomic and molecular spectroscopy is related to the basic aspects of the field, is presented in [3].

We will here focus on environmental monitoring, and numerous examples from the research performed at the Lund Institute of Technology are given. Atmospheric monitoring includes the laser radar, the diode laser spectroscopy and the gas correlation imaging methods, while laser-induced fluorescence (LIF) is applied to hydrospheric pollution measurements, vegetation status assessment and building facade monitoring. Non-intrusive measurements in real time are provided, as is characteristic for remote sensing techniques.

2. ATMOSPHERIC LIDAR

The Lund group has been pursuing atmospheric laser radar monitoring for a long time, including industrial and geophysical measurements. A review of some of the activities is given in Ref. 4. A full integration between plume-transect, integrated-gas-contents measurement and automatic wind assessment for total flux determinations at industries has been achieved [5,6]. Extensive work on the monitoring of mining areas, geothermal fields, and active volcanoes has been performed [7-13]. In particular, the total fluxes from the Italian volcanoes Etna, Stromboli and Vulcano have been measured in shipborne campaigns, where Differential Absorption Lidar (DIAL), Differential Optical Absorption Spectroscopy (DOAS) and Correlation Spectroscopy (COSPEC) were intercompared [14]. Present lidar group activities focus on hydrocarbon monitoring in the IR region. Hydrocarbons play a very important role in modern society, being primary fuels and also very important materials for the chemical and pharmaceutical industries. Monitoring and control of the outflow of gaseous hydrocarbons is important for preventing loss of valuable raw material, for avoiding explosion hazards as well as for environmental protection. The primary C-H stretch molecular vibration occurs in the 3.4 μm region with small and characteristic shifts for different molecules. We use a optical parametric oscillator (OPO) and difference frequency generation in our lidar system to enable IR monitoring of hydrocarbons [15-18].

3. FLUORESCENCE LIDAR

The Lund group has performed extensive fluorescence lidar work regarding water quality, vegetation status, and historical building facades [19-24]. In particular, part of the Lund Cathedral was scanned with a fluorescence lidar system for assessment of algal and lichen growth [25]. Full fluorescence spectra were recorded in each image point and multi-variate analysis was employed for data presentation. A field campaign for similar measurements in Parma, Italy, was recently performed [26].

4. DIODE LASER SPECTROSCOPY

Diode lasers provide convenient means for certain types of spectroscopic diagnostics. Normally, only line-of-sight path-integrated data are obtained, similarly as in DOAS [27]. A particularly high sensitivity in diode laser absorption measurements can be obtained in frequency-modulation (FM) spectroscopy, of which two-tone FM spectroscopy is an especially convenient variety. We have focussed on a full characterisation of the absorption lineshape, which allows the technique to be used under varying gas temperature and pressure conditions. With the fast development of the semiconductor laser technology, new wavelength regions are becoming available. We recently demonstrated first single-mode diode laser spectroscopy in the violet spectral range, frequency-modulation spectroscopy with blue diodes, and also their usefulness in sum-frequency generation to the UV region, also accessing the mercury line at 254 nm [28-30]. We are also, like many other groups, performing diode laser spectroscopy in the IR range, accessible by difference-frequency generation or directly by quantum cascade lasers. Simultaneous monitoring of background atmospheric methane and the oxygen and water vapour concentrations, was recently performed by tuning two diode lasers to appropriate wavelengths [31].

An emerging new field of diode laser spectroscopy, GASMAS (GAs in Scattering Media Absorption Spectroscopy) relates to the study of free gas dispersed as bubbles or pore enclosures in scattering materials such as wood, fruits, ceramics, foams, and insulating materials. Thus, oxygen gas was extensively studied in many materials, with demonstrations also of the possibilities for pressure measurements and diffusion studies [32,33]. By combining CW measurements with pulsed photon propagation studies, which map out the different pathlengths through the scattering medium, it is possible to determine gas concentrations [34]. The new technique uses quite simple equipment, which still allows advanced studies in several novel fields to be performed also in developing countries [35].

5. GAS CORRELATION IMAGING

We have developed a powerful imaging technique for hydrocarbon monitoring. The gas correlation principle combined with passive thermal background illumination is employed for the visualization of gases absorbing in the IR region [36]. The technique was demonstrated at a full-scale polyethylene plant, where ethylene emissions from flares as well as pipes were detected. The method is expected to allow effective surveillance from the production, distribution to the end use phase for petroleum products.

6. CONCLUSIONS

Environmental monitoring using optical techniques has reached a mature state in certain aspects, while other areas of application are emerging. With the fast development of compact and tuneable lasers, sensitive detectors and computers, the techniques can be expected to play a major role in the future surveillance and management of the environment

Acknowledgements. The author gratefully acknowledges a very fruitful collaboration with past and present colleagues and students, as well as generous support from many funding agencies.

REFERENCES

1. C. Fotakis, and S. Svanberg, Applications of Laser Spectroscopy, *Europhysics News* **29**, 151, 1998.
2. S. Svanberg, Chemical Sensing with Laser Spectroscopy, *Sensors and Actuators* **B33**, 1, 1996.
3. S. Svanberg, *Atomic and Molecular Spectroscopy – Basic Aspects and Practical Applications*, 3rd Edition Springer, Heidelberg, Berlin, New York, 2001.
4. S. Svanberg, Differential Absorption Lidar (DIAL), in *Air Monitoring by Spectroscopic Techniques*, M. Sigrist (ed), Wiley, N.Y. 1994
5. P. Weibring, M. Andersson, H. Edner and S. Svanberg, *Remote Monitoring of Industrial Emissions by Combination of Lidar and Plume Velocity Measurements*, *Appl. Phys.* **B66**, 383, 1998.
6. I. Wängberg, H. Edner, R. Ferrara, E. Lanzillotta, J. Munthe, J. Sommar, S. Svanberg, M. Sjöholm, and P. Weibring, Mercury Emissions from a Chlor-alkali Plant in Sweden, *Science of the Total Env.*, in press.
7. H. Edner, G.W. Faris, A. Sunesson, S. Svanberg, J.Ö. Bjarnason, K.H. Sigurdsson and H. Kristmansdottir, Lidar Search for Atomic Mercury in Icelandic Geothermal Fields, *J. Geophys. Res.* **96**, 2977, 1991.
8. H. Edner, P. Ragnarson, S. Svanberg, E. Wallinder, A. de Liso, R. Ferrara and B.E. Maserti, Differential Absorption Lidar Mapping of Atmospheric Atomic Mercury in Italian Geothermal Fields, *J. Geophys. Res.* **97**, 3779, 1992.
9. H. Edner, P. Ragnarson, S. Svanberg, E. Wallinder, R. Ferrara, B.E. Maserti and R. Bargagli, Atmospheric Mercury Mapping in a Cinnabar Mining Area, *The Science of the Total Environment* **133**, 1, 1993.
10. H. Edner, P. Ragnarson, S. Svanberg, E. Wallinder, R. Ferrara, R. Cioni, B. Raco and G. Taddeucci, Total Fluxes of Sulphur Dioxide from the Italian Volcanoes Etna, Stromboli and Vulcano Measured by Differential Absorption Lidar and Passive Differential Optical Absorption Spectroscopy, *J. Geophys. Res.* **99**, 18827, 1994.
11. R. Ferrara, B.E. Maserti, M. Andersson, H. Edner, P. Ragnarson, S. Svanberg and A. Hernandez, Atmospheric Mercury Concentrations and Fluxes in the Almaden District (Spain), *The Science of the Total Environment* **213**, 13, 1998.
12. P. Weibring, H. Edner, S. Svanberg, G. Cecchi, L. Pantani, R. Ferrara and T. Caltabiano, Monitoring of Volcanic Sulphur Dioxide Emissions using Differential Absorption Lidar (DIAL), Differential Optical Absorption Spectroscopy (DOAS) and Correlation Spectroscopy (COSPEC), *Appl. Phys.* **B67**, 419, 1998.
13. S. Svanberg, Geophysical Gas Monitoring using Optical Techniques: Volcanoes, Geothermal Fields and Mines, *Optics and Lasers in Engineering* **37**, 245, 2002.
14. P. Weibring, J. Swartling, H. Edner, S. Svanberg, T. Caltabiano, D. Condarelli, G. Cecchi and L. Pantani, Optical Monitoring of Volcanic Sulphur Dioxide Emissions – Comparison between four Different Remote Sensing Techniques, *Optics and Lasers in Engineering* **37**, 267, 2002.
15. P. Weibring, H. Edner and S. Svanberg, Versatile Mobile Lidar System for Environmental Monitoring, Submitted to *Applied Optics*
16. P. Weibring, J.N. Smith, H. Edner, and S. Svanberg, Development and Testing of a Frequency Agile Optical Parametric Oscillator System for Differential Absorption Lidar, Submitted to *Rev. Scientific Instruments*
17. P. Weibring, Ch. Abrahamsson, J.N. Smith, H. Edner and S. Svanberg, Multi-Component Chemical Analysis of Gas Mixtures Using a Continuously-Tuneable Lidar System, Manuscript
18. Y. Saito, P. Weibring, H. Edner and S. Svanberg, Possibility of Hard-target Lidar Detection of a Biogenic Volatile Organic Compound, α -pinene, over Forest Areas, *Appl. Opt.* **40**, 3572, 2000.
19. H. Edner, J. Johansson, S. Svanberg and E. Wallinder, Fluorescence Lidar Multi-Color Imaging of Vegetation, *Appl. Optics* **33**, 2471, 1994.
20. J. Johansson, M. Andersson, H. Edner, J. Mattsson and S. Svanberg, Remote Fluorescence Measurements of Vegetation Spectrally Resolved and by Multi-Colour Fluorescence Imaging, *J. Plant Physiology* **148**, 632, 1996.
21. S. Svanberg, Fluorescence Lidar Monitoring of Vegetation Status, *Phys. Scripta* **T58**, 79, 1995.
22. P. Weibring, D. Lognoli, R. Chiari, G. Cecchi, H. Edner, T. Johansson, L. Pantani, S. Svanberg, D. Tirelli and M. Trambusti, Lidar Remote Sensing of the Parma Cathedral and Baptistery, *SPIE* **4402-16**, 2001.
23. L. Pantani, G. Ballerini, G. Cecchi, E. Edner, D. Lognoli, T. Johansson, V. Raimondi, S. Svanberg, P. Tiano, L. Tomaselli and P. Weibring, Experiments on stony monument monitoring by laser-induced fluorescence, *J. Cult. Heritage* **1**, 345, 2000.
24. S. Svanberg, Fluorescence lidar, Chapter 7 of T. Fujii (ed.) *Recent Advances in Laser Remote Sensing*, Marcel Dekker, 2003, in press.

25. P. Weibring, Th. Johansson, H. Edner, and S. Svanberg, B. Sundnér, V. Raimondi, G. Cecchi, and L. Pantani, Fluorescence Lidar Imaging of Historical Monuments, *Applied Optics* **40**, 6111, 2001.
26. D. Lognoli, G. Cecchi, L. Pantani, V. Raimondi, R. Chiari, Th. Johansson, P. Weibring, H. Edner and S. Svanberg, Fluorescence Lidar Imaging of the Parma Cathedral and Batistery, *Appl. Physics B*, in press.
27. H. Edner, P. Ragnarson, S. Spännare and S. Svanberg, A Differential Optical Absorption Spectroscopy (DOAS) System for Urban Atmospheric Pollution Monitoring, *Appl. Opt.* **32**, 327, 1993.
28. U. Gustafsson, J. Alnis and S. Svanberg, Atomic Spectroscopy with Violet Laser Diodes, *Am. J. Phys.* **68**, 660, 2000.
29. U. Gustafsson, G. Somesfalean, J. Alnis and S. Svanberg, Frequency Modulation Spectroscopy with Blue Diode Lasers, *Appl. Optics* **39**, 3774, 2000.
30. J. Alnis, U. Gustafsson, G. Somesfalean and S. Svanberg, Sum-frequency Generation with a Blue Diode Laser for Mercury Spectroscopy at 254 nm, *Appl. Phys. Lett.* **76**, 1234, 2000.
31. U. Gustafsson, J. Sandsten, and S. Svanberg, Simultaneous Detection of Methane, Oxygen and Water Vapour Utilizing Near-Infrared Diode Lasers in Conjunction with Difference Frequency Generation, *Appl. Phys. B* **71**, 853 2000.
32. M. Sjöholm, G. Somesfalean, J. Alnis, S. Andersson-Engels, and S. Svanberg, Analysis of Gas Dispersed in Scattering Solids and Liquids, *Opt. Lett.* **26**, 16, 2001.
33. J. Alnis, B. Anderson, G. Somesfalean, M. Sjöholm and S. Svanberg, Laser spectroscopy of molecular oxygen in wood materials, submitted to *Appl. Phys. B*
34. G. Somesfalean, M. Sjöholm, J. Alnis, C. af Klinteberg, S. Andersson-Engels and S. Svanberg, Concentration measurement of gas imbedded in scattering media employing time and spatially resolved techniques, *Appl. Optics* **41**, 3538, 2002.
35. S. Svanberg, Laser Spectroscopy in Development, *Europhysics News* **33**, March/April 2002, p. 52
36. J. Sandsten, P. Weibring, H. Edner and S. Svanberg, Real-time Gas-Correlation Imaging Employing Thermal Background Radiation, *Optics Express* **6**, 92, 2000; <http://www.opticsexpress.org>.