LUND UNIVERSITY

Transcriptional up-regulation in expression of 5-hydroxytryptamine2A and transcriptional down-regulation of angiotensin II type 1 receptors during organ culture of rat mesenteric artery.

Luo, Guogang; Xu, Cang-Bao; Cao, Yong-Xiao; Edvinsson, Lars

Published in:
Basic & Clinical Pharmacology & Toxicology

DOI:
[10.1111/j.1742-7843.2004.t01-1-pto950506.x](https://doi.org/10.1111/j.1742-7843.2004.t01-1-pto950506.x)

2004

[Link to publication](#)

Citation for published version (APA):

Luo, G., Xu, C-B., Cao, Y-X., & Edvinsson, L. (2004). Transcriptional up-regulation in expression of 5-hydroxytryptamine2A and transcriptional down-regulation of angiotensin II type 1 receptors during organ culture of rat mesenteric artery. *Basic & Clinical Pharmacology & Toxicology*, 95(6), 280-287.
<https://doi.org/10.1111/j.1742-7843.2004.t01-1-pto950506.x>

Total number of authors:
4

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Transcriptional Up-Regulation in Expression of 5-Hydroxytryptamine_{2A} and Transcriptional Down-Regulation of Angiotensin II type 1 Receptors during Organ Culture of Rat Mesenteric Artery

Guogang Luo^{1,2}, Cang-Bao Xu^{1,2}, Yong-Xiao Cao^{1,2} and Lars Edvinsson^{1,2}

¹Division of Experimental Vascular Research, Institute of Medicine, Lund University, Sweden and

²Medical College of Xian Jiaotong University, China

(Received March 17, 2004; Accepted August 19, 2004)

Abstract: The purpose of this study was to investigate in rat mesenteric artery if there is up-regulation of 5-hydroxytryptamine (5-HT) receptors and angiotensin II receptors and the potential role of protein kinase C activation in the smooth muscle cells during organ culture. Angiotensin II, 5-HT and potassium induced contraction of ring segments without endothelium, monitored by a sensitive *in vitro* pharmacology method. After the culture of the arterial ring segments for 24 hr, the concentration-contraction curves induced by 5-HT slightly shifted towards the left with pEC₅₀ from 6.64±0.11 to 6.84±0.11 and a significant increase in E_{max} from 147±11% to 246±15% (P<0.05), compared with that obtained in fresh segments. In contrast, the angiotensin II concentration-contraction curve only showed a significant decrease in E_{max} from 99±10% to 37±8%. Specific antagonists for the 5-HT type 2A receptors (5-HT_{2A}) and angiotensin II type 1 receptors (AT₁) demonstrated that the contractions occurred via 5-HT_{2A} and AT₁ receptors, respectively. Real-time PCR revealed that the 5-HT_{2A} receptor mRNA was up-regulated in parallel with the contractile response while there was a down-regulation of AT₁ receptor mRNA. Transcriptional inhibitor actinomycin D and specific protein kinase C inhibitor Ro31-8220 demonstrated that it was a transcriptional mechanism with involvement of protein kinase C that regulated the enhanced expression of 5-HT_{2A} receptors in the mesenteric artery.

5-Hydroxytryptamine (5-HT) and angiotensin II (AngII) are signaling molecules that have important roles in cardiovascular regulation (Turla & Webb 1989; Chester *et al.* 1993; Yildiz *et al.* 1998; Mizuno *et al.* 1989). Both have been seen to be synthesized in vascular endothelial cells (Lincoln *et al.* 1990). Their dominating receptors in smooth muscle cells of rat mesenteric artery are of the 5-HT type 2A (5-HT_{2A}) and Ang II type 1 (AT₁) receptors (Zemin *et al.* 1999). They are considered to be involved both in regulation of tone in the smooth muscle cells and in remodeling via mitogenesis (Florian & Watts 1998; Watts 1998; Ytterberg & Edvinsson 2001; Fang & Marwick 2002). Elevated levels of Ang II and 5-HT are seen in stroke (Nishimura & Suzuki 1995; Walther *et al.* 2002), and in congestive heart failure (Gschwend *et al.* 2003). As a feedback mechanism, there is local release of nitric oxide and prostacyclin, which overweighs the vascular constrictors but these may be particularly important in vascular disease with altered receptor expression. In situations of elevated levels of 5-HT or Ang II their impact on the smooth muscle cells might be pronounced, particularly, when the agonist concentrations in the concentration-response curves are logarithmic. We have previously shown enhanced expression of the endothelin type B receptor

(ET_B) in experimental subarachnoid haemorrhage (Alafaci *et al.* 1990; Hansen-Schwartz *et al.* 2002), in stroke (Stenman *et al.* 2002) and in ischaemic heart disease (Wackenfors *et al.* 2004) which results in markedly enhanced responses to the agonists. The ET_B receptor may not only change its phenotype from a relaxant to a contractile ET_B receptor phenotype but also in increased receptor number as a result of organ culture (Adner *et al.* 1996). The mechanisms behind this phenomenon involve mitogen activated-protein kinase (Uddman *et al.* 2003), protein kinase C (Uddman *et al.* 2002) and transcription factors (Möller *et al.* 1997). However, we do not know if this phenomenon is unique to ET_B receptors or if alterations in other G-protein coupled receptors may occur during organ culture. There are data supporting since there is increased efficacy to Ang II in stroke (Mizuno *et al.* 1999; Stenman & Edvinsson 2004) and to 5-HT in SAH (Saida *et al.* 1997; Miranda *et al.* 1996) which imply a change in receptor expression (Zhu *et al.* 2000), while the elevated circulating level of Ang II in heart failure may results in the down-regulation of Ang II receptors (De Gasparo *et al.* 1994; Regitz-Zagrosek *et al.* 1994; Haywood *et al.* 1997).

The present study was designed to examine if organ culture of arterial segments change vascular smooth muscle cells receptor expression. The rat mesenteric artery has been used as a representative of a peripheral resistance vessel (Ralevic *et al.* 1991), and presently we have examined if there are alterations in the expression of Ang II and 5-HT

Author for correspondence: Cang-Bao Xu, Division of Experimental Vascular Research, Institute of Medicine, Lund University, SE-221 84, Sweden (fax +46 46 222 06 16, e-mail Cang-Bao.Xu@med.lu.se).

receptors in rat mesenteric artery with functional and molecular methods. Actinomycin D (Strohm *et al.* 2002) and the specific protein kinase C inhibitor Ro 31-8220 (Trachsel & Keller 1995) were used to study transcription and involvement of protein kinase C, respectively.

Materials and Methods

Tissue preparation and organ culture procedure. Sprague-Dawley rats (body weight 250–300 g) were anaesthetized with CO₂ and exsanguinated. The superior mesenteric artery was removed gently, immersed into cold buffer solution (for composition, see below) and dissected free of adhering tissue under a microscope. The endothelium was denuded by perfusion of the vessel for 10 sec. with 0.1% Triton X-100 followed by another 10 sec. with a physiologic buffer solution (Hamel *et al.* 1987). The vessels were then cut into 1 mm long cylindrical segments, used directly (fresh group) or incubated for 6 hr–2 days at 37° in humidified 5% CO₂ in O₂ (organ culture group). The segments for organ culture were placed in a 96-well plate, one segment in each well, containing 300 µl Dulbecco's modified Eagle's medium (DMEM) containing L-glutamine (584 mg/l) and supplemented with penicillin (100 U/ml) and streptomycin (100 µg/ml). The medium was changed to fresh daily. The experimental protocol was approved by Lund University Animal Ethic Committee (M217-03).

In vitro pharmacology. Fresh or incubated segments were immersed in temperature-controlled (37°) tissue baths containing a bicarbonate buffer solution. The solution was continuously aerated with 5% CO₂ in O₂ resulting in a pH of 7.4. Each segment was mounted on two L-shaped prongs, one of which was attached to a Grass FT-03 transducer (Grass Instr., Quincy, USA) connected to a PowerLab (ADInstruments, Hastings, UK) unit for continuous recording of isometric tension. A resting tone of about 2.5 mN was applied to each segment and the segments were allowed to stabilize at this tension for at least one hour before being exposed to a potassium-rich (60 mM) buffer solution with the same composition as the standard solution except that NaCl was replaced by an equimolar concentration of KCl. The potassium-induced contraction was used as a reference for the contractile capacity, and the segments was used only if potassium elicited reproducible responses over 1.0 mN. Concentration-response curves for vasoconstrictors were obtained by cumulative administration of the agents.

Buffer solutions and drugs. Standard buffer solution (mM): NaCl 119; NaHCO₃ 15; KCl 4.6; MgCl 1.2; NaH₂PO₄ 1.2; CaCl₂ 1.5; glucose 5.5. Analytical grade chemicals and double distilled water were used for preparing all solutions. Dulbecco's modified Eagle's medium, penicillin and streptomycin were purchased from Gibco BRL, Paisley, Scotland, UK. Ang II, 5-HT, Ro 31-8220, ketanserin, candesartan, cyclohexamide and actinomycin D (Sigma, St. Louis, USA) were dissolved in sterile water with bovine serum albumin (0.1% w/v).

Real time RT-PCR. Smooth muscle isolated from fresh or the cultured segments was homogenated in 1 ml of the RNeasyTM solution (Q-BIOgene, CA, USA) by using a FastPrep[®] instrument (Q-BIOgene, CA, USA). The total RNA was extracted following a protocol from the FastRNA[®] Pro kit supplier. Reverse transcription of total RNA to cDNA was carried out using the Gene Amp RT kit (PE Applied Biosystems) in a Perkin-Elmer 2400 PCR machine at 42° for 30 min. The real-time quantitative PCR was performed with the GeneAmp SYBR Green PCR kit (PE Applied Biosystems) in a Perkin-Elmer real-time PCR machine (PE, GeneAmp 5700 sequence detection system). The system automatically monitors the binding of a fluorescent dye to double-strand DNA by real-time detection of the fluorescence during each cycle of PCR amplifi-

cation. Specific primers for rat AT_{1A} and 5-HT_{2A} receptors were designed based on gene bank data by using primer expression 2 software (PE Applied Biosystems).

5-HT_{2A} receptor primers;

Forward: 5'-CTCAGTCACCATGCGACAGG-3'

Reverse: 5'-TTGTCGCCAGGGTCCG-3'

AT_{1A} receptor primers;

Forward: 5'-3' GGA TGG TTC GCA GAG AGA GTA CAT

Reverse: 5'-3' CCT GCC CTC TTG TAC CTG TTG

The house keeping gene, GAPDH mRNA continuously expressed to a constant amount in the cells, was compared with the house keeping gene EF-1 in a pilot study by real-time PCR (Stenman *et al.* 2002). GAPDH was used as a reference in this study, but both were equally well constant in the tests.

GAPDH primers

Forward: 5'-GGCCTTCCGTGTTCTACC-3'

Reverse: 5'-CGGCATGTCAGATCCACAAC-3'

The PCR reaction was performed in a 50 µl volume and started at a temperature of 50° for 2 min., 95° for 10 min. and the following 40 PCR cycles with 95° for 15 sec. and 60° for 1 min. Dissociation curves were run after the real time PCR to identify the specific PCR products. All primers were designed using the Primer Express 2.0 software (PE Applied Biosystems) and synthesized by TAG Copenhagen A/S (Denmark).

Data were analyzed with the comparative cycle threshold (CT) method. To evaluate the amount of 5-HT_{2A} and AT_{1A} mRNA in a sample, GAPDH mRNA was assessed in the same sample simultaneously. The CT values of GAPDH mRNA were used as a reference to quantify the relative amount of 5-HT_{2A} and AT_{1A} mRNA. The relative amount of mRNA was calculated with the CT values of 5-HT_{2A} and AT_{1A} receptor mRNA in relation to the CT values of GAPDH mRNA in the sample.

Calculation and statistics. Data are expressed as mean values ± S.E.M. Contractile responses in each segment are expressed as a percentage of the potassium-induced contraction. E_{max} represents the maximal contraction induced by an agonist. The pEC₅₀ value was calculated from the line between the concentrations above and below the midpoint of the concentration-response curve. Statistical analysis was performed with unpaired Student's t-test or one-way ANOVA with Dunnet's post test. Differences were considered significant at P<0.05.

Results

5-Hydroxytryptamine receptors.

Contractile response. 5-HT, a general 5-HT receptor agonist, resulted in a concentration-dependent contraction. In fresh arterial segments the E_{max} and pEC₅₀ values were 147±11% and 6.64±0.11, respectively. After 24 hr of incubation, the curves slightly shifted to the left with the pEC₅₀ values from 6.64±0.11 to 6.84±0.11 and the maximal contraction increased significantly from 147±11% to 246±15% (P<0.05) (fig. 1A, table 1). The selective 5-HT_{2A} receptor antagonist ketanserin (10⁻⁹–10⁻⁸ M) shifted the concentration-response in a concentration-dependent manner to the right without any change in maximum contraction (fig. 1B, table 1), indicating that the contraction induced by 5-HT was via a 5-HT_{2A} receptor subtype (Watts 2002). The pEC₅₀ values of 5-HT with ketanserin 10⁻⁹ M and 10⁻⁸ M decreased

from 6.74 ± 0.09 (without ketanserin) to 6.55 ± 0.06 and 5.22 ± 0.07 , respectively. A Schild plot of the data yielded a pA_2 value of 9.43 (slope=1) (Watts 2002). 5-CT, a selective 5-HT₁ agonist, gave rise to a concentration-dependent contraction in the fresh arterial segments. The concentration-contraction curves of 5-CT did not shift obviously following organ culture for 24 or 48 hr and there was no significantly difference in the maximum contraction or the pEC_{50} values (data not shown).

Transcription and translation inhibition experiments. In order to examine the mechanisms of up-regulation of 5-HT_{2A} receptors, we used the transcription inhibitor actinomycin D (Strohm *et al.* 2002) and the translation inhibitor cyclohexamide (Möller *et al.* 1997). The results revealed that

both inhibitors significantly decreased the enhanced 5-HT_{2A} receptor-mediated contraction. The enhanced contraction had returned to that seen in fresh arterial segments in actinomycin D (5×10^{-6} M) treated segments while cyclohexamide (5×10^{-5} M) only partially reduced the up-regulation (fig. 3, table 1). Furthermore, the involvement of protein kinase C was addressed by using its specific inhibitor Ro 31-8220 (10^{-6} M). The presence of Ro31-8220 during organ culture resulted in an attenuated 5-HT concentration-contraction curve with an E_{max} of $158 \pm 12\%$ compared to that seen after organ culture in the absence of Ro31-8220 ($246 \pm 15\%$) ($P < 0.01$) (fig. 4, table 1).

Real-time PCR. To confirm the transcriptional mechanisms and the involvement of protein kinase C, changes in mRNA levels for 5-HT_{2A} receptors were analyzed before and after organ culture by real-time PCR. The mRNA level of 5-HT_{2A} receptors was significantly increased ($P < 0.05$) following organ culture. The presence of the transcription inhibitor actinomycin D or the specific protein kinase C inhibitor Ro31-8220, but not the translation inhibitor cyclohexamide, significantly decreased the up-regulation of mRNA for the 5-HT_{2A} receptors (fig. 5A).

Fig. 1. Concentration response curves for 5-HT in fresh vessels and following 24 hr organ culture (A) or in presence of selective 5-HT_{2A} receptor antagonist ketanserin (10^{-8} M, 10^{-9} M) on rat mesenteric arteries after incubation 24 hr (B). Data are derived from 10–16 identical experiments and shown as mean $\pm$ S.E.M.

Table 1.

Overview of E_{max} and pEC_{50} values of 5-HT and Ang II induced contraction on rat mesenteric arteries.

Agonist	N	E_{max} (%)	pEC_{50}
5-HT			
Fresh	16	147 ± 11	6.64 ± 0.11
Organ culture 24 hr	16	$246 \pm 15^*$	6.84 ± 0.11
Antagonists			
Control	10	248 ± 16	6.74 ± 0.09
Ketanserin 10^{-9} M	10	245 ± 14	$6.11 \pm 0.09^+$
Ketanserin 10^{-8} M	10	248 ± 11	$5.22 \pm 0.07^*$
Inhibitors			
Control	16	248 ± 16	6.61 ± 0.03
Actinomycin D 5×10^{-6} M	11	$147 \pm 12^*$	6.50 ± 0.02
Cyclohexamide 5×10^{-5} M	11	$190 \pm 18^+$	6.52 ± 0.02
Ro 31-8220 10^{-6} M	16	$158 \pm 12^*$	6.44 ± 0.09
Angiotensin II			
Fresh	11	99 ± 10	8.93 ± 0.17
Organ culture 24 hr	8	$37 \pm 8^*$	8.26 ± 0.07
Organ culture 48 hr	10	$33 \pm 12^*$	8.58 ± 0.07
Antagonist			
Control	11	103 ± 7	8.25 ± 0.08
Candesartan 10^{-12} M	12	97 ± 8	8.24 ± 0.03
Candesartan 10^{-10} M	15	$53 \pm 10^+$	$6.25 \pm 0.08^*$
Candesartan 10^{-8} M	15	$17 \pm 8^*$	$5.73 \pm 0.02^*$
Inhibitors			
Control	7	36 ± 8	8.24 ± 0.06
Actinomycin D 5×10^{-6} M	8	0	0
Cyclohexamide 5×10^{-5} M	8	0	0
Ro 31-8220 10^{-6} M	8	0	0

Maximum contractile effect E_{max} are expressed as percent of 60 mM K⁺-induced contraction. pEC_{50} values mean negative logarithm of the molar concentration that produced half maximum contraction. Data are shown as mean $\pm$ S.E.M. N denotes the number of vessel segments. * $P < 0.01$ and $^+P < 0.05$ versus fresh or control segments, respectively.

Angiotensin II receptors.

Contractile response. Ang II caused a concentration-dependent contractile response in fresh mesenteric artery segments. The maximum contraction was $99 \pm 10\%$ and the pEC_{50} value 8.93 ± 0.17 . The contraction response curves were depressed markedly after organ culture for 24 hr or 48 hr (fig. 2A, table 1). The AT₁ receptor antagonist, candesartan, shifted the concentration-contractile curve to right and also reduced the maximum contraction (fig. 2B, table 1). The antagonism was similar to that previously seen in rabbit aorta (Morsing *et al.* 1999).

Fig. 2. Concentration response curves for Angiotensin II in fresh and organ culture for 24, 48 hr (A) or in the presence of AT₁ receptor antagonist candesartan (10^{-8} M, 10^{-10} M, 10^{-12} M) on fresh mesenteric arteries of rat (B). Data are derived from 8–15 identical experiments and shown as mean $\pm$ S.E.M.

Fig. 3. The effects of transcriptional inhibitor actinomycin (5×10^{-6} M) and translation inhibitor cyclohexamide (5×10^{-5} M) exert on 5-HT induced concentration response curve which was co-cultured for 24 hr of rat mesenteric artery segments. Data are derived from 11–16 experiments and shown as mean $\pm$ S.E.M.

Real-time PCR. The presences of mRNA for the AT_{1A} receptors was analyzed before and after organ culture by real-time PCR. In concert with the functional data above, the mRNA level of AT_{1A} was markedly decreased after organ culture of the mesenteric artery for 24 hr ($P < 0.05$) (fig. 5B).

Fig. 4. The specific protein kinase C inhibitor Ro 31-8220 (10^{-6} M) altered the concentration-response curve of 5-HT which was co-cultured for 24 hr of rat mesenteric arteries. Data are derived from 16 identical experiments and shown as mean $\pm$ S.E.M.

Fig. 5. The comparison of mRNA level on rat mesenteric arteries of fresh, organ culture for 24 hr and co-cultured with different inhibitors: translation inhibitor cyclohexamide, transcriptional inhibitor actinomycin and protein kinase C inhibitor Ro 31-8220 (A). The mRNA expression level of AT₁ receptor on fresh rat mesenteric segments or after organ culture for 24 hr (B). Data are derived from 3–5 identical experiments and shown as mean $\pm$ S.E.M. Fresh versus organ culture * P <0.01, ** P <0.001. Organ culture versus inhibitor # P <0.05.

Protein kinase C inhibitor experiments. In order to study if protein kinase C activity was involved in the down-regulation of AT₁ receptors, the specific protein kinase C inhibitor Ro 31-8220 (10^{-6} M) was present during organ culture. Furthermore, inhibitors for transcription and translation, actinomycin D and cyclohexamide, were also studied. The presence of either of the three above inhibitors for 24 hr resulted in the total loss of the Ang II induced contractile responses; it decreased to zero in the organ cultured vessel segments (fig. 6).

Discussion

The present study has for the first time shown that organ culture of rat mesenteric artery results in altered expression of 5-HT_{2A} and AT₁ receptors in the smooth muscle cells. Alteration in the expression of these G-protein-coupled receptors may prove to be a key event in vascular disease resulting in vascular smooth muscle spasm, proliferation and vascular remodeling. We have previously demonstrated that the ET_B receptor shows plasticity by changing from a relax-

Fig. 6. The effects of protein kinase C inhibitor Ro 31-8220 (10^{-6} M), translation inhibitor cyclohexamide 5×10^{-5} M and transcription inhibitor actinomycin D 5×10^{-6} M on angiotensin II induced contraction during organ culture of rat mesenteric artery for 24 hr compared with fresh. Data are derived from 7–8 identical experiments and shown as mean $\pm$ S.E.M..

ant to a contractile phenotype during organ culture of cerebral and peripheral arteries, occurred not only *in vitro* but also *in vivo* e.g. in ischaemic stroke (Stenman *et al.* 2002) and subarachnoid haemorrhage (Hansen-Schwartz *et al.* 2003). The present study has revealed that there are alterations in the expression of 5-HT_{2A} and AT₁ receptors during organ culture of rat mesenteric arteries evident both at mRNA and functional levels. For ET_B receptors in cerebral arteries, this has been shown also at the protein level (Henriksson *et al.* 2004). The up-regulation of the 5-HT_{2A} receptor occurs via a transcriptional mechanism that involves protein kinase C. The results are well in line with our previous studies that have demonstrated that similar mechanisms are involved in the up-regulation of ET_B receptors *in vivo* and *in vitro* (Uddman *et al.* 2002). Protein kinase C activity seems to be a key factor in the transcription of G-protein-coupled receptors and thus blockage of protein kinase C activity during organ culture regulates this. We do not think that smooth muscle cell phenotype changes occurred in the present organ culture of the artery ring segments, since it is known that the contractile response to potassium did not change in the segments cultured for 5 days (Adner *et al.* 1996). In the present study, the segments only cultured for 1–2 days and the culture medium did not contain any serum and growth factors. Smooth muscle cell phenotype changes in primary culture requires longer than 6 days of culture period and also growth factors (Chamley-Campbell *et al.* 1981).

The postsynaptic 5-HT receptors in the peripheral circulation, e.g. the mesenteric artery of the rat, consist mainly

of the 5-HT_{2A} type as based on responses to 5-HT and antagonism by ketanserin with a pA₂ 9.43; this is in concert with published data for 5-HT₂ receptors, pA₂=9.3 for 5-HT_{2A} and 6.5 for 5-HT_{2C} (Hoyer *et al.* 1994; Watts 2002), and the presence of mRNA of 5-HT_{2A} receptors. In fresh rat mesenteric arteries, 5-HT produced a concentration-dependent contraction with an E_{max} of 147±11% which increased after organ culture for 24 hr. The selective 5-HT_{2A} receptor antagonist, ketanserin shifted the concentration-contraction curves of 5-HT to the right in a parallel manner without any change of E_{max} which shows that it was a specific up-regulation of 5-HT_{2A} receptors. The up-regulated receptor has the same pA₂ for ketanserin as that seen in fresh arteries. Real-time PCR is a method that allows the quantitative demonstration of receptor mRNA and here we have showed that the up-regulation is evident also at the mRNA level. This suggests a transcriptional mechanism responsible for the up-regulation. The transcription inhibitor actinomycin and the translation inhibitor cyclohexamide confirmed that the event occurs via a transcriptional mechanism. In agreement with the present study, Adner *et al.* (2002) have shown up-regulation of 5-HT_{2A} receptors in smooth muscle cells during organ culture of mice tracheal segments.

Furthermore, analysis of the mechanisms triggering the up-regulation revealed that protein kinase C has a central role for ET_B receptors (Uddman *et al.* 2002). In our study, the protein kinase C inhibitor Ro 31-8220 abolished both the increase in 5-HT_{2A} receptor mRNA and reduced the contractile effects of 5-HT to that seen in fresh arteries. Thus, protein kinase C activity is involved in the up-regulation of 5-HT_{2A} receptors. Protein kinase C is known to play an important role as an intracellular signal for activation of transcriptional factor nuclear factor-κB (NF-κB) and activation protein-1 (AP-1) (Sliva *et al.* 2002; Chuang *et al.* 2003). Since the protein kinase C inhibitor significantly inhibited the up-regulation of 5-HT_{2A} receptors at both functional and mRNA levels, we have demonstrated that protein kinase C activity is required for the up-regulation. A similar phenomenon with protein kinase C has been demonstrated for ET_B receptors and in addition the downstream extracellular signal-regulated kinase 1/2 (ERK1/2) is involved in the up-regulation. Thus, blockage of protein kinase C activity might be a novel strategy of use in the regulation of the up-regulated G-protein-coupled receptors expression in vascular disease.

Ang II, the major effector molecule in the renin-angiotensin system, plays a central role in the control of sodium excretion and fluid volume as well as of vascular tone. The main actions of Ang II are mediated via a specific membrane-bound G-protein-coupled receptors AT₁ receptor, which plays an important role in the development and progression of many cardiovascular diseases. Available AT receptor blockers have been analysed in details and behave more or less as insurmountable antagonists (Ytterberg & Edvinsson 2001; Pantev *et al.* 2002). In heart failure there is down-regulation of AT receptors (Haywood *et al.* 1997; Touyz & Schiffrin 2000). This has been suggested as a feed-

back loop due to the elevated Ang II level in the circulation of subjects with heart failure (Makita *et al.* 1992; Lassegue *et al.* 1995). The intracellular mechanism behind the regulation of Ang II receptors is unclear, but has sometimes been considered due to enhanced receptor internalization. The contractile response to Ang II was in our study decreased in parallel with down-regulation of AT_{1A} receptor mRNA during organ culture of rat mesenteric arteries for 24 hr. This demonstrates a transcriptional mechanism for the down-regulation of AT_{1A} receptors. We do not know further intracellular mechanisms responsible for the down-regulation of AT₁ receptors, particularly since the experiments with actinomycin D or the protein kinase C inhibitor did not provide any clues. Production of Ang II, nitric oxide (Bucher *et al.* 2001), prostacyclin (Dusting *et al.* 1981; Oriji 1999) and protein kinase A activity (Xu & Murphy 2000) are putatively involved in the down-regulation since they have been shown to suppress G-protein-coupled receptor expression. Down-regulation of AT₁ receptors occur in human heart failure, failing human ventricular myocardium, hypertrophied human myocardium and atrial fibrillation (De Gasparo *et al.* 1994; Regitz-Zagrosek *et al.* 1994; Nozawa *et al.* 1996; Haywood *et al.* 1997; Tsutsumi *et al.* 1998) possibly via a mechanism similar to that seen during organ culture (Doan *et al.* 2004).

In conclusion, the present study demonstrated that 5-HT_{2A} receptors were transcriptionally up-regulated whereas the AT₁ receptors were down-regulated at the transcriptional level in rat mesenteric artery following organ culture. Protein kinase C activity was involved in the up-regulation of 5-HT_{2A} receptors during organ culture. Understanding of the G-protein-coupled receptor alterations that occur in organ culture may provide clues to their functions in cardiovascular disease.

Acknowledgements

This work is supported by Swedish Research Council (grant no. 5958), Council for Swedish Medical Tobacco Research (grant no. 200410) and Swedish Heart and Lung Foundation.

References

- Adner, M., A. C. Rose, Y. Zhang, K. Sward, M. Benson, R. Uddman, N. P. Shankley & L. O. Cardell: An assay to evaluate the long-term effects of inflammatory mediators on murine airway smooth muscle: evidence that TNF_α up-regulates 5-HT(2A)-mediated contraction. *Brit. J. Pharmacol.* 2002, **137**, 971–982.
- Adner, M., L. Cantera, F. Ehlert, L. Nilsson & L. Edvinsson: Plasticity of contractile endothelin-B receptors in human arteries after organ culture. *Brit. J. Pharmacol.* 1996, **119**, 1159–1166.
- Alafaci, C., I. Jansen, M. A. Arbab, Y. Shiokawa, N. A. Svendgaard & L. Edvinsson: Enhanced vasoconstrictor effector of endothelin in cerebral arteries from rats with subarachnoid haemorrhage. *Acta physiol. scand.* 1990, **138**, 317–319.
- Bucher, M., J. Hobbhahn & A. Kurtz: Nitric oxide-dependent down-regulation of angiotensin II type 2 receptors during experimental sepsis. *Crit. Care. Med.* 2001, **29**, 1750–1755.
- Chester, A. H., S. P. Allen, S. Tadjkarimi & M. H. Yacoud: Interaction between thromboxane A₂ and 5-hydroxytryptamine receptor

- subtypes in human coronary arteries. *Circulation* 1993, **87**, 874–880.
- Chuang, S. S., J. K. Lee & P. A. Mathew: Protein kinase C is involved in 2B4 (CD244)-mediated cytotoxicity and AP-1 activation in natural killer cells. *Immunology* 2003, **109**, 432–439.
- Chamley-Campbell, J. H., G. R. Campbell & R. Ross: Phenotype-dependent response of cultured aortic smooth muscle to serum mitogens. *J. Cell. Biol.* 1981, **89**, 379–383.
- De Gasparo, M., H. Rogg, M. Brink, L. Wang, S. Whitebread, G. Bullock & P. Erne: Angiotensin II receptor subtypes and cardiac function. *Eur. Heart. J.* 1994, **15**, Suppl. d98–103.
- Doan, T., P. Farmer, T. Cooney & M. S. Ali: Selective down-regulation of angiotensin II receptor type 1A signaling by protein tyrosine phosphatase SHP-2 in vascular smooth muscle cells. *Cell. Signal.* 2004, **16**, 301–311.
- Dusting, G. J., E. M. Mullins & R. D. Nolan: Prostacyclin (PGI₂) release accompanying angiotensin conversion in rat mesenteric vasculature. *Eur. J. Pharmacol.* 1981, **70**, 129–137.
- Fang, Z. Y. & T. H. Marwick: Vascular dysfunction and heart failure: epiphenomenon or etiologic agent? *Amer. Heart. J.* 2002, **143**, 383–390.
- Florian, J. A. & S. W. Watts: Intergration of mitogen activated protein kinase activation of vascular 5-hydroxytryptamine_{2A} receptor signal transduction. *J. Pharmacol. Exp. Therap.* 1998, **284**, 346–355.
- Gschwend, S., R. H. Henning, Y. M. Pinto, D. de Zeeuw, W. H. van Gilst & H. Buikema: Myogenic constriction is increased in mesenteric resistance arteries from rats with chronic heart failure: instantaneous counteraction by acute AT₁ receptor blockade. *Brit. J. Pharmacol.* 2003, **139**, 1317–1325.
- Hamel, E., C. Assumel-Lurdin, L. Edvinsson, D. Fage & E. T. MacKenzie: Neuronal versus endothelial origin of vasoactive acetylcholine in pial vessels. *Brain Res.* 1987, **420**, 391–396.
- Hansen-Schwartz, J., D. Szok & L. Edvinsson: Expression of ET(A) and ET(B) receptor mRNA in human cerebral arteries. *Brit. J. Neurosurg.* 2002, **16**, 149–153.
- Hansen-Schwartz, J., N. L. Hoel, M. Zhou, C. B. Xu, N. A. Svendgaard & L. Edvinsson: Subarachnoid hemorrhage enhances endothelin receptor expression and function in rat cerebral arteries. *Neurosurgery* 2003, **52**, 1188–1195.
- Haywood, G. A., L. Gullestad, T. Katsuya, H. G. Hutchinson, R. E. Pratt, M. Horiuchi & M. B. Fowler: AT₁ and AT₂ angiotensin receptor gene expression in human heart failure. *Circulation* 1997, **95**, 1201–1206.
- Henriksson M., C. B. Xu & L. Edvinsson: Importance of ERK1/2 in upregulation of endothelin type B receptors in cerebral arteries. *Brit. J. Pharmacol.* 2004, **142**, 1155–1161.
- Hoyer, D., D. E. Clarke, J. R. Fozard, P. R. Hartig, G. R. Martin, E. J. Mylecharane, P. R. Saxena & P. P. Humphrey: International Union of Pharmacology classification of receptors for 5-hydroxytryptamine (Serotonin). *Pharmacol. Rev.* 1994, **46**, 157–203.
- Lassegue, B., R. W. Alexander, G. Nickenig, M. Clark, T. J. Murphy & K. K. Griendling: Angiotensin II down-regulates the vascular smooth muscle AT₁ receptor by transcriptional and post-transcriptional mechanisms: evidence for homologous and heterologous regulation. *Mol. Pharmacol.* 1995, **48**, 601–609.
- Lincoln, J., A. Loesch & G. Burnstock: Localization of vasopressin, serotonin and angiotensin II in endothelial cells of the renal and mesenteric arteries of the rat. *Cell. Tissue. Res.* 1990, **259**, 341–344.
- Makita, N., N. Iwai, T. Inagami & K. F. Badr: Two distinct pathways in the down-regulation of type-1 angiotensin II receptor gene in rat glomerular mesangial cells. *Biochem. Biophys. Res. Commun.* 1992, **185**, 142–146.
- Miranda, F. J., J. A. Alabadi, G. Torregrosa, J. B. Salom, T. Jover, M. D. Barbera & E. Alborch: Modulatory role of endothelial and nonendothelial nitric oxide in 5-hydroxytryptamine-induced contraction in cerebral arteries after subarachnoid hemorrhage. *Neurosurgery* 1996, **39**, 998–1004.
- Mizuno, K., M. Tani, S. Niimura, S. Hashimoto, A. Satoh, K. Shimamoto, T. Inagami & S. Fukuchi: Direct evidence for local generation and release of angiotensin II in human vascular tissue. *Biochem. Biophys. Res. Commun.* 1989, **165**, 457–463.
- Mizuno, H., M. Ikeda, M. Harada, T. Onda & T. Tomita: Sustained contraction to angiotensin II and impaired Ca²⁺-sequestration in the smooth muscle of stroke-prone spontaneously hypertensive rats. *Amer. J. Hypertens.* 1999, **12**, 590–595.
- Morsing, P., G. Adler, U. Brandt-Eliasson, L. Karp, K. Ohlson, L. Renberg, P. O. Sjoquist & T. Abrahamsson: Mechanistic differences of various AT₁-receptor blockers in isolated vessels of different origin. *Hypertension* 1999, **33**, 1406–1413.
- Möller, S., L. Edvinsson & M. Adner: Transcriptional regulated plasticity of vascular contractile endothelin ET(B) receptors after organ culture. *Eur. J. Pharmacol.* 1997, **329**, 69–77.
- Nishimura, Y. & A. Suzuki: Enhanced contractile responses mediated by different 5-HT receptor subtypes in basilar arteries, superior mesenteric arteries and thoracic aortas from stroke-prone spontaneously hypertensive rats. *Clin. Exp. Pharmacol. Physiol. Suppl.* 1995, **22**, S99–101.
- Nozawa, Y., H. Miyake, A. Haruno, S. Yamada, S. Uchida, T. Ohkura, R. Kimura, H. Suzuki & T. Hoshino: Down-regulation of angiotensin II receptors in hypertrophied human myocardium. *Clin. Exp. Pharmacol. Physiol.* 1996, **23**, 514–518.
- Orij, G. K.: Angiotensin II-induced ET and PGI₂ release in rat aortic endothelial cells is mediated by PKC. *Prostaglandins. Leukot. Essent. Fatty Acids* 1999, **61**, 113–117.
- Pantev, E., E. Stenman, A. Wackenfors, L. Edvinsson & M. Malmjö: Comparison of the antagonistic effects of different angiotensin II receptor blockers in human coronary arteries. *Eur. J. Heart Fail.* 2002, **4**, 699–705.
- Ralevic, V., R. T. Mathie, B. Alexander & G. Burnstock: NG-nitro-L-arginine methyl ester attenuates vasodilator responses to acetylcholine but enhances those to sodium nitroprusside. *J. Pharm. Pharmacol.* 1991, **43**, 871–874.
- Regitz-Zagrosek, V., W. Auch-Schwelk, M. Neuss & E. Fleck: Regulation of the angiotensin receptor subtypes in cell cultures, animal models and human diseases. *Eur. Heart J.* 1994, **15**, Suppl D, 92–97.
- Saida, A., H. Ito, T. Shibuya & Y. Watanabe: Time-course alterations of monoamine levels and cerebral blood flow in brain regions after subarachnoid hemorrhage in rats. *Brain Res. Bull.* 1997, **43**, 69–80.
- Sliva, D., D. English, D. Lyons & F. P. Lloyd: Protein kinase C induces motility of breast cancers by upregulating secretion of urokinase-type plasminogen activator through activation of AP-1 and NF-kappaB. *Biochem. Biophys. Res. Commun.* 2002, **290**, 552–557.
- Stenman, E., M. Malmjö, E. Uddman, G. Gido, T. Wieloch & L. Edvinsson: Cerebral ischemia upregulates vascular endothelin ET(B) receptors in rat. *Stroke* 2002, **33**, 2311–2316.
- Stenman, E. & L. Edvinsson: Cerebral ischemia enhances vascular angiotensin AT₁ receptor-mediated contraction in rats. *Stroke* 2004, **35**, 970–974.
- Strohm, C., M. Barancik, M. von Bruehl, M. Strniskova, C. Ullmann, R. Zimmermann & W. Schaper: Transcription inhibitor actinomycin-D abolishes the cardioprotective effect of ischemic preconditioning. *Cardiovasc. Res.* 2002, **55**, 602–618.
- Touyz, R. M. & E. L. Schiffrin: Signal transduction mechanisms mediating the physiological and pathophysiological actions of angiotensin II in vascular smooth muscle cells. *Pharmacol. Rev.* 2000, **52**, 639–672.
- Trachsel, S. & H. U. Keller: Selective responses (actin polymerization, shape changes, locomotion, pinocytosis) to the PKC inhibitor Ro 31-8220 suggest that PKC discriminately regulates functions of human blood lymphocytes. *J. Leukoc. Biol.* 1995, **57**, 587–591.
- Tsutsumi, Y., H. Matsubara, N. Ohkubo, Y. Mori, Y. Nozawa, S. Murasawa, K. Kijima, K. Maruyama, H. Masaki, Y. Moriguchi,

- Y. Shibasaki, H. Kamihata, M. Inada & T. Iwasaka: Angiotensin II type 2 receptor is up-regulated in human heart with interstitial fibrosis, and cardiac fibroblasts are the major cell type for its expression. *Circ. Res.* 1998, **83**, 1035–1046.
- Turla, M. B. & R. C. Webb: Vascular responsiveness to 5-hydroxytryptamine in experimental hypertension. In: *The peripheral actions of 5-hydroxytryptamine*. Ed.: J. R. Fozard. Oxford University Press, 1989, 327–353.
- Uddman, E., M. Adner & L. Edvinsson: Protein kinase C inhibitors decrease endothelin ET_B receptor mRNA expression and contraction during organ culture of rat mesenteric artery. *Eur. J. Pharmacol.* 2002, **452**, 215–222.
- Uddman, E., M. Henriksson, K. Eskesen & L. Edvinsson: Role of mitogen-activated protein kinases in endothelin ET_B receptor up-regulation after organ culture of rat mesenteric artery. *Eur. J. Pharmacol.* 2003, **482**, 39–47.
- Wackenfors, A., M. Emilson, R. Ingemansson, T. Hortobagyi, D. Szok, J. Tajti, L. Vecsei, L. Edvinsson & M. Malmjö: Ischemic heart disease induce upregulation of endothelin receptor mRNA in human coronary arteries. *Eur. J. Pharmacol.* 2004, **484**, 103–109.
- Walther, T., L. Olah, C. Harms, B. Maul, M. Bader, H. Hortnagl, H. P. Schultheiss & G. Mies: Ischemic injury in experimental stroke depends on angiotensin II. *FASEB. J.* 2002, **16**, 169–176.
- Watts, S. W.: Activation of the mitogen-activated protein kinase pathway via the 5-HT_{2A} receptor. *Ann. N. Y. Acad. Sci.* 1998, **861**, 162–168.
- Watts, S. W.: Serotonin-induced contraction in mesenteric resistance arteries: signaling and changes in deoxycorticosterone acetate-salt hypertension. *Hypertension* 2002, **39**, 825–829.
- Xu, K. & T. J. Murphy: Reconstitution of angiotensin receptor mRNA down-regulation in vascular smooth muscle. Post-transcriptional control by protein kinase A but not mitogenic signaling directed by the 5'-untranslated region. *J. Biol. Chem.* 2000, **275**, 7604–7611.
- Yildiz, O., J. R. Smith & R. E. Purdy: Serotonin and vasoconstrictor synergism. *Life Sci.* 1998, **62**, 1723–1732.
- Ytterberg, H. & L. Edvinsson: Evidence for a cyclic AMP-dependent pathway in angiotensin AT₁-receptor activation of human omental arteries. *JRAAS* 2001, **2**, 42–47.
- Zemin, C., D. Rachael, W. Leonard, C. David & E. C. Mark: Role of angiotensin receptor subtypes in mesenteric vascular proliferation and hypertrophy. *Hypertension* 1999, **34**, 408–414.
- Zhu, Y. Z., G. N. Chimon, Y. C. Zhu, Q. Lu, B. Li, H. Z. Hu, E. H. Yap, H. S. Lee & P. T. Wong: Expression of angiotensin II AT₂ receptor in the acute phase of stroke in rats. *Neuroreport* 2000, **11**, 1191–1194.