LUND UNIVERSITY

Female gender increases stiffness of elastic but not of muscular arteries in type I diabetic patients.

Rydén Ahlgren, Åsa; Sundkvist, Göran; Sandgren, T; Länne, T

Published in:
Clinical Physiology and Functional Imaging

DOI:
[10.1046/j.1475-097X.2002.00451.x](https://doi.org/10.1046/j.1475-097X.2002.00451.x)

2002

[Link to publication](#)

Citation for published version (APA):
Rydén Ahlgren, Å., Sundkvist, G., Sandgren, T., & Länne, T. (2002). Female gender increases stiffness of elastic but not of muscular arteries in type I diabetic patients. *Clinical Physiology and Functional Imaging*, 22(6), 409-415. <https://doi.org/10.1046/j.1475-097X.2002.00451.x>

Total number of authors:
4

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Female gender increases stiffness of elastic but not of muscular arteries in type I diabetic patients

Å. Rydén Ahlgren¹, G. Sundkvist², T. Sandgren³ and T. Länne⁴

Departments of ¹Clinical Physiology, ²Endocrinology, and ³Surgery, Lund University, Malmö University Hospital, Malmö, Sweden, and ⁴Division of Vascular Surgery, Linköping University, Jönköping Hospital, Jönköping, Sweden

Summary

Correspondence

Dr Åsa Rydén Ahlgren,
Department of Clinical Physiology, Lund University,
Malmö University Hospital, S-205 02 Malmö,
Sweden
E-mail: asa.ryden-ahlgren@klinfys.mas.lu.se

Accepted for publication

Received 8 April 2002;
accepted 2 September 2002

Key words

arterial stiffness; distensibility; femoral artery;
gender difference; type 1 diabetes; ultrasound

The reason for the particularly increased risk for cardiovascular complications in diabetic women is still unclear. We have previously found decreased distensibility of elastic arteries in type I diabetic women, indicating increased cardiac load, not seen in type I diabetic men, which might be one contributing factor. Whether the effect of gender is different in muscular arteries in type I diabetic patients has not been assessed. As estimates of arterial distensibility we measured stiffness (β) and pressure strain elastic modulus (E_p) in the muscular common femoral artery using echo-tracking sonography in 30 women (mean age 34 years, range 20–61) and 26 men (mean age 38 years, range 22–56) with type I diabetes. The results were compared with those of 89 healthy individuals of corresponding age and gender and with previously published results from elastic arteries in these patients obtained at the same occasion. The internal common femoral diameter was significantly decreased in both diabetic men and women. In sharp contrast to the highly significant decreased distensibility of the elastic abdominal aorta and common carotid artery in the type I diabetic women, the distensibility of the common femoral artery did not clearly differ between patients and controls, neither for women nor for men. Thus, the gender difference in changes of arterial distensibility found in elastic arteries was absent or far less obvious in the femoral artery. In conclusion, female gender seems to affect the mechanical properties of elastic, but not of large muscular arteries in type I diabetic patients. Thus, putative gender differences in arterial changes in type I diabetes are to be sought in elastic rather than muscular arteries.

Introduction

The risk for cardiovascular complications is particularly increased in women with diabetes (Kannel & McGee, 1979; Barret-Conner & Wingard, 1983; Lerner & Kannel, 1986). In addition, diabetic women have consistently been observed to have a worse prognosis than diabetic men after myocardial infarction (Garcia et al., 1974; Kannel, 1985; Tofler et al., 1987; Stone et al., 1989). Further, the relative risk for stroke is increased in diabetic women compared with diabetic men (Bell, 1994). The reason for this association between cardiovascular and cerebrovascular complications and female gender in diabetic patients is still unclear and inadequately explained by classical risk factors. Alterations of the mechanical properties of large arteries are increasingly recognized as a major factor for cardiovascular morbidity and mortality. Studies of the mechanical properties of arteries in type I diabetes have not been conclusive; both decreased (Pillsbury et al., 1962; Woolam et al., 1962; Christensen and Neubauer 1987; Oxlund et al., 1989;

Airaksinen et al., 1993; Lambert et al., 1998; Giannattasio et al., 1999), increased (Lehmann et al., 1992) and unchanged (Kool et al., 1995; Zenere et al., 1995) distensibility have been reported. The discrepancy may partly be due to the fact that gender differences have rarely been considered. We have previously reported gender-related changes with decreased distensibility of central elastic arteries in type I diabetic women, indicating increased cardiac load, not seen in diabetic men (Rydén Ahlgren et al., 1995). In healthy subjects the mechanical properties of elastic arteries are age- (Kawasaki et al., 1987; Länne et al., 1992a) and gender-dependent (Sonesson et al., 1993, 1994; Hansen et al., 1995). In contrast, muscular arteries, like the common femoral artery, are not clearly affected by either age (Kawasaki et al., 1987; Benetos et al., 1993; Rydén Ahlgren et al., 2001) or gender (Rydén Ahlgren et al., 2001). There is also a possibility that diabetes has different effects on muscular and elastic arteries. The aim of this study was to evaluate the mechanical properties of the common femoral artery in type I diabetic patients to clarify whether the gender-related

differences found in central elastic arteries are also present in this muscular artery.

Materials and methods

Patients and control subjects

Thirty women (mean age 34 years, range 20–61) and 26 men (mean age 38, range 22–56) with type I diabetes mellitus were investigated. Data regarding the mechanical properties of the abdominal aorta and the common carotid artery used for comparison have previously been published (Rydén Ahlgren et al., 1995, 1999). All subjects are followed-up regularly at the Department of Endocrinology at Malmö University Hospital, Sweden. The mean duration of diabetes at the time of this study was 19 years for both women and men (women: range 9–36 years, men: 9–38 years). None of the patients had a history of acute myocardial infarction, cerebrovascular events, or intermittent claudication. Moreover, none of the patients had angina pectoris or were being treated for coronary artery disease. Clinical details of the patients are given in Table 1; besides gender there were no significant differences in clinical features between women and men. One woman and four men showed clinical nephropathy (Table 1); two women were treated for hypertension with furosemide and metoprolol, respectively. Smoking was reported in 12 women and eight men. All patients gave informed consent according to the Helsinki declaration. To obtain age- and gender-matched reference values for common femoral artery diameter and distensibility indices, a control group of 158 healthy subjects (80 women: age range 7–81 years; 78 men: age range 8–78

years), was used (Rydén Ahlgren et al., 2001). Of these, 42 men and 47 women were in the age range 20–63 years, i.e. in the age range of the diabetic patients and these 89 subjects were used as controls in this study. None of the control subjects reported previous cardiopulmonary disease, intermittent claudication, diabetes or smoking and all were free from current medication. The ratio between ankle and brachial arterial pressure was >1 in all subjects indicating absence of obliterative atherosclerotic lesions in the arteries to the lower limb. The studies of normal subjects and diabetic patients were partly overlapping in time.

Ultrasonic measurements of arterial diameter and distensibility

The pulsatile changes in vessel diameter were registered with an ultrasound echo-tracking system (Diamove, Teltec AB, Lund, Sweden) (Lindström et al., 1987), interfaced with a 3.5 and a 5-MHz B-mode real time linear scanner (EUB 240, Hitachi, Tokyo, Japan), capable of detecting vessel wall movements of less than 10 µm and with a time resolution of 1.2 ms (Lindström et al., 1987; Benthin et al., 1991). In combination with blood pressure measurements the pulsatile diameter changes form the basis for calculation of vessel wall distensibility. Details of the study technique have previously been described (Länne et al., 1992b; Hansen et al., 1993; Sonesson et al., 1993).

All measurements were performed in a quiet room with the subject in the supine position after at least 15-min rest. An experienced ultrasound technician performed the examinations. The right common femoral artery was examined with the

Table 1 Demographical and clinical data in diabetic patients. All data, except frequencies, are presented as mean ± SD.

	Diabetic women	Diabetic men	Control women	Control men
Number	30	26	47	42
Duration (years)	19 ± 7	19 ± 8		
Height (cm)	165 ± 6	179 ± 7	166 ± 6	179 ± 5
Weight (kg)	64 ± 9	76 ± 7	67 ± 10	81 ± 10
Body surface area (m ²)	1.7 ± 0.1	1.9 ± 0.1	1.7 ± 0.1	2.0 ± 0.1
Body mass index (kg m ⁻²)	23.4 ± 2.5	23.7 ± 2.2	24 ± 4.4	25.4 ± 3.0
Albuminuria >0.5 g per 24 h	1	4		
Microalbuminuria 30–300 mg per 24 h	5	3		
Retinopathy				
Background	11	8		
More severe than background	3	4		
HbA1c (%)	8.56 ± 1.76	7.85 ± 1.41		
Systolic blood pressure (mmHg)	121 ± 12	127 ± 14	117 ± 13	125 ± 11
Diastolic blood pressure (mmHg)	76 ± 6	77 ± 6	74 ± 8	79 ± 8
Mean arterial pressure (mmHg)	91 ± 7	94 ± 7	89 ± 9	94 ± 8
Mean common femoral diameter (mm)	7.2 ± 0.65*	8.8 ± 1.0**	8.2 ± 0.8	9.7 ± 1.0
Common femoral artery strain (%)	6.7 ± 3.5	6.0 ± 3.2	6.1 ± 2.4	5.6 ± 2.1
Common femoral artery stiffness	9.2 ± 4.8	10.7 ± 5.8	8.6 ± 3.2	9.8 ± 4.7
Common femoral artery Ep (10 ⁵ N m ⁻²)	1.2 ± 0.7***	1.4 ± 0.8	1.1 ± 0.4	1.3 ± 0.6

*P < 0.00001 versus control women, **P = 0.002 versus control men, ***P = 0.04 versus control women.

5 MHz transducer and measurements were performed at the inguinal fossa with the hip joint as a landmark. The artery was visualized in the longitudinal section. Care was taken to minimize the pressure of the transducer to the skin. From each registration a sequence of at least five representative consecutive diameter cycles was manually chosen and the femoral artery diameters and the diameter changes calculated as mean of the selected diameter cycles. Blood pressure was measured by the auscultatory method using a sphygmomanometer on the upper arm immediately after measurement of the pulsatile common femoral artery diameter changes. Mean arterial blood pressure (MAP) was taken as the diastolic pressure plus one-third of the pulse pressure. Each vessel was examined three times with calculation of mean diameter, strain, stiffness (β), and the pressure strain elastic modulus (Ep) from the corresponding diameters, pulsatile diameter changes and blood pressures. The average of the values for each subject was then calculated.

The arterial strain, or the fractional diameter change, was defined as:

$$\text{Strain} = \frac{D_{\text{systolic}} - D_{\text{diastolic}}}{D_{\text{diastolic}}}$$

where D_{systolic} and $D_{\text{diastolic}}$ are the end-systolic and end-diastolic diameters.

The mean vessel diameter was defined as $(D_{\text{max}} + D_{\text{min}})/2$.

Distensibility characteristics of large arteries depend on the extent to which they are expanded, the vessels being very distensible at low pressures and small diameters, but gradually stiffer with increasing pressure and diameter. Thus, there is a non-linear pressure–diameter relationship of the arterial wall. It was observed that the relation between the logarithm of relative pressure and strain (fractional diameter change) was linear *in vitro*. Based on this logarithmic relation, and thus independent of pressure in the physiological range, the index stiffness (β) was established as an estimate of vascular distensibility. It was later modified and used *in vivo* by Kawasaki et al. (1987). This index has been used in this study and is expressed as:

$$\text{Stiffness } (\beta) = \frac{\ln(P_{\text{systolic}}/P_{\text{diastolic}})}{(D_{\text{systolic}} - D_{\text{diastolic}})/D_{\text{diastolic}}}$$

where P_{systolic} and $P_{\text{diastolic}}$ are the systolic and diastolic blood pressure levels in mmHg.

We also calculated Ep, defined according to Peterson et al. (1960) as:

$$\text{Ep} = K \times \frac{P_{\text{systolic}} - P_{\text{diastolic}}}{(D_{\text{systolic}} - D_{\text{diastolic}})/D_{\text{diastolic}}}$$

Ep is measured in N m^{-2} . $K = 133.3$ is the factor for converting mmHg to N m^{-2} .

Using this technique the intraobserver variability for repeated measurements have previously, at this laboratory, been shown to be 15% for the pulsatile diameter change, 5% for the mean diameter and 18% for stiffness (β) and Ep for the vessel investigated (Hansen et al., 1993).

Statistics

Unpaired t-test or analyses of covariance were used to compare the vessel diameters, strain, stiffness and Ep in diabetic men and women with control subjects in corresponding age ranges. The effects of age, duration and HbA1c on the studied parameters were tested by correlation analysis (Pearson or Spearman). Differences between smoking and non-smoking patients and differences in clinical characteristics (HbA1c, blood pressure) were statistically tested with Mann–Whitney U-test. Multiple regression analysis, correcting for age, were used to sought correlations between the arterial indices and blood pressure. Statistical significance was defined as $P < 0.05$. Data are presented as the mean value $\pm$ SD, unless otherwise stated.

Results

Blood pressures in the diabetic patients did not differ significantly from those in the control group of corresponding age, either for women or for men (Table 1). There was no significant difference in HbA1c levels between diabetic men and women (Table 1). Body mass index (BMI) was slightly less in diabetic patients than in controls (Table 1).

The common femoral artery diameter

The internal common femoral artery mean diameter (Table 1) was significantly decreased in both diabetic men and diabetic women compared with the corresponding control subjects (men: $P = 0.002$, women: $P < 0.00001$). The difference in diameter between diabetic patients and controls could not be explained by differences in body surface area. No increase in diameter with ageing could be detected in either diabetic men or diabetic women. No significant correlation between the common femoral artery diameter and the duration of diabetes or the HbA1c levels, respectively, could be detected.

The common femoral artery strain

There were no significant differences in strain of the common femoral artery between men with type I diabetes and control men ($P = 0.5$) or between women with type I diabetes and control women ($P = 0.3$). Further, no significant correlation between strain and age was seen (diabetic men $r = -0.21$, $P = 0.3$; control men $r = -0.14$, $P = 0.4$, diabetic women $r_s = -0.32$, $P = 0.09$; control women, $r = 0.18$, $P = 0.2$).

The common femoral artery stiffness (β)

Figure 1a shows stiffness (β) of the common femoral artery in men with type I diabetes and in control men. There was no significant difference in stiffness between diabetic men and control men ($P = 0.5$) and no correlation between stiffness and age was seen (diabetics $r = 0.08$, $P = 0.7$; controls $r = 0.15$, $P = 0.4$).

Figure 1 (a) Stiffness (β) of the common femoral artery related to age in men with type I diabetes (black circles) and control men (white circles). There is no significant difference in stiffness between men with diabetes and control men ($P = 0.5$) and no correlation with age is seen. (b) Stiffness (β) of the common femoral artery related to age in women with type I diabetes (black circles, solid line) and control women (white circles, dotted line). There is no significant difference in the level of stiffness values between women with diabetes and control women ($P = 0.1$). In diabetic women a correlation between age and stiffness is found, which can not be detected in control women. The difference in slope between diabetic women and control women reach significance ($P = 0.048$).

Figure 1b shows stiffness (β) of the common femoral artery in women with type I diabetes and in control women. In diabetic women a significant positive correlation between age and stiffness was seen ($r_s = 0.45$, $P = 0.01$), not detected in control women ($r = 0.11$, $P = 0.5$). In agreement, the difference in slope between diabetic women and control women reached significance ($P = 0.048$). On the other hand no significant difference in the level of stiffness between diabetic women and control women could be detected ($P = 0.1$).

The common femoral artery pressure strain elastic modulus (Ep)

As for stiffness (β), there was no significant difference in Ep between men with type I diabetes and control men ($P = 0.4$) and similarly there was no correlation between Ep and age (diabetic men $r = 0.11$, $P = 0.6$; control men $r = 0.17$, $P = 0.3$).

As for stiffness (β), in diabetic women there was a correlation between age and Ep ($r_s = 0.48$, $P = 0.007$), not seen in control women ($r = 0.20$, $P = 0.2$). In agreement, the difference in slope between diabetic women and control women reached significance ($P = 0.046$) and, in fact, the level of Ep tended to be increased in diabetic women, although the difference was low ($P = 0.04$).

The arterial indices and clinical characteristics

The strain, stiffness and Ep values in the few patients with nephropathy and/or proliferative retinopathy, did not differ from those without complications and our results were the same if these patients were excluded. No significant correlation between common femoral artery strain, stiffness (β) or Ep and the duration of diabetes could be detected in either men or women. Further, no correlation between strain, stiffness or Ep and HbA1c could be detected. Using multiple regression analyses, correcting for age, no correlation between the arterial indices and blood pressure, analysed as MAP, systolic and diastolic pressure, was seen. No significant differences in strain, stiffness or Ep between smoking and non-smoking patients for either women or men could be detected.

Comparison between the common femoral artery and the elastic arteries

Figure 2 shows schematically common femoral artery stiffness (β) data in diabetic women and men in comparison with previously published results from the abdominal aorta and the common carotid artery in the same patients examined at the same occasion (Rydén Ahlgren et al., 1995). The clear increase in stiffness (β) in the abdominal aorta (mean 64%) and the common carotid artery (mean 22%) in diabetic women is in sharp contrast to the lack of increase in stiffness (β) in the common femoral artery in the same diabetic women.

Figure 2 Schematic illustration of the percentage increase in stiffness (β) in men ($n = 26$, white bars) and women ($n = 30$, black bars) with type I diabetes compared with predicted mean normal values according to age and gender in the abdominal aorta (AO), the common carotid artery (CCA) and the common femoral artery (CFA). The original data from the abdominal aorta and the common carotid artery have been published (Rydén Ahlgren et al., 1995).

Discussion

Alterations of the mechanical properties of large arteries are increasingly recognized as an important factor for cardiovascular morbidity and mortality. In a previous study of patients with type I diabetes (Rydén Ahlgren et al., 1995) we reported gender-related changes with decreased distensibility of the elastic abdominal aorta and the elastic common carotid artery in diabetic women compared with control women, whereas in diabetic men no difference compared with control men was seen, thus indicating increased cardiac load in diabetic women. We have now evaluated the muscular common femoral artery in the same patients, examined at the same occasion as the elastic arteries, i.e. with the same duration and metabolic control, to clarify whether the gender-related differences in changes of arterial distensibility are also present in the muscular common femoral artery.

In agreement with our findings from the abdominal aorta in the same patients (Rydén Ahlgren et al., 1995) we found the common femoral artery mean internal diameter to be decreased in both diabetic men and women. This is in agreement with a previous study (Christensen & Neubauer, 1988), but in contrast to Kool et al. (1995), who did not find a decreased common femoral artery diameter in type I diabetic patients. The patient material in the latter study was, however, small and the duration of diabetes was shorter. Our finding of a decreased inner vessel diameter suggests a thicker vessel wall in the diabetic patients than in the control subjects. Intima-media thickness has been shown to be increased in several studies of arteries in diabetic patients (Yamasaki et al., 1994; Frost & Beischer, 1998; Giannattasio et al., 1999). Another possible explanation for found decreased diameter might be an effect of reduced vasodilatory function in type I diabetic patients, which has been described for the common femoral artery (Zenere et al., 1995).

In sharp contrast to the findings from the abdominal aorta, where a 60–70% increase in stiffness (β) was noted in diabetic women, no clear-cut differences in distensibility of the common femoral artery were seen in either diabetic women or men compared with control subjects. Accordingly, the pronounced gender-related differences in changes of arterial distensibility, found in the central elastic arteries in these patients, are far less obvious or absent in the common femoral artery, indicating that the diabetic state has different influence on elastic and muscular arteries in type I diabetic women, probably because of differences in histological and histochemical structure. The common femoral artery is a predominantly muscular artery with large content of smooth muscles, while the abdominal aorta and the common carotid are predominantly elastic arteries with a large content of elastin and collagen and only sparse smooth muscle cells.

Studies of the mechanical properties of the common femoral artery in type I diabetic patients are sparse and putative gender differences have not been evaluated. Kool et al. (1995), in a study of 30 type I diabetic patients, found the distensibility of

the femoral artery to be decreased. Zenere et al. (1995), however, in a smaller number of patients, including those with microalbuminuria, did not find changes in the mechanical properties of the femoral artery, thus supporting our findings.

Decreased distensibility of central arteries increases central pulse pressure and left ventricular load (Nichols & O'Rourke, 1998). Thus, if further confirmed, our finding of a gender difference with more decreased distensibility in central arteries in diabetic women than in diabetic men, is likely to be an important factor for the particularly increased risk for cardiovascular complications in diabetic women, including their poor prognosis after myocardial infarction with increased incidence of congestive heart failure, reinfarction and death (Garcia et al., 1974; Kannel, 1985; Tofler et al., 1987). Studies regarding putative gender-related differences in arterial changes in diabetic patients are few. Our findings of gender-related differences in changes of arterial distensibility in elastic arteries in type I diabetic patients have, however, support from Hu et al. (1997) in a study of adolescents with type I diabetes. Further, recently, Lambert et al. (1998) reported female gender to be a determinant of decreased distensibility of the common carotid artery in type I diabetic patients. Indeed, gender-related differences have also been reported in type II diabetic patients with findings of increased aorto-femoral pulse wave velocity in women, but not in men (Lehmann, 1996). These studies are all based on findings from predominantly elastic arteries, thus supporting our initial finding. The present study with no clear cut gender-related changes in the common femoral artery, despite clear gender-related changes in the central elastic arteries, thus shows that gender-related changes in type I diabetic patients can be present in one type of arteries, but not in others. Further, our study shows that putative major gender-related differences in arterial changes in type I diabetes are to be sought in central arteries rather than in large muscular arteries. As a factor for cardiac load the elastic arteries are also those of the most importance.

Measures of arterial distensibility or stiffness have been proposed as surrogate markers for atherosclerosis (Arnett et al., 1994; Lehmann et al., 1996), at least in central arteries. At present the putative relation between early atherosclerosis and changes in the mechanical properties of the arterial wall are, however, far from clear (Nichols & O'Rourke, 1998). In this study we found the common femoral inner diameter to be decreased in both diabetic men and women, which might be due to early atherosclerosis. Thus, our finding that the mechanical properties of the common femoral artery, a vessel prone to atherosclerotic disease, are not clearly affected in patients with long-term type I diabetes, may seem surprising. It is well known that the structure of arterial walls change with ageing with an increase in collagen and thickening of the vessel wall and diabetic patients have in several studies been shown to have increased intima-media thickness (Yamasaki et al., 1994; Frost & Beischer, 1998; Giannattasio et al., 1999), a factor likely to increase vessel wall stiffness. This leads to the hypothesis that the smooth muscle within the common femoral artery has a

larger impact on vessel wall mechanics than age- and putative diabetes-related histological changes *per se*. Further, it gives support to the opinion that there is no clear relation between atherosclerosis and vessel wall motion. One must, however, keep in mind that in this study all patients investigated were without clinical signs of vascular disease and the vast majority of the patients were free of diabetic late complications. Further, although the relation between femoral artery distensibility and atherosclerosis is not clear, it might still be possible that our finding of a decrease in femoral artery distensibility with age in diabetic women, but not in men also indicates an increased vulnerability related to gender in the common femoral artery.

In several studies the arterial wall properties of central elastic arteries in healthy, non-diabetic subjects have been shown to differ between men and women. This suggests that these differences may result from hormonal influence. In contrast, in the common femoral artery, a large muscular artery, gender- and age-related changes, seem to be sparse or lacking (Rydén Ahlgren *et al.*, 2001). Thus, our finding that the normal gender-related differences in arterial distensibility of the common carotid artery and the abdominal aorta are absent in patients with type I diabetes, whereas no clear cut differences in changes of the mechanical properties of the common femoral artery are seen, favours the assumption that the sex hormonal balance is affected in type I diabetic women. Indeed, women with type I diabetes have recently been shown to have a high prevalence of hyperandrogenic disorders (Escobar-Morreale *et al.*, 2000). Future studies will show whether an altered sex hormone balance contributes to the alterations in arterial distensibility seen in type I diabetic women.

In attempting to measure indices of arterial distensibility the arterial diameter change and the intra-arterial blood pressure ideally should be measured simultaneously at the same location, as it is well known that the arterial pressure waves undergo transformation in the arterial tree. Although widely used, the method of using the auscultatory blood pressure in the upper arm, when calculating distensibility indices for, e.g. the femoral artery, may be questioned. Comparison between intra-arterial pressure in the common femoral artery and the brachial blood pressure obtained by the auscultatory method in healthy subjects has, however, shown that the pulse pressure is systematically underestimated when the auscultatory method is used without obvious differences between gender or between young and elderly (Rydén Ahlgren *et al.*, 2001). Hence, our use of auscultatory blood pressure cannot explain our findings.

In conclusion, this study shows that the internal common femoral diameter significantly decreases in both type I diabetic men and women without clinical signs of vascular disease. Further, it shows that female gender has a major effect on the mechanical properties of elastic, but not on muscular arteries in type I diabetic patients. Accordingly, the obvious gender-related differences found in elastic arteries with decreased arterial distensibility in women, but not in men, are far less evident or absent in the muscular common femoral artery. This indicates that

the metabolic state has different effects on elastic and muscular arteries in diabetic women. Further, it suggests that putative gender-related differences in arterial changes in type I diabetes are to be sought in central elastic arteries rather than in muscular arteries, which are also of most importance for cardiac load.

Acknowledgements

This study was supported by the Swedish Medical Research Council (no. 12661 and no. 7507), the Swedish Diabetes Association, The Lundström Foundation, the Novo-Nordic Foundation, the Swedish Heart-Lung Foundation and the Medical Faculty, Lund University, Lund.

References

- Airaksinen KEJ, Salmela PI, Linnaluoto MK, Ikäheimo MJ, Ahola K, Ryhänen LJ. Diminished arterial elasticity in diabetes: association with fluorescent advanced glycosylation end products in collagen. *Cardiovasc Res* (1993); **27**: 942–945.
- Arnett DK, Evans GW, Riley WA. Arterial stiffness: a new cardiovascular risk factor? *Am J Epidemiol* (1994); **140**: 669–682.
- Barret-Conner E, Wingard DL. Sex differential in ischemic heart disease mortality in diabetics: A prospective population based study. *Am J Epidemiol* (1983); **118**: 489–496.
- Bell D. Stroke in the diabetic patient. *Diabetes Care* (1994); **17**: 213–219.
- Benetos A, Laurent S, Hoeks AP, Boutouyrie PH, Safar ME. Arterial alterations with aging and high blood pressure. A non-invasive study of carotid and femoral arteries. *Arteriosclerosis and Thrombosis* (1993); **13**: 90–97.
- Benthin M, Dahl P, Ruzicka R, Lindström K. Calculation of pulse wave velocity using cross correlations-effects of reflexes in the arterial tree. *Ultrasound Med Biol* (1991); **17**: 461–469.
- Christensen T, Neubauer B. Arterial wall stiffness in insulin-dependent diabetes mellitus. *Acta Radiol* (1987); **28**: 207–209.
- Christensen T, Neubauer B. Internal diameter of the common femoral artery in patients with insulin-dependent diabetes mellitus. *Acta Radiol* (1988); **29**: 423–425.
- Escobar-Morreale HF, Roldan B, Barrio R *et al.* High prevalence of the polycystic ovary syndrome and hirsutism in women with type I diabetes mellitus. *J Clin Endocrinol Metab* (2000); **85**: 4182–4187.
- Frost D, Beischer W. Determinants of carotid artery wall thickening in young patients with type 1 diabetes mellitus. *Diabetic Medicine* (1998); **15**: 851–857.
- Garcia MJ, McNamara PM, Gordon T, Kannel WB. Morbidity and mortality in diabetics in Framingham population: sixteen year follow-up study. *Diabetes* (1974); **23**: 105–111.
- Giannattasio C, Failla M, Piperno A *et al.* Early impairment of large artery structure and function in Type I diabetes mellitus. *Diabetologia* (1999); **42**: 987–994.
- Hansen F, Bergqvist D, Mangell P, Rydén Å, Sonesson B, Länne T. Non-invasive measurement of pulsatile vessel diameter changes and elastic properties in human arteries: a methodological study. *Clin Physiol* (1993); **13**: 631–643.
- Hansen F, Mangell P, Sonesson B, Länne T. Diameter and compliance in the human common carotid artery – variations with age and sex. *Ultrasound Med Biol* (1995); **21**: 1–9.
- Hu J, Norman M, Wallenstein M, Gennser G. Dynamic properties of the aorta and of the foot microcirculation in adolescents with diabetes mellitus. *Acta Paediatr* (1997); **86**: 620–625.

- Kannel W. Lipids, diabetes, and coronary heart disease: insights from the Framingham study. *Am Heart J* (1985); **110**: 1110–1117.
- Kannel WB, McGee DL. Diabetes and cardiovascular disease: the Framingham study. *JAMA* (1979); **241**: 2035–2038.
- Kawasaki T, Sasayama S, Yagi S-I, Asakawa T, Hirai T. Non-invasive assessment of the age related changes in stiffness of major branches of the human arteries. *Cardiovasc Res* (1987); **21**: 678–687.
- Kool MJ, Hoeks AP, Lambert J, Struijker Boudier HA, Stehouwer CD, Van Bortel LM. Vessel wall properties of large arteries in uncomplicated IDDM. *Diabetes care* (1995); **18**: 618–624.
- Lambert J, Smulders RA, Aarsen M, Donker JM, Stehouwer CD. Carotid artery stiffness is increased in microalbuminuric IDDM patients. *Diabetes Care* (1998); **21**: 99–103.
- Länne T, Sonesson B, Bergqvist D, Bengtsson H, Gustafsson D. Diameter and compliance in the male human abdominal aorta: influence of age and aortic aneurysm. *Eur J Vasc Surg* (1992a); **6**: 178–184.
- Länne T, Stale H, Bengtsson H et al. Noninvasive measurement of diameter changes in the distal abdominal aorta in man. *Ultrasound Med Biol* (1992b); **18**: 451–457.
- Lehmann ED. Pulse wave velocity as a marker of vascular disease. *Lancet* (1996); **348**: 744.
- Lehmann ED, Gosling RG, Sönksen PH. Arterial wall compliance in diabetes. *Diabet Med* (1992); **9**: 114–119.
- Lehmann ED, Hopkins KD, Gosling RG. Increased aortic stiffness in women with NIDDM. *Diabetologia* (1996) **39**: 870–871.
- Lerner DJ, Kannel WB. Patterns of coronary heart disease morbidity and mortality in the sexes: a 24-year follow-up of the Framingham population. *Am Heart J* (1986); **111**: 383–390.
- Lindström K, Gennser G, Sindberg Eriksson P, Benthin M, Dahl P. An improved echo-tracker for studies on pulse-waves in the fetal aorta. In: *Fetal Physiological Measurements* (ed. Rolfe, P.) (1987) pp. 217–226. Butterworths, London.
- Nichols WW, O'Rourke MF. (1998) *McDonald's Blood Flow in Arteries. Theoretical, Experimental and Clinical Principles* (eds. Nichols W. W., O'Rourke M. F.), (1998), 4th edn. Edward Arnold, London.
- Oxlund H, Rasmussen LM, Andreassen TT, Heickendorff L. Increased aortic stiffness in patients with type 1 (insulin-dependent) diabetes mellitus. *Diabetologia* (1989); **3**: 748–752.
- Peterson LH, Jensen RE, Parnell J. Mechanical properties of arteries in vivo. *Circ Res* (1960); **8**: 622–633.
- Pillsbury HC, Hung W, Kyle MC, Freis ED. Arterial pulse waves and systolic time intervals in diabetic children. *Am Heart J* (1962); **87**: 783–790.
- Rydén Ahlgren Å, Länne T, Wollmer P, Hansen F, Sundkvist G. Increased arterial stiffness in women, but not in men, with IDDM. *Diabetologia* (1995); **38**: 1082–1089.
- Rydén Ahlgren Å, Sundkvist G, Wollmer P, Sonesson B, Länne T. Increased aortic stiffness in women with type 1 diabetes mellitus is associated with diabetes duration and autonomic nerve function. *Diabetic Med* (1999); **16**: 291–297.
- Rydén Ahlgren Å, Åstrand H, Sandgren T, Vernerström E, Sonesson B, Länne T. Dynamic behavior of the common femoral artery – age and gender of minor importance. *Ultrasound Med Biol* (2001); **27**: 181–188.
- Sonesson B, Hansen F, Stahle H, Länne T. Compliance and diameter in the human abdominal aorta – the influence of age and sex. *Eur J Vasc Surg* (1993); (1993) **7**: 690–697.
- Sonesson B, Vernerström E, Hansen F, Länne T. Sex differences in the mechanical properties of the abdominal aorta in man. *J Vasc Surg* (1994); **20**: 959–969.
- Stone PH, Muller JE, Hartwell T. The effect of diabetes on prognosis and serial left ventricular function after acute myocardial infarction: contribution of both coronary artery disease and diastolic left ventricular dysfunction to the adverse prognosis. *J Am Coll Cardiol* (1989); **14**: 49–57.
- Toffer GH, Stone PH, Muller JE et al. Effects of gender and race on prognosis after myocardial infarction: adverse prognosis for women, particularly black women. *J Am Coll Cardiol* (1987); **9**: 473–482.
- Woolam GL, Schnur PL, Valbona C, Hoff HE. The pulse wave velocity as an early indicator of atherosclerosis in diabetic subjects. *Circulation* (1962); **25**: 533–539.
- Yamasaki Y, Kawamori R, Matsushima H et al. Atherosclerosis in carotid artery of young IDDM patients monitored by ultrasound high resolution B-mode imaging. *Diabetes* (1994); **43**: 634–639.
- Zenere BM, Rossi L, Arcaro G, Muggeo M, Saggiani F, Lechi A. of functional alterations of the arterial wall in IDDM patients with and without microalbuminuria. *Diabetes Care* (1995); **18**: 975–982.