LUND UNIVERSITY

Phosphatase activities of arbuscular mycorrhizal intraradical

van Aarle, Ingrid; Rouhier, Hervé; Saito, M

Published in:
Mycological Research

DOI:
[10.1017/S0953756202006470](https://doi.org/10.1017/S0953756202006470)

2002

[Link to publication](#)

Citation for published version (APA):

van Aarle, I., Rouhier, H., & Saito, M. (2002). Phosphatase activities of arbuscular mycorrhizal intraradical. *Mycological Research*, 106(10), 1224. <https://doi.org/10.1017/S0953756202006470>

Total number of authors:
3

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Phosphatase activities of arbuscular mycorrhizal intraradical and extraradical mycelium, and their relation to phosphorus availability

Ingrid M. van AARLE¹, Hervé ROUHIER² and Masanori SAITO³

¹Department of Microbial Ecology, Ecology Building, Lund University, SE-223 62 Lund, Sweden.

²Laboratory of Chemistry, Teikyo University, School of Medicine, 359 Ohtsuka, Hachioji, 192–0395, Japan.

³Laboratory of Soil Ecology, Department of Grassland Ecology, National Institute of Livestock and Grassland Science, 329–2793 Nishi-nasuno, Tochigi, Japan.

E-mail: Ingrid.van_Aarle@mbioekol.lu.se

Received 3 March 2002; accepted 29 July 2002.

We investigated the influence of changes in external phosphorus (P) concentration on the proportion of phosphatase-active structures of the arbuscular mycorrhizal fungus *Gigaspora margarita* associated with *Allium cepa*. The P treatment was started when mycorrhizal colonisation had been established, and plant systems were harvested three times after the start of the P treatment. Higher shoot dry weights and P contents were observed in the high-P treated plants at the last harvest. We did not find any change in the proportion of phosphatase-active extraradical mycelium following P treatment. However, the proportion of alkaline phosphatase-active mycelium was positively correlated for extraradical and intraradical mycelium. Also, the proportion of alkaline phosphatase-active arbuscules seemed to increase with the shoot fresh weight, whereas the proportion of acid phosphatase-active arbuscules decreased with higher shoot P concentration and dry weight. We have shown experimentally that the intraradical mycelium of *G. margarita*, but not the extraradical mycelium, responds metabolically to plant P concentration, and possibly also to external P availability.

INTRODUCTION

Decreased soil phosphorus (P) may lead to an increase in the colonisation of plants by arbuscular mycorrhizal (AM) fungi (Abbott, Robson & de Boer 1984, Thomson, Robson & Abbott 1991, Tawarayama *et al.* 1994). The AM fungal association often improves plant P uptake under P-limited conditions (Smith & Read 1997, Dickson, Smith & Smith 1999a, b). It has recently been shown that a phosphate transporter gene in the extraradical mycelium of *Glomus intraradices* is expressed at low phosphate concentration in the growth medium (Maldonado-Mendoza, Dewbre & Harrison 2001). When grown under P-limited conditions the production of phosphatase enzymes by many fungi increases (Grotelueschen *et al.* 1994, Kaffman *et al.* 1994, O'Neill *et al.* 1996), although such regulation has not been demonstrated for mycorrhizal fungi. The acid phosphatase (ACP) of intraradical AM fungal hyphae is located mainly in the vacuoles and sometimes in arbuscules (Gianinazzi, Gianinazzi-Pearson & Dexheimer 1979, Ezawa, Saito & Yoshida 1995, Saito 1995). It has been hypothesized that ACP could be involved in the hydrolysis of vacuolar polyphosphate after which inorganic

P could be released by the fungus into the plant–fungal interface (Ezawa, Smith & Smith 2001). The presence of ACP activity has also been demonstrated in extraradical mycelium (van Aarle, Olsson & Söderström 2001) and alkaline phosphatase (ALP) activity has been found in both intraradical (Gianinazzi *et al.* 1979, Smith & Gianinazzi-Pearson 1990, Tisserant *et al.* 1993) and extraradical AM fungal mycelium (Boddington & Dodd 1999, van Aarle *et al.* 2001). It is assumed that the ALP of the intraradical mycelium is involved in the transfer of P from the fungus to the plant (Tisserant *et al.* 1993), since a relationship has been found between the ALP of the intraradical mycelium and P transfer (Tisserant, Gianinazzi & Gianinazzi-Pearson 1996).

It has been suggested that the location of phosphatases may reflect the site for P transfer (Ezawa *et al.* 1995), but no experimental data on the relationship between external P concentration and the proportion of phosphatase-active extraradical and intraradical AM fungal mycelium are available. In the present work we studied the influence of external P availability on the relative amount of phosphatase-active structures of an AM fungus, in order to illustrate the P metabolism of an AM fungal symbiosis. We compared extraradical and

intraradical mycelium associated with plants established under P-limited conditions, and then exposed these plants either to increased P availability or to P depletion. We then examined whether the proportions of phosphatase-active mycelium and arbuscules were related to P availability or to fungal P transfer.

MATERIALS AND METHODS

Experimental set-up and harvesting

Seeds of *Allium cepa* (cv. 'Senshuchukou') were surface-sterilized and incubated for 12 d in the dark at 23 °C during the daytime and 20 ° during the night. 50 ml tubes (disposable polypropylene syringes without a plunger: Iwaki, Japan) were painted black to prevent light transmission and filled with 75 g of washed autoclaved river sand. One seedling was transplanted into each tube and, for the mycorrhizal treatment, 30 spores of *Gigaspora margarita* (MAFF 520054) were placed below the root at the time of transplantation. Spores were collected from a soil culture with *Sorghum bicolor* as host. To all tubes, 10 ml of a nutrient solution with 1 mg P l⁻¹ (low-P treatment) was added containing (mg l⁻¹): NH₄NO₃ (114.3), NaNO₃ (121.3), NaHPO₄·12H₂O (11.6), K₂SO₄ (133.7), CaCl₂ (221.5), MgSO₄·7H₂O (202.8), Fe-EDTA (15.1), MnCl₂·4H₂O (3.6), CuSO₄·5H₂O (0.039), Na₂MoO₄·2H₂O (0.0126), H₃BO₃ (2.29) and ZnCl₂ (0.417), and the pH was adjusted to approx. 6.5. The tubes were kept in a growth cabinet (23/20 °, 14 h photoperiod and a photosynthetic photon flux density of 465 µmol m⁻² s⁻¹) and watered twice a week with 10 ml of the nutrient solution.

Three mycorrhizal and three non-mycorrhizal plants were harvested 37 d after transplantation. The fresh weight of the plant and the dry weight of the shoot were recorded. For the mycorrhizal plants, the extraradical mycelium was collected from the sand. After staining with trypan blue (adapted from Phillips & Hayman 1970), root colonisation and arbuscule formation were assessed with the magnified intersections method (McGonigle *et al.* 1990).

The remaining mycorrhizal plants were randomly divided into two groups, each group receiving a different kind of P treatment. One group was watered twice a week with the nutrient solution containing no P (no-P treatment), while the other group was watered with the nutrient solution containing a five times increased P concentration (5 mg P l⁻¹; high-P treatment). At 42, 49 and 56 d after transplantation, 3 mycorrhizal plants from each P treatment were harvested. Two non-mycorrhizal plants were watered with the low-P solution and harvested 56 d after transplantation.

Collection of intraradical mycelium

Fresh root samples (cleaned from debris, extraradical mycelium and spores) were kept on ice, cut into pieces 5 mm in length and placed in 10 ml of a freshly prepared

filtered enzyme digestion solution, as described by Saito (1995). The samples were placed under vacuum for 8 min, covered with Parafilm, and then incubated at 30 ° for 2 h on a reciprocal shaker at 120 strokes min⁻¹. The digested root samples were collected on a 45 µm nylon mesh, washed clean with a washing buffer (Saito 1995) and kept on ice. The root cortex was opened under a dissecting microscope and removed with two pairs of fine forceps. The intraradical mycelium was carefully separated from the stele and transferred to fresh washing buffer.

Enzyme histochemical staining

The intraradical and extraradical mycelium was subjected to histochemical assessment of ACP and ALP activity. For ALP activity a 0.1 M Tris-HCl buffer (pH 8.5) with 1.1 mg ml⁻¹ α-naphthyl phosphate Na salt (Sigma) and 1.0 mg ml⁻¹ fast blue RR (Sigma) was used, while for ACP activity the Tris-HCl buffer was replaced by a 0.1 M sodium acetate buffer (pH 5.5). The samples were incubated for 1 h at 30 °, washed with deionised water, counterstained with acid fuchsin in lactic acid and stored in the stain (Saito, Stribley & Hepper 1993). The samples were mounted on microscope slides with lactoglycerol. Fast blue RR precipitation of the samples was assessed with a compound light microscope (Nikon Eclipse E800) with a bright field view at 400 times magnification. For each sample, approx. 200 hyphal intersections were assessed. Intersections of hyphae or arbuscules were classified as active or non-active, depending on whether they exhibited fast blue RR precipitation or not (see Fig. 1). The proportion (%) of ACP- and ALP-active hyphae and arbuscules was determined. Images were taken with a SPOT RT camera (Diagnostic Instruments, Burrough, MN).

P determination

Dried shoot material (10–50 mg) was digested (with H₂SO₄ and H₂O₂) at 200 ° according to the method of Mizuno & Minami (1980). Colorimetric determination of phosphate was carried out according to Watanabe & Olsen (1965) with a spectrophotometer by measuring the absorbance at 710 nm 15 min after mixing the samples with the chromophoric solution (ammonium molybdate, potassium antimony tartrate and ascorbic acid).

Statistical analysis

Means, standard errors and linear regression coefficients were calculated. To test for effects of P treatment (no-P and high-P) the data were subjected to a 1-tailed *t* test with unequal variances, for shoot dry weight and shoot P content, and a 2-tailed *t* test with unequal variances for the proportion of phosphatase-active structures. To compare the proportion of ACP-active structures with that of ALP-active structures, the data were subjected to a 2-tailed paired *t* test with unequal variances.

Fig. 1. Extraradical and intraradical mycelium of *Gigaspora margarita* collected from enzymatically digested roots of *Allium cepa*. The mycelium was stained with fast blue RR to visualise alkaline (ALP) and acid (ACP) phosphatase activity (dark precipitate) and counterstained with acid fuchsin (magenta). In (a) intraradical hyphae with low and an arbuscule with high ALP activity are shown; in (b) the intraradical hyphae show high and the arbuscule low ACP activity. Extraradical mycelium with some ACP activity is shown in (c). Bars = 20 μ m.

RESULTS

Growth and phosphatases

At the final harvest (56 d), the shoot dry weight of the high-P treated plants was significantly higher than that of the no-P treated plants (Table 1). High-P treated plants also had a higher shoot weight than the

non-mycorrhizal plants at 56 d. The P content of the shoot increased with time, and a greater ($P=0.056$) increase was observed for the high-P treated plants (Table 1).

Root colonisation and arbuscule formation by *Gigaspora margarita* were 33 and 24 %, respectively, at the start of the P treatment. After a further slight increase in colonisation, both root colonisation and arbuscule formation seemed to stabilise. Mean values for root colonisation were between 40 and 55 % of the total root length, while arbuscules were present in 29–49 % of the root length. No significant differences were found between the kind of P treatment or time of harvest.

The proportions of phosphatase-active extraradical and intraradical mycelium were estimated after staining with the fast blue RR substrate (Table 2). The staining of the intraradical mycelium was not affected by enzymatic digestion of the roots. The fast blue RR precipitation in arbuscules was mostly confined to the trunk hyphae. Arbuscules with fast blue RR precipitation in the fine arbuscular branches (Fig. 1a) were rarely seen.

Apart from the proportion of ACP-active arbuscules at 49 d, no difference was found in the proportions of phosphatase-active intraradical mycelium between the two kinds of P treatment. The proportions of both ACP- and ALP-active intraradical mycelium decreased with time. In the intraradical mycelium, the proportion of ACP-active mycelium was significantly higher than the ALP-active proportion (Table 2), and the hyphae were more intensely stained (compare hyphae in Figs 1a and 1b).

No difference could be seen in the proportion of phosphatase-active extraradical mycelium, between the two kinds of P treatment. Extraradical mycelium did not show strong fast blue RR precipitation (Fig. 1c), and only a small proportion of the mycelium showed any precipitation. In contrast to the intraradical mycelium, the proportion of ALP-active mycelium was significantly higher than the ACP-active proportion (Table 2).

The proportion of ALP-active extraradical mycelium was correlated with the proportion of ALP-active intraradical mycelium (Fig. 2). This correlation was positive for both kinds of P treatment (significant correlations), and was higher for the no-P than for the high-P treated plants. Due to the low proportion of ACP-active extraradical mycelium, no correlation with ACP in the intraradical mycelium could be observed.

Response to phosphorus availability

The proportion of ACP-active intraradical mycelium in the no-P treated plants showed a significant negative correlation to both the shoot P concentration (Fig. 3) and to the shoot dry weight (data not shown). The proportion of ACP-active arbuscules also showed a significant negative correlation with the shoot P concentration, but no correlation with shoot dry weight was found. For the high-P treated plants no clear trends were observed. No correlations were found regarding the

Table 1. Shoot dry weight and P content at sequential harvests of mycorrhizal and non-mycorrhizal *Allium cepa*. Standard errors (SE) are given in parentheses. The no-P (watered with nutrient solution without P) and high-P (watered with nutrient solution with 5 mg P l⁻¹) treatment was initiated at 37 days on plants previously treated with low-P (1 mg P l⁻¹) solution. NM, non-mycorrhizal plants; Myc, mycorrhizal plants.

		Shoot					
		Dry weight (mg)			P content (µg)		
	Days	No-P	Low-P	High-P	No-P	Low-P	High-P
NM	37		22.1 (2.4)			10.5 (0.9)	
	56		31.2 (0.8)			16.4 (1.4)	
Myc	37		29.7 (3.6)			20.6 (3.2)	
	42	33.2 (1.8)	ns	28.3 (5.3)	22.3 (0.9)	ns	17.1 (6.2)
	49	41.9 (1.7)	ns	42.0 (4.8)	31.5 (4.1)	ns	36.3 (6.6)
	56	44.6 (4.7)	*	62.4 (4.6)	37.5 (9.4)	[*]	61.2 (5.6)

* Significant difference at the 5% level between the no-P and the high-P treated plants (1-tailed *t* test with unequal variances).

[*] Significant difference at the 10% level.

ns, No significant difference.

Table 2. Proportion (%) of phosphatase-active hyphae, arbuscules and ERM as indicated by fast blue RR staining. The intraradical hyphae and arbuscules were collected after enzymatic digestion of the roots. The SE is given in parentheses. P treatment as described in Table 1. Arbs, arbuscules; ERM, extraradical mycelium; IRM, intraradical mycelium.

		Proportion of phosphatase active structures ¹ (%)					
		ACP ²			ALP		
	Days	No-P	Low-P	High-P	No-P	Low-P	High-P
IRM-Hyphae ***	37		80.4 (3.8)			39.7 (17.0)	
	42	88.0 (4.4)		88.5 (4.6)	53.6 (8.8)		58.8 (10.5)
	49	79.5 (2.4)		67.6 (3.1)	53.0 (10.7)		38.1 (17.3)
	56	69.8 (10.2)		70.5 (7.1)	26.3 (3.2)		28.7 (4.4)
IRM-Arbs ***	37		92.1 (6.2)			40.5 (24.3)	
	42	89.3 (3.8)		94.0 (2.3)	35.0 (17.5)		50.9 (7.9)
	49	71.1 (6.9)		64.9 (10.5)	44.6 (13.2)		36.2 (4.3)
	56	80.6 (10.0)		75.7 (5.4)	9.8 (7.4)		31.5 (10.0)
ERM **	37		0.4 (0.4)			1.7 (0.9)	
	42	3.7 (3.5)		0.0 (0.0)	07.6 (1.5)		12.6 (3.7)
	49	0.3 (0.3)		0.8 (0.3)	11.0 (2.3)		4.1 (2.4)
	56	1.1 (0.5)		2.2 (0.8)	4.9 (1.6)		4.2 (1.8)

¹ Proportion (%) of phosphatase-active hyphal intersections calculated as the percentage of active hyphae from the total amount of hyphae counted in the specific subsample that was stained for the specified phosphatase activity. A similar procedure was followed for arbuscules and ERM.

² ACP and ALP indicate staining for acid phosphatase and alkaline phosphatase, respectively. Bold numbers indicate that a significant difference (5% level) was found between the P treatments.

** Significant difference between ACP and ALP staining at the 1% level (2-tailed paired *t* test).

*** Significant difference at the 0.1% level.

proportion of ALP-active intraradical mycelium or arbuscules, or the proportion of ACP- and ALP-active extraradical mycelium.

The shoot P content was positively correlated to the shoot fresh weight (significant correlation, Fig. 4) after 19 d of P treatment (56 d of plant growth), and the proportion of ALP-active intraradical mycelium appeared to increase with higher shoot fresh weight (not significant; data not shown). Also, the proportion of ALP-active arbuscules seemed to increase with higher shoot fresh weight (not significant; Fig. 4). In earlier harvests (at 37, 42 and 49 d), the proportion of ALP-active structures was constant and no correlation with shoot fresh weight was observed.

DISCUSSION

We have presented experimental data on the metabolic response of an AM fungal symbiosis, established under P-limited conditions, to altered external P concentrations. This is the first study in which the relative amount of ACP- and ALP-active structures in both the extraradical and intraradical mycelium has been assayed in relation to a change in external P availability.

Except in one case, we were not able to show any direct change in the relative amount of phosphatase-active mycelium following P treatment for *G. margarita* (Table 2). However, the correlation between the proportion of ALP-active arbuscules and shoot fresh

Fig. 2. The relationship between the proportion of ALP-active intraradical (IRM) and extraradical (ERM) mycelium. ---●---●---, no-P treatment; —▲—▲—, high-P treatment; triplicate samples were used. P treatment as described in Table 1.

Fig. 3. Proportion of ACP-active intraradical mycelium (IRM) in relation to shoot P concentration. ---●---●---, no-P treatment; ▲, high-P treatment ($R^2=0.25$; no regression line shown); triplicate samples were used. P treatment as described in Table 1.

weight was positive, as was that for shoot P content and shoot fresh weight (Fig. 4). This suggests that the proportion of ALP-active arbuscules is related to the fungal P status or to fungal P transfer to the plant. However, there was a lag phase before the response in ALP was observed. Correlations between ALP and plant P content have been demonstrated previously (Tisserant *et al.* 1996). The proportion of ALP-active intraradical mycelium was not as strongly correlated to shoot fresh weight as the arbuscular proportion, which could be an effect of the central role of arbuscules for P transfer.

Under P-deprived conditions, the proportion of ACP-active intraradical mycelium was correlated to shoot P concentration (Fig. 3) and plant growth. This may suggest that under P limitation, in *G. margarita*, the host P status affected the proportion of ACP-active intraradical mycelium. Ezawa *et al.* (2001) suggested that a non-specific ACP could be involved in the hydrolysis of polyphosphate in intraradical hyphae before inorganic P is released into the plant–fungal interface by

Fig. 4. Shoot P content and the proportion of ACP- and ALP-active arbuscules as a function of the shoot fresh weight of mycorrhizal plants at 56 d. The mycorrhizal plants had been exposed to no-P (open symbols) or to high-P (solid symbols) for 19 d. —○—○—, shoot P content; △ △, proportion of ACP-active arbuscules ($R^2=0.12$; no regression line shown), ---□---□---, proportion of ALP-active arbuscules; triplicate samples were used. P treatments as described in Table 1.

the fungus. In this case, increased ACP activity in the intraradical mycelium could lead to an increase in P transfer from fungus to plant.

All these results indicate that the intraradical mycelium of *G. margarita* responded metabolically to P treatment. The influence of the plant P concentration on the fungal metabolic activity may also be of importance. Such an influence has been found earlier by Thomson *et al.* (1991) and Tisserant *et al.* (1996) who showed that increased plant P led to a decrease in AM fungal growth and activity. However, the lack of correlation between the proportion of phosphatase-active extraradical mycelium and shoot growth or P content supports observations by Boddington & Dodd (1999) that the proportion of ALP-active extraradical mycelium was unaffected by P treatment.

In the extraradical mycelium a large part of the phosphatase activity is associated with the cell walls (Joner & Johansen 2000), whereas in the intraradical mycelium it is mainly located in the acid vacuoles (Gianinazzi *et al.* 1979, Ezawa *et al.* 1995, 2001). A different location and proportion of ALP- and ACP-active structures could indicate that these enzymes have different roles in P metabolism. In the present study we showed that the intraradical mycelium had a higher proportion of phosphatase-active hyphae than the extraradical mycelium (Fig. 1, Table 2). This has previously been observed for ALP in a *Glomus* species (Zhao *et al.* 1997, Kjoller & Rosendahl 2000). The proportion of ALP-active extraradical mycelium could be correlated to the proportion of ALP-active intraradical mycelium (Fig. 2). Furthermore, while in the intraradical mycelium the proportion of ACP-active structures was highest, in the extraradical mycelium the proportion of ALP-active hyphae was highest. This suggests that in the extraradical mycelium the ALPs might be involved in hyphal P acquisition. The high proportion of ACP-active structures in the intraradical

mycelium indicates that one or several ACPs might be involved in P transfer processes.

Our results indicate that mycorrhizal plants, which have previously been limited in P, can respond metabolically to either the external P availability or the plant P concentration. This study emphasizes the importance of studying not only the amount of AM fungal mycelium but also the proportion of active mycelium. Furthermore, we showed that the extraradical and the intraradical mycelium should not be considered as one entity when studying the transfer of nutrients.

ACKNOWLEDGEMENTS

The financial support of the Swedish Council for Forestry and Agricultural Research, the Sweden–Japan Foundation and the Royal Physiographic Society in Lund is gratefully acknowledged. This work has also, in part, been supported by the Promotion of Basic Research Activities for Innovation of Biosciences (PROBRAIN), Bio-oriented Technology Research Advances Institution, Japan. The initial work was performed at the National Institute of Livestock and Grassland Science (Nishinasuno) and the Faculty of Medicine, Chemistry Laboratory, Teikyo University (Tokyo). We wish to thank Pål Axel Olsson for critical reading of the manuscript. I. v. A. is indebted to the research and technical members of the Laboratory of Soil Ecology (National Institute of Livestock and Grassland Science, Japan).

REFERENCES

- Abbott, L. K., Robson, A. D. & de Boer, G. (1984) The effect of phosphorus on the formation of hyphae in soil by the vesicular-arbuscular mycorrhizal fungus, *Glomus fasciculatum*. *New Phytologist* **97**: 437–446.
- Boddington, C. L. & Dodd, J. C. (1999) Evidence that differences in phosphate metabolism in mycorrhizas formed by species of *Glomus* and *Gigaspora* might be related to their life-cycle strategies. *New Phytologist* **142**: 531–538.
- Dickson, S., Smith, S. E. & Smith, F. A. (1999a) Characterization of two arbuscular mycorrhizal fungi in symbiosis with *Allium porrum*: colonization, plant growth and phosphate uptake. *New Phytologist* **144**: 163–172.
- Dickson, S., Smith, S. E. & Smith, F. A. (1999b) Characterization of two arbuscular mycorrhizal fungi in symbiosis with *Allium porrum*: inflow and flux of phosphate across the symbiotic interface. *New Phytologist* **144**: 173–181.
- Ezawa, T., Saito, M. & Yoshida, T. (1995) Comparison of phosphatase localization in the intraradical hyphae of arbuscular mycorrhizal fungi, *Glomus* spp. and *Gigaspora* spp. *Plant and Soil* **176**: 57–63.
- Ezawa, T., Smith, S. E. & Smith, F. A. (2001) Differentiation of polyphosphate metabolism between the extra- and intraradical hyphae of arbuscular mycorrhizal fungi. *New Phytologist* **149**: 555–563.
- Gianinazzi, S., Gianinazzi-Pearson, V. & Dexheimer, J. (1979) Enzymatic studies on the metabolism of vesicular-arbuscular mycorrhiza. III. Ultrastructural localization of acid and alkaline phosphatase in onion roots infected by *Glomus mosseae* (Nicol. & Gerd.). *New Phytologist* **82**: 127–132.
- Grotelueschen, J., Peleg, Y., Glass, N. L. & Metzenberg, R. L. (1994) Cloning and characterization of the *pho-2⁺* gene encoding a repressible alkaline phosphatase in *Neurospora crassa*. *Gene* **144**: 147–148.
- Joner, E. J. & Johansen, A. (2000) Phosphatase activity of external hyphae of two arbuscular mycorrhizal fungi. *Mycological Research* **104**: 81–86.
- Kaffman, A., Herskowitz, I., Tjian, R. & O'Shea, E. K. (1994) Phosphorylation of the transcription factor PHO4 by a cyclin-CDK complex, PHO80-PHO85. *Science* **263**: 1153–1156.
- Kjøller, R. & Rosendahl, S. (2000) Effects of fungicides on arbuscular mycorrhizal fungi: differential responses in alkaline phosphatase activity of external and internal hyphae. *Biology and Fertility of Soils* **31**: 361–365.
- McGonigle, T. P., Miller, M. H., Evans, D. G., Fairchild, G. L. & Swan, J. A. (1990) A new method which gives an objective measure of colonization of roots by vesicular-arbuscular mycorrhizal fungi. *New Phytologist* **115**: 495–501.
- Maldonado-Mendoza, I. E., Dewbre, G. R. & Harrison, M. J. (2001) A phosphate transporter gene from the extra-radical mycelium of an arbuscular mycorrhizal fungus *Glomus intraradices* is regulated in response to phosphate in the environment. *Molecular Plant–Microbe Interactions* **14**: 1140–1148.
- Mizuno, N. & Minami, M. (1980) [The use of H₂SO₄–H₂O₂ for destruction of plants matters as a preliminary determination of N, K, Mg, Ca, Fe, Mn.] *Journal of Science of Soil and Manure, Japan* **51**: 418–420 [In Japanese].
- O'Neill, E. M., Kaffman, A., Jolly, E. R. & O'Shea, E. K. (1996) Regulation of PHO4 nuclear localization by the PHO80-PHO85 cyclin-CDK complex. *Science* **271**: 209–212.
- Phillips, J. M. & Hayman, D. S. (1970) Improved procedures for clearing roots and staining parasitic and vesicular-arbuscular mycorrhizal fungi for rapid assessment of infection. *Transactions of the British Mycological Society* **55**: 158–161.
- Saito, M. (1995) Enzyme activities of the internal hyphae and germinated spores of an arbuscular mycorrhizal fungus, *Gigaspora margarita* Becker & Hall. *New Phytologist* **129**: 425–431.
- Saito, M., Stribley, D. P. & Hepper, C. M. (1993) Succinate dehydrogenase activity of external and internal hyphae of a vesicular-arbuscular mycorrhizal fungus, *Glomus mosseae* (Nicol. & Gerd.) Gerdmann and Trappe, during mycorrhizal colonization of roots of leek (*Allium porrum* L.), as revealed by *in situ* histochemical staining. *Mycorrhiza* **4**: 59–62.
- Smith, S. E. & Gianinazzi-Pearson, V. (1990) Phosphate uptake and arbuscular activity in mycorrhizal *Allium cepa* L.: effects of photon irradiance and phosphate nutrition. *Australian Journal of Plant Physiology* **17**: 177–188.
- Smith, S. E. & Read, D. J. (1997) *Mycorrhizal Symbiosis*. 2nd edn. Academic Press, London.
- Tawaray, K., Saito, M., Morioka, M. & Wagatsuma, T. (1994) Effect of phosphate application to arbuscular mycorrhizal onion on the development and succinate dehydrogenase activity of internal hyphae. *Soil Science and Plant Nutrition* **40**: 667–673.
- Thomson, B. D., Robson, A. D. & Abbott, L. K. (1991) Soil mediated effects of phosphorus supply on the formation of mycorrhizas by *Scutellispora calospora* (Nicol. & Gerd.) Walker & Sanders on subterranean clover. *New Phytologist* **118**: 463–469.
- Tisserant, B., Gianinazzi, S. & Gianinazzi-Pearson, V. (1996) Relationships between lateral root order, arbuscular mycorrhiza development, and the physiological state of the symbiotic fungus in *Platanus acerifolia*. *Canadian Journal of Botany* **74**: 1947–1955.
- Tisserant, B., Gianinazzi-Pearson, V., Gianinazzi, S. & Gollotte, A. (1993) *In planta* histochemical staining of fungal alkaline phosphatase activity for analysis of efficient arbuscular mycorrhizal infections. *Mycological Research* **97**: 245–250.
- van Aarle, I. M., Olsson, P. A. & Söderström, B. (2001) Microscopic detection of phosphatase activity of saprophytic and arbuscular mycorrhizal fungi using a fluorogenic substrate. *Mycologia* **93**: 17–24.
- Watanabe, F. S. & Olsen, S. R. (1965) Test of an ascorbic acid method for determining phosphorus in water and NaHCO₃ extracts from soil. *Soil Science Society of American Proceeding* **29**: 677–678.
- Zhao, B., Trouvelot, A., Gianinazzi, S. & Gianinazzi-Pearson, V. (1997) Influence of two legume species on hyphal production and activity of two arbuscular mycorrhizal fungi. *Mycorrhiza* **7**: 179–185.

Corresponding Editor: J. R. Leake