



LUND UNIVERSITY

Fetal growth predicts stress susceptibility independent of parental education in 161 991 adolescent Swedish male conscripts.

Nilsson, P M; Nilsson, J-Å; Östergren, Per-Olof; Rasmussen, F

Published in:
Journal of Epidemiology and Community Health

DOI:
[10.1136/jech.2003.015495](https://doi.org/10.1136/jech.2003.015495)

2004

[Link to publication](#)

Citation for published version (APA):
Nilsson, P. M., Nilsson, J.-Å., Östergren, P.-O., & Rasmussen, F. (2004). Fetal growth predicts stress susceptibility independent of parental education in 161 991 adolescent Swedish male conscripts. *Journal of Epidemiology and Community Health*, 58(7), 571-573. <https://doi.org/10.1136/jech.2003.015495>

Total number of authors:
4

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00



Fetal growth predicts stress susceptibility independent of parental education in 161 991 adolescent Swedish male conscripts

P M Nilsson, J-Å Nilsson, P-O Östergren and F Rasmussen

J. Epidemiol. Community Health 2004;58;571-573
doi:10.1136/jech.2003.015495

Updated information and services can be found at:
<http://jech.bmjjournals.com/cgi/content/full/58/7/571>

These include:

References

This article cites 5 articles, 3 of which can be accessed free at:
<http://jech.bmjjournals.com/cgi/content/full/58/7/571#BIBL>

Rapid responses

You can respond to this article at:
<http://jech.bmjjournals.com/cgi/eletter-submit/58/7/571>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

Topic collections

Articles on similar topics can be found in the following collections

[Pregnancy](#) (777 articles)
[Psychology](#) (333 articles)
[Adolescent medicine](#) (358 articles)

Notes

To order reprints of this article go to:
<http://www.bmjjournals.com/cgi/reprintform>

To subscribe to *Journal of Epidemiology and Community Health* go to:
<http://www.bmjjournals.com/subscriptions/>

SHORT REPORT

Fetal growth predicts stress susceptibility independent of parental education in 161 991 adolescent Swedish male conscripts

P M Nilsson, J-Å Nilsson, P-O Östergren, F Rasmussen

J Epidemiol Community Health 2004;**58**:571–573. doi: 10.1136/jech.2003.015495

Psychosocial stress could lead to a wide range of possible physiological reactions, due to both the total burden of stress as well as individual susceptibility. Two useful Swedish registers to investigate early life influences on stress susceptibility are the Swedish Medical Birth Register (MBR) and the Military Service Conscription Register (MSCR). In a previous study we showed a positive relation between fetal growth and psychological functioning (PF) including an assessment of stress susceptibility.¹ However, in that study we did not adjust for family social class—nor did another related study.² We have therefore now carried out such an analysis in an expanded cohort study, by adding parental educational level as a marker of family social class. The aim was to investigate independent associations between fetal growth and stress susceptibility in young men.

METHODS

We selected all Swedish men born in 1973–1979 and registered in the MBR ($n = 306\,497$). Birth characteristics recorded in MBR were linked to data on PF in the MSCR during 1990–1997. Excluded from the conscript test are men with severe physical or mental handicap ($n = 43\,890$). In all, 161 991 young men had complete birth data and full data on PF using the same standardised methods. The study was approved by the ethical committee, University of Lund.

Recorded MBR variables were: birth weight (g), birth length (cm), head circumference (cm), gestational age (weeks), maternal age (years), and parity (n). We used data only from subjects with a birth weight between 1500–4499

grams, born at term (\geq week 37), and with head circumference 20–60 cm, to avoid potential bias.

At conscription (MSCR) each young man undergoes a standard examination for PF, which is based on a personal standardised interview with a military psychologist.¹ This test (graded 1–9 scores; 9 highest) includes an evaluation of ability to cope with stress.

Data on parental education (years) were derived from the Population and Housing Census conducted by Statistics Sweden in 1970 and 1990.

Statistics

The three registers were linked by using unique personal 10 digit identification number. Birth weight was subdivided into centiles of aberration from expected birth weight adjusted for gestational week. We calculated the odds ratio (OR; 95%CI) for the risk of being assessed for a low PF (≤ 4 scores) in relation to impaired fetal growth, after adjustment for other birth characteristics and parental education, by use of the SAS program version 8 (Cary, NC, 1995). A p value less than 0.05 was considered significant.

RESULTS

Mean birth weight was 3576 (SD 436) grams after a mean of 39.9 (1.4) gestational weeks, as compared with a birth weight

Abbreviations: MBR, Medical Birth Register; MSCR, Military Service Conscription Register; PF, psychological functioning

Table 1 Psychological functioning (PF) scores in relation to categories of birth weight (registered birth weight in relation to expected birth weight) and paternal education (1970). Means (SD) and 95% confidence intervals (CI)

Birth weight category (centiles)	Paternal educational category	Number	PF score mean (SD)	95%CI
Lowest (1%–5%)	lowest	2041	4.61 (1.81)	4.53 to 4.69
	median-low	3394	5.18 (1.78)	5.12 to 5.23
	median-high	188	5.57 (1.73)	5.33 to 5.82
	highest	372	5.54 (1.75)	5.36 to 5.72
Second lowest (6%–10%)	lowest	2093	4.73 (1.84)	4.66 to 4.81
	median-low	3348	5.22 (1.77)	5.16 to 5.28
	median-high	212	5.39 (1.87)	5.13 to 5.64
	highest	369	5.74 (1.71)	5.57 to 5.91
Medium (11%–90%)	lowest	32176	4.94 (1.76)	4.92 to 4.96
	median-low	56630	5.40 (1.74)	5.39 to 5.42
	median-high	3919	5.77 (1.71)	5.72 to 5.83
	highest	8109	5.78 (1.76)	5.74 to 5.82
Second highest (91%–95%)	lowest	2177	4.98 (1.80)	4.91 to 5.06
	median-low	3563	5.45 (1.76)	5.39 to 5.51
	median-high	289	5.82 (1.71)	5.63 to 6.02
	highest	596	5.93 (1.76)	5.79 to 6.07
Highest (96%–100%)	lowest	2169	4.94 (1.79)	4.86 to 5.01
	median-low	3545	5.45 (1.80)	5.39 to 5.51
	median-high	252	5.83 (1.63)	5.63 to 6.03
	highest	598	5.81 (1.77)	5.67 to 5.96

Key points

- Impaired fetal growth has been associated with various adult disease manifestations in epidemiological studies from different populations.
- Neuropsychological development could be negatively influenced by reduced fetal growth, one possible consequence being impaired cognitive function and increased stress susceptibility in young adult life.
- In 161 991 Swedish male conscripts we found inverse associations between birth weight and psychological assessment scores including an evaluation of stress susceptibility, adjusted for other birth variables and parental educational level as a measure of childhood social class.
- Stress susceptibility in early adult life is more strongly associated with parental educational level than fetal growth, but both represent independent associations.
- Fetal growth thus seems to influence not only cognitive function, but also assessed susceptibility to stress—a factor of possible importance for adverse cardiovascular reactions with long term detrimental effects.

of 3448 (661) grams in the subjects excluded from conscript testing because of mental or physical handicap ($p < 0.001$). Mean value of the PF was 5.20 (1.80) scores. The PF score continuously increased with paternal educational level within each birth weight category (table 1) and with increasing paternal educational level. The odds ratio for receiving a low PF score (≤ 4 ; 34% of all) was higher for subjects belonging to the two lowest birth weight categories, OR 1.27 (1.21 to 1.33) for 1–5th centile of birth weight in relation to expected birth weight for gestational age, and OR 1.25 (1.19 to 1.31) for 6–10th centile, as compared with the reference category (10–90th centile). This was still evident, after full adjustment for parental educational level and age, as well as other birth characteristics (maternal parity, head circumference, and birth length). Low paternal educational level was slightly more important, OR 1.28 (1.24 to 1.33), as a risk factor for obtaining a low PF score (≤ 4) than birth weight alone in the fully adjusted statistical model (fig 1).

DISCUSSION

This study showed an increased risk of stress susceptibility among growth retarded male military conscripts. This




Figure 1 Psychological functioning score results, including an assessment of stress susceptibility, in relation to birth weight deviation from expected.

Policy implications

- Preventive efforts to ensure normal fetal growth and to strengthen family socioeconomic position in childhood may decrease the risk of boys growing up with increased stress susceptibility.

association was still evident after adjustment for parental educational level and other birth characteristics. This is in accordance with what has previously been reported in two similar Swedish studies, both however lacking adjustment for parental educational level.^{1,2} A positive association between size at birth and psychological function in women and men 42 years of age has recently been reported from a British population based cohort study, also after adjustment for childhood social class.³ Thus, both our studies consistently support the notion that fetal growth is important for PF in early adulthood.

We are aware of the fact that our findings are limited to young men and that the assessment of stress susceptibility by a military psychologist could potentially be influenced by day to day variation in mood, mental concentration, and cooperativeness of the conscripts. We also lack data on a number of subjects who were either not invited to the conscript test because of physical or mental abnormalities, or did not have a complete set of data. This potential bias is most probably diluting our findings, and not exaggerating them, as it would be suspected that an increased proportion of stress susceptible subjects belong to the excluded categories with handicaps. The correlations we obtained were weak, which is often the case in large scale epidemiological studies because of imprecision of measures, random variability, etc. Cognitive function may be a useful marker for long term health consequences as shown in other epidemiological studies.^{4,5}

In conclusion, impaired fetal growth is independently associated with increased stress susceptibility in young men.

ACKNOWLEDGEMENTS

We are grateful to Johan Lothigius, chief psychologist at the National Service Administration, Karlstad, Sweden, for fruitful discussions and advice as regards the psychological test methods used at conscript examinations.

CONTRIBUTORS

PMN and FR developed the research questions, study design, and analytical strategies. PMN drafted the paper and all authors contributed to the final version. FR created the database and JÄN conducted the statistical analyses. PMN is the guarantor.

Authors' affiliations

P M Nilsson, J-Å Nilsson, Department of Medicine, University Hospital, Malmö, Sweden

P-O Östergren, Department of Community Medicine, University Hospital, Malmö, Sweden

F Rasmussen, Department of Public Health Sciences, Karolinska Institute, Stockholm, Sweden

Funding: PMN was supported by grants from the Swedish Medical Association.

Competing interests: none declared.

Correspondence to: Dr P Nilsson, Department of Medicine, University Hospital, S-205 02 Malmö, Sweden; Peter.Nilsson@medforsk.mas.lu.se

Accepted for publication 11 November 2003

REFERENCES

- 1 Nilsson P, Nyberg P, Östergren P-O. Susceptibility to a psychological stress test in young males is associated with low birth weight. *Int J Epidemiol* 2001;**30**:75–80.
- 2 Lundgren EM, Cnattingius S, Jonsson B, *et al.* Intellectual and psychological performance in males born small for gestational age with and without catch-up growth. *Pediatr Res* 2001;**50**:91–6.
- 3 Cheung YB, Khoo KS, Karlberg J, *et al.* Association between psychological symptoms in adults and growth in early life: longitudinal follow up study. *BMJ* 2002;**325**:749–53.
- 4 Jiang GX, Rasmussen F, Wasserman D. Short stature and poor psychological performance: risk factors for attempted suicide among Swedish male conscripts. *Acta Psychiatr Scand* 1999;**100**:433–40.
- 5 Breslau N. Psychiatric sequelae of low birth weight. *Epidemiol Rev* 1995;**17**:96–106.

THE JECH GALLERY

doi: 10.1136/jech.2003.008821

Influential women in occupational health

Andrea Kidd Taylor—Ensuring safety for labour



January 1955–, Country of birth: USA.

Andrea Kidd Taylor, DrPH, MSPH, embodies the personification of the American Dream. Born of a middle class Southern family in the late 20th century, she was one of the first African-American children to attend, and thereby, de-segregate an all white school.

Reflecting her family's strong background in organised labour, Taylor worked as an industrial hygienist and occupational health policy consultant for the International Union, United Automobile, Aerospace, and Agricultural Implement Workers of America (UAW), playing a seminal part in many Right-to-Know programmes throughout the USA.

“To achieve gains in health, public policies must be implemented that overcome discrimination and reduce exploitation of labor, thereby promoting socioeconomic equity and improved health status for all people.”

Unafraid to tackle major health policy questions, she served on the Presidential Advisory Committee on Gulf War Veterans' Illnesses, and has represented organised labour on the National Advisory Committee on Occupational Safety and Health. In 1998, Taylor was appointed to the US Chemical Safety and Hazard Investigation Board (CSB), empowered to investigate major chemical accidents that involve deaths or serious harm to workers and the public.

Ilise L Feitshans

Adjunct Faculty, Cornell University, School of Industrial and Labor Relations, Albany, NY, USA

Deborah F Salerno

Clinical Communications Scientist, Pfizer Global Research and Development—Michigan Laboratories, Ann Arbor, MI, USA

Correspondence to: Deborah F Salerno, 2800 Plymouth Road, Ann Arbor, MI 48105, USA; deborah.salerno@pfizer.com