



# LUND UNIVERSITY

## Intralipid decreases apolipoprotein m levels and insulin sensitivity in rats.

Zheng, Lu; Feng, Yuehua; Shi, Yuanping; Zhang, Jun; Mu, Qinfeng; Qin, Li; Berggren Söderlund, Maria; Nilsson-Ehle, Peter; Zhang, Xiaoying; Luo, Guanghua; Xu, Ning

*Published in:*  
PLoS ONE

*DOI:*  
[10.1371/journal.pone.0105681](https://doi.org/10.1371/journal.pone.0105681)

2014

[Link to publication](#)

*Citation for published version (APA):*

Zheng, L., Feng, Y., Shi, Y., Zhang, J., Mu, Q., Qin, L., Berggren Söderlund, M., Nilsson-Ehle, P., Zhang, X., Luo, G., & Xu, N. (2014). Intralipid decreases apolipoprotein m levels and insulin sensitivity in rats. *PLoS ONE*, 9(8), Article e105681. <https://doi.org/10.1371/journal.pone.0105681>

*Total number of authors:*  
11

### General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00



# Intralipid Decreases Apolipoprotein M Levels and Insulin Sensitivity in Rats

Lu Zheng<sup>1</sup>, Yuehua Feng<sup>1</sup>, Yuanping Shi<sup>1</sup>, Jun Zhang<sup>1</sup>, Qinfeng Mu<sup>1</sup>, Li Qin<sup>1</sup>, Maria Berggren-Söderlund<sup>3</sup>, Peter Nilsson-Ehle<sup>3</sup>, Xiaoying Zhang<sup>2</sup>, Guanghua Luo<sup>1\*</sup>, Ning Xu<sup>3\*</sup>

**1** Comprehensive Laboratory, the Third Affiliated Hospital of Soochow University, Changzhou, P.R. China, **2** Department of Cardiothoracic Surgery, the Third Affiliated Hospital of Soochow University, Changzhou, P.R. China, **3** Division of Clinical Chemistry and Pharmacology, Department of Laboratory Medicine, Lunds University, Lund, Sweden

## Abstract

**Background:** Apolipoprotein M (ApoM) is a constituent of high-density lipoproteins (HDL). It plays a crucial role in HDL-mediated reverse cholesterol transport. Insulin resistance is associated with decreased ApoM levels.

**Aims:** To assess the effects of increased free fatty acids (FFAs) levels after short-term Intralipid infusion on insulin sensitivity and hepatic ApoM gene expression.

**Methods:** Adult male Sprague-Dawley (SD) rats infused with 20% Intralipid solution for 6 h. Glucose infusion rates (GIR) were determined by hyperinsulinemic-euglycemic clamp during Intralipid infusion and plasma FFA levels were measured by colorimetry. Rats were sacrificed after Intralipid treatment and livers were sampled. Human embryonic kidney 293T cells were transfected with a lentivirus mediated human apoM overexpression system. Goto-Kakizaki (GK) rats were injected with the lentiviral vector and insulin tolerance was assessed. Gene expression was assessed by real-time RT-PCR and PCR array.

**Results:** Intralipid increased FFAs by 17.6 folds and GIR was decreased by 27.1% compared to the control group. ApoM gene expression was decreased by 40.4% after Intralipid infusion. PPAR<sub>β/δ</sub> expression was not changed by Intralipid. Whereas the mRNA levels of *Acaca*, *Acox1*, *Akt1*, *V-raf murine sarcoma 3611 viral oncogene homolog*, *G6pc*, *Irs2*, *Ldlr*, *Map2k1*, *pyruvate kinase* and *RBC* were significantly increased in rat liver after Intralipid infusion. The Mitogen-activated protein kinase 8 (MAPK8) was significantly down-regulated in 293T cells overexpressing ApoM. Overexpression of human ApoM in GK rats could enhance the glucose-lowering effect of exogenous insulin.

**Conclusion:** These results suggest that Intralipid could decrease hepatic ApoM levels. ApoM overexpression may have a potential role in improving insulin resistance *in vivo* and modulating apoM expression might be a future therapeutic strategy against insulin resistance in type 2 diabetes.

**Citation:** Zheng L, Feng Y, Shi Y, Zhang J, Mu Q, et al. (2014) Intralipid Decreases Apolipoprotein M Levels and Insulin Sensitivity in Rats. PLoS ONE 9(8): e105681. doi:10.1371/journal.pone.0105681

**Editor:** Victor Sanchez-Margalet, Virgen Macarena University Hospital, School of Medicine, University of Seville, Spain

**Received:** March 7, 2014; **Accepted:** July 22, 2014; **Published:** August 21, 2014

**Copyright:** © 2014 Zheng et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This research project was supported by the National Natural Science Foundation of China (NSFC) (81071414, 81201352, <http://www.nsf.gov.cn/>), the Natural Science Foundation of Jiangsu Province (BK2011245, BK2012154, <http://www.jskj.gov.cn/>) and a research grant from the Changzhou Science & Technology Bureau (CJ20122012, <http://new.czstb.gov.cn/>). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* Email: [shineroar@163.com](mailto:shineroar@163.com) (GL); [ning.xu@med.lu.se](mailto:ning.xu@med.lu.se) (NX)

## Introduction

Apolipoprotein M (ApoM) is a constituent of plasma high-density lipoproteins (HDL) and most plasma ApoM are bound to HDL, which plays an important role on lipid and lipoprotein metabolism [1,2]. ApoM could influence pre-β HDL formation and cholesterol efflux, which is thought to be one of key regulators of HDL metabolism and reverse cholesterol transport [1,2]. It has been demonstrated that ApoM expression could be directly regulated by the hepatic nuclear factor-1α (HNF-1α) [3], liver receptor homolog-1 (LRH-1) [4], forkhead box A2 (Foxa2) [5], and liver X receptor (LXR) [5]. And all of these transcription factors are also involved in hepatic lipid and glucose metabolism [3–6].

Type 2 diabetes is a major health problem and its prevalence increased dramatically in the last decades, mostly due to obesity and sedentary lifestyle [7,8]. Furthermore, insulin resistance, a key feature of type 2 diabetes, induces major metabolic abnormalities, resulting in high free fatty acids (FFA) plasma levels, hypertriglyceridemia, low HDL levels and small dense LDL particles [9,10]. In addition, size and composition of HDL particles are abnormal in diabetic patients [11]. Indeed, serum/plasma ApoM levels are significantly reduced in diabetic and metabolic syndrome patients [12–14].

Since insulin resistance is one of the key features of type 2 diabetes, finding new ways to improve insulin resistance is important for the management of these patients. *In vitro* insulin and insulin-like growth factor I (IGF-I) could significantly inhibit

**Table 1.** Fatty acid composition of soybean oil in 20% Intralipid.

Fatty acid	Ratio (% w/w)
<C14:0	≤0.1
Myristic acid	≤0.2
Palmitic acid	9.0–13.0
Palmitoleic Acid	≤0.3
Stearic acid	3.5–5.0
Oleic acid	17.0–30.0
Linoleic acid	48.0–58.0
Linolenic acid	5.0–11.0
Arachidic acid	≤0.1
Eicosenoic acid	≤0.1
Docosanoic acid	≤0.1

doi:10.1371/journal.pone.0105681.t001

apoM expression with a dose- and time-dependent manner [15,16]. Moreover, both *in vivo* and *in vitro* observations suggested that ApoM may also be associated with diabetes and obesity [12–14]. Exogenous insulin administration could partially reverse abnormal ApoM expression in diabetic rats [17]. ApoM levels were significantly decreased in hyperglycemic rats, and high glucose and insulin concentrations inhibited ApoM expression in cultured cells [16].

Intralipid is a solution of soybean oil, phosphatidylcholine, glycerol and water, and is used to increase FFA levels. It contains significant amounts of ω-6 polyunsaturated fatty acids (PUFA) that are easily oxidized to generate reactive oxygen species [18]. Short-term Intralipid infusion significantly increases FFA levels and insulin resistance [19,20] by decreasing peripheral glucose uptake [21] and down-regulating intracellular insulin signaling [22,23]. Elevated FFA levels decrease insulin sensitivity in trained and sedentary humans [24], and induce insulin resistance in both skeletal and cardiac muscles [25]. FFAs are ligands for ApoM in plasma, which could contribute to FFA removal from the circulation, preventing their ill effects [26].

We hypothesized that downregulation of ApoM expression by hyperglycemia may be associated with insulin resistance. In the present study, we studied the effects of artificially increasing FFAs on ApoM expression and insulin sensitivity in rats. We showed that increased FFA levels decreased both ApoM levels and insulin sensitivity. Therefore, modulating ApoM expression might be a future therapeutic strategy against insulin resistance in type 2 diabetes.

## Materials and Methods

### Animals

Each experimental group contained 5–6 adult male Sprague-Dawley (SD) rats (286.2±18.3 g) or, as a model for insulin resistance, aged male Goto-Kakizaki (GK) rats (416.1±40.0 g). In the present study, 10 male SD rats (8 weeks old) underwent a hyperinsulinemic-euglycemic clamp (HEC) and 10 aged male GK rats (32 weeks old) were obtained from the Shanghai Slac Laboratory Animal Co., China. Another 12 male SD rats (8 weeks old) were obtained from the Changzhou Cavens Laboratory Animal Co., China.

Rats were kept in a temperature-controlled (22°C) room with 12-hrs light-dark cycle, and were provided with standard rodent chow and water ad libitum. Rats were acclimatized for one week before placing catheters. All procedures and animal experiments were approved by the Animal Care and Use Committee of the Soochow University (Suzhou, China) (permit number SYXK(Su)2002–0045)).

### Surgical preparation

After being acclimated to their new environment, SD rats underwent surgery to place catheters 7 days before experiments. Rats were anesthetized with 10% chloral hydrate (4 ml/kg). Two catheters were placed, one in each jugular vein: one for Intralipid infusion, and the other for 20% glucose infusion during hyperinsulinemic euglycemic clamp (HEC). An additional catheter was placed in a carotid artery for blood sampling. The free ends of the catheters were attached to steel tubing and tunneled subcutaneously on the back of the neck. The catheters were flushed with isotonic saline containing 50 IU/ml heparin (Qianhong Bio-pharma Co., Ltd., Changzhou, China) and filled with a viscous solution of heparin (500 IU/ml) and 300 g/L polyvinyl pyrrolidone (PVP-10; Sigma, St Louis, MO, USA) to prevent blood reflux into the catheter lumen.

### Intralipid infusion

20% Intralipid (Sino-Swed Pharmaceutical Corp, Ltd., Jiangsu, China) containing 20% soybean oil, 1.2% lecithin and 2.2% glycerin, is a triglyceride emulsion, which releases fatty acids (Table 1) with the concomitant infusion of heparin, a stimulant of the lipoprotein lipase enzyme. On the day of the experiment, the catheters were carefully connected to infusion pumps (Smiths Medical, Lower Pemberton, UK). A 20% Intralipid solution (10 ml·kg<sup>-1</sup>·h<sup>-1</sup>) combined with heparin (0.0975 IU/min) was infused for 6 h. In addition, Intralipid/heparin was infused via the tail vein in another parallel experiments, without HEC test, to avoid the interference of the 20% glucose infusion on liver gene expression. All control rats received 5% glucose solution (10 ml·kg<sup>-1</sup>·h<sup>-1</sup>) combined with heparin (0.0975 IU/min). After Intralipid treatment for 6 h, SD rats were anesthetized using 10% chloral hydrate (4 ml/kg) and sacrificed. Blood samples were obtained from the inferior vena cava. Plasma was separated by

**Table 2.** Sequences of primers and probes.

Gene	Primer/Probe	Sequence (5'-3')	Product (bp)
Rat ApoM	Forward	acaagagacccccagagccc	67
	Reverse	tccatggtgggagccg	
	Probe	FAM-acctgggctgtggtactttattgctgg-TAMRA	
Rat $\beta$ -actin	Forward	gccactgccgcatcctct	108
	Reverse	ctggaagagagcctcgggg	
	Probe	FAM-agctgcctgacggtcaggtcatcactatc-TAMRA	
Rat PPAR $\beta/\delta$	Forward	gaagaaccgcaacaagtgtcagta	85
	Reverse	cctccaagcgatagcgt	
	Probe	FAM-cttcagaagtgcctggcctcggc-TAMRA	

doi:10.1371/journal.pone.0105681.t002

centrifugation and stored at  $-70^{\circ}\text{C}$ . Livers were removed, sectioned, and stored in liquid nitrogen.

### Hyperinsulinemic-euglycemic clamp

To assess insulin sensitivity, rats underwent a primed-constant infusion of 10 mU/kg/min of insulin to achieve a steady state. A 20% D-glucose solution (AMRESCO Inc., Solon, OH, USA) was then infused. Blood glucose was measured at 5-min intervals using an ACCU-CHECK Active glucometer (Roche Diagnostics, Basel, Switzerland). The glucose infusion rate (GIR) was adjusted in order to maintain a blood glucose level of approximately 5.5 mmol/L. The mean glucose infusion rate in the last 30 min was used for analysis [27].

### Determinations of FFAs


FFAs were determined with the nonesterified fatty acid (NEFA) colorimetric method (Applygen Technologies Inc, Beijing, China). Briefly, total plasma FFAs were extracted with a chloroform:N-heptane:methanol solution (56:42:2), coupled with copper, reacted with color reagent and measured with a UV-2401PC UV-visible

spectrophotometer (Shimadzu, Tokyo, Japan) at 550 nm. The standard curve was created using a series of dilution of palmitic acid.


### Lentiviral expression system for overexpression of human ApoM gene

A lentiviral expression system for overexpression of the human ApoM gene was constructed by Shanghai GenePharma Co., Ltd (Shanghai, China). A 564-bp fragment of the ApoM gene (GenBank accession number: AF118393) was cloned into a lentivirus transfection vector. In brief, a plasmid containing the ApoM gene, a plasmid encoding the Gag/Pol gene, a plasmid encoding the rev gene, and a plasmid encoding the vesicular stomatitis virus G glycoprotein gene were co-transferred into human embryonic kidney 293T cells (HEK 293T) using lipofectamine 2000 (Invitrogen Inc., Carlsbad, CA, USA). A control lentivirus vector expressing green fluorescent protein (GFP) was constructed in the same manner, but without the ApoM gene.

HEK 293T cells (ATCC, Manassas, VA, USA) were cultured in RPMI1640 supplemented with 10% fetal bovine serum (GIBCO,



**Figure 1. Effects of Intralipid on insulin sensitivity, free fatty acids (FFA) levels and hepatic ApoM gene expression in SD rats.** (A) Plasma FFA levels in rats after infusion of 5% glucose solution (controls,  $n=6$ ) or 20% Intralipid solution ( $n=6$ ) for 6 h, and glucose infusion rate (GIR) during hyperinsulinemic euglycemic clamp (HEC) in control rats ( $n=5$ ) and Intralipid-infused rats ( $n=5$ ). (B) Hepatic ApoM mRNA levels in control rats and in Intralipid-infused rats with ( $n=5$  for each group) or without ( $n=6$  for each group) HEC were determined by real-time RT-PCR. ApoM mRNA levels in control rats were set as 100%. Data are presented as means  $\pm$  standard error of the mean (SEM).  $*P<0.05$ ,  $**P<0.01$ ,  $***P<0.001$  vs. controls. doi:10.1371/journal.pone.0105681.g001




**Figure 2. Effects of Intralipid on genes related to insulin signaling in liver tissues of SD rats.** Genes related to insulin signaling were determined by PCR array. Fold changes of certain liver genes in Intralipid-infused SD rats (experimental group) compared with 5% glucose solution (control group). \* $P < 0.05$  and \*\* $P < 0.01$  vs. control group; (n = 6 for each group, total of 12 PCR arrays, all genes normalized to  $\beta$ -actin individually). Numbers in bold represent fold-changes. Genes are identified using their abbreviated name (in red). A (in blue): Average threshold cycle was relatively high (>30) in the control or the experimental sample, and was reasonably low in the other sample (<30). B (in blue): Average threshold cycle was relatively high (>30), meaning that its relative expression level was low, in both control and experimental samples, and the  $P$ -value for the fold-change was either unavailable or high ( $P > 0.05$ ). C (in blue): Average threshold cycle was either not determined or greater than the predefined cut-off value (default value of 35) in both samples, meaning that its expression was undetected, making the result erroneous and non-interpretable. "OKAY" indicates that threshold cycle in the control and the experimental sample was accepted (<30). doi:10.1371/journal.pone.0105681.g002

Invitrogen Inc., Carlsbad, CA, USA), 100 U/ml of penicillin, 100  $\mu$ g/ml of streptomycin and 2 mM of L-glutamine (Invitrogen, Carlsbad, CA, USA) at 37°C in a 5% CO<sub>2</sub> humidified incubator. Cells were seeded in 6-well cell culture plates, and were grown to 50–70% confluence. Prior to experiments, cells were washed once with phosphate buffered saline (PBS), and once with serum-free RPMI1640 without antibiotics. In each well, the experimental medium contained 3 ml of RPMI1640 with 1.0% human serum albumin (HSA), 50  $\mu$ l of  $1 \times 10^9$  TU lentivirus expressing GFP and ApoM simultaneously (n = 6) or lentivirus only expressing GFP (control group, n = 6), and 15  $\mu$ g of polybrene (Invitrogen, Carlsbad, CA, USA).

### Insulin tolerance test

An insulin tolerance test (ITT) was performed in GK rats (n = 5 for each group) 14 days after being transfected with  $5 \times 10^8$  TU of lentiviral vectors with or without the human ApoM gene via tail vein injection. After a 5-h fast, rats were injected with 1 IU/kg of insulin (Wanbang Biopharmaceuticals, China) intra-peritoneally. Blood was sampled from the tail vein 0, 30, 60, 90, 120, 150, 180 and 210 min after insulin injection. Blood glucose levels were determined using a glucosimeter (ACCU-CHEK Active, Roche Diagnostics, Basel, Switzerland). Blood glucose half-time ( $t_{1/2}$ ) was calculated from the slope of the least squares regression line of the blood glucose concentration during the linear phase of decrease.





**Figure 3. Effects of Intralipid on hepatic PPAR $\beta/\delta$  mRNA expression in SD rats.** Total RNA was obtained from the liver of rats after infusion of 5% glucose solution (controls, n=6) or rats infused with 20% Intralipid solution (n=6). mRNA expression was determined by real-time PCR. mRNA levels in control rats were set as 100%. Data are presented as means  $\pm$  SEM. doi:10.1371/journal.pone.0105681.g003

After ITT, GK rats were anesthetized using 10% chloral hydrate (4 ml/kg) and sacrificed. Lungs, livers and kidneys were removed, sectioned, and stored in liquid nitrogen.

#### Real-time Reverse transcription (RT)-PCR and PCR array

Total RNA from 293T cells, liver tissues from SD rats and liver, kidney and lung tissues from GK rats was extracted using a total RNA purification kit (Omega Bio-Tek Inc., Norcross, GA, USA), according to the manufacturer's instructions. RNA quality was determined by absorbance at 260/280 nm. Using the first strand cDNA synthesis kit (Qiagen, Venlo, Netherlands), 2  $\mu$ g of total RNA was reverse transcribed to cDNA. PCR primers were designed according to the data available in GenBank (Table 2).  $\beta$ -actin was used as the reference gene. Relative standard curves were created to compensate for PCR efficiency. mRNA levels are expressed in relation to  $\beta$ -actin mRNA level. The real-time PCR reaction for each gene was performed in a 25  $\mu$ L volume, in a glass capillary containing 0.4  $\mu$ M of each primer and probe, 2  $\mu$ L of cDNA, 1  $\times$  buffer, 1.5 mM of MgCl<sub>2</sub>, 200  $\mu$ M of dNTPs, and 1.25 U of Taq DNA polymerase. Thermal cycling conditions were: initial denaturation at 95°C for 3 min, followed by 40 cycles at 95°C for 5 sec and 60°C for 15 sec for Rat PPAR $\beta/\delta$  (rat ApoM, 58°C for 12 sec; rat  $\beta$ -actin, 61°C for 10 sec). All PCRs were performed on a LightCycler real-time PCR system (Roche Diagnostics, Basel, Switzerland). The human ApoM gene quantification assay was performed according to a published method [28].

To scan genes of interest in liver of SD rats administrated with 20% Intralipid solution and in 293T cells transfected with a lentivirus mediated human apoM overexpression system, we used the PCR Array analyses for genes related to Rat Insulin Signaling Pathway (PARN-030Z) (SABiosciences, Qiagen, Venlo, Netherlands) in rats, and Human Type 2 Diabetes Mellitus (PA2) (CT bioscience, Jiangsu, China) in 293T cells, according to the manufacturer's instructions. PCR array data were calculated by the comparative cycle threshold method, normalized against multiple housekeeping genes, and expressed as mean fold change in experimental samples relative to control samples.

#### Statistical analysis

Data are expressed as means  $\pm$  standard error of the mean (SEM). Statistical analyses were performed using GraphPad Prism 6.0 (GraphPad Software, Inc., San Diego, California, USA).

Comparisons between two groups were evaluated by unpaired *t*-tests. *P*-values <0.05 were considered significant.

## Results

### Intralipid decreased insulin sensitivity and inhibited ApoM gene expression


In SD rats infused with 20% Intralipid, plasma FFA levels increased by 17.6-folds and GIR was reduced by 27.1% compared with SD rats infused with 5% glucose solution (both *P*<0.01) (Figure 1A). As shown in Figure 1B, liver ApoM gene expression was significantly inhibited by 40.4% after Intralipid infusion followed by HEC, compared with control SD rats. When the HEC test was omitted to avoid the potentially confounding effect of 20% glucose, ApoM mRNA levels were still significantly decreased by 40.4% in SD rats infused with Intralipid compared with control SD rats (Figure 1B). There was no significant difference in ApoM mRNA levels between rats infused with Intralipid with or without HEC (*P*>0.05).

### Effects of short-term Intralipid infusion on genes related to the insulin signaling pathway in rat liver

Using PCR array, we examined 84 genes reportedly related to the insulin response in SD rats. We observed that the mRNA levels of acetyl-CoA carboxylase alpha (Acaca), acyl-CoA oxidase 1 (Acox1), v-akt murine thymoma viral oncogene homolog 1 (Akt1), V-raf murine sarcoma 3611 viral oncogene homolog (Araf, component of MAPK pathway), catalytic subunit (G6pc), insulin receptor substrate 2 (Irs2), low density lipoprotein receptor (Ldlr), mitogen-activated protein kinase kinase 1 (Map2k1) and pyruvate kinase in liver and RBC (Pklr, target gene for SREBP1) were significantly increased in SD rats treated with Intralipid (Figure 2).

### Effect of Intralipid on PPAR $\beta/\delta$ mRNA expression in rat liver

We previously reported [15] that activation of PPAR $\beta/\delta$ , but not PPAR $\alpha$  and PPAR $\gamma$ , might be involved in the down-regulation of ApoM. Since the gene chip did not contain the PPAR $\beta/\delta$  gene, we detected the mRNA levels of PPAR $\beta/\delta$  by real-time RT-PCR after infusion of 20% Intralipid in SD rats. Intralipid solution did not



**Figure 4. Human ApoM overexpression increased MAPK8 mRNA expression in 293T cells.** Lentiviral vectors (LV) were used as delivery vehicles. 293T cells were treated with LV4(GFP)-NC that expressed GFP alone (control group, n = 6) or LV4(GFP)-ApoM that expressed GFP and ApoM (n = 6). (A) MAPK8 mRNA levels. (B) ApoM mRNA levels. mRNA expression was determined by real-time RT-PCR. mRNA levels in control rats were set as 100%. Data are presented as means ± SEM. \*\**P*<0.01 and \*\*\**P*<0.001 vs. controls. doi:10.1371/journal.pone.0105681.g004

significantly influence the hepatic PPAR<sub>β/δ</sub> mRNA expression in rats (*P* = 0.21) (Figure 3).

**Overexpression of human ApoM gene on insulin sensitivity in 293T cells**

PCR array analyses using the Human Type 2 Diabetes Mellitus (PA2) array in 293T cells overexpressing human ApoM (Table S1) demonstrated that the mitogen-activated protein kinase 8 (MAPK8) gene, which is related to insulin resistance [29], was down-regulated by 2.1 folds (*P* = 0.0029) (Figure 4A), while ApoM mRNA levels in 293T cells transfected with the human ApoM gene were increased by 80 folds (*P* = 0.0001) (Figure 4B).


**Overexpression of human ApoM gene on insulin sensitivity in GK rats**

We investigated the effects of overexpressing the human ApoM gene on insulin sensitivity in GK rats, which is a non-obese Wistar substrain rat characterized by mild hyperglycemia, insulin resistance and hyperinsulinemia [30,31]. As shown in Figure 5A,

human ApoM mRNA levels were significantly increased in the lungs (No human ApoM gene expression in kidneys and livers) of GK rats after injecting  $5 \times 10^8$  TU of lentiviral vectors integrating the human ApoM gene. There was no expression of the human ApoM gene in the lungs of GK rats transfected with the LV4(GFP)-NC vector. After 14 days, there were no obvious differences in fasting blood glucose levels between control rats and rats transfected with the human ApoM gene ( $10.82 \pm 0.92$  mmol/L vs.  $11.34 \pm 1.06$  mmol/L, *P* = 0.72). Interestingly, IIT analysis (Figure 5B) showed that the blood glucose t<sub>1/2</sub> of GK rats transfected with the human ApoM gene (61.5 min) was shorter than in control rats (89.0 min), although the difference did not reach statistical significance (*P* = 0.15), which may be due to the limited number of animals.

**Discussion**

The aim of the present study was to assess the effects of increased FFAs levels after short-term Intralipid infusion on insulin sensitivity and hepatic ApoM gene expression. Intralipid could



**Figure 5. Human ApoM overexpression in Goto-Kakizaki (GK) rats improves insulin resistance.** GK rats were injected with  $5 \times 10^8$  TU of LV4(GFP)-NC that expressed GFP alone (control group, n = 5) or LV4(GFP)-ApoM that expressed GFP and ApoM (n = 5) via the tail vein. (A) Human ApoM mRNA expressions in lungs of GK rats. mRNA expression was determined by real-time RT-PCR. Copy number of ApoM in the lungs of GK rats treated with the LV4(GFP)-NC or LV4(GFP)-ApoM, respectively. (B) Insulin tolerance test (ITT) was performed on day 14. Blood was sampled from the tail vein at 0, 30, 60, 90, 120, 150, 180 and 210 min after insulin administration, and blood glucose half-time (t<sub>1/2</sub>) was calculated from the slope of the least squares regression during the linear phase of decline. Data are presented as means ± SEM. doi:10.1371/journal.pone.0105681.g005

increase FFAs by 17.6 folds and decrease GIR by 27.1%. This further confirms previously findings [32] that Intralipid infusion increases plasma FFA levels and simultaneously decreases insulin sensitivity.

ApoM is the acceptor of HDL-carrying S1P [33], apoM could enhance HDL-mediated anti-oxidation effects [34], and it plays an important role for the pre- $\beta$  HDL formation [35]. In the present study, we demonstrated that ApoM expression was also associated with insulin sensitivity. Artificially increased FFA levels after short-term Intralipid infusion could decrease ApoM expression. In order to minimize the experiment bias, the rats in the control group received glucose to avoid the confounding effect of starving on normal physiological processes. As expected, infusion of 20% Intralipid solution resulted in elevated plasma FFAs levels and decreased insulin sensitivity [5,32,36,37]. Intralipid infusion could also decrease hepatic apoM expression in rats. To explore the mechanisms of down-regulation of ApoM by FFA, we monitored the expression of 84 genes related to the insulin response in liver tissues on rats after Intralipid infusion. The mRNA levels of *Acaca*, *Acox1*, *Akt1*, *Araf*, *G6pc*, *Irs2*, *Ldlr*, *Map2k1* and pyruvate kinase, which are all proteins involved in energy metabolism and insulin signaling in the liver, were all significantly increased. A number of transcription factors are involved in the liver response to increased FFA levels and in insulin signaling, such as HNF-1 $\alpha$ , LRH-1, Foxa2, and LXR [38–40]. These transcriptions factors are also involved in ApoM transcription, suggesting that ApoM may directly or indirectly involved in energy source metabolism and insulin signaling [3–6].

We have previously reported [15] that activation of PPAR $_{\beta/\delta}$ , but not PPAR $\alpha$  and PPAR $\gamma$ , might be involved in the down-regulation of ApoM, suggesting that the FFA-induced down-regulation of ApoM expression may be mediated, at least in part, by the PPAR $_{\beta/\delta}$  pathway. However, in the present study, Intralipid infusion did not influence hepatic PPAR $_{\beta/\delta}$  mRNA expression, despite marked elevations in plasma FFA levels. This discrepancy may be explained by the fatty acid composition of Intralipid (limoleic acid 44–62%, oleic acid 19–30%, palmitic acid 7–14%, linolenic acid 4–11% and stearic acid 1.4–5.5%). The effects of different FFA on PPAR $_{\beta/\delta}$  expression warrant further studies.

## References

- Xu N, Dahlback B (1999) A novel human apolipoprotein (apoM). *J Biol Chem* 274: 31286–31290.
- Christoffersen C, Nielsen LB, Axler O, Andersson A, Johnsen AH, et al. (2006) Isolation and characterization of human apolipoprotein M-containing lipoproteins. *J Lipid Res* 47: 1833–1843.
- Skupien J, Kepka G, Gorczynska-Kosiorz S, Gebaska A, Klupa T, et al. (2007) Evaluation of Apolipoprotein M Serum Concentration as a Biomarker of HNF-1 $\alpha$  MODY. *Rev Diabet Stud* 4: 231–235.
- Venteclef N, Haroniti A, Tousaint JJ, Talianidis I, Delerive P (2008) Regulation of anti-atherogenic apolipoprotein M gene expression by the orphan nuclear receptor LRH-1. *J Biol Chem* 283: 3694–3701.
- Zhang X, Zhu Z, Luo G, Zheng L, Nilsson-Ehle P, et al. (2008) Liver X receptor agonist downregulates hepatic apoM expression in vivo and in vitro. *Biochem Biophys Res Commun* 371: 114–117.
- Hu YW, Zheng L, Wang Q, Zhong TY, Yu X, et al. (2012) Vascular endothelial growth factor downregulates apolipoprotein M expression by inhibiting Foxa2 in a Nur77-dependent manner. *Rejuvenation Res* 15: 423–434.
- Engelgau MM, Geiss LS, Saaddine JB, Boyle JP, Benjamin SM, et al. (2004) The evolving diabetes burden in the United States. *Ann Intern Med* 140: 945–950.
- Sullivan PW, Morratto EH, Ghushchyan V, Wyatt HR, Hill JO (2005) Obesity, inactivity, and the prevalence of diabetes and diabetes-related cardiovascular comorbidities in the U.S., 2000–2002. *Diabetes Care* 28: 1599–1603.
- Garg A, Grundy SM (1990) Nicotinic acid as therapy for dyslipidemia in non-insulin-dependent diabetes mellitus. *JAMA* 264: 723–726.
- Garvey WT, Kwon S, Zheng D, Shaughnessy S, Wallace P, et al. (2003) Effects of insulin resistance and type 2 diabetes on lipoprotein subclass particle size and concentration determined by nuclear magnetic resonance. *Diabetes* 52: 453–462.
- Adiels M, Olofsson SO, Taskinen MR, Boren J (2008) Overproduction of very low-density lipoproteins is the hallmark of the dyslipidemia in the metabolic syndrome. *Arterioscler Thromb Vasc Biol* 28: 1225–1236.
- Richter S, Shih DQ, Pearson ER, Wolfrum C, Fajans SS, et al. (2003) Regulation of apolipoprotein M gene expression by MODY3 gene hepatocyte nuclear factor-1 $\alpha$ : haploinsufficiency is associated with reduced serum apolipoprotein M levels. *Diabetes* 52: 2989–2995.
- Ooi EM, Watts GF, Chan DC, Nielsen LB, Plomgaard P, et al. (2010) Association of apolipoprotein M with high-density lipoprotein kinetics in overweight-obese men. *Atherosclerosis* 210: 326–330.
- Dullaart RP, Plomgaard P, de Vries R, Dahlback B, Nielsen LB (2009) Plasma apolipoprotein M is reduced in metabolic syndrome but does not predict intima media thickness. *Clin Chim Acta* 406: 129–133.
- Xu N, Nilsson-Ehle P, Ahren B (2006) Suppression of apolipoprotein M expression and secretion in alloxan-diabetic mouse: Partial reversal by insulin. *Biochem Biophys Res Commun* 342: 1174–1177.
- Zhang X, Jiang B, Luo G, Nilsson-Ehle P, Xu N (2007) Hyperglycemia down-regulates apolipoprotein M expression in vivo and in vitro. *Biochim Biophys Acta* 1771: 879–882.
- Xu N, Ahren B, Jiang J, Nilsson-Ehle P (2006) Down-regulation of apolipoprotein M expression is mediated by phosphatidylinositol 3-kinase in HepG2 cells. *Biochim Biophys Acta* 1761: 256–260.
- Waitzberg DL, Torrinhas RS, Jacintho TM (2006) New parenteral lipid emulsions for clinical use. *JPEN J Parenter Enteral Nutr* 30: 351–367.
- Kroggh-Madsen R, Plomgaard P, Akerstrom T, Moller K, Schmitz O, et al. (2008) Effect of short-term intralipid infusion on the immune response during low-dose endotoxemia in humans. *Am J Physiol Endocrinol Metab* 294: E371–379.

Lentiviruses can permanently integrate genetic material into the genome of host cells, are able to maintain long-term expression of integrated material, and can be used to express transgenes and to suppress the expression of endogenous genes by RNA interference (RNAi) [41]. In the present study, we constructed lentiviral vectors to overexpress the human ApoM gene in order to increase ApoM expression in 293T cells and in GK rats. Our results demonstrated that MAPK8 was significantly down-regulated, while ApoM mRNA levels in 293T cells transfected with human ApoM gene were dramatically increased. MAPK8 is known as the c-Jun NH $_2$ -terminal kinase 1 (JNK1), and is involved in the mechanism of obesity-induced insulin resistance [29]. Feeding a high fat diet can cause activation of the JNK1 signaling pathway, insulin resistance, and obesity in mice [29]. Interestingly, ITT analysis showed that human ApoM expression in GK rats had a tendency to enhance the glucose-lowering effects of exogenous insulin, suggesting that overexpression of ApoM might improve insulin sensitivity. However, further studies are needed to confirm this observation and to clarify the mechanism.

In conclusion, the present study demonstrated that Intralipid could increase plasma FFA levels, decreased insulin sensitivity and suppressed ApoM expression. Moreover, Intralipid could enhance a number of genes involved in insulin signaling. Our results suggest that ApoM overexpression may have a potential role in improving insulin resistance in *vivo*, and could be considered as a future therapeutic target against insulin resistance and type 2 diabetes.

## Supporting Information

**Table S1.**  
(DOC)

## Author Contributions

Conceived and designed the experiments: GHL NX. Performed the experiments: LZ YHF YPS JZ QFM. Analyzed the data: LQ MBS PNE XYZ. Contributed reagents/materials/analysis tools: XYZ. Wrote the paper: GHL NX LZ.



20. Choi CS, Lee FN, Youn JH (2001) Free fatty acids induce peripheral insulin resistance without increasing muscle hexosamine pathway product levels in rats. *Diabetes* 50: 418–424.
21. Brehm A, Krssak M, Schmid AI, Nowotny P, Waldhausl W, et al. (2010) Acute elevation of plasma lipids does not affect ATP synthesis in human skeletal muscle. *Am J Physiol Endocrinol Metab* 299: E33–38.
22. Itani SI, Ruderman NB, Schmieder F, Boden G (2002) Lipid-induced insulin resistance in human muscle is associated with changes in diacylglycerol, protein kinase C, and IkappaB-alpha. *Diabetes* 51: 2005–2011.
23. Gosmanov AR, Smiley DD, Peng L, Siquiera J, Robalino G, et al. (2012) Vascular effects of intravenous intralipid and dextrose infusions in obese subjects. *Metabolism* 61: 1370–1376.
24. Chow LS, Seaquist ER, Eberly LE, Mashek MT, Schimke JM, et al. (2012) Acute free fatty acid elevation eliminates endurance training effect on insulin sensitivity. *J Clin Endocrinol Metab* 97: 2890–2897.
25. Liu J, Jahn LA, Fowler DE, Barrett EJ, Cao W, et al. (2011) Free fatty acids induce insulin resistance in both cardiac and skeletal muscle microvasculature in humans. *J Clin Endocrinol Metab* 96: 438–446.
26. Sevvana M, Ahnstrom J, Egerer-Sieber C, Lange HA, Dahlback B, et al. (2009) Serendipitous fatty acid binding reveals the structural determinants for ligand recognition in apolipoprotein M. *J Mol Biol* 393: 920–936.
27. Kim JK (2009) Hyperinsulinemic-euglycemic clamp to assess insulin sensitivity in vivo. *Methods Mol Biol* 560: 221–238.
28. Luo G, Zhang X, Mu Q, Chen L, Zheng L, et al. (2010) Expression and localization of apolipoprotein M in human colorectal tissues. *Lipids Health Dis* 9: 102.
29. Sabio G, Davis RJ (2010) cJun NH2-terminal kinase 1 (JNK1): roles in metabolic regulation of insulin resistance. *Trends Biochem Sci* 35: 490–496.
30. Goto Y, Suzuki K, Sasaki M, Ono T, Abe S (1988) GK rat as a model of non-obese non-insulin-dependent diabetes: selective breeding over 35 generations. In: Shafir E, Renold A, editors. *Frontiers in Diabetes Research Lessons from Animal Diabetes*. London: John Libbey. pp.301–303.
31. Ogata M, Uchimura T, Iizuka Y, Murata R, Suzuki S, et al. (1997) Effect of non-insulin dependent diabetes on cyclosporin A disposition in Goto-Kakizaki (GK) rats. *Biol Pharm Bull* 20: 1026–1029.
32. Griffin ME, Marcucci MJ, Cline GW, Bell K, Barucci N, et al. (1999) Free fatty acid-induced insulin resistance is associated with activation of protein kinase C theta and alterations in the insulin signaling cascade. *Diabetes* 48: 1270–1274.
33. Christoffersen C, Obinata H, Kumaraswamy SB, Galvani S, Ahnstrom J, et al. (2011) Endothelium-protective sphingosine-1-phosphate provided by HDL-associated apolipoprotein M. *Proc Natl Acad Sci U S A* 108: 9613–9618.
34. Else S, Ahnstrom J, Christoffersen C, Hoofnagle AN, Plomgaard P, et al. (2012) Apolipoprotein M binds oxidized phospholipids and increases the antioxidant effect of HDL. *Atherosclerosis* 221: 91–97.
35. Wolfrum C, Poy MN, Stoffel M (2005) Apolipoprotein M is required for prebeta-HDL formation and cholesterol efflux to HDL and protects against atherosclerosis. *Nat Med* 11: 418–422.
36. Umpierrez GE, Smiley D, Robalino G, Peng L, Kitabchi AE, et al. (2009) Intravenous intralipid-induced blood pressure elevation and endothelial dysfunction in obese African-Americans with type 2 diabetes. *J Clin Endocrinol Metab* 94: 609–614.
37. Kashyap SR, Ioachimescu AG, Gornik HL, Gopan T, Davidson MB, et al. (2009) Lipid-induced insulin resistance is associated with increased monocyte expression of scavenger receptor CD36 and internalization of oxidized LDL. *Obesity (Silver Spring)* 17: 2142–2148.
38. Kim MY, Bac JS, Kim TH, Park JM, Ahn YH (2012) Role of transcription factor modifications in the pathogenesis of insulin resistance. *Exp Diabetes Res* 2012: 716425.
39. Mounier C, Posner BI (2006) Transcriptional regulation by insulin: from the receptor to the gene. *Can J Physiol Pharmacol* 84: 713–724.
40. Zhang Y, Xiao M, Niu G, Tan H (2005) Mechanisms of oleic acid deterioration in insulin secretion: role in the pathogenesis of type 2 diabetes. *Life Sci* 77: 2071–2081.
41. Dissen GA, Lomniczi A, Neff TL, Hobbs TR, Kohama SG, et al. (2009) In vivo manipulation of gene expression in non-human primates using lentiviral vectors as delivery vehicles. *Methods* 49: 70–77.