



LUND UNIVERSITY

Complete genome sequences of three novel human papillomavirus types, 175, 178, and 180.

Johansson, Hanna K; Forslund, Ola

Published in:
Genome Announcements

DOI:
[10.1128/genomeA.00443-14](https://doi.org/10.1128/genomeA.00443-14)

2014

[Link to publication](#)

Citation for published version (APA):
Johansson, H. K., & Forslund, O. (2014). Complete genome sequences of three novel human papillomavirus types, 175, 178, and 180. *Genome Announcements*, 2(3). <https://doi.org/10.1128/genomeA.00443-14>

Total number of authors:
2

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Complete Genome Sequences of Three Novel Human Papillomavirus Types, 175, 178, and 180

Hanna Johansson, Ola Forslund

Department of Laboratory Medicine, Medical Microbiology, Lund University, Malmö, Sweden

We report the characterization of three novel human papillomavirus (HPV) types of the genus *Gammapapillomavirus*. HPV175 and HPV180 were isolated from a condyloma. HPV178 was isolated from healthy skin adjacent to an actinic keratosis.

Received 22 April 2014 Accepted 5 May 2014 Published 22 May 2014

Citation Johansson H, Forslund O. 2014. Complete genome sequences of three novel human papillomavirus types, 175, 178, and 180. *Genome Announc.* 2(3):e00443-14. doi:10.1128/genomeA.00443-14.

Copyright © 2014 Johansson and Forslund. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Ola Forslund, ola.forslund@med.lu.se.

We report the complete genome sequences of three novel human papillomavirus (HPV) types of the genus *Gammapapillomavirus*. HPV175 (SE87) and HPV180 (FA69) were originally detected as complete genomes in a condyloma swab sample from a 30-year-old male (1). HPV178 was isolated from a swab of healthy skin, next to an actinic keratosis of an 86-year-old male, and discovered after an attempt to amplify a closely related virus (SE46 [GenBank accession number JX198657]).

Briefly, multiple-displacement amplification (MDA)-amplified HPV DNA (1) was reamplified using the PrimeSTAR GXL DNA polymerase kit (TaKaRa Bio, Shiga, Japan). Primers were designed using the Primer Express Software v. 3.0 (Applied Biosystems). HPV175 (7,226 bp) was amplified in three parts, amplicon 1 (3,677 bp) (forward primer [fwd] 5'-ACAAATCTCTGGGAGGA CTAATGC-3' and reverse primer [rev] 5'-GCCTCCAGTTCTTC TATAGTTCCCTT-3'), amplicon 2 (2,397 bp) (fwd 5'-CGCATG CCATGTTTGTCTG-3' and rev 5'-GGCCTGATTCATCTTGG ATATCTT-3'), and amplicon 3 (2,601 bp) (fwd 5'-ATTTGGAA ATGTTGTGATGAGTGAAA-3' and rev 5'-TTGAGCCTCTCA ATAGGTGTG-3') (Eurofins MWG Operon, Germany). HPV178 (7,314 bp) was obtained as a single amplicon (fwd 5'-GTTGGTG GTGACCGAGTTTACTTT-3' and rev 5'-TTCATTAATACAGG CCATTATAATCTACAAGT-3') (Eurofins). HPV180 (7,356 bp) was obtained as a single amplicon (fwd 5'-TATTTGCACAAGGT GCACCAG-3' and rev 5'-AAGGAAAGGTCAGAAAAGAGAAG CT-3') (DNA Technology, Denmark).

Amplicons were cloned using a TOPO TA cloning kit and the pCR 2.1-TOPO vector (Invitrogen, Carlsbad, CA) and sequenced using primer walking (Eurofins).

Pairwise comparisons of the L1 open reading frames (ORF) of HPV175, HPV178, and HPV180 demonstrated closest nucleotide homology to HPV60 (68.7%, species 4), HPV65 (67.6%, species 1), and HPV121 (81.8%, species 10), respectively.

HPV175, HPV178, and HPV180 all demonstrate a typical genome organization of cutaneous HPVs, with an upper regulatory region (URR), five early genes (E1, E2, E4, E6, and E7), and two late genes (L1 and L2). In the URR of HPV175 (488 bp) we identified five consensus E2-binding sites (ACC-N₆-GGT), one putative TATA box (TATAA), and one putative polyadenylation site

(ATATAA). In the URR of HPV178 (456 bp), three E2-binding sites were present, as well as one putative TATA box and one putative polyadenylation site. In the URR of HPV180 (544 bp), we identified four consensus E2-binding sites, three putative TATA boxes, and one putative polyadenylation site.

The putative E6 proteins contained two zinc finger domains [CxxC(x)₂₉CxxC] (2) separated by 36 amino acids. One zinc finger domain was also present in the E7 proteins. The LxCxE motif (binding site for the tumor suppressor retinoblastoma protein) (3) was observed in E7 of HPV178, whereas serine was substituted for cysteine in the corresponding domain (LxSxE) of HPV175 and HPV180. Among 55 E7 proteins of representative HPV types of the genus *Gammapapillomavirus*, 19 demonstrated the LxCxE motif and 27 the LxSxE motif.

Only the putative E1 protein of HPV178 had the conserved ATP-binding site (GPPDTGKS) (4, 5). For HPV175 we identified GPSDTGKS and for HPV180 GKPNTGKS. An initiation codon of the putative start of the E4 ORF was present in HPV175 and HPV178, whereas the corresponding codon was absent in the E4 ORF of HPV180. This is in agreement with the HPV types of species 10 of the genus *Gammapapillomavirus* (6).

Nucleotide sequence accession numbers. The complete genomic sequences are available in GenBank under these accession numbers: HPV175 (KC108721), HPV178 (KJ130020), and HPV180 (KC108722).

ACKNOWLEDGMENTS

This project was supported by BioCARE, a Strategic Research Program at Lund University, Sweden, and by the Swedish Cancer Society.

REFERENCES

- Johansson H, Bzhalava D, Ekström J, Hultin E, Dillner J, Forslund O. 2013. Metagenomic sequencing of "HPV-negative" condylomas detects novel putative HPV types. *Virology* 440:441–447. <http://dx.doi.org/10.1016/j.virol.2013.01.023>.
- Ullman CG, Haris PI, Galloway DA, Emery VC, Perkins SJ. 1996. Predicted alpha-helix/beta-sheet secondary structures for the zinc-binding motifs of human papillomavirus E7 and E6 proteins by consensus prediction averaging and spectroscopic studies of E7. *Biochem. J.* 319(Pt 1): 229–239.

3. Dahiya A, Gavin MR, Luo RX, Dean DC. 2000. Role of the LXCXE binding site in Rb function. *Mol. Cell. Biol.* **20**:6799–6805. <http://dx.doi.org/10.1128/MCB.20.18.6799-6805.2000>.
4. Iyer LM, Leipe DD, Koonin EV, Aravind L. 2004. Evolutionary history and higher order classification of AAA+ ATPases. *J. Struct. Biol.* **146**: 11–31. <http://dx.doi.org/10.1016/j.jsb.2003.10.010>.
5. Liu X, Schuck S, Stenlund A. 2010. Structure-based mutational analysis of the bovine papillomavirus E1 helicase domain identifies residues involved in the nonspecific DNA binding activity required for double trimer formation. *J. Virol.* **84**:4264–4276. <http://dx.doi.org/10.1128/JVI.02214-09>.
6. Ure AE, Forslund O. 2014. Characterization of human papillomavirus type 154 and tissue tropism of gammapapillomaviruses. *PLoS One* **9**:e89342. <http://dx.doi.org/10.1371/journal.pone.0089342>.