



LUND UNIVERSITY

HIV prevalence and risk behaviours among people who inject drugs in Iran: the 2010 National Surveillance Survey

Khajehkazemi, Razieh; Osooli, Mehdi; Sajadi, Leily; Karamouzian, Mohammad; Sedaghat, Abbas; Fahimfar, Noushin; Safaie, Afshin; Mostafavi, Ehsan; Haghdoost, Ali-Akbar

Published in:
Sexually Transmitted Infections

DOI:
[10.1136/sextrans-2013-051204](https://doi.org/10.1136/sextrans-2013-051204)

2013

[Link to publication](#)

Citation for published version (APA):

Khajehkazemi, R., Osooli, M., Sajadi, L., Karamouzian, M., Sedaghat, A., Fahimfar, N., Safaie, A., Mostafavi, E., & Haghdoost, A.-A. (2013). HIV prevalence and risk behaviours among people who inject drugs in Iran: the 2010 National Surveillance Survey. *Sexually Transmitted Infections*, 89, 29-32. <https://doi.org/10.1136/sextrans-2013-051204>

Total number of authors:
9

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

HIV Prevalence and Risk Behaviors among People Who Inject Drugs in Iran: The 2010 National Surveillance Survey

Razieh Khajehkazemi¹, Mehdi Osooli^{1,2}, Leily Sajadi¹, Mohammad Karamouzian¹, Abbas Sedaghat³, Noushin Fahimfar³, Afshin Safaie⁴, Ehsan Mostafavi^{5, 1}, Ali-Akbar Haghdoost^{6, 1}

¹Regional Knowledge Hub, and WHO Collaborating Centre for HIV Surveillance, Institute for Futures Studies in Health, Kerman University of Medical Sciences, Kerman, Iran

²Centre for Haemostasis and Thrombosis, Skane University Hospital, Lund University, Sweden

³HIV/AIDS Control Office, Center for Disease Control (CDC), Ministry of Health and Medical Education, Tehran, Iran

⁴Director of Laboratories, Alborz Medical University, Karaj, Iran

⁵Department of Epidemiology, Pasteur Institute of Iran, Tehran, Iran

⁶Research Center for Modeling in Health, , Institute for Futures Studies in Health, Kerman University of Medical Sciences, Kerman, Iran

Key Words: HIV, People Who Inject Drugs, Surveillance, Risk Behaviors, Iran.

Word count: Abstract: 167, Text: 1425 words.

Number of tables: 1

Number of figures: 0

Corresponding author: Ali-Akbar Haghdoost, MD, PhD of Epidemiology and Biostatistics, Research Center for Modeling in Health, Institute for Futures Studies in Health, Kerman University of Medical Sciences, Avicenna Ave., Jahad Blvd., Postal Code: 7619813159, Kerman, Iran.

Telephone: + (98) 341-2263787. Fax: + (98) 341-2263725. Emails: ahaghdoost@kmu.ac.ir, Ali-Akbar.Haghdoost@lshtm.ac.uk

Abstract

Objectives: To assess the prevalence of HIV and related risk behaviors among People Who Inject Drugs (PWID) in Iran.

Methods: We conducted a national cross-sectional bio-behavioral surveillance survey between March and July 2010, interviewing male PWID from a geographically dispersed sample through a facility-based sampling method.

Results: We recruited 2480, and tested 2290 PWID. The overall prevalence of HIV was 15.2% (95% CI 9.7-23.1). Among those who had injected drugs over the last month, 36.9% had used a non-sterile needle, and 12.6% had practiced shared injection. Over the past 12 months preceding the interview, 30.4% had sold sex for money, drugs, goods or a favor. In the multivariate analysis, the prevalence of HIV had a positive association with age, while having above high school education, and permanent job were protective.

Conclusion: Unsafe injection, and sexual risk behaviors were still frequent and the prevalence of HIV among PWID remains high. Intensified efforts are needed to prevent the further spread of HIV among Iranian PWID and their sexual partners.

Background

It is estimated that around 300,000 People Who Inject Drugs (PWID) are living in Iran.[1] National statistics show injection drug use is still the principal mode of HIV transmission in Iran.[2] From 1986 to 2006, around 64% of new HIV cases were reported as acquiring the infection through injecting drug use. Recent evidence show that the prevalence of HIV infection might be on a downward trend among in this group.[3] However, the first Iranian national Bio-Behavioral Surveillance Survey (BBSS1) of HIV among PWID conducted in 2008, found an alarmingly high HIV prevalence of 15.3% with high levels of risky injection and sexual behaviors.[4]

More effective prevention and care response requires tracking the key features of the HIV epidemic. This paper, therefore presents the results of the second round of BBSS (BBSS2) among male PWID in Iran thereby establishing the current status of HIV infection and gauging the potential for further spread due to injection and sexual risk behaviors.

Methods

Setting and Participants

This cross-sectional survey was conducted in the same ten provinces (out of 31) as in BBSS1. Based on the trend of identified HIV cases among PWID during 2000- 2003, provinces were categorized in two strata: one stratum included the provinces with an upward trend and the other with a downward or stable trend. From each stratum, five provinces were purposefully selected to provide the most disperse geographic representativeness.

In the current survey, between March and July 2010, using a facility-based sampling approach, PWID were recruited from drop-in centers, shelters, drug treatment centers, voluntary counseling and testing centers, and outreach spots in each selected province. Having the inputs from the officers in charge of the HIV surveillance system in each province, we selected minimum five different facilities in the capital city and neighborhood cities (if feasible). Participants were recruited by convenience sampling at the center regardless of their known or unknown HIV status and their verbal informed consent was requested to participate in the study. Eligible participants were men ≥ 18 years of age, who had injected drug at least once during the past 12 months.

Data Collection

A structured questionnaire was used to collect data on the demographic characteristics, and the key indicators of injection and sexual risk behaviors. Provincial supervisors trained and monitored the facility's staff to recruit, interview, and collect Dried Blood Samples (DBS) based on a standard protocol. Upon completion of the behavioral survey, each subject was given an incentive of 1.5 USD (Tehran 2.5 USD) in cash, and if agreed to be tested for HIV, they would

get an additional 0.5 USD. DBS samples were tested for HIV antibodies by ELISA (using bioMérieux Vironostika Uni-Form II Ag/Ab). All positives and 10% of the negative samples were rechecked in the Pasteur reference laboratory (using Bio-Rad Genscreen Plus HIV Ag-Ab); the Kappa coefficient was 95.8% . .

Statistical Analysis

Multivariate logistic regression models were applied to determine the factors associated with HIV infection. Variables with a p-value less than 0.2 in the bivariate analysis were entered into the multivariate model. In order to adjust for the clustering effects within facilities and provinces, and also to weight for the size of facilities, the survey function in stata (version 10) was used.

Results

A total of 2518 PWID were asked to participate in this study. Of whom 27 individuals did not meet the eligibility criteria, and 11 refused to provide informed consent (Overall response rate 97.5%). Out of 2480 PWID, 2417 provided informed consent for HIV testing (response rate 97.5%). We were unable to complete the HIV test on 127 DBS as the specimen was not sufficient for HIV testing.

The overall prevalence of HIV was 15.2% (95% CI: 9.7-23.1); the highest and the lowest prevalence in different provinces were 31.9% and 2.2%, respectively. Among subgroups, unemployed PWID had the highest prevalence (20.1%), and those with above high school education had the lowest (4.6%). The mean (standard deviation) age at the time of study, age at first drug use and age at first drug injection were 34.6 (8.9), 18.7 (5.2), and 25.9 (7.2) years, respectively. Over the past month, among those who had injected drugs, 36.9% had used a non-sterile needle and, 12.6% had practiced shared injection. Over the past 12 months, 30.4% had sold sex for money, drugs, goods or a favor. In the multivariate model, HIV was significantly associated with current age, the level of education, and job status. Compared to 18-25 age group, the adjusted odds ratio [AOR] of infection were 3.1 and 4.1 in 26-35 and >35 years age groups, respectively. Regarding the education level, $AOR_{(\text{secondary and high school/ above high school})}$, $AOR_{(\text{primary/ above high school})}$, and $AOR_{(\text{illiterate/ above high school})}$ were 3.2, 3.1, and 2.7, respectively. Having temporary job, and unemployment increased the AORs as well (3.1 and 4.6, respectively)(Table 1).

Discussion

This survey showed that the prevalence of HIV among PWID in Iran is still alarmingly high, and behaviors for acquisition and transmission of HIV are common. The prevalence of HIV in BBSS1(2008) and what we observed in this survey was remarkably similar (15.3% versus 15.2%).[4] However, the apparently stable prevalence of HIV must be considered as a dynamic balance of several competing or offsetting factors, including new infections against mortality of HIV-infected PWID, in against out migration, initiation against discontinuation of injection and other factors such as incarceration and improved survival with anti-retroviral therapy. Nonetheless, our data make a strong case that HIV transmission will remain a major health problem throughout Iran for the years to come.

In other high risk groups in Iran such as female sex workers, still rather low levels of HIV prevalence (below 5%) are found compared to PWID.[5] Our findings confirm that PWID are still the most affected high risk group by HIV in Iran with a prevalence of about 15%. Though this level of prevalence among PWID is rather high compared to other countries in the Middle East and North Africa, it is still lower than those found in other countries including Libya and Pakistan.[6-7] This might be partly explained by our method of sampling recruiting PWID from the facilities; however, other studies which recruited individuals through respondent driven sampling have also reported high prevalence of HIV (25%) among PWID.[8]

Although there has been a history of prevention efforts among PWID in Iran, our findings potentially point to a sustained level of risky injection among PWID. While a large proportion of participants reported the use of sterile needles and syringes in their last injection, a considerable

fraction still practice shared injections. Indeed, almost 40% of participants reported a history of using previously used needles during the month before the interview. Possible explanations may be peer pressure among PWID[9-10] or having poor access to harm reduction services where and when they are needed the most.

In addition to risky injecting behaviors, sexual risk behaviors were relatively common among PWID. We observed selling sex in exchange with money, drugs, goods or a favor during the previous year were common; an issue that is difficult to address in the Iranian context. Iranian law outlaws male-male sex which makes any explicit intervention regarding this behavior even harder. Compounding the situation is that more than 60% of the participants reported not having used a condom in their last sexual encounter with a client. We think practical approaches are needed to slow down this potential mode of transmission among this population.

We acknowledge the limitations of our study. Our main sampling method (convenience sampling of those who seek services at the facilities) may limit the generalization of the findings to all PWID in Iran, particularly those who are hidden and not linked to services. However, we tried to address such bias by recruiting from venues and hotspots of PWID through out-reach activities. This method of sampling and recruitment is feasible given the geography and time constraints (more than 10 provinces over 4 months). It is also similar to that used in BBSS1 in 2008, thereby facilitating comparison with the results of that round.[4]. As is common in behavioral surveys, social desirability may have affected the reporting of sensitive behaviors. Some sort of calibration may be needed to address this issue.[11]

In conclusion, injection and sexual risks are common among PWID and they still have the highest prevalence of HIV infection among the key populations in Iran. This high prevalence of

HIV and its associated risky behaviors among PWID raise an alarm for the health authorities in Iran to design and implement effective and timely interventions to prevent further transmission among this population and to their partners.

Key messages

- ▶ PWID continue to have the highest HIV prevalence in Iran.
- ▶ Unsafe injection and sex are still common among Iranian PWID.
- ▶ Targeted HIV interventions tailored for the specific needs of PWID are critical to prevent further transmission of HIV among this high risk group and to other sub-populations.

Acknowledgements

The authors are grateful to the following people for their scientific and administrative contributions to the study: Dr. Omid Zamani, Dr. Ali Mirzazadeh, Ms. Soodeh Arabnejad, Dr. Soodabeh Navadeh, Dr. Kianoosh Kamali, Dr. Mohammad Reza Aghasadeghi, Mr. Farhang Vahabpour, Dr. Kayhan Azadmanesh, and Dr. Farzaneh Zolala. Special thanks are also due to the field supervisors and interviewers for their valuable work and to Dr. Majid Rezazadeh from the State Welfare Organization of Iran and the staff of the AIDS and Sexually Transmitted Diseases Office of the Ministry of health for providing logistics support and expert opinions. In addition, we highly appreciate the significant contributions of Dr Laith Abu-Raddad and Dr Willi McFarland in editing the manuscript.

Conflict of interest

None.

Contributors list

AH supervised the project and led the analysis and manuscript development. MO participated in writing the proposal, project management, and drafted the first version of the manuscript. RK and MK carried out the statistical analysis and contributed to the development of the manuscript. All authors had contribution in the data collection process. In addition, all authors read and approved the final version of this manuscript.

Funding

The study was funded by the United Nations Development Programme's (UNDP) the primary recipient of funds from the Global Fund for AIDS, Tuberculosis, and Malaria projects in Iran (grant number IC/10/26).

Ethics approval

The ethics committee of Kerman University of Medical Sciences reviewed and approved the study's protocol.

References:

1. Razzaghi EM, Movaghar AR, Green TC, et al. Profiles of risk: a qualitative study of injecting drug users in Tehran, Iran. *Harm Reduct J* 2006;**3**:12.
2. Nasirian M, Doroudi F, Gouya MM, et al. Modeling of Human Immunodeficiency Virus Modes of Transmission in Iran. *J Res Health Sci* 2012;**12**:81-87.
3. Rahimi-Movaghar A, Amin-Esmaeili M, Haghdoost AA, et al. HIV prevalence amongst injecting drug users in Iran: a systematic review of studies conducted during the decade 1998-2007. *Int J Drug Policy* 2012;**23**:271-8.
4. Islamic Republic of Iran, AIDS Progress Report on Monitoring of the United Nations General Assembly Special Session on HIV and AIDS, March 2012. Available online: http://www.unaids.org/en/dataanalysis/knowyourresponse/countryprogressreports/2012countries/IRIran%20AIDS%20Progress%20Report%202012%20English%20final1_1.pdf.
5. Mirzazadeh A, Nedjat S, Navadeh S, et al. HIV and related risk behaviors among female sex workers in Iran: bias-adjusted estimates from the 2010 national bio-behavioral survey. *AIDS Behav* 2013:1-6.
6. Emmanuel F, Salim M, Akhtar N, et al. Second-generation surveillance for HIV/AIDS in Pakistan: results from the 4th round of integrated behavior and biological survey 2011–2012. *Sex Transm Infect* Published Online First: 2 August 2013.
7. Mirzoyan L, Berendes S, Jeffery C, et al. New evidence on the HIV epidemic in Libya: Why countries must implement prevention programs among people who inject drugs. *J Acquir Immune Defic Syndr* 2013;**62**:577-583.
8. Malekinejad M, Mohraz M, Razani N, et al. HIV and related risk behaviors of injecting drug users (IDU) in Iran: findings from the first respondent-driven sampling (RDS) survey of IDU in Tehran in 2006-2007. *AIDS 2008 - XVII International AIDS Conference: Abstract no. THAC0202*. <http://www.iasociety.org/Abstracts/A200720902.aspx> (accessed 7 Aug 2013).

9. Ball AL. HIV, injecting drug use and harm reduction: a public health response. *Addiction* 2007;**102**:684-690.
10. Gjeruldsen S, Myrvang B, and Opjordsmoen S. Risk factors for drug addiction and its outcome. A follow-up study over 25 years. *Nord J Psychiatry* 2003;**57**:373-6.
11. Mirzazadeh A, Mansournia M-A, Nedjat S, et al. Bias analysis to improve monitoring an HIV epidemic and its response: approach and application to a survey of female sex workers in Iran. *J Epidemiol Community Health* 2013.

Table 1. Socio-demographic characteristics, risk behaviors, and HIV infection among people who inject drugs in Iran, 2010

	Percent	HIV prevalence N (%)	Bivariate Model		Multivariate Model	
			OR (95%CI)	p-value	OR (95%CI)	p-value
Current age (years)						
18-25	13.9 (11.6- 16.6)	7.8 (3.3- 17.4)	Ref	----	Ref	----
26-35	46.6 (43.9- 49.3)	15.8 (10.1- 23.8)	2.3 (1.2- 4.4)	0.017	3.1 (1.6- 6.1)	0.001
>=36	39.5 (35.8- 43.3)	17.0 (10.5- 26.4)	2.4 (1.2- 5.1)	0.018	4.1 (1.7- 9.5)	0.002
Age at first drug use (years)						
<= 18	56.7 (51.5- 61.8)	14.2 (9.9- 19.9)	Ref	----	b	b
>18	43.3 (38.2- 48.5)	16.1 (8.5- 28.4)	1.2 (0.6- 2.1)	0.617	b	b
Age at first drug injection (years)						
<= 25	57.0 (52.9- 60.9)	16.4 (10.1- 25.6)	Ref	----	Ref	----
>25	43.0 (39.0- 47.1)	13.2 (8.6- 19.6)	0.8 (0.6- 1.1)	0.164	0.7 (0.5- 1.1)	0.106
Marital Status						
Currently married	31.2 (27.4- 35.3)	13.2 (7.1- 23.2)	Ref	----	Ref	----
Ever been married (divorced and widowed)	21.5 (18.7- 24.6)	16.7 (10.4- 25.9)	1.3 (0.1-1.9)	0.164	1.1 (0.7- 1.9)	0.605
Single	47.3 (43.6- 51.0)	15.9 (10.2- 23.8)	1.3 (0.8-2.1)	0.339	1.4(0.7- 2.6)	0.259
Education						
Above high school	4.5 (3.1- 6.6)	4.6 (1.4- 13.7)	Ref	----	Ref	----
Secondary and High school	65.2 (60.6- 69.6)	15.6 (9.6- 24.4)	3.8 (1.2-12.1)	0.024	3.2 (1.1- 9.4)	0.039
Primary school	24.6 (21.3- 28.2)	16.2 (9.6- 26.0)	3.9 (1.2-12.6)	0.024	3.1 (1.1- 8.7)	0.036
Illiterate	5.7 (3.5- 9.2)	14.7 (7.9- 25.6)	3.4 (0.9-12.6)	0.063	2.7 (0.7-9.8)	0.125
Job status						
Permanent job	12.3 (7.5- 19.7)	5.1 (1.9- 12.5)	Ref	----	Ref	----
Temporary job	48.9 (43.7- 54.1)	13.8 (9.6- 19.6)	3.3 (1.2-8.7)	0.020	3.1 (1.2- 8.1)	0.025
Unemployed	38.8 (31.9- 46.2)	20.1 (10.9- 33.9)	5.2 (1.6-16.4)	0.006	4.6 (1.6- 13.6)	0.007
Having at least one injection in the last month						
No	38.2 (29.2- 48.1)	11.0 (6.8- 17.2)	Ref	----	Ref	----
Yes	61.8 (51.8- 70.8)	17.8 (10.1- 29.4)	1.7 (0.8- 3.8)	0.178	1.7 (0.8- 3.6)	0.184
Using a used needle or syringe for injection in the last month *						
No	63.1 (55.0- 70.6)	17.9 (9.6- 30.9)	Ref	----	b	b
Yes	36.9 (29.4- 44.9)	17.6 (10.1- 28.9)	0.9 (0.6- 1.5)	0.903	b	b
Using a sterile needle or syringe at last injection						
No	11.1 (8.3- 14.6)	19.0 (10.6- 31.5)	Ref	----	b	b
Yes	88.9 (85.4- 91.7)	14.7 (9.1- 22.9)	0.8 (0.5- 1.3)	0.368	b	b
Sharing injection tools with others over the last month*						
No	87.4 (83.2- 90.7)	18.5 (10.2- 31.4)	Ref	----	b	b
Yes	12.6 (9.3- 16.8)	13.5 (7.9- 22.2)	0.7 (0.3-1.5)	0.353	b	b
Sold sex during last 12 months						
No	69.6	195 (15.1)	Ref	----	b	b
Yes	30.4	78 (13.6)	0.9 (0.6- 1.2)	0.392	b	b
Condom use at last sexual intercourse**						
No	61.7	47 (13.4)	Ref	----	b	b
Yes	38.3	29 (13.2)	0.9 (0.5- 2.1)	0.976	b	b

* Among those who had at least one injection in the last month.

** Among those who sold sex in exchange with money, drug, goods, or any favor during the last 12 months.

b: Not entered into the multivariate model (p-value in bivariate model was >0.2).