



# LUND UNIVERSITY

## Functional Modeling of Genes Upregulated in Chronic Myeloid Leukemia

Hansen, Nils

2013

[Link to publication](#)

*Citation for published version (APA):*

Hansen, N. (2013). *Functional Modeling of Genes Upregulated in Chronic Myeloid Leukemia*. [Doctoral Thesis (compilation), Division of Clinical Genetics]. Division of Clinical Genetics, Lund University.

*Total number of authors:*

1

### General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00

# Functional Modeling of Genes Upregulated in Chronic Myeloid Leukemia

Nils Hansen



**LUND**  
UNIVERSITY

DOCTORAL DISSERTATION

With the approval of the Lund University Faculty of Medicine, this thesis will be defended on November 22, 2013 at 13.00 in Segerfalksalen, Lund, Sweden.

*Faculty opponent*


Professor Ravi Bhatia, MD, PhD,  
City of Hope National Medical Center,  
California, USA

Organization LUND UNIVERSITY		Document name DOCTORAL DISSERTATION	
Department of Clinical Genetics BMC B13, Klinikgatan 26 SE-221 84 Lund Sweden		Date of issue November 22, 2013	
Author(s) Nils Hansen		Sponsoring organization	
Title and subtitle Functional modeling of genes upregulated in chronic myeloid leukemia			
Abstract Chronic myeloid leukemia (CML) is caused by the transformation of a primitive hematopoietic cell by the BCR/ABL1 fusion gene that is formed through the chromosomal translocation t(9;22). CML is currently successfully treated with tyrosine kinase inhibitors targeting the ABL1 kinase domain. However, the CML stem cells are insensitive to this drug and a large fraction of patients will have relapse following discontinuation of the drug. Thus, improved therapeutic strategies are needed towards the ultimate goal of curing CML. In Article I, the functional role of suppressor of cytokine signaling 2 (SOCS2) in CML and in normal hematopoiesis was investigated. Mice deficient for the Socs2 gene displayed normal steady-state hematopoiesis and hematopoietic stem cell (HSC) function. Transduction of bone marrow (BM) cells from Socs2-deficient mice with BCR/ABL1 and subsequent transplantation resulted in a CML-like disease indistinguishable from the disease in control mice, suggesting that SOCS2 is dispensible for normal hematopoiesis and CML pathogenesis. In Article II, global gene expression analysis and subsequent flow cytometric analysis of normal and CML BM cells showed that interleukin-1 receptor accessory protein (IL1RAP) was highly upregulated on the cell surface of CML cells, allowing prospective separation of candidate CML stem cells from normal hematopoietic stem cells. The use of a polyclonal IL1RAP antibody enabled antibody-dependent cellular cytotoxicity (ADCC) of CD34+CD38- CML cells in vitro. In Article III, the function of IL1RAP in normal hematopoiesis and in CML was investigated. Mice lacking IL1rap displayed lower myeloid steady-state cell counts. Overexpression of IL1RAP in cord blood cells, followed by transplantation into immunodeficient mice, resulted in increased levels of myelopoeisis. Finally, CD34+CD38- CML cells were found to respond strongly to IL1B stimulation in vitro. Together, the results suggest that IL1RAP is a promising target for novel therapeutic approaches in CML.			
Key words: Chronic myeloid leukemia, SOCS2, IL1RAP, BCR/ABL1, leukemia models			
Classification system and/or index terms (if any):			
Supplementary bibliographical information:		Language English	
ISSN and key title: 1652-8220		ISBN 978-91-87449-97-0	
Recipient's notes		Number of pages 101	Price
		Security classification	

Distribution by (name and address)

I, the undersigned, being the copyright owner of the abstract of the above-mentioned dissertation, hereby grant to all reference sources permission to publish and disseminate the abstract of the above-mentioned dissertation.

Signature



Date October 17, 2013

# Functional Modeling of Genes Upregulated in Chronic Myeloid Leukemia

Nils Hansen



**LUND**  
UNIVERSITY

Department of Clinical Genetics  
Faculty of Medicine

Lund University  
2013

Copyright © Nils Hansen

LUND UNIVERSITY

Faculty of Medicine, Doctoral Dissertation Series 2013:124

ISBN 978-91-87449-97-0

ISSN 1652-8220

Printed in Sweden by Media-Tryck, Lund University

Lund 2013



**CLIMATE  
COMPENSATED  
PAPER**



**REPA**<sup>®</sup>  
A part of FNI (the Packaging and  
Newspaper Collection Service)

"A model is a lie that helps you see the truth."

Howard Skipper



# Table of contents

Table of contents	7
Abbreviations	9
Articles included in the thesis	11
Other related publications	13
Preface	15
Summary in Swedish	17
Introduction	21
Hematopoiesis	21
Cytokine signaling in hematopoiesis	23
Leukemia	25
Leukemia stem cells	25
Chronic myeloid leukemia	27
Clinical characteristics of chronic myeloid leukemia and current treatment options	27
Blast phase and known secondary genetic changes	30
CML stem cells	31
Signaling pathways in chronic myeloid leukemia	31
Experimental models of chronic myeloid leukemia	35
The present investigation	39
Aims of the study	39
Results and discussion	40
Article I	40
Article II	40
Article III	41



Conclusions	42
Article I	42
Article II	42
Article III	42
Concluding remarks and future perspectives	43
Acknowledgements	47
References	49
Articles I–III	71

# Abbreviations

ALL	Acute lymphoblastic leukemia
AML	Acute myeloid leukemia
AP	Advanced phase
BM	Bone marrow
BP	Blast phase
CLL	Chronic lymphoblastic leukemia
CLP	Common lymphoid progenitor
CML	Chronic myeloid leukemia
CP	Chronic phase
CSC	Cancer stem cell
FTI	Farnesyl protein transferase inhibitor
GH	Growth hormone
GMP	Granulocyte macrophage progenitor
HSC	Hematopoietic stem cell
LIC	Leukemia-initiating cell
LMPP	Lymphoid multi-potential progenitor
LSC	Leukemic stem cell
MEP	Megakaryocyte erythrocyte progenitor

MPP	Multi-potent progenitor
PB	Pheripheral blood
Ph chromosome	Philadelphia chromosome
QPCR	Quantitative polymerase chain reaction
SCT	Stem cell transplantation
TKI	Tyrosine kinase inhibitor
WBC	White blood cell counts

# Articles included in the thesis

This thesis is based on the following articles, which are referred to in the text by their Roman numerals (I–III).

- I **Hansen N**, Ågerstam H, Wahlestedt M, Landberg N, Askmyr M, Ehinger M, Rissler M, Lilljebjörn H, Johnels P, Ishiko J, Melo JV, Alexander WS, Bryder D, Järås M, Fioretos T. SOCS2 is dispensable for BCR/ABL1-induced chronic myeloid leukemia-like disease and for normal hematopoietic stem cell function. *Leukemia* 2013;27(1):130-135.
- II Järås M, Johnels P, **Hansen N**, Ågerstam H, Tsapogas P, Rissler M, Lassen C, Olofsson T, Bjerrum OW, Richter J, Fioretos T. Isolation and killing of candidate chronic myeloid leukemia stem cells by antibody targeting of IL-1 receptor accessory protein. *Proc Natl Acad Sci* 2010;107(37):16280–16285.
- III **Hansen N**, Reckzeh K, Ågerstam H, Askmyr M, Högberg C, Gordon S, Rissler M, Richter J, Järås M, Fioretos T. IL1RAP promotes myelopoiesis and its upregulation on candidate CML stem cells is associated with increased IL1 sensitivity. *Manuscript*.



## Other related publications

Askmyr M, Ågerstam H, Hansen N, Gordon S, Arvanitakis A, Rissler M, Juliusson G, Richter J, Järås M, Fioretos T. Selective killing of candidate AML stem cells by antibody targeting of IL1RAP. *Blood* 2013; 2;121(18):3709-3713.

Ågerstam H, Järås M, Andersson A, Johnels P, Hansen N, Lassen C, Rissler M, Gisselsson D, Olofsson T, Richter J, Fan X, Ehinger M, Fioretos T. Modeling the human 8p11-myeloproliferative syndrome in immunodeficient mice. *Blood* 2010;116(12): 2103-2111.



# Preface

Chronic myeloid leukemia (CML) is a myeloproliferative disorder arising from a hematopoietic stem cell that has acquired the Philadelphia chromosome, which was discovered in 1960 as the first chromosomal abnormality consistently associated with a specific type of cancer. The fusion gene *BCR/ABL1* is transcribed from the breakpoint of the Philadelphia chromosome and 40 years after its discovery, a targeted therapy (imatinib), which inhibits the BCR/ABL1 fusion protein, was introduced in the clinic. Despite the remarkable success of imatinib and other tyrosine kinase inhibitors as therapy for CML, the CML stem cells are insensitive to the treatment and following drug discontinuation, a large proportion of the patients will experience relapse. Thus, an improved understanding of the disease mechanisms and new therapeutic strategies will be needed for complete cure of CML.

The aim of the work in this thesis was to study molecular mechanisms responsible for the emergence of CML. For this, we used various models of CML and normal hematopoiesis to assay the function of specific genes found to be upregulated in CML. This thesis is divided into three sections, the first of which provides a background to the subject by discussing normal hematopoiesis, CML, and the different experimental models available for the study of CML. The second part describes the studies performed during the work for the thesis and the conclusions we have drawn from them. The third and last part constitutes the articles that this thesis is based upon.

Lund

October, 2013





# Summary in Swedish

Kroppens blodceller nybildas kontinuerligt ifrån blodstamceller (hematopoetiska stamceller) i benmärgen i en process som kallas hematopoies. Detta sker genom att stamcellerna ger upphov till både nya stamceller och till celler som mognar ut till olika typer av blodceller. Blodstamcellerna bildar en rad olika mogna celltyper med olika funktioner, så som syresättning av kroppens vävnader (röda blodkroppar, erythrocyter), förmåga att bekämpa infektioner (vita blodkroppar, leukocyter) och för att hejda blödningar (blodplättar, trombocyter). Cancer i den blodbildande benmärgen (leukemi) drabbar varje år ca 1200 personer i Sverige. Vid leukemi uppstår en okontrollerad delning av omogna blodceller, ofta i kombination med att dessa celler inte mognar ut eller dör undan på ett kontrollerat sätt. När detta sker uppstår brist på normala blodceller, vilket i sin tur bland annat kan leda till trötthet, blödningar, förstörd mjälte och ökad känslighet för infektioner. När celler delar sig organiseras allt DNA i 46 kromosomer och en typ av genetisk förändring som är vanlig i leukemier är så kallade translokationer, som uppstår när två kromosomer går sönder och felaktigt läker samman med varandra. I brytpunkterna mellan de olika kromosomdelarna, kan två olika gener sammanfogas och bilda en fusionsgen. En translokation mellan kromosomerna 9 och 22 kan skapa en fusionsgen där generna *BCR* och *ABL1* sammanfogas till *BCR/ABL1*, vilken cellen översätter till ett fusionsprotein som felaktigt aktiverar flera av cellens signalvägar. Om fusionsgenen *BCR/ABL1* uppstår i en blodstamcell kan den orsaka kronisk myeloisk leukemi (KML) som kännetecknas av en förhöjd produktion av en typ av vita blodkroppar benämnda "myeloida celler". Ungefär 80 personer om året insjuknar i KML i Sverige och behandlas med läkemedlet imatinib som blockerar BCR/ABL1. Imatinib har inneburit dramatiskt förbättrade utsikter för patienter med KML och framhålls idag som ett av de bästa exemplen där biologisk kunskap om cancerceller har möjliggjort utvecklingen av en målstyrd behandling. Behandlingen av KML är dock fortfarande förknippat med problem eftersom alla patienter inte svarar på behandlingen med imatinib. Dessutom krävs i de flesta fallen en livslång behandling eftersom en majoritet av patienterna återfaller i KML om medicineringen avbryts. Detta beror sannolikt på att leukemistamcellen i KML överlever behandlingen.

Den övergripande målsättningen med den här avhandlingen har varit att öka kunskapen om hur KML uppstår och hitta nya angreppspunkter för målstyrd terapi. För att göra detta har vi använt olika leukemimodeller som möjliggör detaljerade studier av sjukdomsbiologin vid KML. Avhandlingens första delarbete (Article I) syftade till att undersöka effekten av genen *SOCS2* som är uppreglerad hos KML-patienter. *SOCS2*-

proteinet fungerar som en broms i cellens signalsystem och har till uppgift att reglera svaret på specifika signalmolekyler. I delarbetet undersökte vi specifikt ifall SOCS2 även påverkade signaleringen som BCR/ABL1 startar i cellen. För att undersöka detta använde vi benmärgsceller från en mus som saknar *Socs2* genen (*Socs2* KO). Genom att transplantera dessa celler till normala möss, fann vi att även benmärgsstamceller utan SOCS2 ger upphov till alla de celltyper som normalt bildas i benmärgen. Dessa resultat tyder på att SOCS2 inte är viktigt för normal hematopoes eller att andra gener kan kompensera förlusten av SOCS2. Genom att uttrycka fusionsproteinet BCR/ABL1 i normala och i *Socs2* KO celler och sedan transplantera cellerna till normala möss, fann vi att mössen utvecklade samma KML-liknande sjukdomsbild även om de saknade SOCS2. Därför kunde vi dra slutsatsen att SOCS2 sannolikt inte har någon central betydelse för utvecklingen av KML.

I det andra delarbetet (Article II) var målet att hitta ett protein på cellytan som går att använda som markör för att skilja KML-stamceller från normala stamceller och potentiellt även som måltavla för behandling riktad mot KML-stamcellerna. För att göra det använde vi en metod (microarray) som gör det möjligt att jämföra uttrycket av ett stort antal gener, till att undersöka vilka gener som var högt uttryckta på cellytan av KML-celler jämfört med normala celler. I arbetet identifierade vi ett ökat uttryck av proteinet IL1RAP på cellytan av KML-celler jämfört med normala celler. IL1RAP är ett protein som utgör en viktig del av det komplex på cellytan som signalmolekylen IL1 binder till. Genom att använda antikroppar som binder till IL1RAP kunde vi även visa att denna cellytemarkör kan användas för att sortera ut och därmed särskilja KML-stamceller från normala stamceller. I arbetet kunde vi också visa att KML-stamceller selektivt kunde avdödas genom inbindning av en polyklonal antikropp riktad mot IL1RAP.

I det tredje delarbetet (Article III) undersökte vi vilken betydelse IL1RAP har för normal hematopoes och för celler som uttrycker BCR/ABL1. När vi undersökte blodet i möss som saknar *Il1rap*-genen (*Il1rap* KO möss), upptäckte vi lägre nivåer än normalt av myeloida celltyper i blodet. När vi sedan använde virus för att uttrycka IL1RAP i normala mänskliga celler som sedan transplanterades till en typ av möss med mycket försvagat immunförsvar (immundefekta möss), såg vi att fler av de omogna humana cellerna med IL1RAP utvecklades till myeloida typer av blodceller, än kontrollcellerna gjorde. Därav kunde vi dra slutsatsen att om omogna blodceller får förhöjt uttryck av IL1RAP, kan detta påverka cellens förmåga att bli myeloida celler. Vi upptäckte även att de allra mest primitiva cellerna hos KML patienter, som har mycket högre nivåer av IL1RAP på cellytan än normala celler, reagerade kraftigt på signalmolekylen IL1 medan normala celler inte märkbart påverkades. Detta tyder på att de höga nivåerna av IL1RAP på KML-stamceller har betydelse för sjukdomsutvecklingen av KML genom att påverka hur cellen reagerar på IL1. Därför är det också sannolikt att behandlingsstrategier som blockerar IL1RAP kan komplettera imatinib i behandlingen av KML-patienter.

Sammanfattningsvis har studierna i den här avhandlingen bidragit till en ökad förståelse av de cellulära förändringarna som orsakar KML. Genom att undersöka två proteiner som är uppreglerade i KML-celler, har vi kunnat visa att ett den ena av dessa (IL1RAP) sannolikt har betydelse för utvecklingen av KML, genom att bl.a. öka leukemicellernas svar på signalmolekylen IL1. Därför kan IL1RAP vara ett lämpligt cellyteprotein att blockera vid utvecklingen av nya behandlingsmetoder i KML.




# Introduction

## Hematopoiesis

The gradual development of all mature blood cells from hematopoietic stem cells (HSCs) is called hematopoiesis and a trillion (1,000,000,000,000) new cells of different types migrate from the bone marrow (BM) into the peripheral blood (PB) every day (Ogawa 1993). During their lifetime, these cells are responsible for a wide variety of functions such as oxygen transport by erythrocytes, blood clotting after vessel damage by thrombocytes, production of antibodies by B-cells, killing of cells infected by virus or potentially damaged cells by T-cells and NK-cells, and innate immunity from macrophage/monocytes and granulocytes. In order to produce this variety of cells, the hematopoietic system is hierarchically organized with the self-renewing multi-potent HSCs at the top of the hierarchy. According to the classical model of hematopoiesis, HSCs differentiate into multi-potent progenitors (MPPs) that increase their proliferative rate while losing self-renewal capacity (Reya et al. 2001). More lineage-committed progenitors are gradually formed, and they have been defined based on the expression of different transcription factors and surface markers. These progenitors are generally named according to their potential to differentiate into different mature cell types, and some well-studied progenitor types are: common lymphoid progenitors (CLPs), which differentiate into B-cells, T-cells, and NK-cells; and granulocyte macrophage progenitors (GMPs). These give rise to innate immune cells. The megakaryocyte erythrocyte progenitors (MEPs) are precursors for thrombocytes and erythrocytes (Kondo et al. 1997; Akashi et al. 2000; Reya et al. 2001).

The differentiation pathways and surface markers needed to identify the various progenitor populations were first defined in the mouse, in the classical hematopoietic model described in Figure 1 (Kondo et al. 1997; Akashi et al. 2000; Reya et al. 2001). However, this model has been challenged after the identification of a lymphoid multi-potential progenitor (LMPP), which can give rise to the lymphoid, granulocytic, and monocytic lineages but not to erythroid cells (Adolfsson et al. 2005). This spurred further detailed investigations and proposal of alternative models for murine hematopoiesis (Giebel & Punzel 2008; Kawamoto et al. 2010). Human hematopoiesis was first believed to be consistent with the classical model. However, the discovery of a human cell population corresponding to the LMPP in the mouse prompted a revision of this model (Doulatov et al. 2010; Doulatov et al. 2012). Furthermore, it has recently been suggested

that the human hematopoietic model, established after the discovery of LMPPs, needs to be revised further, as LMPPs only give rise to neutrophilic granulocytes, while other types of granulocytes, macrophages, and erythrocytes develop from myeloid progenitors (Görgens et al. 2013).



**Figure 1. The classical model of hematopoiesis and a revised model with the recently described MLLPs and their possible differentiation pathways indicated.** In the classical model of hematopoiesis, self-renewing hematopoietic stem cells (HSCs) differentiate into multi-potent progenitors MPPs, which lack self-renewal potential and produce common lymphoid progenitors (CLPs) and common myeloid progenitors (CMPs). The CLPs give rise to B-cells, T-cells, and NK-cells, while the CMPs give rise to megakaryocyte erythrocyte progenitors (MEPs) and granulocyte macrophage progenitors (GMPs). MEPs give rise to erythrocytes and megakaryocytes while monocytes and neutrophils are produced by GMPs. The discovery of lymphoid multi-potent progenitors (LMPPs) with lymphoid but also a limited myeloid lineage potential, caused a revision of the hematopoietic model. However, it is still not fully understood which subsets of myeloid cells LMPPs can give rise to and how these differentiation pathways are arranged.

A major limitation of the hematopoietic models is that they are constructed through functional evaluation of progenitor populations defined by expression of specific surface markers. However, these surface markers are not necessarily relevant for the cellular function of the progenitors and may not fully reflect gradual processes in the differentiating progenitor cells. Thus, the apparently homogeneous progenitor populations may be more heterogeneous than implied by the model and contain cells with slightly different properties and lineage potentials. Characterization of global molecular programs in order to better understand the differentiation process may be a feasible way to overcome this limitation, but the efforts so far have still been hampered by our inability to prospectively separate HSCs and progenitor cells without relying on individual surface markers and flow cytometry.

## **Cytokine signaling in hematopoiesis**

Hematopoietic progenitor cells are guided in their decisions to proliferate, differentiate, and survive by external stimuli from cytokines. Each cytokine binds its specific receptor, which is expressed differentially among cell types, and can thereby activate downstream signaling pathways in a cell-specific manner. Despite numerous different receptors and several downstream pathways, cell signaling is, however, characterized by crosstalk between pathways, resulting in complex networks of intracellular signal transduction. A few pathways relevant to this thesis are outlined below.

### *KIT receptor*

One of the best-known cytokines that regulate hematopoietic stem and progenitor cells is the stem cell factor (SCF), which signals through the KIT receptor (c-KIT) and can, by itself, regulate maintenance of HSCs, although a proportion of HSCs are generated and survive even in the absence of functional KIT (Li & Johnson 1994; Linnekin et al. 1995; Thoren et al. 2008). Stimulation of hematopoietic stem and progenitor cells with SCF is synergistic in combination with one of a number of other cytokines, for example interleukin-3 (IL3), granulocyte colony-stimulating factor (GCSF), and thrombopoietin (TPO) (Bernstein et al. 1991; Metcalf & Nicola 1991; Kobayashi et al. 1996), allowing *in vitro* expansion of progenitor cells and development of several myeloid cell types *in vivo* (Broudy 1997).

### *JAK/STAT*

Janus kinase/signal transducer and activator of transcription (JAK/STAT) signaling, is another important pathway that can be activated by a number of different cytokines, among others interleukin-6 (IL6) and erythropoietin (EPO). The JAK/STAT pathway is normally activated by cytokines that bind and oligomerize cell-surface receptors, allowing them to bind and facilitate cross-phosphorylation of JAK proteins (Dorritie et al. 2013). Once JAK is activated, it phosphorylates STATs, which are then able to enter the nucleus



and initiate transcription of a number of target genes, including the suppressor of cytokine signaling (*SOCS*) family. The *SOCS* family act like classical feedback inhibitors of signaling (Krebs & Hilton 2001; Aaronson & Horvath 2002). The importance of JAK/STAT proteins can be exemplified by *STAT3*, which is activated by several cytokines including *IL3* and *GCSF* (Duarte & Frank 2000). Furthermore, truncation of both *STAT5A* and *STAT5B* genes in mice results in lower B-cell counts, reduces the cellular response to erythropoietin, and may cause anemia (Teglund et al. 1998; Socolovsky et al. 1999; Sexl et al. 2000; Socolovsky et al. 2001), while complete deletion of the *STAT5* genes blocks development of several hematopoietic cell types and results in embryonic lethality (Hoelbl et al. 2006).

#### *Interleukin-1 receptor*

The Interleukin-1 receptor (*IL1R*)/Toll-like receptor (*TLR*) family includes both *TLRs*, which are mainly expressed on immune cells, and the *IL1Rs*, which have mainly been studied in the context of inflammation (O'Neill 2008). The two *IL1* isoforms, *IL1A/B*, can bind *IL1R* type 1, which then forms a complex with the *IL1R* accessory protein (*IL1RAP*), resulting in recruitment of the adaptor protein *MyD88* (Brikos et al. 2007). Following this, the proteins *IRAK1* and *IRAK4* bind to this complex and become phosphorylated, which in turn activates several downstream processes, leading to the activation of the nuclear factor kappa-light-chain-enhancer of activated B-cells 1 (*NFKB* or *NF- $\kappa$ B*) complex (O'Neill 2008). Although *NFKB* is a well-known transcription factor mainly studied in the context of immune response and T-cell development, only a few studies have been published on the role of *NFKB* signaling in hematopoiesis (Gerondakis et al. 2012). At the HSC level, *NFKB* has been suggested to regulate HSC self-renewal (Zhao et al. 2012a). Myelopoiesis may also be enhanced by *NFKB* through cooperation with *C/EBPA* (Friedman 2007).

# Leukemia

Cancers, including leukemia, are caused by genetic aberrations that alter normal mechanisms regulating self-renewal, cellular proliferation, differentiation, and programmed cell death (apoptosis) (Hanahan & Weinberg 2011). The term leukemia refers to malignancies affecting immature hematopoietic cells, and it has traditionally been divided into acute and chronic leukemias, depending on various morphological and clinical characteristics. Acute leukemias are typically manifested by an accumulation of immature blast cells due to a block in differentiation, while chronic leukemias mainly show an increase in mature cells in PB and BM due to increased proliferation among the progenitors. Leukemias are subdivided further, based on which lineage(s) the cells differentiate into. Thus, patients with expansion of myeloid cells can be divided into having acute myeloid leukemia (AML) with immature myeloid blasts or chronic myeloid leukemia (CML) with an expansion of myeloid cells at all stages of maturation and, for the definition of CML, the presence of the *BCR/ABL1* fusion gene. The corresponding classification for lymphoid (B- and T-cell) leukemia is acute lymphoblastic leukemia (ALL) and chronic lymphoid leukemia (CLL). Currently, the classification of myeloid leukemias and neoplasms into clinically and scientifically relevant entities is based on genetic, morphological, cytochemical, and immuno-phenotypic characteristics with the genetic alterations playing an increasingly important role for the diagnosis of these malignancies (Vardiman et al. 2009). Myeloproliferative neoplasms and myelodysplastic syndromes are other important disease entities that mainly affect the myeloid lineage and that have the capacity to progress into AML (Vardiman et al. 2009).

## Leukemia stem cells

It is still debated from which cell types different leukemias arise and are maintained. Leukemic stem cells (LSCs) can be defined as cells that have the capacity to both self-renew and propagate cell populations expanded in the leukemia (Reya et al. 2001; Dick & Lapidot 2005). The discoveries leading to the current LSC concept were made by transplantation experiments where primary AML cells were transplanted into immunodeficient mice (Lapidot et al. 1994; Bonnet & Dick 1997). Different AML cell populations were sorted and transplanted into non-obese diabetic/severe combined immunodeficiency disease (NOD/SCID) immune-incompetent mice to test whether only specific cell populations were able to propagate the leukemia. The most primitive CD34<sup>+</sup>CD38<sup>-</sup> fraction of cells was the only one that could reconstitute the leukemia, and it was proposed that the LSCs in AML were transformed normal primitive cells (Bonnet & Dick 1997). LSCs are not only found in AML, but they have also been found in various other leukemias and malignant disorders. In MDS with 5q deletions, the cell of

origin and the subsequent LSCs are found in the CD34+CD38- compartment (Nilsson et al. 2000). In B-ALL, various B-cell progenitor and blast populations, including the abnormal CD34+CD38-CD19+ population, have LSC activity (Cobaleda et al. 2000; Castor et al. 2005; le Viseur et al. 2008). The nature of the LSCs in AML was questioned a decade after their discovery, when Taussig et al. (2008) showed that the more mature CD34+CD38+ cell population in AML patients also contained LSCs. Furthermore, AML patients with NPM1 mutations even had their LSC activity in the CD34- cell population (Taussig et al. 2010). Although controversy remains, many of the diverse results could be explained by variations in the experimental systems used to study the LSCs; for example, treating recipient mice with immunosuppressing antibodies allows engraftment of CD38+ AML cells (Taussig et al. 2008). In the end, it is still uncertain which cells propagate AML, but there is a correlation between high engraftment potential and worse clinical outcome (van Rhenen et al. 2005; Pearce et al. 2006; Gentles et al. 2010; Eppert et al. 2011). Interestingly, NF $\kappa$ B is activated in primitive AML LSCs, and inhibition of this transcription pathway has led to successful eradication of CD34+CD38- cells in vitro (Guzman et al. 2001; Guzman et al. 2002; Guzman et al. 2005; Kuo et al. 2013).

When the LSC concept was first used to define the cells propagating a leukemia, they were termed LSCs, as they, like normal stem cells, have the potential for unlimited self-renewal and the ability to drive the leukemogenesis (Reya et al. 2001). The LSC concept does not, however, imply that the leukemic clone necessarily has to arise or share its surface marker profile with a normal stem cell. In fact, there are several examples where differentiated progenitors have been proven to regain self-renewal and become LSCs (Cozzio et al. 2003; Huntly et al. 2004; Krivtsov et al. 2006). Perhaps much confusion could have been avoided if another term, such as leukemia-propagating or leukemia-initiating cells (LICs), had been used instead of LSCs.

## Chronic myeloid leukemia

In 1960, CML was the first cancer to be associated with a causative chromosomal aberration, when a minute chromosome named the Philadelphia chromosome (Ph chromosome), was discovered (Nowell & Hungerford 1960). The Ph chromosome is the result of a reciprocal translocation, t(9;22), which forms the *BCR/ABL1* fusion gene at its breakpoints (Rowley 1973; Heisterkamp et al. 1985; Shtivelman et al. 1985; Grosveld et al. 1986; Mes-Masson et al. 1986). Later mouse studies established that the expression of the *BCR/ABL1* fusion gene causes a CML-like disease in mice, and that the tyrosine kinase activity of BCR/ABL1 is indispensable (Daley et al. 1990; Lugo et al. 1990). Compared to AML, which is a heterogeneous disease caused by several different genetic aberrations, CML is always initiated by the *BCR/ABL1* fusion gene. Since its original discovery, CML and the disease-causing *BCR/ABL1* fusion gene have not only provided a framework for the study of leukemia, but have also provided an important basis for the study and dissection of several other tumor types. Several molecular and cellular characteristics of CML are discussed in the sections below.

### **Clinical characteristics of chronic myeloid leukemia and current treatment options**

When diagnosed, CML often manifests in symptoms such as tiredness and clinical findings of an enlarged spleen accompanied by elevated white blood cell counts (WBC). However, today several cases are discovered accidentally from an elevated WBC count at a routine check-up performed for other medical reasons (Randolph 2005). The actual diagnosis of CML requires the presence of a t(9;22), or a confirmation that the *BCR/ABL1* fusion gene is present<sup>1</sup>. If left untreated, CML typically progresses through three phases. First, there is a chronic phase (CP), which manifests in an abnormal increase in mature granulocytes in the peripheral blood. After 3–5 years, the CP progresses into an accelerated phase (AP), with increasing numbers of immature myeloid progenitors. Finally, a blast phase (BP) (also called blast crisis or BC) will occur, with a block in differentiation that causes massive accumulation of immature blasts in both PB and BM (Sawyers 1999; Jabbour & Kantarjian 2012).

---

<sup>1</sup> According to the Swedish national treatment recommendations from 2010. <http://www.sfhem.se/>

Depending on the accumulation of either myeloid blasts (representing about 70% of all CML BP cases) or lymphoid blasts (30% of the cases), the BP is referred to as either 'myeloid BP' or 'lymphatic BP' (Randolph 2005; Jabbour & Kantarjian 2012). The blast accumulation, with its severe side effects, will be lethal within a few weeks if left untreated.

#### *First-generation tyrosine kinase inhibitors*

In 1996, a novel tyrosine kinase inhibitor (TKI) that blocks ABL1 activity was described (Druker et al. 1996), and it was later shown to be a very effective new treatment for CML (Druker et al. 2001). In long-term evaluations, this TKI called imatinib, has proved to effectively suppress the disease and induce remission in most patients (Druker et al. 2006). Since then, imatinib (or other TKIs) has become the recommended standard for the treatment of CP CML (Baccarani et al. 2013). The development of imatinib was made possible by a detailed characterization of the BCR/ABL1 fusion protein and the ability of different chemical compounds to block the ABL1 kinase activity. To date, imatinib most likely provides the best example of a truly targeted and successful therapy of any malignant disorder. Despite the remarkable success of imatinib, the majority of patients will experience relapse of the disease if the TKI treatment is discontinued, even after achieving undetectable levels of *BCR/ABL1* in BM and PB (Merante et al. 2005; Rousselot et al. 2007; Mahon et al. 2010). The most likely explanation for the recurrence of CML in patients stopping treatment with imatinib after achieving complete molecular remission, is a reservoir of residual CML stem cells that survive TKI treatment (Diamond & Melo 2011).

So far, it has not been possible to predict when the TKI therapy can be safely discontinued, but a fast decrease in *BCR/ABL1*-positive cells after initiation of therapy is believed to indicate a lower risk of relapse (Hughes et al. 2010). Other trials have indicated that in patients who achieve molecularly undetectable leukemia, which is defined as less than 0.0032% (4.5 log) *BCR/ABL1* transcripts per ABL1 transcript (as detected by quantitative polymerase chain reaction (QPCR)) (Baccarani et al. 2013), for a period of 2 years, between 50% and 60% will relapse after withdrawal of imatinib (Mahon et al. 2010; Ross et al. 2013). Thus, despite the revolutionary improvements in outcome following the introduction of TKIs in CML, additional therapeutic strategies are needed to meet the ultimate goal of a cure.

### *Second-generation tyrosine kinase inhibitors*

Despite the success of imatinib in the treatment of CML, about 5% of patients in CP and virtually all of the responding patients in BP, will relapse with disease (Druker et al. 2006; Hochhaus et al. 2009). Relapses can have a number of explanations including upregulation of *BCR/ABL1*, low intracellular concentration of imatinib due to poor bioavailability or poor cellular uptake, other activated signaling pathways, or mutations at the active site of ABL1 (Diamond & Melo 2011). Imatinib binds competitively to the adenosine triphosphate (ATP) binding site on ABL1, thereby inhibiting the kinase activity of the BCR/ABL1 fusion protein (Schindler et al. 2000). Different point mutations at the active site of ABL1 can therefore prevent binding of imatinib, and are often found in relapse patients who have acquired resistance to therapy (Hochhaus et al. 2002). Today, several new small molecules have been developed with the aim of inhibiting ABL1, and they enable treatment of patients with ABL1 mutations. Using one of the second-generation TKIs (nilotinib or dasatinib) to treat relapsing patients, it is now possible to overcome most mutations in BCR/ABL1 except for T315I (Shah et al. 2004; Golemovic et al. 2005; Talpaz et al. 2006; Kantarjian et al. 2006). However, with the introduction of ponatinib, it has recently become possible to successfully treat also patients with the T315I mutation (O'Hare et al. 2009; Cortes et al. 2012).

### *Allogeneic stem cell transplantation*

Before the introduction of TKI treatment in CML, allogeneic stem cell transplantation (SCT) remained the only curative treatment option (Sawyers 1999). As referred to earlier, this is likely to be still true in most cases, although a fraction of CML patients have been off drug treatment for several years without detectable BCR/ABL1 transcripts (Mahon et al. 2010). In patients diagnosed with BP CML or patients who develop resistance to second-generation TKIs, allogeneic SCT is still the recommended treatment of choice (Jain & van Besien 2011; Jabbour & Kantarjian 2012;). It is worth mentioning, however, that allogeneic SCT is associated with high treatment-related mortality and risk of graft-versus-host disease which, combined with the occurrence of relapse after transplant, results in a five-year survival ranging from 30% to 90%, depending on various risk factors (Jabbour et al. 2011; Kantarjian et al. 2012). Due to the risks associated with allogeneic SCT, and the challenge to find suitable HSC donors, it is still difficult to treat CML at advanced stages.

## **Blast phase and known secondary genetic changes**

The exact mechanism and cellular effects leading to progression of CML into BP remain elusive, despite the fact that there have been several studies. Upregulation of *BCR/ABL1* has been suggested as one factor contributing to disease progression, and *BCR/ABL1* is known to increase the rate of mutagenesis (Nowicki et al. 2004). As might be expected, CML cells in BC generally have a higher number of genetic aberrations than in CP (Johansson et al. 2002; Mullighan et al. 2008). As a consequence, several genes (e.g. *RUNX1*, *ASXL1*, *WT1*, *NRAS*, *KRAS*, and *TET2*) are deleted or mutated in a proportion of BP patients (Feinstein et al. 1991; Sill et al. 1995; Grossmann et al. 2011; Makishima et al. 2011; Zhao et al. 2012b). In addition, deletion of the transcription factors *IKZF1* and *CEBPA*, which are essential for differentiation into lymphoid and myeloid lineages, respectively, provide examples of genetic aberrations that may contribute to induction of the differentiation block required for lymphoid or myeloid blast formation (Wagner et al. 2006; Mullighan et al. 2008). Genetic events leading to gain of function of affected proteins may also contribute to progression of CML. For instance, formation of a secondary *NUP98/HOXA9* or *MSI2/HOXA9* fusion gene have been found important for the disease progression in CML (Yamamoto et al. 2000; Dash et al. 2002; Barbouti et al. 2003).

CML cells have an increased level of non-homologous end-joining, leading to increased susceptibility to chromosomal aberrations (Chakraborty et al. 2012). As a possible result, aberrations like +8, i(17q), and also an extra Philadelphia chromosome, are associated with progression to BP (Mitelman et al. 1976). In addition to the existence of an extra Philadelphia chromosome in patients with BP, increased *BCR/ABL1* expression in both the bulk leukemic cells and in cells with a GMP phenotype is associated with progression to BP (Gaiger et al. 1995; Jamieson et al. 2004). Cell line experiments have also shown that higher *BCR/ABL1* expression reduces cell adherence and increases transforming effects such as cytokine independence and migration (Barnes et al. 2005). These findings indicate that upregulation of *BCR/ABL1* may be involved in the transformation of CML to BP. Despite the finding of several genetic events that are associated with progression of CML to BP, this process is not yet fully understood, but the introduction of massive parallel sequencing will most likely give a more comprehensive view of the genetic landscape of CML in BP. Moreover, it may well be true that the most frequent cytogenetic changes observed in CML BP, i.e. trisomy 8 and isochromosome 17q, affect the expression levels of multiple genes that contribute to the progression of CML CP into BP.

## **CML stem cells**

The first indication that the LSC in CML is created by transformation of a normal HSC was the finding that the *BCR/ABL1* gene is present not only in leukemic cells but also in multiple lineages of hematopoietic cells (Fialkow et al. 1977; Fialkow et al. 1978). Furthermore, experiments using transgenic mouse models closely mimic CML when *BCR/ABL1* is expressed specifically in HSCs and progenitor cells (Koschmieder et al. 2005). It is generally believed that the CML stem cells, which are insensitive to imatinib treatment, are the source of disease relapse in CML CP patients after withdrawal of imatinib. Several mechanisms for the insensitivity of CML stem cells to TKIs have been suggested, including high baseline expression of *BCR/ABL1*, protection of CML stem cells by the BM microenvironment, and insensitivity because the CML stem cells are kinase-independent, possibly due to cell cycle quiescence (Jiang et al. 2007; Corbin et al. 2011; B. Zhang et al. 2013). Recent studies using transgenic mice where *BCR/ABL1* expression was turned on and off after establishment of disease, strongly suggested that CML stem cells are not addicted to BCR/ABL1, as leukemia reappears after re-induction of the *BCR/ABL1* expression (Hamilton et al. 2012). Thus, it seems unlikely that TKIs by themselves would be able to provide an efficient cure for CML. In myeloid BP, CML stem cells appear to be present both in the CD34<sup>+</sup>CD38<sup>-</sup> population and in the GMP compartment (Jamieson et al. 2004), while in lymphoid BP, CML stem cells have been found even among CD34<sup>-</sup>CD38<sup>+</sup> BM cells (Tanizaki et al. 2010).


## **Signaling pathways in chronic myeloid leukemia**

Fusion of BCR to ABL1 results in a constitutive activation of the ABL1 tyrosine kinase activity and leads to phosphorylation and activation of several downstream signaling pathways. The network of signaling pathways activated by BCR/ABL1 is very complex, but a few main pathways that are very relevant to this thesis are described below and summarized in Figure 2.

### *RAS pathway*

Autophosphorylation of the Tyr177 residue of BCR/ABL1 allows binding of the GRB2 adaptor protein (Pendergast et al. 1993), which enables subsequent binding of SOS and activation of the RAS/MAPK pathway (Druker et al. 1992; Tauchi et al. 1994; Cortez et al. 1997). Activated RAS can also induce signaling through the JUN amino-terminal kinase pathway (Raitano et al. 1995). RAS activity is essential in CML, and blocking of the RAS pathway reverses transformation by BCR/ABL1 (Sakai et al. 1994; Gishizky et al. 1995; Sawyers et al. 1995). Moreover, the cell transforming activity of RAS is dependent on a posttranslational modification, the addition of an isoprenoid group catalyzed by farnesyl protein transferase, which allows RAS to associate with the cell





**Figure 2. Schematic overview of signaling pathways activated by BCR/ABL1.** The tyrosine kinase domain of ABL1 activates JAK2 and STAT5, both leading to translocation of phosphorylated STAT5 to the nucleus where it induces transcription of target genes. Phosphorylation of the Tyr177 in BCR/ABL1 allows binding of GRB2, which enables complex formation with SOS and GAB2 leading to activation of the RAS/MAPK pathway and transcriptional activation. GAB2 also activates PI3K, a signaling mediator with potential to cross-activate NFKB through AKT. The NFKB-complex is also activated downstream from the IL1R, which is dependent on IL1RAP for its activation. The focus of Article I, in which the role of SOCS2 was investigated in normal and malignant BCR/ABL1-induced hematopoiesis, is indicated with an asterisk (\*) in the figure. In Article II, IL1RAP was identified as upregulated on candidate CML stem cells, indicated by \*\*. The effects of the signaling through IL1RAP were investigated in Article III, indicated by \*\*\*.

membrane (Hancock et al. 1990; Gibbs et al. 1994). Thus, farnesyl protein transferase inhibitors (FTIs) were developed to inhibit both mutated RAS and normal RAS (Kato et al. 1992; Nagasu et al. 1995; Sepp-Lorenzino et al. 1995). It was, however, established later that RAS function could be partially maintained by alternative pathways and that FTIs had unspecific effects on a number of other farnesylated proteins (Downward 2003). Nevertheless, these compounds had some anti-tumorigenic activity, and FTIs have been tested in clinical trials as therapy for CML in CP, AP, and BP, but they only resulted in transient responses in a minority of patients tested (Cortes et al. 2003; Borthakur et al. 2006)

### *JAK2/STAT5 pathway*

In CML, the JAK2/STAT5 pathway is directly activated by the BCR/ABL1 fusion protein, but the question of whether only STAT5 or also JAK2 is important for the BCR/ABL1-induced transformation is still being debated (Ilaria & Van Etten 1996; Carlesso et al. 1996; Shuai et al. 1996). However, STAT5 was believed to be dispensable in CML, given that *BCR/ABL1* transduction-transplantation experiments, using BM from mice with a truncated form of *STAT5*, was still able to induce a CML-like disease (Sexl et al. 2000). Although *STAT5* was truncated, these cells retained residual STAT5 activity and when the experiment was repeated with BM cells from mice lacking the complete *STAT5A* or *A/B* genes, BCR/ABL1 was shown to be dependent on STAT5 (Hoelbl et al. 2006; Ye et al. 2006; Hoelbl et al. 2010).

Normally, STAT5 is activated through phosphorylation of tyrosine residues by the upstream protein JAK2. Thus, it can be reasoned that because of a direct activation of STAT5 by BCR/ABL1 (Chai et al. 1997; Wilson-Rawls et al. 1997), JAK2 is redundant in CML. However, it has been argued that inhibition of JAK2 reduces the transforming effects of BCR/ABL1 and that JAK2 either cooperates with BCR/ABL1 to activate STAT5 or mediates cross-activation of other signaling pathways (Warsch et al. 2013). One of the main proofs supporting involvement of JAK2 in CML is that inhibition of JAK2 with a small-molecule inhibitor was shown to induce apoptosis in CML cells (Samanta et al. 2009). However, several of the inhibitors used in similar experiments were later shown to inhibit BCR/ABL1 as well (Hantschel et al. 2012). Hence, although this is still controversial, JAK2 now appears to be non-essential for the development of CML but has been suggested to be essential in mediating cytokine signaling to allow CML stem cells to survive TKI therapy (Warsch et al. 2013). In addition, it is quite possible that JAK2 has a different role in CML BP.

Another member of the JAK2/STAT5 pathway is the suppressor of cytokine signaling 2 (SOCS2), whose main function is feedback inhibition of cytokine signaling (Endo et al. 1997; Naka et al. 1997). *SOCS2* is upregulated upon STAT5 activation and although it is a feedback inhibitor, high levels of SOCS2 have been reported to stimulate signaling, possibly by inhibiting other inhibitors (Tannahill et al. 2005; Piessevaux et al. 2006). We and other investigators have previously found that *SOCS2* expression is upregulated in CML (Schultheis et al. 2002; Håkansson et al. 2008). In Article 1, we used a *Socs2*-deficient mouse strain to evaluate whether SOCS2 is a critical regulator of BCR/ABL1 signaling and important for the pathogenesis of CML. We showed that cells from *Socs2*-deficient mice develop leukemia indistinguishable from wild-type cells upon transplantation after transduction with *BCR/ABL1*. Our results indicate that SOCS2 is not critical for the induction of CML and STAT5 phosphorylation by BCR/ABL1. In myeloproliferative neoplasms, STAT5 phosphorylation is often achieved by mutations activating JAK2, so SOCS2 may be of more importance in that setting (Quentmeier et al. 2008).

### *NFKB pathway*

BCR/ABL1 activates NFKB in a tyrosine kinase- dependent fashion (Hamdane et al. 1997; Kirchner et al. 2003). However, it has also been suggested that RAS activation is involved in transactivation of NFKB (Reuther et al. 1998). Thus, it is still possible that multiple co-operating pathways contribute to the activation of NFKB in CML. NFKB is known to enhance resistance to apoptosis and to increase proliferation in some forms of cancer, and it is known to be activated in both CML and AML (Hamdane et al. 1997; Reuther et al. 1998; Guzman et al. 2001; Karin et al. 2002; Kirchner et al. 2003), possibly through crosstalk from the PI3K/AKT signaling pathway (Zhu et al. 2011). Furthermore, inhibition of the NFKB complex by blocking of various proteins in the signaling pathway suggests that NFKB may be a possible therapeutic target in CML (Cilloni et al. 2006; Lounnas et al. 2009; Lu et al. 2010; Wei et al. 2013). IL1 is one cytokine known to activate NFKB in several types of immune cells (Fitzgerald & O'Neill 2000). In addition, a study of microenvironmental effects in CML showed that IL1 and GCSF induce more proliferation of primitive CD34+CD38- CML cells than of corresponding normal cells (Zhang et al. 2012). Furthermore, altered cytokine expression in CML cells with an increase in IL1A, GCSF, and other cytokines provides a selective advantage for the CML stem cells (Zhang et al. 2012). The intracellular mechanisms underlying the growth advantage of primitive CML cells over normal HSCs after IL1 stimulation are unclear, but they could be an effect of additional NFKB activation. In Article II, we found that expression of the IL1 receptor accessory protein (IL1RAP) was upregulated in primitive CD34+CD38- CML cells compared to the corresponding normal cells. IL1RAP is an essential component of the IL1 signaling pathway which (among other effects) results in NFKB activation (Cullinan et al. 1998). Thus, in Article III, we continued to investigate whether IL1RAP has a role in hematopoiesis and whether upregulation of *IL1RAP* is sufficient to induce features of a myeloproliferative disease. We demonstrated that *Il1rap*-deficient mice have a reduction in myeloid cells in PB. We also showed that CML CD34+CD38- BM cells, the population with the highest *IL1RAP* expression compared to corresponding normal cells, are hypersensitive to IL1B, while IL1B has limited effect on more mature CD34+ cells. It may be speculated that since IL1RAP is essential for activation of the IL1R, the higher levels of IL1RAP on the cell surface of CD34+CD38- CML cells may help to sensitize the receptor complex and downstream signaling pathways to activation, leading to a CML-specific cell expansion.

## **Experimental models of chronic myeloid leukemia**

### *Cell culture*

Immortalized leukemic cell lines are widely used in research and offer a system that is easy to work with, is renewable, and has the benefit that experiments can be carried out at multiple sites using the same cell model. Several CML cell lines are available, all of which were established from PB or BM of patients in advanced BP, but there are no cell lines that faithfully represent CML CP. Despite the convenience of using immortalized cell lines, they have several drawbacks that limit their use. Cell lines have usually been cultured extensively, leading to the acquisition and selection of new genetic alterations and to genetic drift, when the cells adapt to *in vitro* culture (Drexler et al. 1999; Dirks et al. 2010).

To overcome the drawbacks of immortalized cell lines, retrovirally transduced primary cells are often used when studying the cellular and molecular effects of BCR/ABL1 expression. These cell cultures are believed to reflect leukemic cells more closely, as they have not been immortalized and retain much of their intrinsic differentiation potential and cytokine dependency. Compared to *in vitro* culture of primary leukemic cells, retrovirally transduced cells have both benefits and drawbacks. While retroviral transduction is often a way of ensuring access to sufficient numbers of cells for a successful study (especially when rare genetic aberrations are studied), primary leukemia cells often contain additional genetic and epigenetic changes that are difficult to mimic in transduced cells.

### *Mouse models*

Instead of culturing *BCR/ABL1*-transduced cells *in vitro*, it is possible to transplant transduced mouse cells into mice of the same genetic background in order to study the cellular functions *in vivo*. Daley et al. (1990) established a mouse model of CML by transducing mouse BM cells with a retrovirus expressing *BCR/ABL1* and transplanting cells into irradiated recipients. These mice developed several different hematopoietic malignancies, including macrophage tumors and lymphoid leukemia. Subsequently, the transduction and transplantation model was gradually improved and better characterized to give a disease more similar to CML (Zhang & Ren 1998; Li et al. 1999). These models have been used successfully combined with KO strains, to investigate the role of specific genes in CML, such as *Stat5* and *Alox5* (Hoelbl et al. 2006; Ye et al. 2006; Chen et al. 2009). However, the transduction-transplantation model of CML still has a number of limitations. The retroviral constructs target not only HSCs but also progenitor cells, leading to massive progenitor proliferation that complicates the evaluation of experiments. The irradiation of recipient mice and the variation in *BCR/ABL1* expression levels may also increase the variability of the experiments. In addition, the retroviral models typically show a very aggressive disease, with death occurring around three weeks

after transplantation. Thus, studies investigating the contributions by various ‘associated genes’ such as *SOCS2* (Article 1) may not necessarily exclude the possibility that these genes play a role in the pathogenesis of CML, as, due to its limitations, this mouse model may fail to account for the function of that specific gene. An alternative way to model CML that has been widely used is to generate transgenic *BCR/ABL1* mice (Koschmieder & Schemionek 2011). Currently, the most elegant one is transgenic mice that express *BCR/ABL1* from an HSC-specific tetracycline-inducible promoter (Huettner et al. 2003; Koschmieder et al. 2005). This approach makes it possible to both induce and to reverse a CML-like disease by removal or re-administration of tetracycline to the drinking water of the mice, thereby allowing detailed *in vivo* studies of CML pathogenesis (Zhang et al. 2012). To date, this transgenic model is probably the most elegant and pure murine model of CML.

### *Immunodeficient mouse models*

Some immunodeficient recipient mice, such as NOD/SCID mice, have impaired T-cell and B-cell development and can therefore be engrafted by human cells (Prochazka et al. 1992; Lowry et al. 1996). Xenotransplantation of human cells into these mice have been of great importance for the functional characterization of normal human hematopoiesis and HSCs (Lowry et al. 1996; Doulatov et al. 2012). However, not only can normal hematopoietic cells be transplanted into NOD/SCID mice, but CML cells also give long-term engraftment (Lewis et al. 1998, Wang et al. 1998). Although they can be robustly engrafted by *BCR/ABL1*-positive cells, none of the transplanted mice were reported to have succumbed from development of a disease similar to CML.

Immunodeficient mouse models for xenotransplantation of human cells are still being improved to allow higher engraftment levels and to better support differentiation of cells into lineages. The NOD/SCID model has been improved by deletion of the IL2R gamma chain, which leads to lack of functional NK-cells in these mice, commonly referred to as NSG (Shultz et al. 2005). In Article III of this thesis, we used NSG mice to study the effect of IL1RAP overexpression in human HSCs and progenitor cells. Clearance of human donor cells by recipient immune cells is, however, not the only limiting factor for xenotransplantation efficiency. Lack of cross-reactivity of some niche factors and cytokines is also likely to limit the development of leukemic cells after xenotransplantation. Mice transgenically expressing human SCF, granulocyte macrophage colony-stimulating factor (GM-CSF), and IL3 have increased development of both normal human hematopoietic cells and leukemic cells after transplantation (Feuring-Buske et al. 2003; Nicolini et al. 2004).

Recently, Wunderlich et al. (2010) crossed these mice with NSG mice to generate an NSG strain expressing human SCF, granulocyte macrophage colony-stimulating factor (GM-CSF), and IL3 (referred to as NSGS mice), which even better supports the engraftment of human leukemic cells. As it has already been reported that CML cells transplanted into NOD/SCID mice achieve higher engraftment after injections of human SCF into the recipients (Lewis et al. 1998), it is quite possible that NSGS mice are more permissive in allowing engraftment of primary CML cells.



# The present investigation

This section describes the aims of the present study and the articles included, a summary and discussion of the main findings, and the conclusions arrived at. The experimental procedures are based on well-known methods and will not be discussed in this chapter. Instead, the reader is referred to the materials and methods sections of the original articles (I–III), which are presented in the third section of this thesis.

## Aims of the study

The overall aim of this study was to identify genes upregulated by the BCR/ABL1 fusion protein and to use *in vitro* and *in vivo* models to study their function in normal and malignant hematopoiesis. More specifically, the aims were:

- To determine the function of SOCS2 in normal hematopoiesis and its role in BCR/ABL1-induced cell signaling (Article I).
- To identify a cell-surface marker for primitive CML cells, allowing prospective isolation of candidate CML stem cells (Article II).
- To study the function of IL1RAP in normal hematopoiesis and the consequences of upregulation of *IL1RAP* in malignant hematopoiesis (Article III).



## Results and discussion

### Article I

#### **SOCS2 is dispensable for BCR/ABL1-induced chronic myeloid leukemia-like disease and for normal hematopoietic stem cell function.**

Signaling cascades activated by BCR/ABL1 elicit transcriptional gene expression signatures that are thought to contribute to CML pathogenesis. Thus, not only may the primary downstream mediators of BCR/ABL1 be important for the establishment of CML, but also genes becoming upregulated as a result of signaling by BCR/ABL1. The STAT5 signaling pathway is one important mediator of BCR/ABL1-induced transformation. Activation of STAT5 is well known to induce expression of its feedback inhibitor SOCS2, which had already been found to be highly expressed in primary BM cells from patients with CML. However, at the outset of this study it had not been established whether SOCS2 is involved in the pathogenesis of CML or whether it is important for normal HSC function. In Article I, we demonstrated that although *Socs2* was found to be preferentially expressed in long-term HSCs, *Socs2*-deficient HSCs were indistinguishable from wild-type HSCs when challenged in competitive BM transplantation experiments. Furthermore, by using a retroviral BCR/ABL1-induced mouse model of CML, we showed that SOCS2 is dispensable for the induction and propagation of the disease. The phosphorylation of STAT5 following *BCR/ABL1* transduction was unaffected by SOCS2, suggesting that the SOCS2-mediated feedback regulation of the JAK/STAT pathway is deficient in BCR/ABL1-induced CML.

### Article II

#### **Isolation and killing of candidate chronic myeloid leukemia stem cells by antibody targeting of IL-1 receptor accessory protein.**

Therapeutic strategies aimed at cure of CML will most likely require full eradication of the CML stem cells. In an attempt to find cell-surface markers useful for identification and isolation of CML stem cells, we performed global gene expression analysis both on primary CD34<sup>+</sup> CML cells and umbilical cord blood (CB) cells transduced with BCR/ABL1. The IL1R1 co-receptor IL1RAP was found to be selectively expressed on CML cells and to become upregulated following expression of BCR/ABL in CB cells. To determine whether IL1RAP expression would allow a prospective isolation of Ph-positive and normal residual Ph-negative cells within the CML CD34<sup>+</sup>CD38<sup>-</sup> cell compartment, we established a new protocol for FISH analysis on small numbers of sorted cells.

Interestingly, we found that the CML CD34+CD38-IL1RAP+ cells were preferentially Ph-positive, whereas CML CD34+CD38-IL1RAP- cells were almost exclusively normal. We hypothesized that, being exposed on the cell surface of candidate CML stem cells, IL1RAP could provide a target for a novel antibody-based therapy of CML. After generating a polyclonal anti-IL1RAP antibody, we found that IL1RAP could be used to induce antibody-dependent cell-mediated cytotoxicity (ADCC) of CD34+CD38- CML cells. In summary, this study identified IL1RAP as the first specific marker for candidate CML stem cells and also demonstrated that targeting of IL1RAP by an antibody-based approach may open up new avenues in the treatment of CML.

### **Article III**

#### **IL1RAP promotes myelopoiesis and its upregulation on candidate CML stem cells is associated with increased IL1 sensitivity.**

In addition to several progenitor populations, IL1RAP is expressed on candidate CML stem cells and other myeloid malignancies (Article II; Barreyro et al. 2012; Askmyr et al. 2013). IL1RAP is an essential co-receptor for IL1R1 and IL33R (ST2), mediating signals from IL1 and IL33, respectively. However, little is known about the cellular consequences of IL1RAP-mediated signaling in these cell types. In Article III, we found that *Il1rap*-deficient mice have reduced myeloid cell counts in peripheral blood at steady state, but that IL1rap is dispensable for normal hematopoietic stem cell function, as assessed by competitive SCTs. In xenotransplantation studies, enforced expression of IL1RAP in CB CD34+ cells resulted in a myeloid lineage skewing with expansion of a CD33<sup>low</sup>CD15+ cell population. We also determined whether CB cells with enforced IL1RAP expression and primary CML cells showed a different sensitivity for cytokines signaling through receptors known to associate with IL1RAP in other cell types. Whereas IL1B did not support *in vitro* proliferation of normal CB CD34+ cells, not even upon enforced IL1RAP expression, primary CML CD34+CD38- cells expanded significantly in response to IL1B stimulation. Furthermore, we found that CD34+ CML cells under stimulation by SCF expanded more than the primitive CD34+CD38- population. This is in accordance with a recent report showing differential sensitivity of CML cells to SCF (Corbin et al. 2013). In summary, these results suggest that upregulation of IL1RAP may contribute to the pathogenesis of myeloid neoplasms by increasing the sensitivity to IL1 signaling and promoting myelopoiesis.

## Conclusions

The main findings of this thesis are presented below:

### **Article I**

- SOCS2 is not required for normal hematopoiesis and HSC function.
- SOCS2 is not critical for BCR/ABL1 to initiate disease.
- The direct activation of STAT5 by BCR/ABL1 probably by-passes the SOCS2-mediated feedback inhibition of the JAK2/STAT5 pathway.

### **Article II**

- IL1RAP is a cell-surface marker for candidate CML stem cells and can be used to prospectively isolate such cells.
- Antibodies to IL1RAP can be used to induce ADCC in candidate CD34+CD38-CML stem cells, while sparing the corresponding normal cells.

### **Article III**

- IL1RAP is dispensable for HSCs but promotes the myeloid lineage *in vivo*.
- Forced IL1RAP expression is insufficient to increase the sensitivity of CB cells to IL1B, while primitive CML cells are hypersensitive and expand massively *in vitro* upon IL1B stimulation.

# Concluding remarks and future perspectives

Cancer is caused by a variety of genetic aberrations, resulting in loss or abnormal activation of protein functions. Recent technological advancements have facilitated genetic studies, with the sequencing of the human genome as an important landmark (Lander et al. 2001; Venter et al. 2001). Since then, sequencing technology has improved and has reduced costs further. Currently, massive parallel sequencing technology provides large amounts of data describing the genetic landscape of cancer and leukemia (Cancer Genome Atlas Research Network 2013a; Cancer Genome Atlas Research Network 2013b). The human genome project had great hopes of giving benefit to patients in the clinic, but it has proven to be more difficult than expected to understand the functional relevance, at the cellular and tissue levels, of the genetic findings. Several new functional methods such as shRNA technology and refined mouse models are being developed to facilitate experiments to understand the functional effects of the genetic findings.

The aim of this thesis was to improve our understanding of the functional effects of the upregulation of *SOCS2* and *IL1RAP* seen in CML cells. *SOCS2* is known to be involved in intracellular regulation of cytokine signaling, but as described in this thesis, *SOCS2* is dispensable for BCR/ABL1 inducing disease *in vivo*, as assessed in mice receiving transplants with BCR/ABL1-transduced mouse BM cells. As referred to earlier, it is worth noting that this retroviral CML model has several limitations; for example, the disease observed is very aggressive with a short latency period, and as such it is difficult to entirely rule out a role of *SOCS2* in the pathogenesis of CML. A relevant follow-up study would be to cross *Socs2*-deficient mice with transgenic mice expressing *BCR/ABL1* from an HSC-specific inducible promoter (Koschmieder et al. 2005). Interestingly, there have been several publications indicating that *SOCS2* is important in other tumor types, including prostate cancer, and in MPDs with *JAK2* mutation (Quentmeier et al. 2008; Iglesias-Gato et al. 2013).

The current CML therapy is based on inhibition of BCR/ABL1 or stem cell transplantation. However, as LSCs appear to be independent of the BCR/ABL1 tyrosine kinase activity, it is questionable whether ABL1 inhibitors would have the potential to be highly curative in CML. In addition, TKIs have proven to be ineffective in advanced stages of the disease (Pellicano et al. 2011). One way to overcome the TKI insensitivity is to find alternative targets in the CML cells, such as proteins with a function that becomes

critical after transformation by BCR/ABL1. One example of such a target in CML is RAD52, which is involved in the repair of double-strand breaks (Cramer-Morales et al. 2013). TKI treatment of CML cells has also been shown to induce autophagy, a process that leads to TKI resistance (Mishima et al. 2008; Kamitsuji et al. 2008). Inhibition of autophagy increases leukemic cell death and significantly increases the elimination of candidate CML stem cells in combination with a TKI (Mishima et al. 2008; Bellodi et al. 2009). Thus, autophagy may be a promising novel therapeutic target in CML.

The *IL1RAP* protein is an alternative target in CML that was discovered in a quest to find a cell-surface marker specific for CML stem cells (Article II). Cell-surface proteins can be used both to identify leukemic cells and as targets for different antibody-based therapies (Morris & Waldmann 2009). One alternative to the ADCC-based approach using NK-cells in Article II is the cancer vaccine concept, where synthetic peptides from leukemia-specific proteins induce T-cell-mediated cell killing (Dao & Scheinberg 2008). Based on this concept, there have been several trials using various BCR/ABL1 fusion peptides as vaccines in CML treatment (Maslak et al. 2008; Casnici et al. 2011). Peptides of WT1 are another target of vaccines tested for AML, but they have recently also been used for antibodies targeting the peptide when presented by human leukocyte antigen (HLA) in Ph chromosome-positive ALL (Maslak et al. 2010; Dao et al. 2013). These approaches might prove to be feasible ways to achieve an improved therapy of CML and Ph chromosome-positive ALL. Most likely, however, targeting of cell-surface proteins will turn out to be an easier and more effective approach, as exemplified by the successful therapy of B-cell malignancies using anti-CD20 antibodies (e.g. Rituximab) (Barth et al. 2012).

In Article III, we studied the function of IL1RAP and found that mice deficient for *Il1rap* had fewer myeloid cells in PB and that forced expression of *IL1RAP* in human CB cells resulted in myeloid skewing after transplantation into NSG mice. These findings indicate that IL1RAP has a functional role in promoting myelopoiesis by enabling cytokine signaling in normal hematopoiesis. It should, however, be noted that *in vitro* cultures of cells with forced expression of IL1RAP did not yield higher cell numbers after cytokine stimulation with IL1RAP-dependent cytokines. Thus, the exact mechanism of IL1RAP function in normal hematopoiesis is not yet understood, and should be further investigated. In contrast, primitive CML CD34+CD38- cells with increased expression of IL1RAP expanded significantly in response to IL1B stimulation (Article III; Zhang et al. 2012). This effect of IL1 in the primitive CML cell population supports the idea that extrinsic microenvironmental regulation, and in particular IL1, may be of importance to the LSCs. IL1 signals through the IL1R1/IL1RAP complex and is known to result in NFkB activation (Cullinan et al. 1998). It is also known that NFkB activation increases the self-renewal of normal HSCs (Zhao et al. 2012a), which may indicate that upregulation of IL1RAP combined with expression of IL1R leads to increased self-renewal of the LSCs. However, our results with cell expansions after liquid cultures suggest more of a proliferative expansion of the CD34+CD38- CML fraction. A detailed

analysis of the consequences of IL1 signaling, regarding both gene expression response and in terms of cellular response of primitive CML cells, will therefore be needed to clarify how primitive CML cells respond to IL1. It may very well be that an IL1RAP-based therapy would both inhibit IL1 signaling in the CML cells and target them for immune-mediated killing in order to be effective.

Despite the fact that CML is always initiated by the formation of the *BCR/ABL1* fusion gene, CML cells may evolve into separate heterogeneous sub-clones during disease progression, which is indicated by the variation in mutations and genetic aberrations found in CML cells (Grossmann et al. 2011; Makishima et al. 2011). Thus, it may be not be enough to use only one therapeutic approach, but combinational therapy may be needed in order to find cures for CML patients in the future. Given the heterogeneity of most cancers, it may be that therapy could be based on the presence of specific target molecules rather than the type of cancer. For example, IL1RAP is expressed in CML, and also in a proportion of AML patients, and antibodies against IL1RAP are also capable of inducing *in vitro* ADCC in cells from these patients (Barreyro et al. 2012; Askmyr et al. 2013). Perhaps a large panel of immune-based therapy targets, like anti-IL1RAP agents, cancer vaccines, or therapies inhibiting the key driver mutations (like imatinib) will eventually become the basis of cancer therapy, where treatment is selected by genome-wide analysis to identify target proteins, enabling individually selected drug combinations irrespective of the tumor type.



# Acknowledgements

I would like to express my sincere gratitude to many people for their help and support during my thesis work. In particular, I would like to thank:

My excellent supervisor **Thoas Fioretos**, who is always kind and generous when enthusiastically sharing his wisdom, knowledge, and experience in science and of life. Thank you for letting me take part in your exciting research, and for encouraging me whenever I needed it.

My co-supervisor **Marcus Järås**, who has taught me so much and has always pushed me to take one step further, improving my work a little more and never giving up or leaving things only half finished. I would not have made it this far without your help.

**Jonas Larsson** and **Johan Richter** who have been my co-supervisors, for giving good advice and sharing opinions and expert knowledge that helped me move the project forward.

The founding father of the research environment at the Department of Clinical Genetics, **Felix Mitelman** and all the senior scientists **Bertil Johansson**, **Fredrik Mertens**, **Nils Mandahl**, **David Gisselsson Nord**, **Kajsa Paulsson**, and **Anna Andersson**, for creating this excellent environment with its open and kind intellectual atmosphere.

The postdocs **Helena**, **Kristian**, **Maria**, and **Henrik**, who worked with me and have been very supportive both regarding the projects and in understanding the research field in general, with countless discussions at meetings, in the lunch room, and at the lab bench. The very skilled and helpful technicians **Marianne Rissler**, **Sandra Gordon**, **Carl Högberg**, and **Carin Lassen**, who made my work in the lab so much easier. Also, thanks to **Nina** and the postdoctoral fellows **Christina** and **Christine**, who recently joined the lab.



Present and former PhD students of the department: **Alexandros, Anders, Axel, Charles, Daniel, Elsa, Emely, Hammurabi, Gisela, Josef, Karolin, Kristina, Linda, Mia, Niklas, Pablo, Setareh,** and **Ylva**, who have been great companions at coffee breaks and at conferences. I also want to thank you for all the good times and interesting discussions. It would not have been the same without you.

**Anette Welin**, for your admirable work in keeping this department running and for efficiently sorting out all kinds of administrative matters.

Also, thanks to **Linda Magnusson, Jenny Nilsson, Andrea Biloglav, Helena Sturesson, Eleanor Woodward, Jenny Karlsson, Linda Holmquist Mengelbier, Rebeqa Gunnarsson, Kristina Lundin, Johnbosco Tayebwa, Catarina Lundin** and **Margreth Isaksson** for always being helpful.

All my co-workers at the Department of Clinical Genetics at the University Hospital, and at BMC B13.

A big thank you to my wonderful friends **Carolina, Kalle,** and **Ällen**, who have always been there for some rock climbing, skiing, “fika”, or just a “walk and talk”. You have all been a great support throughout the years. **Daniel** and **Marianne**, for sharing dorm with me and staying good friends after we moved out, even though we may not be able to meet as often as we used to.

Finally, I especially want to thank my entire family for their love and support. My parents **Bodil** and **Bosse**, my brothers **Erik** and **Olof**, and my grandmother **Disa**, for always encouraging me and believing in me. **Nora**, for all your support and for letting me enjoy life with you!

# References

- Aaronson, D.S., and Horvath, C.M. (2002). A road map for those who don't know JAK-STAT. *Science* 296, 1653–1655.
- Adolfsson, J., Månsson, R., Buza-Vidas, N., Hultquist, A., Liuba, K., Jensen, C.T., Bryder, D., Yang, L., Borge, O.-J., Thoren, L.A.M., Anderson, K., Sitnicka, E., Sasaki, Y., Sigvardsson, M., and Jacobsen, S.E.W. (2005). Identification of Flt3+ lympho-myeloid stem cells lacking erythro-megakaryocytic potential a revised road map for adult blood lineage commitment. *Cell* 121, 295–306.
- Akashi, K., Traver, D., Miyamoto, T., and Weissman, I.L. (2000). A clonogenic common myeloid progenitor that gives rise to all myeloid lineages. *Nature* 404, 193–197.
- Askmyr, M., Agerstam, H., Hansen, N., Gordon, S., Arvanitakis, A., Rissler, M., Juliusson, G., Richter, J., Järås, M., and Fioretos, T. (2013). Selective killing of candidate AML stem cells by antibody targeting of IL1RAP. *Blood* 121, 3709-3713.
- Baccarani, M., Deininger, M.W., Rosti, G., Hochhaus, A., Soverini, S., Apperley, J.F., Cervantes, F., Clark, R.E., Cortes, J.E., Guilhot, F., Hjorth-Hansen, H., Hughes, T.P., Kantarjian, H.M., Kim, D.-W., Larson, R.A., Lipton, J.H., Mahon, F.-X., Martinelli, G., Mayer, J., Müller, M.C., Niederwieser, D., Pane, F., Radich, J.P., Rousset, P., Saglio, G., Sauße, S., Schiffer, C., Silver, R., Simonsson, B., Steegmann, J.-L., Goldman, J.M., and Hehlmann, R. (2013). European LeukemiaNet recommendations for the management of chronic myeloid leukemia: 2013. *Blood* 122, 872–884.
- Barboudi, A., Höglund, M., Johansson, B., Lassen, C., Nilsson, P.-G., Hagemeijer, A., Mitelman, F., and Fioretos, T. (2003). A novel gene, MSI2, encoding a putative RNA-binding protein is recurrently rearranged at disease progression of chronic myeloid leukemia and forms a fusion gene with HOXA9 as a result of the cryptic t(7;17)(p15;q23). *Cancer Research*. 63, 1202–1206.
- Barnes, D.J., Schultheis, B., Adedeji, S., and Melo, J.V. (2005). Dose-dependent effects of Bcr-Abl in cell line models of different stages of chronic myeloid leukemia. *Oncogene* 24, 6432–6440.
- Barreyro, L., Will, B., Bartholdy, B., Zhou, L., Todorova, T.I., Stanley, R.F., Ben-Neriah, S., Montagna, C., Parekh, S., Pellagatti, A., Boultonwood, J., Paietta, E., Ketterling, R.P., Cripe, L., Fernandez, H.F., Greenberg, P.L., Tallman, M.S., Steidl, C., Mitsiades, C.S., Verma,

- A., and Steidl, U. (2012). Overexpression of IL-1 receptor accessory protein in stem and progenitor cells and outcome correlation in AML and MDS. *Blood* 120, 1290–1298.
- Barth, M., Raetz, E., and Cairo, M.S. (2012). The future role of monoclonal antibody therapy in childhood acute leukaemias. *British journal of haematology* 159, 3–17.
- Bellodi, C., Lidonnici, M.R., Hamilton, A., Helgason, G.V., Soliera, A.R., Ronchetti, M., Galavotti, S., Young, K.W., Selmi, T., Yacobi, R., Van Etten, R.A., Donato, N., Hunter, A., Dinsdale, D., Tirrò, E., Vigneri, P., Nicotera, P., Dyer, M.J., Holyoake, T., Salomoni, P., and Calabretta, B. (2009). Targeting autophagy potentiates tyrosine kinase inhibitor-induced cell death in Philadelphia chromosome-positive cells, including primary CML stem cells. *The Journal of clinical investigation* 119, 1109–1123.
- Bernstein, I.D., Andrews, R.G., and Zsebo, K.M. (1991). Recombinant human stem cell factor enhances the formation of colonies by CD34+ and CD34+lin- cells, and the generation of colony-forming cell progeny from CD34+lin- cells cultured with interleukin-3, granulocyte colony-stimulating factor, or granulocyte-macrophage colony-stimulating factor. *Blood* 77, 2316–2321.
- Bonnet, D., and Dick, J.E. (1997). Human acute myeloid leukemia is organized as a hierarchy that originates from a primitive hematopoietic cell. *Nature medicine* 3, 730–737.
- Borthakur, G., Kantarjian, H., Daley, G., Talpaz, M., O'Brien, S., Garcia-Manero, G., Giles, F., Faderl, S., Sugrue, M., and Cortes, J. (2006). Pilot study of lonafarnib, a farnesyl transferase inhibitor, in patients with chronic myeloid leukemia in the chronic or accelerated phase that is resistant or refractory to imatinib therapy. *Cancer* 106, 346–352.
- Brikos, C., Wait, R., Begum, S., O'Neill, L.A.J., and Saklatvala, J. (2007). Mass spectrometric analysis of the endogenous type I interleukin-1 (IL-1) receptor signaling complex formed after IL-1 binding identifies IL-1RAcP, MyD88, and IRAK-4 as the stable components. *Molecular & cellular proteomics* 6, 1551–1559.
- Broudy, V.C. (1997). Stem cell factor and hematopoiesis. *Blood* 90, 1345–1364.
- Cancer Genome Atlas Research Network (2013a). Genomic and epigenomic landscapes of adult de novo acute myeloid leukemia. *The New England journal of medicine* 368, 2059–2074.
- Cancer Genome Atlas Research Network (2013b). The Cancer Genome Atlas Pan-Cancer analysis project. *Nature genetics* 45, 1113–1120.
- Carlesso, N., Frank, D.A., and Griffin, J.D. (1996). Tyrosyl phosphorylation and DNA binding activity of signal transducers and activators of transcription (STAT) proteins in hematopoietic cell lines transformed by Bcr/Abl. *The Journal of experimental medicine* 183, 811–820.

- Casnici, C., Volpe, G., Crotta, K., Panuzzo, C., Lattuada, D., Cabras, C.A., Longhi, R., Saglio, G., and Marelli, O. (2011). Characterization of a monoclonal antibody specific for novel Bcr/Abl out-of-frame fusion proteins. *Hybridoma* 30, 261–269.
- Castor, A., Nilsson, L., Astrand-Grundström, I., Buitenhuis, M., Ramirez, C., Anderson, K., Strömbeck, B., Garwicz, S., Békássy, A.N., Schmiegelow, K., Lausen, B., Hokland, P., Lehmann, S., Juliusson, G., Johansson, B., and Jacobsen, S.E.W. (2005). Distinct patterns of hematopoietic stem cell involvement in acute lymphoblastic leukemia. *Nature medicine* 11, 630–637.
- Chai, S.K., Nichols, G.L., and Rothman, P. (1997). Constitutive activation of JAKs and STATs in BCR-Abl-expressing cell lines and peripheral blood cells derived from leukemic patients. *Journal of immunology* 159, 4720–4728.
- Chakraborty, S., Stark, J.M., Sun, C.-L., Modi, H., Chen, W., O'Connor, T.R., Forman, S.J., Bhatia, S., and Bhatia, R. (2012). Chronic myelogenous leukemia stem and progenitor cells demonstrate chromosomal instability related to repeated breakage-fusion-bridge cycles mediated by increased nonhomologous end joining. *Blood* 119, 6187–6197.
- Chen, Y., Hu, Y., Zhang, H., Peng, C., and Li, S. (2009). Loss of the Alox5 gene impairs leukemia stem cells and prevents chronic myeloid leukemia. *Nature genetics* 41, 783–792.
- Cilloni, D., Messa, F., Arruga, F., Defilippi, I., Morotti, A., Messa, E., Carturan, S., Giugliano, E., Pautasso, M., Bracco, E., Rosso, V., Sen, A., Martinelli, G., Baccarani, M., and Saglio, G. (2006). The NF-kappaB pathway blockade by the IKK inhibitor PS1145 can overcome imatinib resistance. *Leukemia* 20, 61–67.
- Cobaleda, C., Gutiérrez-Cianca, N., Pérez-Losada, J., Flores, T., García-Sanz, R., González, M., and Sánchez-García, I. (2000). A primitive hematopoietic cell is the target for the leukemic transformation in human philadelphia-positive acute lymphoblastic leukemia. *Blood* 95, 1007–1013.
- Corbin, A.S., Agarwal, A., Loriaux, M., Cortes, J., Deininger, M.W., and Druker, B.J. (2011). Human chronic myeloid leukemia stem cells are insensitive to imatinib despite inhibition of BCR-ABL activity. *The Journal of clinical investigation* 121, 396–409.
- Corbin, A.S., O'Hare, T., Gu, Z., Kraft, I.L., Eiring, A.M., Khorashad, J.S., Pomictter, A.D., Zhang, T.Y., Eide, C.A., Manley, P.W., Cortes, J.E., Druker, B.J., and Deininger, M.W. (2013). KIT signaling governs differential sensitivity of mature and primitive CML progenitors to tyrosine kinase inhibitors. *Cancer research* 73, 5775-86.
- Cortes, J., Albitar, M., Thomas, D., Giles, F., Kurzrock, R., Thibault, A., Rackoff, W., Koller, C., O'Brien, S., Garcia-Manero, G., Talpaz, M., and Kantarjian, H. (2003). Efficacy of the farnesyl transferase inhibitor R115777 in chronic myeloid leukemia and other hematologic malignancies. *Blood* 101, 1692–1697.

- Cortes, J.E., Kantarjian, H., Shah, N.P., Bixby, D., Mauro, M.J., Flinn, I., O'Hare, T., Hu, S., Narasimhan, N.I., Rivera, V.M., Clackson, T., Turner, C.D., Haluska, F.G., Druker, B.J., Deininger, M.W.N., and Talpaz, M. (2012). Ponatinib in refractory Philadelphia chromosome-positive leukemias. *The New England journal of medicine* 367, 2075–2088.
- Cortez, D., Reuther, G., and Pendergast, A.M. (1997). The Bcr-Abl tyrosine kinase activates mitogenic signaling pathways and stimulates G1-to-S phase transition in hematopoietic cells. *Oncogene* 15, 2333–2342.
- Cozzio, A., Passegué, E., Ayton, P.M., Karsunky, H., Cleary, M.L., and Weissman, I.L. (2003). Similar MLL-associated leukemias arising from self-renewing stem cells and short-lived myeloid progenitors. *Genes & development* 17, 3029–3035.
- Cramer-Morales, K., Nieborowska-Skorska, M., Scheibner, K., Padgett, M., Irvine, D.A., Sliwinski, T., Haas, K., Lee, J., Geng, H., Roy, D., Slupianek, A., Rassool, F.V., Wasik, M.A., Childers, W., Copland, M., Müschen, M., Civin, C.I., and Skorski, T. (2013). Personalized synthetic lethality induced by targeting RAD52 in leukemias identified by gene mutation and expression profile. *Blood* 122, 1293–1304.
- Cullinan, E.B., Kwee, L., Nunes, P., Shuster, D.J., Ju, G., McIntyre, K.W., Chizzonite, R.A., and Labow, M.A. (1998). IL-1 receptor accessory protein is an essential component of the IL-1 receptor. *Journal of immunology* 161, 5614–5620.
- Daley, G.Q., Van Etten, R.A., and Baltimore, D. (1990). Induction of chronic myelogenous leukemia in mice by the P210bcr/abl gene of the Philadelphia chromosome. *Science* 247, 824–830.
- Dao, T., and Scheinberg, D.A. (2008). Peptide vaccines for myeloid leukaemias. *Best practice & research. Clinical haematology* 21, 391–404.
- Dao, T., Yan, S., Veomett, N., Pankov, D., Zhou, L., Korontsvit, T., Scott, A., Whitten, J., Maslak, P., Casey, E., Tan, T., Liu, H., Zakhaleva, V., Curcio, M., Doubrovina, E., O'Reilly, R.J., Liu, C., and Scheinberg, D.A. (2013). Targeting the intracellular WT1 oncogene product with a therapeutic human antibody. *Science translational medicine* 176ra33.
- Dash, A.B., Williams, I.R., Kutok, J.L., Tomasson, M.H., Anastasiadou, E., Lindahl, K., Li, S., Van Etten, R.A., Borrow, J., Housman, D., Druker, B., and Gilliland, D.G. (2002). A murine model of CML blast crisis induced by cooperation between BCR/ABL and NUP98/HOXA9. *Proceedings of the National Academy of Sciences of the United States of America* 99, 7622–7627.
- Diamond, J.M., and Melo, J.V. (2011). Mechanisms of resistance to BCR-ABL kinase inhibitors. *Leukemia & lymphoma* 52 Suppl 1, 12–22.

- Dick, J.E., and Lapidot, T. (2005). Biology of normal and acute myeloid leukemia stem cells. *International journal of hematology* 82, 389–396.
- Dirks, W.G., MacLeod, R.A.F., Nakamura, Y., Kohara, A., Reid, Y., Milch, H., Drexler, H.G., and Mizusawa, H. (2010). Cell line cross-contamination initiative: an interactive reference database of STR profiles covering common cancer cell lines. *International journal of cancer* 126, 303–304.
- Dorritie, K.A., McCubrey, J.A., and Johnson, D.E. (2013). STAT transcription factors in hematopoiesis and leukemogenesis: opportunities for therapeutic intervention. *Leukemia* Epub ahead of print.
- Doulatov, S., Notta, F., Eppert, K., Nguyen, L.T., Ohashi, P.S., and Dick, J.E. (2010). Revised map of the human progenitor hierarchy shows the origin of macrophages and dendritic cells in early lymphoid development. *Nature immunology* 11, 585–593.
- Doulatov, S., Notta, F., Laurenti, E., and Dick, J.E. (2012). Hematopoiesis: a human perspective. *Cell Stem Cell* 10, 120–136.
- Downward, J. (2003). Targeting RAS signalling pathways in cancer therapy. *Nature reviews Cancer* 3, 11–22.
- Drexler, H.G., Dirks, W.G., and MacLeod, R.A. (1999). False human hematopoietic cell lines: cross-contaminations and misinterpretations. *Leukemia* 13, 1601–1607.
- Druker, B., Okuda, K., Matulonis, U., Salgia, R., Roberts, T., and Griffin, J.D. (1992). Tyrosine phosphorylation of rasGAP and associated proteins in chronic myelogenous leukemia cell lines. *Blood* 79, 2215–2220.
- Druker, B.J., Guilhot, F., O'Brien, S.G., Gathmann, I., Kantarjian, H., Gattermann, N., Deininger, M.W.N., Silver, R.T., Goldman, J.M., Stone, R.M., Cervantes, F., Hochhaus, A., Powell, B.L., Gabrilove, J.L., Rouselot, P., Reiffers, J., Cornelissen, J.J., Hughes, T., Agis, H., Fischer, T., Verhoef, G., Shepherd, J., Saglio, G., Gratwohl, A., Nielsen, J.L., Radich, J.P., Simonsson, B., Taylor, K., Baccarani, M., So, C., Letvak, L., Larson, R.A., IRIS investigators (2006). Five-year follow-up of patients receiving imatinib for chronic myeloid leukemia. *The New England journal of medicine* 355, 2408–2417.
- Druker, B.J., Talpaz, M., Resta, D.J., Peng, B., Buchdunger, E., Ford, J.M., Lydon, N.B., Kantarjian, H., Capdeville, R., Ohno-Jones, S., and Sawyers, C.L. (2001). Efficacy and safety of a specific inhibitor of the BCR-ABL tyrosine kinase in chronic myeloid leukemia. *The New England journal of medicine* 344, 1031–1037.
- Druker, B.J., Tamura, S., Buchdunger, E., Ohno, S., Segal, G.M., Fanning, S., Zimmermann, J., and Lydon, N.B. (1996). Effects of a selective inhibitor of the Abl tyrosine kinase on the growth of Bcr-Abl positive cells. *Nature medicine* 2, 561–566.

- Duarte, R.F., and Frank, D.A. (2000). SCF and G-CSF lead to the synergistic induction of proliferation and gene expression through complementary signaling pathways. *Blood* 96, 3422–3430.
- Endo, T.A., Masuhara, M., Yokouchi, M., Suzuki, R., Sakamoto, H., Mitsui, K., Matsumoto, A., Tanimura, S., Ohtsubo, M., Misawa, H., Miyazaki, T., Leonor, N., Taniguchi, T., Fujita, T., Kanakura, Y., Komiya, S., and Yoshimura, A. (1997). A new protein containing an SH2 domain that inhibits JAK kinases. *Nature* 387, 921–924.
- Eppert, K., Takenaka, K., Lechman, E.R., Waldron, L., Nilsson, B., van Galen, P., Metzeler, K.H., Poepl, A., Ling, V., Beyene, J., Canty, A.J., Danska, J.S., Bohlander, S.K., Buske, C., Minden, M.D., Golub, T.R., Jurisica, I., Ebert, B.L., and Dick, J.E. (2011). Stem cell gene expression programs influence clinical outcome in human leukemia. *Nature medicine* 17, 1086–1093.
- Feinstein, E., Cimino, G., Gale, R.P., Alimena, G., Berthier, R., Kishi, K., Goldman, J., Zaccaria, A., Berrebi, A., and Canaani, E. (1991). p53 in chronic myelogenous leukemia in acute phase. *Proceedings of the National Academy of Sciences of the United States of America* 88, 6293–6297.
- Feuring-Buske, M., Gerhard, B., Cashman, J., Humphries, R.K., Eaves, C.J., and Hogge, D.E. (2003). Improved engraftment of human acute myeloid leukemia progenitor cells in beta 2-microglobulin-deficient NOD/SCID mice and in NOD/SCID mice transgenic for human growth factors. *Leukemia* 17, 760–763.
- Fialkow, P.J., Denman, A.M., Jacobson, R.J., and Lowenthal, M.N. (1978). Chronic myelocytic leukemia. Origin of some lymphocytes from leukemic stem cells. *The Journal of clinical investigation* 62, 815–823.
- Fialkow, P.J., Jacobson, R.J., and Papayannopoulou, T. (1977). Chronic myelocytic leukemia: clonal origin in a stem cell common to the granulocyte, erythrocyte, platelet and monocyte/macrophage. *The American journal of medicine* 63, 125–130.
- Fitzgerald, K.A., and O'Neill, L.A. (2000). The role of the interleukin-1/Toll-like receptor superfamily in inflammation and host defence. *Microbes and infection* 2, 933–943.
- Friedman, A.D. (2007). Transcriptional control of granulocyte and monocyte development. *Oncogene* 26, 6816–6828.
- Gaiger, A., Henn, T., Hörth, E., Geissler, K., Mitterbauer, G., Maier-Dobersberger, T., Greinix, H., Mannhalter, C., Haas, O.A., Lechner, K., and Lion, T. (1995). Increase of bcr-abl chimeric mRNA expression in tumor cells of patients with chronic myeloid leukemia precedes disease progression. *Blood* 86, 2371–2378.

- Gentles, A.J., Plevritis, S.K., Majeti, R., and Alizadeh, A.A. (2010). Association of a leukemic stem cell gene expression signature with clinical outcomes in acute myeloid leukemia. *The Journal of the American Medical Association* 304, 2706–2715.
- Gerondakis, S., Banerjee, A., Grigoriadis, G., Vasanthakumar, A., Gugasyan, R., Sidwell, T., and Grumont, R.J. (2012). NF- $\kappa$ B subunit specificity in hemopoiesis. *Immunological reviews* 246, 272–285.
- Gibbs, J.B., Oliff, A., and Kohl, N.E. (1994). Farnesyltransferase inhibitors: Ras research yields a potential cancer therapeutic. *Cell* 77, 175–178.
- Giebel, B., and Punzel, M. (2008). Lineage development of hematopoietic stem and progenitor cells. *Biological chemistry* 389, 813–824.
- Gishizky, M.L., Cortez, D., and Pendergast, A.M. (1995). Mutant forms of growth factor-binding protein-2 reverse BCR-ABL-induced transformation. *Proceedings of the National Academy of Sciences of the United States of America* 92, 10889–10893.
- Golemovic, M., Verstovsek, S., Giles, F., Cortes, J., Manshour, T., Manley, P.W., Mestan, J., Dugan, M., Alland, L., Griffin, J.D., Arlinghaus, R.B., Sun, T., Kantarjian, H., and Beran, M. (2005). AMN107, a novel aminopyrimidine inhibitor of Bcr-Abl, has in vitro activity against imatinib-resistant chronic myeloid leukemia. *Clinical cancer research* 11, 4941–4947.
- Görgens, A., Radtke, S., Möllmann, M., Cross, M., Dürig, J., Horn, P.A., and Giebel, B. (2013). Revision of the human hematopoietic tree: granulocyte subtypes derive from distinct hematopoietic lineages. *Cell Reports* 3, 1539–1552.
- Grossmann, V., Kohlmann, A., Zenger, M., Schindela, S., Eder, C., Weissmann, S., Schnittger, S., Kern, W., Müller, M.C., Hochhaus, A., Haferlach, T., and Haferlach, C. (2011). A deep-sequencing study of chronic myeloid leukemia patients in blast crisis (BC-CML) detects mutations in 76.9% of cases. *Leukemia* 25, 557–560.
- Grosveld, G., Verwoerd, T., van Agthoven, T., de Klein, A., Ramachandran, K.L., Heisterkamp, N., Stam, K., and Groffen, J. (1986). The chronic myelocytic cell line K562 contains a breakpoint in bcr and produces a chimeric bcr/c-abl transcript. *Molecular and cellular biology* 6, 607–616.
- Guzman, M.L., Neering, S.J., Upchurch, D., Grimes, B., Howard, D.S., Rizzieri, D.A., Luger, S.M., and Jordan, C.T. (2001). Nuclear factor-kappaB is constitutively activated in primitive human acute myelogenous leukemia cells. *Blood* 98, 2301–2307.
- Guzman, M.L., Rossi, R.M., Karnischky, L., Li, X., Peterson, D.R., Howard, D.S., and Jordan, C.T. (2005). The sesquiterpene lactone parthenolide induces apoptosis of human acute myelogenous leukemia stem and progenitor cells. *Blood* 105, 4163–4169.



- Guzman, M.L., Swiderski, C.F., Howard, D.S., Grimes, B.A., Rossi, R.M., Szilvassy, S.J., and Jordan, C.T. (2002). Preferential induction of apoptosis for primary human leukemic stem cells. *Proceedings of the National Academy of Sciences of the United States of America* 99, 16220–16225.
- Hamdane, M., David-Cordonnier, M.H., and D'Halluin, J.C. (1997). Activation of p65 NF-kappaB protein by p210BCR-ABL in a myeloid cell line (P210BCR-ABL activates p65 NF-kappaB). *Oncogene* 15, 2267–2275.
- Hamilton, A., Helgason, G.V., Schemionek, M., Zhang, B., Myssina, S., Allan, E.K., Nicolini, F.E., Müller-Tidow, C., Bhatia, R., Brunton, V.G., Koschmieder, S., and Holyoake, T.L. (2012). Chronic myeloid leukemia stem cells are not dependent on Bcr-Abl kinase activity for their survival. *Blood* 119, 1501–1510.
- Hanahan, D., and Weinberg, R.A. (2011). Hallmarks of cancer: the next generation. *Cell* 144, 646–674.
- Hancock, J.F., Paterson, H., and Marshall, C.J. (1990). A polybasic domain or palmitoylation is required in addition to the CAAX motif to localize p21ras to the plasma membrane. *Cell* 63, 133–139.
- Hantschel, O., Warsch, W., Eckelhart, E., Kaupe, I., Grebien, F., Wagner, K.-U., Superti-Furga, G., and Sexl, V. (2012). BCR-ABL uncouples canonical JAK2-STAT5 signaling in chronic myeloid leukemia. *Nature chemical biology* 8, 285–293.
- Håkansson, P., Nilsson, B., Andersson, A., Lassen, C., Gullberg, U., and Fioretos, T. (2008). Gene expression analysis of BCR/ABL1-dependent transcriptional response reveals enrichment for genes involved in negative feedback regulation. *Genes Chromosomes Cancer* 47, 267–275.
- Heisterkamp, N., Stam, K., Groffen, J., de Klein, A., and Grosveld, G. (1985). Structural organization of the bcr gene and its role in the Ph' translocation. *Nature* 315, 758–761.
- Hochhaus, A., Kreil, S., Corbin, A.S., La Rosée, P., Müller, M.C., Lahaye, T., Hanfstein, B., Schoch, C., Cross, N.C.P., Berger, U., Gschaidmeier, H., Druker, B.J., and Hehlmann, R. (2002). Molecular and chromosomal mechanisms of resistance to imatinib (STI571) therapy. *Leukemia* 16, 2190–2196.
- Hochhaus, A., O'Brien, S.G., Guilhot, F., Druker, B.J., Branford, S., Foroni, L., Goldman, J.M., Müller, M.C., Radich, J.P., Rudoltz, M., Mone, M., Gathmann, I., Hughes, T.P., Larson, R.A., IRIS investigators (2009). Six-year follow-up of patients receiving imatinib for the first-line treatment of chronic myeloid leukemia. *Leukemia* 23, 1054–1061.
- Hoelbl, A., Kovacic, B., Kerenyi, M.A., Simma, O., Warsch, W., Cui, Y., Beug, H., Hennighausen, L., Moriggl, R., and Sexl, V. (2006). Clarifying the role of Stat5 in lymphoid development and Abelson-induced transformation. *Blood* 107, 4898–4906.

- Hoelbl, A., Schuster, C., Kovacic, B., Zhu, B., Wickre, M., Hoelzl, M.A., Fajmann, S., Grebien, F., Warsch, W., Stengl, G., Hennighausen, L., Poli, V., Beug, H., Moriggl, R., and Sexl, V. (2010). Stat5 is indispensable for the maintenance of bcr/abl-positive leukaemia. *EMBO molecular medicine* 2, 98–110.
- Huettner, C.S., Koschmieder, S., Iwasaki, H., Iwasaki-Arai, J., Radomska, H.S., Akashi, K., and Tenen, D.G. (2003). Inducible expression of BCR/ABL using human CD34 regulatory elements results in a megakaryocytic myeloproliferative syndrome. *Blood* 102, 3363–3370.
- Hughes, T.P., Hochhaus, A., Branford, S., Müller, M.C., Kaeda, J.S., Foroni, L., Druker, B.J., Guilhot, F., Larson, R.A., O'Brien, S.G., Rudoltz, M.S., Mone, M., Wehrle, E., Modur, V., Goldman, J.M., Radich, J.P., IRIS investigators (2010). Long-term prognostic significance of early molecular response to imatinib in newly diagnosed chronic myeloid leukemia: an analysis from the International Randomized Study of Interferon and STI571 (IRIS). *Blood* 116, 3758–3765.
- Huntly, B.J.P., Shigematsu, H., Deguchi, K., Lee, B.H., Mizuno, S., Duclos, N., Rowan, R., Amaral, S., Curley, D., Williams, I.R., Akashi, K., and Gilliland, D.G. (2004). MOZ-TIF2, but not BCR-ABL, confers properties of leukemic stem cells to committed murine hematopoietic progenitors. *Cancer Cell* 6, 587–596.
- Iglesias-Gato, D., Chuan, Y.-C., Wikström, P., Augsten, S., Jiang, N., Niu, Y., Seipel, A., Danneman, D., Vermeij, M., Fernandez-Perez, L., Jenster, G., Egevad, L., Norstedt, G., and Flores-Morales, A. (2013). SOCS2 mediates the crosstalk between androgen and growth hormone signaling in prostate cancer. *Carcinogenesis* Epub ahead of print.
- Ilaria, R.L., and Van Etten, R.A. (1996). P210 and P190(BCR/ABL) induce the tyrosine phosphorylation and DNA binding activity of multiple specific STAT family members. *The Journal of biological chemistry* 271, 31704–31710.
- Jabbour, E., and Kantarjian, H. (2012). Chronic myeloid leukemia: 2012 update on diagnosis, monitoring, and management. *American journal of hematology* 87, 1037–1045.
- Jabbour, E., Cortes, J., Santos, F.P.S., Jones, D., O'Brien, S., Rondon, G., Popat, U., Giral, S., Kebriaei, P., Jones, R.B., Kantarjian, H., Champlin, R., and de Lima, M. (2011). Results of allogeneic hematopoietic stem cell transplantation for chronic myelogenous leukemia patients who failed tyrosine kinase inhibitors after developing BCR-ABL1 kinase domain mutations. *Blood* 117, 3641–3647.
- Jain, N., and van Besien, K. (2011). Chronic myelogenous leukemia: role of stem cell transplant in the imatinib era. *Hematology/oncology clinics of North America* 25, 1025–1048.
- Jamieson, C.H.M., Ailles, L.E., Dylla, S.J., Muijtjens, M., Jones, C., Zehnder, J.L., Gotlib, J., Li, K., Manz, M.G., Keating, A., Sawyers, C.L., and Weissman, I.L. (2004). Granulocyte-macrophage progenitors as candidate leukemic stem cells in blast-crisis CML. *The New England journal of medicine* 351, 657–667.

- Järås, M., Johnels, P., Hansen, N., Agerstam, H., Tsapogas, P., Rissler, M., Lassen, C., Olofsson, T., Bjerrum, O.W., Richter, J., and Fioretos, T. (2010). Isolation and killing of candidate chronic myeloid leukemia stem cells by antibody targeting of IL-1 receptor accessory protein. *Proceedings of the National Academy of Sciences of the United States of America* 107, 16280–16285.
- Jiang, X., Zhao, Y., Smith, C., Gasparetto, M., Turhan, A., Eaves, A., and Eaves, C. (2007). Chronic myeloid leukemia stem cells possess multiple unique features of resistance to BCR-ABL targeted therapies. *Leukemia* 21, 926–935.
- Johansson, B., Fioretos, T., and Mitelman, F. (2002). Cytogenetic and molecular genetic evolution of chronic myeloid leukemia. *Acta Haematol.* 107, 76–94.
- Kamitsuji, Y., Kuroda, J., Kimura, S., Toyokuni, S., Watanabe, K., Ashihara, E., Tanaka, H., Yui, Y., Watanabe, M., Matsubara, H., Mizushima, Y., Hiraumi, Y., Kawata, E., Yoshikawa, T., Maekawa, T., Nakahata, T., and Adachi, S. (2008). The Bcr-Abl kinase inhibitor INNO-406 induces autophagy and different modes of cell death execution in Bcr-Abl-positive leukemias. *Cell death and differentiation.* 15, 1712–1722.
- Kantarjian, H., Giles, F., Wunderle, L., Bhalla, K., O'Brien, S., Wassmann, B., Tanaka, C., Manley, P., Rae, P., Mietlowski, W., Bochinski, K., Hochhaus, A., Griffin, J.D., Hoelzer, D., Albitar, M., Dugan, M., Cortes, J., Alland, L., and Ottmann, O.G. (2006). Nilotinib in imatinib-resistant CML and Philadelphia chromosome-positive ALL. *The New England journal of medicine* 354, 2542–2551.
- Kantarjian, H., O'Brien, S., Jabbour, E., Garcia-Manero, G., Quintás-Cardama, A., Shan, J., Rios, M.B., Ravandi, F., Faderl, S., Kadia, T., Borthakur, G., Huang, X., Champlin, R., Talpaz, M., and Cortes, J. (2012). Improved survival in chronic myeloid leukemia since the introduction of imatinib therapy: a single-institution historical experience. *Blood* 119, 1981–1987.
- Karin, M., Cao, Y., Greten, F.R., and Li, Z.-W. (2002). NF-kappaB in cancer: from innocent bystander to major culprit. *Nature reviews Cancer* 2, 301–310.
- Kato, K., Cox, A.D., Hisaka, M.M., Graham, S.M., Buss, J.E., and Der, C.J. (1992). Isoprenoid addition to Ras protein is the critical modification for its membrane association and transforming activity. *Proceedings of the National Academy of Sciences of the United States of America* 89, 6403–6407.
- Kawamoto, H., Ikawa, T., Masuda, K., Wada, H., and Katsura, Y. (2010). A map for lineage restriction of progenitors during hematopoiesis: the essence of the myeloid-based model. *Immunological reviews*, 238, 23–36.
- Kirchner, D., Duyster, J., Ottmann, O., Schmid, R.M., Bergmann, L., and Munzert, G. (2003). Mechanisms of Bcr-Abl-mediated NF-kappaB/Rel activation. *Experimental hematology* 31, 504–511.

- Kobayashi, M., Laver, J.H., Kato, T., Miyazaki, H., and Ogawa, M. (1996). Thrombopoietin supports proliferation of human primitive hematopoietic cells in synergy with steel factor and/or interleukin-3. *Blood* 88, 429–436.
- Kondo, M., Weissman, I.L., and Akashi, K. (1997). Identification of clonogenic common lymphoid progenitors in mouse bone marrow. *Cell* 91, 661–672.
- Koschmieder, S., and Schemionek, M. (2011). Mouse models as tools to understand and study BCR-ABL1 diseases. *American journal of blood research* 1, 65–75.
- Koschmieder, S., Göttgens, B., Zhang, P., Iwasaki-Arai, J., Akashi, K., Kutok, J.L., Dayaram, T., Geary, K., Green, A.R., Tenen, D.G., and Huettner, C.S. (2005). Inducible chronic phase of myeloid leukemia with expansion of hematopoietic stem cells in a transgenic model of BCR-ABL leukemogenesis. *Blood* 105, 324–334.
- Krebs, D.L., and Hilton, D.J. (2001). SOCS proteins: negative regulators of cytokine signaling. *Stem Cells* 19, 378–387.
- Krivtsov, A.V., Twomey, D., Feng, Z., Stubbs, M.C., Wang, Y., Faber, J., Levine, J.E., Wang, J., Hahn, W.C., Gilliland, D.G., Golub, T.R., and Armstrong, S.A. (2006). Transformation from committed progenitor to leukaemia stem cell initiated by MLL-AF9. *Nature* 442, 818–822.
- Kuo, H.-P., Wang, Z., Lee, D.-F., Iwasaki, M., Duque-Afonso, J., Wong, S.H.K., Lin, C.-H., Figueroa, M.E., Su, J., Lemischka, I.R., and Cleary, M.L. (2013). Epigenetic Roles of MLL Oncoproteins Are Dependent on NF- $\kappa$ B. *Cancer Cell*.
- Lander, E.S., Linton, L.M., Birren, B., Nusbaum, C., Zody, M.C., Baldwin, J., Devon, K., Dewar, K., Doyle, M., FitzHugh, W., Funke, R., Gage, D., Harris, K., Heaford, A., Howland, J., Kann, L., Lehoczy, J., LeVine, R., McEwan, P., McKernan, K., Meldrim, J., Mesirov, J.P., Miranda, C., Morris, W., Naylor, J., Raymond, C., Rosetti, M., Santos, R., Sheridan, A., Sougnez, C., Stange-Thomann, N., Stojanovic, N., Subramanian, A., Wyman, D., Rogers, J., Sulston, J., Ainscough, R., Beck, S., Bentley, D., Burton, J., Clee, C., Carter, N., Coulson, A., Deadman, R., Deloukas, P., Dunham, A., Dunham, I., Durbin, R., French, L., Grafham, D., Gregory, S., Hubbard, T., Humphray, S., Hunt, A., Jones, M., Lloyd, C., McMurray, A., Matthews, L., Mercer, S., Milne, S., Mullikin, J.C., Mungall, A., Plumb, R., Ross, M., Shownkeen, R., Sims, S., Waterston, R.H., Wilson, R.K., Hillier, L.W., McPherson, J.D., Marra, M.A., Mardis, E.R., Fulton, L.A., Chinwalla, A.T., Pepin, K.H., Gish, W.R., Chissoe, S.L., Wendl, M.C., Delehaunty, K.D., Miner, T.L., Delehaunty, A., Kramer, J.B., Cook, L.L., Fulton, R.S., Johnson, D.L., Minx, P.J., Clifton, S.W., Hawkins, T., Branscomb, E., Predki, P., Richardson, P., Wenning, S., Slezak, T., Doggett, N., Cheng, J.F., Olsen, A., Lucas, S., Elkin, C., Uberbacher, E., Frazier, M., Gibbs, R.A., Muzny, D.M., Scherer, S.E., Bouck, J.B., Sodergren, E.J., Worley, K.C., Rives, C.M., Gorrell, J.H., Metzker, M.L., Naylor, S.L., Kucherlapati, R.S., Nelson, D.L., Weinstock, G.M., Sakaki, Y., Fujiiyama, A., Hattori, M., Yada, T., Toyoda, A., Itoh, T., Kawagoe, C., Watanabe, H., Totoki, Y., Taylor, T., Weissenbach, J., Heilig,

- R., Saurin, W., Artiguenave, F., Brottier, P., Bruls, T., Pelletier, E., Robert, C., Wincker, P., Smith, D.R., Doucette-Stamm, L., Rubenfield, M., Weinstock, K., Lee, H.M., Dubois, J., Rosenthal, A., Platzer, M., Nyakatura, G., Taudien, S., Rump, A., Yang, H., Yu, J., Wang, J., Huang, G., Gu, J., Hood, L., Rowen, L., Madan, A., Qin, S., Davis, R.W., Federspiel, N.A., Abola, A.P., Proctor, M.J., Myers, R.M., Schmutz, J., Dickson, M., Grimwood, J., Cox, D.R., Olson, M.V., Kaul, R., Raymond, C., Shimizu, N., Kawasaki, K., Minoshima, S., Evans, G.A., Athanasiou, M., Schultz, R., Roe, B.A., Chen, F., Pan, H., Ramser, J., Lehrach, H., Reinhardt, R., McCombie, W.R., la Bastide, de, M., Dedhia, N., Blöcker, H., Hornischer, K., Nordsiek, G., Agarwala, R., Aravind, L., Bailey, J.A., Bateman, A., Batzoglou, S., Birney, E., Bork, P., Brown, D.G., Burge, C.B., Cerutti, L., Chen, H.C., Church, D., Clamp, M., Copley, R.R., Doerks, T., Eddy, S.R., Eichler, E.E., Furey, T.S., Galagan, J., Gilbert, J.G., Harmon, C., Hayashizaki, Y., Haussler, D., Hermjakob, H., Hokamp, K., Jang, W., Johnson, L.S., Jones, T.A., Kasif, S., Kasprzyk, A., Kennedy, S., Kent, W.J., Kitts, P., Koonin, E.V., Korf, I., Kulp, D., Lancet, D., Lowe, T.M., McLysaght, A., Mikkelsen, T., Moran, J.V., Mulder, N., Pollara, V.J., Ponting, C.P., Schuler, G., Schultz, J., Slater, G., Smit, A.F., Stupka, E., Szustakowski, J., Thierry-Mieg, D., Thierry-Mieg, J., Wagner, L., Wallis, J., Wheeler, R., Williams, A., Wolf, Y.I., Wolfe, K.H., Yang, S.P., Yeh, R.F., Collins, F., Guyer, M.S., Peterson, J., Felsenfeld, A., Wetterstrand, K.A., Patrinos, A., Morgan, M.J., de Jong, P., Catanese, J.J., Osoegawa, K., Shizuya, H., Choi, S., Chen, Y.J., Szustakowki, J., International Human Genome Sequencing Consortium (2001). Initial sequencing and analysis of the human genome. *Nature* 409, 860–921.
- Lapidot, T., Sirard, C., Vormoor, J., Murdoch, B., Hoang, T., Caceres-Cortes, J., Minden, M., Paterson, B., Caligiuri, M.A., and Dick, J.E. (1994). A cell initiating human acute myeloid leukaemia after transplantation into SCID mice. *Nature* 367, 645–648.
- le Viseur, C., Hotfilder, M., Bomken, S., Wilson, K., Röttgers, S., Schrauder, A., Rosemann, A., Irving, J., Stam, R.W., Shultz, L.D., Harbott, J., Jürgens, H., Schrappe, M., Pieters, R., and Vormoor, J. (2008). In childhood acute lymphoblastic leukemia, blasts at different stages of immunophenotypic maturation have stem cell properties. *Cancer Cell* 14, 47–58.
- Lewis, I.D., McDiarmid, L.A., Samels, L.M., To, L.B., and Hughes, T.P. (1998). Establishment of a reproducible model of chronic-phase chronic myeloid leukemia in NOD/SCID mice using blood-derived mononuclear or CD34+ cells. *Blood* 91, 630–640.
- Li, C.L., and Johnson, G.R. (1994). Stem cell factor enhances the survival but not the self-renewal of murine hematopoietic long-term repopulating cells. *Blood* 84, 408–414.
- Li, S., Ilaria, R.L., Million, R.P., Daley, G.Q., and Van Etten, R.A. (1999). The P190, P210, and P230 forms of the BCR/ABL oncogene induce a similar chronic myeloid leukemia-like syndrome in mice but have different lymphoid leukemogenic activity. *The Journal of experimental medicine* 189 1399–1412.

- Linnekin, D., Keller, J.R., Ferris, D.K., Mou, S.M., Broudy, V., and Longo, D.L. (1995). Stem cell factor induces phosphorylation of a 200 kDa protein which associates with c-kit. *Growth Factors* 12, 57–67.
- Lounnas, N., Frelin, C., Gonthier, N., Colosetti, P., Sirvent, A., Cassuto, J.-P., Berthier, F., Sirvent, N., Rousselot, P., Dreano, M., Peyron, J.-F., and Imbert, V. (2009). NF-kappaB inhibition triggers death of imatinib-sensitive and imatinib-resistant chronic myeloid leukemia cells including T315I Bcr-Abl mutants. *International journal of cancer* 125, 308–317.
- Lowry, P.A., Shultz, L.D., Greiner, D.L., Hesselton, R.M., Kittler, E.L., Tiarks, C.Y., Rao, S.S., Reilly, J., Leif, J.H., Ramshaw, H., Stewart, F.M., and Quesenberry, P.J. (1996). Improved engraftment of human cord blood stem cells in NOD/LtSz-scid/scid mice after irradiation or multiple-day injections into unirradiated recipients. *Biology of blood and marrow transplantation* 2, 15–23.
- Lu, Z., Jin, Y., Chen, C., Li, J., Cao, Q., and Pan, J. (2010). Pristimerin induces apoptosis in imatinib-resistant chronic myelogenous leukemia cells harboring T315I mutation by blocking NF-kappaB signaling and depleting Bcr-Abl. *Molecular cancer* 9, 112.
- Lugo, T.G., Pendergast, A.M., Muller, A.J., and Witte, O.N. (1990). Tyrosine kinase activity and transformation potency of bcr-abl oncogene products. *Science* 247, 1079–1082.
- Mahon, F.-X., Rea, D., Guilhot, J., Guilhot, F., Huguier, F., Nicolini, F., Legros, L., Charbonnier, A., Guerci, A., Varet, B., Etienne, G., Reiffers, J., Rousselot, P., Intergroupe Français des Leucémies Myéloïdes Chroniques (2010). Discontinuation of imatinib in patients with chronic myeloid leukaemia who have maintained complete molecular remission for at least 2 years: the prospective, multicentre Stop Imatinib (STIM) trial. *The lancet oncology* 11, 1029–1035.
- Makishima, H., Jankowska, A.M., McDevitt, M.A., O'Keefe, C., Dujardin, S., Cazzolli, H., Przychodzen, B., Prince, C., Nicoll, J., Siddaiah, H., Shaik, M., Szpurka, H., Hsi, E., Advani, A., Paquette, R., and Maciejewski, J.P. (2011). CBL, CBLB, TET2, ASXL1, and IDH1/2 mutations and additional chromosomal aberrations constitute molecular events in chronic myelogenous leukemia. *Blood* 117, e198–e206.
- Manz, M.G., Miyamoto, T., Akashi, K., and Weissman, I.L. (2002). Prospective isolation of human clonogenic common myeloid progenitors. *Proceedings of the National Academy of Sciences of the United States of America* 99, 11872–11877.
- Maslak, P.G., Dao, T., Gomez, M., Chanel, S., Packin, J., Korontsvit, T., Zakhaleva, V., Pinilla-Ibarz, J., Berman, E., and Scheinberg, D.A. (2008). A pilot vaccination trial of synthetic analog peptides derived from the BCR-ABL breakpoints in CML patients with minimal disease. *Leukemia* 22, 1613–1616.

- Maslak, P.G., Dao, T., Krug, L.M., Chanel, S., Korontsvit, T., Zakhaleva, V., Zhang, R., Wolchok, J.D., Yuan, J., Pinilla-Ibarz, J., Berman, E., Weiss, M., Jurcic, J., Frattini, M.G., and Scheinberg, D.A. (2010). Vaccination with synthetic analog peptides derived from WT1 oncoprotein induces T-cell responses in patients with complete remission from acute myeloid leukemia. *Blood* 116, 171–179.
- Merante, S., Orlandi, E., Bernasconi, P., Calatroni, S., Boni, M., and Lazzarino, M. (2005). Outcome of four patients with chronic myeloid leukemia after imatinib mesylate discontinuation. *Haematologica* 90, 979–981.
- Mes-Masson, A.M., McLaughlin, J., Daley, G.Q., Paskind, M., and Witte, O.N. (1986). Overlapping cDNA clones define the complete coding region for the P210c-abl gene product associated with chronic myelogenous leukemia cells containing the Philadelphia chromosome. *Proceedings of the National Academy of Sciences of the United States of America* 83, 9768–9772.
- Metcalf, D., and Nicola, N.A. (1991). Direct proliferative actions of stem cell factor on murine bone marrow cells in vitro: effects of combination with colony-stimulating factors. *Proceedings of the National Academy of Sciences of the United States of America* 88, 6239–6243.
- Mishima, Y., Terui, Y., Mishima, Y., Taniyama, A., Kuniyoshi, R., Takizawa, T., Kimura, S., Ozawa, K., and Hatake, K. (2008). Autophagy and autophagic cell death are next targets for elimination of the resistance to tyrosine kinase inhibitors. *Cancer Science*. 99, 2200–2208.
- Mitelman, F., Levan, G., Nilsson, P.G., and Brandt, L. (1976). Non-random karyotypic evolution in chronic myeloid leukemia. *International journal of cancer* 18, 24–30.
- Morris, J.C., and Waldmann, T.A. (2009). Antibody-based therapy of leukaemia. *Expert reviews in molecular medicine* 11, e29.
- Mullighan, C.G. (2013). Genome sequencing of lymphoid malignancies. *Blood* Epub ahead of print.
- Mullighan, C.G., Miller, C.B., Radtke, I., Phillips, L.A., Dalton, J., Ma, J., White, D., Hughes, T.P., Le Beau, M.M., Pui, C.-H., Relling, M.V., Shurtleff, S.A., and Downing, J.R. (2008). BCR-ABL1 lymphoblastic leukaemia is characterized by the deletion of Ikaros. *Nature* 453, 110–114.
- Nagasu, T., Yoshimatsu, K., Rowell, C., Lewis, M.D., and Garcia, A.M. (1995). Inhibition of human tumor xenograft growth by treatment with the farnesyl transferase inhibitor B956. *Cancer research* 55, 5310–5314.

- Naka, T., Narazaki, M., Hirata, M., Matsumoto, T., Minamoto, S., Aono, A., Nishimoto, N., Kajita, T., Taga, T., Yoshizaki, K., Akira, S., and Kishimoto, T. (1997). Structure and function of a new STAT-induced STAT inhibitor. *Nature* 387, 924–929.
- Nicolini, F.E., Cashman, J.D., Hogge, D.E., Humphries, R.K., and Eaves, C.J. (2004). NOD/SCID mice engineered to express human IL-3, GM-CSF and Steel factor constitutively mobilize engrafted human progenitors and compromise human stem cell regeneration. *Leukemia* 18, 341–347.
- Nilsson, L., Astrand-Grundström, I., Arvidsson, I., Jacobsson, B., Hellström-Lindberg, E., Hast, R., and Jacobsen, S.E. (2000). Isolation and characterization of hematopoietic progenitor/stem cells in 5q-deleted myelodysplastic syndromes: evidence for involvement at the hematopoietic stem cell level. *Blood* 96, 2012–2021.
- Nowell, P.C., and Hungerford, D. (1960). A minute chromosome in human granulocytic leukemia. *Science* 132, 1497.
- Nowicki, M.O., Falinski, R., Koptyra, M., Slupianek, A., Stoklosa, T., Gloc, E., Nieborowska-Skorska, M., Blasiak, J., and Skorski, T. (2004). BCR/ABL oncogenic kinase promotes unfaithful repair of the reactive oxygen species-dependent DNA double-strand breaks. *Blood* 104, 3746–3753.
- O'Hare, T., Shakespeare, W.C., Zhu, X., Eide, C.A., Rivera, V.M., Wang, F., Adrian, L.T., Zhou, T., Huang, W.-S., Xu, Q., Metcalf, C.A., Tyner, J.W., Loriaux, M.M., Corbin, A.S., Wardwell, S., Ning, Y., Keats, J.A., Wang, Y., Sundaramoorthi, R., Thomas, M., Zhou, D., Snodgrass, J., Commodore, L., Sawyer, T.K., Dalgarno, D.C., Deininger, M.W.N., Druker, B.J., and Clackson, T. (2009). AP24534, a pan-BCR-ABL inhibitor for chronic myeloid leukemia, potently inhibits the T315I mutant and overcomes mutation-based resistance. *Cancer Cell* 16, 401–412.
- O'Neill, L.A.J. (2008). The interleukin-1 receptor/Toll-like receptor superfamily: 10 years of progress. *Immunological reviews*. 226, 10–18.
- Ogawa, M. (1993). Differentiation and proliferation of hematopoietic stem cells. *Blood* 81, 2844–2853.
- Pearce, D.J., Taussig, D., Zibara, K., Smith, L.-L., Ridler, C.M., Preudhomme, C., Young, B.D., Rohatiner, A.Z., Lister, T.A., and Bonnet, D. (2006). AML engraftment in the NOD/SCID assay reflects the outcome of AML: implications for our understanding of the heterogeneity of AML. *Blood* 107, 1166–1173.
- Pellicano, F., Sinclair, A., and Holyoake, T.L. (2011). In search of CML stem cells' deadly weakness. *Current hematologic malignancy reports* 6, 82–87.
- Pendergast, A.M., Quilliam, L.A., Cripe, L.D., Bassing, C.H., Dai, Z., Li, N., Batzer, A., Rabun, K.M., Der, C.J., and Schlessinger, J. (1993). BCR-ABL-induced oncogenesis is mediated



- by direct interaction with the SH2 domain of the GRB-2 adaptor protein. *Cell* 75, 175–185.
- Piessevaux, J., Lavens, D., Montoye, T., Wauman, J., Catteeuw, D., Vandekerckhove, J., Belsham, D., Peelman, F., and Tavernier, J. (2006). Functional cross-modulation between SOCS proteins can stimulate cytokine signaling. *The Journal of biological chemistry* 281, 32953–32966.
- Prochazka, M., Gaskins, H.R., Shultz, L.D., and Leiter, E.H. (1992). The nonobese diabetic scid mouse: model for spontaneous thymomagenesis associated with immunodeficiency. *Proceedings of the National Academy of Sciences of the United States of America*. 89, 3290–3294.
- Quentmeier, H., Geffers, R., Jost, E., Macleod, R.A.F., Nagel, S., Röhrs, S., Romani, J., Scherr, M., Zaborski, M., and Drexler, H.G. (2008). SOCS2: inhibitor of JAK2V617F-mediated signal transduction. *Leukemia* 22, 2169–2175.
- Raitano, A.B., Halpern, J.R., Hambuch, T.M., and Sawyers, C.L. (1995). The Bcr-Abl leukemia oncogene activates Jun kinase and requires Jun for transformation. *Proceedings of the National Academy of Sciences of the United States of America* 92, 11746–11750.
- Randolph, T.R. (2005). Chronic myelocytic leukemia--Part I: History, clinical presentation, and molecular biology. *Clinical laboratory science* 18, 38–48.
- Reuther, J.Y., Reuther, G.W., Cortez, D., Pendergast, A.M., and Baldwin, A.S. (1998). A requirement for NF-kappaB activation in Bcr-Abl-mediated transformation. *Genes & development*. 12, 968–981.
- Reya, T., Morrison, S.J., Clarke, M.F., and Weissman, I.L. (2001). Stem cells, cancer, and cancer stem cells. *Nature* 414, 105–111.
- Ross, D.M., Branford, S., Seymour, J.F., Schwazer, A.P., Arthur, C., Yeung, D.T., Dang, P., Goyne, J.M., Slader, C., Filshie, R.J., Mills, A.K., Melo, J.V., White, D.L., Grigg, A.P., and Hughes, T.P. (2013). Safety and efficacy of imatinib cessation for CML patients with stable undetectable minimal residual disease: results from the TWISTER study. *Blood* 122, 515–522.
- Rousselot, P., Huguette, F., Rea, D., Legros, L., Cayuela, J.M., Maarek, O., Blanchet, O., Marit, G., Gluckman, E., Reiffers, J., Gardembas, M., and Mahon, F.-X. (2007). Imatinib mesylate discontinuation in patients with chronic myelogenous leukemia in complete molecular remission for more than 2 years. *Blood* 109, 58–60.
- Rowley, J.D. (1973). Letter: A new consistent chromosomal abnormality in chronic myelogenous leukaemia identified by quinacrine fluorescence and Giemsa staining. *Nature* 243, 290–293.

- Sakai, N., Ogiso, Y., Fujita, H., Watari, H., Koike, T., and Kuzumaki, N. (1994). Induction of apoptosis by a dominant negative H-RAS mutant (116Y) in K562 cells. *Experimental cell research* 215, 131–136.
- Samanta, A.K., Chakraborty, S.N., Wang, Y., Kantarjian, H., Sun, X., Hood, J., Perrotti, D., and Arlinghaus, R.B. (2009). Jak2 inhibition deactivates Lyn kinase through the SET-PP2A-SHP1 pathway, causing apoptosis in drug-resistant cells from chronic myelogenous leukemia patients. *Oncogene* 28, 1669–1681.
- Sawyers, C.L. (1999). Chronic myeloid leukemia. *The New England journal of medicine* 340, 1330–1340.
- Sawyers, C.L., McLaughlin, J., and Witte, O.N. (1995). Genetic requirement for Ras in the transformation of fibroblasts and hematopoietic cells by the Bcr-Abl oncogene. *The Journal of experimental medicine*. 181, 307–313.
- Schindler, T., Bornmann, W., Pellicena, P., Miller, W.T., Clarkson, B., and Kuriyan, J. (2000). Structural mechanism for STI-571 inhibition of abelson tyrosine kinase. *Science* 289, 1938–1942.
- Schultheis, B., Carapeti-Marootian, M., Hochhaus, A., Weisser, A., Goldman, J.M., and Melo, J.V. (2002). Overexpression of SOCS-2 in advanced stages of chronic myeloid leukemia: possible inadequacy of a negative feedback mechanism. *Blood* 99, 1766–1775.
- Sepp-Lorenzino, L., Ma, Z., Rands, E., Kohl, N.E., Gibbs, J.B., Oliff, A., and Rosen, N. (1995). A peptidomimetic inhibitor of farnesyl:protein transferase blocks the anchorage-dependent and -independent growth of human tumor cell lines. *Cancer research* 55, 5302–5309.
- Sexl, V., Piekorz, R., Moriggl, R., Rohrer, J., Brown, M.P., Bunting, K.D., Rothhammer, K., Roussel, M.F., and Ihle, J.N. (2000). Stat5a/b contribute to interleukin 7-induced B-cell precursor expansion, but abl- and bcr/abl-induced transformation are independent of stat5. *Blood* 96, 2277–2283.
- Shah, N.P., Tran, C., Lee, F.Y., Chen, P., Norris, D., and Sawyers, C.L. (2004). Overriding imatinib resistance with a novel ABL kinase inhibitor. *Science* 305, 399–401.
- Shtivelman, E., Lifshitz, B., Gale, R.P., and Canaani, E. (1985). Fused transcript of abl and bcr genes in chronic myelogenous leukaemia. *Nature* 315, 550–554.
- Shuai, K., Halpern, J., Hoeve, ten, J., Rao, X., and Sawyers, C.L. (1996). Constitutive activation of STAT5 by the BCR-ABL oncogene in chronic myelogenous leukemia. *Oncogene* 13, 247–254.
- Shultz, L.D., Lyons, B.L., Burzenski, L.M., Gott, B., Chen, X., Chaleff, S., Kotb, M., Gillies, S.D., King, M., Mangada, J., Greiner, D.L., and Handgretinger, R. (2005). Human lymphoid and myeloid cell development in NOD/LtSz-scid IL2R gamma null mice

- engrafted with mobilized human hemopoietic stem cells. *Journal of immunology* 174, 6477–6489.
- Sill, H., Goldman, J.M., and Cross, N.C. (1995). Homozygous deletions of the p16 tumor-suppressor gene are associated with lymphoid transformation of chronic myeloid leukemia. *Blood* 85, 2013–2016.
- Socolovsky, M., Fallon, A.E., Wang, S., Brugnara, C., and Lodish, H.F. (1999). Fetal anemia and apoptosis of red cell progenitors in Stat5a<sup>-/-</sup>5b<sup>-/-</sup> mice: a direct role for Stat5 in Bcl-X(L) induction. *Cell* 98, 181–191.
- Socolovsky, M., Nam, H., Fleming, M.D., Haase, V.H., Brugnara, C., and Lodish, H.F. (2001). Ineffective erythropoiesis in Stat5a<sup>-/-</sup>5b<sup>-/-</sup> mice due to decreased survival of early erythroblasts. *Blood* 98, 3261–3273.
- Talpaz, M., Shah, N.P., Kantarjian, H., Donato, N., Nicoll, J., Paquette, R., Cortes, J., O'Brien, S., Nicaise, C., Bleickardt, E., Blackwood-Chirchir, M.A., Iyer, V., Chen, T.-T., Huang, F., Decillis, A.P., and Sawyers, C.L. (2006). Dasatinib in imatinib-resistant Philadelphia chromosome-positive leukemias. *The New England journal of medicine* 354, 2531–2541.
- Tanizaki, R., Nomura, Y., Miyata, Y., Minami, Y., Abe, A., Hanamura, A., Sawa, M., Murata, M., Kiyoi, H., Matsushita, T., and Naoe, T. (2010). Irrespective of CD34 expression, lineage-committed cell fraction reconstitutes and re-establishes transformed Philadelphia chromosome-positive leukemia in NOD/SCID/IL-2R $\gamma$ mmac<sup>-/-</sup> mice. *Cancer science*. 101, 631–638.
- Tannahill, G.M., Elliott, J., Barry, A.C., Hibbert, L., Cacalano, N.A., and Johnston, J.A. (2005). SOCS2 can enhance interleukin-2 (IL-2) and IL-3 signaling by accelerating SOCS3 degradation. *Molecular and cellular biology* 25, 9115–9126.
- Tauchi, T., Boswell, H.S., Leibowitz, D., and Broxmeyer, H.E. (1994). Coupling between p210bcr-abl and Shc and Grb2 adaptor proteins in hematopoietic cells permits growth factor receptor-independent link to ras activation pathway. *The Journal of experimental medicine* 179, 167–175.
- Taussig, D.C., Miraki-Moud, F., Anjos-Afonso, F., Pearce, D.J., Allen, K., Ridler, C., Lillington, D., Oakervee, H., Cavenagh, J., Agrawal, S.G., Lister, T.A., Gribben, J.G., and Bonnet, D. (2008). Anti-CD38 antibody-mediated clearance of human repopulating cells masks the heterogeneity of leukemia-initiating cells. *Blood* 112, 568–575.
- Taussig, D.C., Vargaftig, J., Miraki-Moud, F., Griessinger, E., Sharrock, K., Luke, T., Lillington, D., Oakervee, H., Cavenagh, J., Agrawal, S.G., Lister, T.A., Gribben, J.G., and Bonnet, D. (2010). Leukemia-initiating cells from some acute myeloid leukemia patients with mutated nucleophosmin reside in the CD34<sup>-</sup> fraction. *Blood* 115, 1976–1984.

- Teglund, S., McKay, C., Schuetz, E., van Deursen, J.M., Stravopodis, D., Wang, D., Brown, M., Bodner, S., Grosveld, G., and Ihle, J.N. (1998). Stat5a and Stat5b proteins have essential and nonessential, or redundant, roles in cytokine responses. *Cell* 93, 841–850.
- Thoren, L.A., Liuba, K., Bryder, D., Nygren, J.M., Jensen, C.T., Qian, H., Antonchuk, J., and Jacobsen, S.E.W. (2008). Kit regulates maintenance of quiescent hematopoietic stem cells. *Journal of immunology* 180, 2045–2053.
- van Rhenen, A., Feller, N., Kelder, A., Westra, A.H., Rombouts, E., Zweegman, S., van der Pol, M.A., Waisfisz, Q., Ossenkoppele, G.J., and Schuurhuis, G.J. (2005). High stem cell frequency in acute myeloid leukemia at diagnosis predicts high minimal residual disease and poor survival. *Clinical cancer research* 11, 6520–6527.
- Vardiman, J.W., Thiele, J., Arber, D.A., Brunning, R.D., Borowitz, M.J., Porwit, A., Harris, N.L., Le Beau, M.M., Hellström-Lindberg, E., Tefferi, A., and Bloomfield, C.D. (2009). The 2008 revision of the World Health Organization (WHO) classification of myeloid neoplasms and acute leukemia: rationale and important changes. *Blood* 114, 937–951.
- Venter, J.C., Adams, M.D., Myers, E.W., Li, P.W., Mural, R.J., Sutton, G.G., Smith, H.O., Yandell, M., Evans, C.A., Holt, R.A., Gocayne, J.D., Amanatides, P., Ballew, R.M., Huson, D.H., Wortman, J.R., Zhang, Q., Kodira, C.D., Zheng, X.H., Chen, L., Skupski, M., Subramanian, G., Thomas, P.D., Zhang, J., Gabor Miklos, G.L., Nelson, C., Broder, S., Clark, A.G., Nadeau, J., McKusick, V.A., Zinder, N., Levine, A.J., Roberts, R.J., Simon, M., Slayman, C., Hunkapiller, M., Bolanos, R., Delcher, A., Dew, I., Fasulo, D., Flanigan, M., Florea, L., Halpern, A., Hannenhalli, S., Kravitz, S., Levy, S., Mobarry, C., Reinert, K., Remington, K., Abu-Threideh, J., Beasley, E., Biddick, K., Bonazzi, V., Brandon, R., Cargill, M., Chandramouliswaran, I., Charlab, R., Chaturvedi, K., Deng, Z., Di Francesco, V., Dunn, P., Eilbeck, K., Evangelista, C., Gabrielian, A.E., Gan, W., Ge, W., Gong, F., Gu, Z., Guan, P., Heiman, T.J., Higgins, M.E., Ji, R.R., Ke, Z., Ketchum, K.A., Lai, Z., Lei, Y., Li, Z., Li, J., Liang, Y., Lin, X., Lu, F., Merkulov, G.V., Milshina, N., Moore, H.M., Naik, A.K., Narayan, V.A., Neelam, B., Nusskern, D., Rusch, D.B., Salzberg, S., Shao, W., Shue, B., Sun, J., Wang, Z., Wang, A., Wang, X., Wang, J., Wei, M., Wides, R., Xiao, C., Yan, C., Yao, A., Ye, J., Zhan, M., Zhang, W., Zhang, H., Zhao, Q., Zheng, L., Zhong, F., Zhong, W., Zhu, S., Zhao, S., Gilbert, D., Baumhueter, S., Spier, G., Carter, C., Cravchik, A., Woodage, T., Ali, F., An, H., Awe, A., Baldwin, D., Baden, H., Barnstead, M., Barrow, I., Beeson, K., Busam, D., Carver, A., Center, A., Cheng, M.L., Curry, L., Danaher, S., Davenport, L., Desilets, R., Dietz, S., Dodson, K., Doup, L., Ferreira, S., Garg, N., Gluecksmann, A., Hart, B., Haynes, J., Haynes, C., Heiner, C., Hladun, S., Hosten, D., Houck, J., Howland, T., Ibegwam, C., Johnson, J., Kalush, F., Kline, L., Koduru, S., Love, A., Mann, F., May, D., McCawley, S., McIntosh, T., McMullen, I., Moy, M., Moy, L., Murphy, B., Nelson, K., Pfannkoch, C., Pratts, E., Puri, V., Qureshi, H., Reardon, M., Rodriguez, R., Rogers, Y.H., Romblad, D., Ruhfel, B., Scott, R., Sitter, C., Smallwood, M., Stewart, E., Strong, R., Suh, E., Thomas, R., Tint, N.N., Tse, S., Vech, C., Wang, G., Wetter, J., Williams, S., Williams, M., Windsor, S., Winn-Deen, E., Wolfe, K., Zaveri, J., Zaveri, K., Abril, J.F., Guigó, R., Campbell, M.J., Sjolander, K.V., Karlak, B., Kejariwal, A., Mi, H., Lazareva, B., Hatton, T., Narechania,

- A., Diemer, K., Muruganujan, A., Guo, N., Sato, S., Bafna, V., Istrail, S., Lippert, R., Schwartz, R., Walenz, B., Yoosheph, S., Allen, D., Basu, A., Baxendale, J., Blick, L., Caminha, M., Carnes-Stine, J., Caulk, P., Chiang, Y.H., Coyne, M., Dahlke, C., Mays, A., Dombroski, M., Donnelly, M., Ely, D., Esparham, S., Fosler, C., Gire, H., Glanowski, S., Glasser, K., Glodek, A., Gorokhov, M., Graham, K., Gropman, B., Harris, M., Heil, J., Henderson, S., Hoover, J., Jennings, D., Jordan, C., Jordan, J., Kasha, J., Kagan, L., Kraft, C., Levitsky, A., Lewis, M., Liu, X., Lopez, J., Ma, D., Majoros, W., McDaniel, J., Murphy, S., Newman, M., Nguyen, T., Nguyen, N., Nodell, M., Pan, S., Peck, J., Peterson, M., Rowe, W., Sanders, R., Scott, J., Simpson, M., Smith, T., Sprague, A., Stockwell, T., Turner, R., Venter, E., Wang, M., Wen, M., Wu, D., Wu, M., Xia, A., Zandieh, A., and Zhu, X. (2001). The sequence of the human genome. *Science* 291, 1304–1351.
- Wagner, K., Zhang, P., Rosenbauer, F., Drescher, B., Kobayashi, S., Radoska, H.S., Kutok, J.L., Gilliland, D.G., Krauter, J., and Tenen, D.G. (2006). Absence of the transcription factor CCAAT enhancer binding protein alpha results in loss of myeloid identity in bcr/abl-induced malignancy. *Proceedings of the National Academy of Sciences of the United States of America* 103, 6338–6343.
- Wang, J.C., Lapidot, T., Cashman, J.D., Doedens, M., Addy, L., Sutherland, D.R., Nayar, R., Laraya, P., Minden, M., Keating, A., Eaves, A.C., Eaves, C.J., and Dick, J.E. (1998). High level engraftment of NOD/SCID mice by primitive normal and leukemic hematopoietic cells from patients with chronic myeloid leukemia in chronic phase. *Blood* 91, 2406–2414.
- Warsch, W., Walz, C., and Sexl, V. (2013). JAK of all trades: JAK2-STAT5 as novel therapeutic targets in BCR-ABL1+ chronic myeloid leukemia. *Blood* 122, 2167-2175.
- Wei, W., Huang, H., Zhao, S., Liu, W., Liu, C.-X., Chen, L., Li, J.-M., Wu, Y.-L., and Yan, H. (2013). Alantolactone induces apoptosis in chronic myelogenous leukemia sensitive or resistant to imatinib through NF-κB inhibition and Bcr/Abl protein deletion. *Apoptosis* 18, 1060–1070.
- Wilson-Rawls, J., Liu, J., Laneville, P., and Arlinghaus, R.B. (1997). P210 Bcr-Abl interacts with the interleukin-3 beta c subunit and constitutively activates Jak2. *Leukemia* 11 Suppl 3, 428–431.
- Yamamoto, K., Nakamura, Y., Saito, K., and Furusawa, S. (2000). Expression of the NUP98/HOXA9 fusion transcript in the blast crisis of Philadelphia chromosome-positive chronic myelogenous leukaemia with t(7;11)(p15;p15). *British journal of haematology* 109, 423–426.
- Wunderlich, M., Chou, F.-S., Link, K.A., Mizukawa, B., Perry, R.L., Carroll, M., and Mulloy, J.C. (2010). AML xenograft efficiency is significantly improved in NOD/SCID-IL2RG mice constitutively expressing human SCF, GM-CSF and IL-3. *Leukemia* 24, 1785–1788.

- Ye, D., Wolff, N., Li, L., Zhang, S., and Ilaria, R.L. (2006). STAT5 signaling is required for the efficient induction and maintenance of CML in mice. *Blood* 107, 4917–4925.
- Zhang, B., Ho, Y.W., Huang, Q., Maeda, T., Lin, A., Lee, S.-U., Hair, A., Holyoake, T.L., Huettner, C., and Bhatia, R. (2012). Altered microenvironmental regulation of leukemic and normal stem cells in chronic myelogenous leukemia. *Cancer Cell* 21, 577–592.
- Zhang, B., Li, M., McDonald, T., Holyoake, T.L., Moon, R.T., Campana, D., Shultz, L., and Bhatia, R. (2013). Microenvironmental protection of CML stem and progenitor cells from tyrosine kinase inhibitors through N-cadherin and Wnt- $\beta$ -catenin signaling. *Blood* 121, 1824–1838.
- Zhang, X., and Ren, R. (1998). Bcr-Abl efficiently induces a myeloproliferative disease and production of excess interleukin-3 and granulocyte-macrophage colony-stimulating factor in mice: a novel model for chronic myelogenous leukemia. *Blood* 92, 3829–3840.
- Zhao, C., Xiu, Y., Ashton, J., Xing, L., Morita, Y., Jordan, C.T., and Boyce, B.F. (2012a). Noncanonical NF- $\kappa$ B signaling regulates hematopoietic stem cell self-renewal and microenvironment interactions. *Stem Cells* 30, 709–718.
- Zhao, L.-J., Wang, Y.-Y., Li, G., Ma, L.-Y., Xiong, S.-M., Weng, X.-Q., Zhang, W.-N., Wu, B., Chen, Z., and Chen, S.-J. (2012b). Functional features of RUNX1 mutants in acute transformation of chronic myeloid leukemia and their contribution to inducing murine full-blown leukemia. *Blood* 119, 2873–2882.
- Zhu, X., Wang, L., Zhang, B., Li, J., Dou, X., and Zhao, R.C. (2011). TGF- $\beta$ 1-induced PI3K/Akt/NF- $\kappa$ B/MMP9 signalling pathway is activated in Philadelphia chromosome-positive chronic myeloid leukaemia hemangioblasts. *Journal of biochemistry*, 149, 405–414.