

Molecular subtyping of serous ovarian tumors reveals multiple connections to intrinsic breast cancer subtypes.

Jönsson, Jenny-Maria; Johansson, Ida; Dominguez, Mev; Kimbung, Siker; Jönsson, Mats; Bonde, Jesper Hansen; Kannisto, Päivi; Måsbäck, Anna; Malander, Susanne; Nilbert, Mef; Hedenfalk, İngrid

Published in: PLoS ONE

2014

Link to publication

Citation for published version (APA):

10.1371/journal.pone.0107643

Jönsson, J.-M., Johansson, I., Dominguez, M., Kimbung, S., Jönsson, M., Bonde, J. H., Kannisto, P., Måsbäck, A., Malander, S., Nilbert, M., & Hedenfalk, I. (2014). Molecular subtyping of serous ovarian tumors reveals multiple connections to intrinsic breast cancer subtypes. *PLoS ONE*, *9*(9), Article e107643. https://doi.org/10.1371/journal.pone.0107643

Total number of authors:

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- · Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

 • You may not further distribute the material or use it for any profit-making activity or commercial gain

 • You may not further distribute the LIPL identifying the publication in the public portal

Read more about Creative commons licenses: https://creativecommons.org/licenses/

Take down policyIf you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

Download date: 29. Oct. 2025



Molecular Subtyping of Serous Ovarian Tumors Reveals Multiple Connections to Intrinsic Breast Cancer Subtypes CrossMark



Jenny-Maria Jönsson^{1*}, Ida Johansson^{1,2}, Mev Dominguez-Valentin^{1,5}, Siker Kimbung^{1,2}, Mats Jönsson¹, Jesper Hansen Bonde³, Päivi Kannisto⁴, Anna Måsbäck¹, Susanne Malander¹, Mef Nilbert^{1,5}, Ingrid Hedenfalk^{1,2}

1 Division of Oncology-Pathology, Department of Clinical Sciences, Skåne University Hospital, Lund University, Lund, Sweden, 2 CREATE Health Strategic Center for Translational Cancer Research, Lund University, Lund, Sweden, 3 Department of Pathology, Clinical Research Centre, Hvidovre Hospital, Copenhagen University, Hvidovre, Denmark, 4 Division of Obstetrics and Gynecology, Department of Clinical Sciences, Skåne University Hospital, Lund University, Lund, Sweden, 5 Clinical Research Centre, Hvdiovre Hospital, Copenhagen University, Hvidovre, Denmark

Abstract

Objective: Transcriptional profiling of epithelial ovarian cancer has revealed molecular subtypes correlating to biological and clinical features. We aimed to determine gene expression differences between malignant, benign and borderline serous ovarian tumors, and investigate similarities with the well-established intrinsic molecular subtypes of breast cancer.

Methods: Global gene expression profiling using Illumina's HT12 Bead Arrays was applied to 59 fresh-frozen serous ovarian malignant, benign and borderline tumors. Nearest centroid classification was performed applying previously published gene profiles for the ovarian and breast cancer subtypes. Correlations to gene expression modules representing key biological breast cancer features were also sought. Validation was performed using an independent, publicly available dataset.

Results: 5,944 genes were significantly differentially expressed between benign and malignant serous ovarian tumors, with cell cycle processes enriched in the malignant subgroup. Borderline tumors were split between the two clusters. Significant correlations between the malignant serous tumors and the highly aggressive ovarian cancer signatures, and the basal-like breast cancer subtype were found. The benign and borderline serous tumors together were significantly correlated to the normal-like breast cancer subtype and the ovarian cancer signature derived from borderline tumors. The borderline tumors in the study dataset, in addition, also correlated significantly to the luminal A breast cancer subtype. These findings remained when analyzed in an independent dataset, supporting links between the molecular subtypes of ovarian cancer and breast cancer beyond those recently acknowledged.

Conclusions: These data link the transcriptional profiles of serous ovarian cancer to the intrinsic molecular subtypes of breast cancer, in line with the shared clinical and molecular features between high-grade serous ovarian cancer and basallike breast cancer, and suggest that biomarkers and targeted therapies may overlap between these tumor subsets. The link between benign and borderline ovarian cancer and luminal breast cancer may indicate endocrine responsiveness in a subset of ovarian cancers.

Citation: Jönsson J-M, Johansson I, Dominguez-Valentin M, Kimbung S, Jönsson M, et al. (2014) Molecular Subtyping of Serous Ovarian Tumors Reveals Multiple Connections to Intrinsic Breast Cancer Subtypes. PLoS ONE 9(9): e107643. doi:10.1371/journal.pone.0107643

Editor: Goli Samimi, Garvan Institute of Medical Research, Australia

Received June 5, 2014; Accepted August 12, 2014; Published September 16, 2014

Copyright: © 2014 Jönsson et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability: The authors confirm that all data underlying the findings are fully available without restriction. All gene expression data files are available in the National Center for Biotechnology Information Gene Expression Omnibus [GEO accession number: GSE57477].

Funding: Financial support was granted from a governmental funding of clinical research within the National Health Services (JMJ), the Swedish Cancer Society (MN, IH), the Swedish Research Council (MN, IH) and an unrestricted educational grant from the Swedish Society of Gynecologic Oncology sponsored by Roche (JMJ). Funding has neither influenced study design, data collection nor analysis or manuscript writing.

Competing Interests: The authors have declared that no competing interests exist.

* Email: jenny-maria.jonsson@med.lu.se

Introduction

Epithelial ovarian tumors constitute a heterogeneous group of neoplasms that differ in epidemiology, genetic risk factors, precursor lesions and clinical behavior. The different histopathologic subtypes, i.e. serous, mucinous, endometrioid, clear cell and transitional carcinomas and carcinosarcomas, likely have different origins and appear to evolve along distinct pathways [1–3]. Alongside with the standard taxane and platinum based agents

used for ovarian cancer [4-6], multiple targeted agents are being evaluated, with e.g. bevacizumab recently being included in the therapeutic arsenal [7]. Personalized therapy is called for in ovarian cancer particularly since the histopathologic subtypes, as well as tumors with different malignant potential and tumor grade, can be viewed as separate diseases with differences related to both prognosis and treatment response [8-12]. Refined molecular subtyping and recognition of key genetic mechanisms constitutes

an encouraging basis for further development of subtype-specific targeted therapies.

Previous efforts to characterize ovarian cancers at the molecular level have identified distinct profiles related to the histologic subtypes and have suggested predictive gene signatures [13–17]. Tothill et al. suggested six different subtypes, referred to as C1–C6, based on serous and endometrioid ovarian, primary peritoneal and fallopian tube tumors. The C1–C2 and C4–C5 subtypes, in general, are thought to characterize high-grade serous tumors. The C1 signature is characterized by a high degree of desmoplasia, C5 by mesenchymal genes and overexpression of proliferation genes and the C2 and C4 signatures by high numbers of intra-tumoral and stroma associated CD3+ cells. The signatures are outcome predictive, with the C1 signature corresponding to a considerably worse outcome than the other signatures. The C3 signature represents low-grade serous and borderline tumors and the C6 signature low-grade, early-stage endometrioid tumors; in general they show good response to treatment and long-time survival [18]. Likewise, molecular subtyping in breast cancer is well established and recent reports have recognized similarities between high-grade serous ovarian cancer and basal-like breast cancer [19].

We performed global gene expression profiling of serous ovarian tumors, including serous cystadenomas, serous borderline tumors and serous adenocarcinomas, and applied previously described gene signatures including the well-known intrinsic breast cancer subtypes [18,20–22] to outline further possible similarities between these tumor types. Since mutations in the MAPK/ERK pathways are common in both borderline and low-grade ovarian cancer and luminal breast cancers the presence of *KRAS* and *BRAF* mutations was investigated among the ovarian tumors [23,24]. Shared common features between ovarian and breast cancer may be useful for future development of predictive biomarkers and tailored treatments in both tumor types, and in this study we present interesting connections between the molecular subtypes of ovarian and breast cancer.

Materials and Methods

Tumor samples

In total, 37 serous ovarian adenocarcinomas, 17 serous cystadenomas/adenofibromas and 5 serous borderline tumors were obtained from the Skåne University Hospital ovarian tumor biobank (table 1). A total of 13 biological replicates (6 omental metastases, 1 pelvic metastasis and 2 metastases to the contralateral ovary as well as 3 benign and 1 borderline ovarian tumors) were included to account for intra-tumor heterogeneity. All tumor samples were collected at primary surgery (2003-2011) and the patients had not received chemotherapy prior to surgery. Histologic subtype and grade were determined according to Silverberg and WHO [25,26] and all tumors were staged according to the International Federation of Gynecology and Obstetrics (FIGO) criteria. Hematoxylin & Eosin stained slides were used to assess tumor grade. This was performed by a senior pathologist (AM). Ethical approval for the study was granted from the Lund University ethics committee, Sweden, waiving the requirement for informed consent for the study.

RNA extraction and gene expression analysis

Total RNA was extracted using the Allprep kit (Qiagen, Heidelberg, Germany) according to the manufacturer's instructions. RNA concentration was determined using a NanoDrop Spectrophotometer (NanoDrop Technologies, Wilmington, DE) and samples with ≥200 ng RNA with 260/280 ratios ≥1.8 were

used for further analysis. RNA quality was assessed using a Bioanalyzer (Agilent technologies, Santa Clara, CA), and RNA integrity numbers (RIN) >6 were regarded as sufficient.

Gene expression profiling analyses were performed at the SCIBLU Genomics Centre, Lund University, Sweden. The cDNA synthesis, labeling, and subsequent hybridization to the HumanHT-12 v4 Expression BeadChips (Illumina Inc., San Diego, CA) was performed according to the manufacturer's instructions. The Illumina HumanHT-12 v4 Expression BeadChips allow genome-wide expression profiling of more than 47,000 gene transcripts and splice variants. The 59 samples and 13 biological replicates were randomized on the chips. The BeadChips were then scanned on an i-Scan (Illumina Inc.), during which fluorescence intensities were read and images extracted. The gene expression data are available in the National Center for Biotechnology Information Gene Expression Omnibus [GEO accession number: GSE57477] [27].

Data analysis

Gene expression data were uploaded to the GenomeStudio software (Illumina Inc.), quantile normalized, background corrected and log2 transformed. Probes with a mean intensity <2.5 and variance <0.1 were excluded, leaving a total of 16,024 probes corresponding to 12,313 unique genes. Thereafter the data were uploaded to the MeV v4 software, an application used for identification of genes and expression patterns in microarray data [28], mean centered and a variance filter was applied to select the 20% of the probes with the greatest variation of expression across the dataset. Unsupervised hierarchical clustering was performed using complete linkage and Pearson distant metric. Two-class unpaired significance analysis of microarrays (SAM) was performed based on all 16,024 probes to identify differentially expressed probes between the different tumor subgroups (benign, borderline, malignant) at a false discovery rate (FDR) < 0.01 [29]. Hierarchical clustering, supervised by the SAM analysis results and thereby identifying significantly differentially expressed genes between the tumor subgroups, was performed using the same methods as for unsupervised clustering. Gene ontology analyses based on the significant genes were performed in the gene ontology enrichment analysis and visualization tools GOrilla and ToppGene for identification of possible gene enrichment with biological or functional differences separating the subgroups [30,31].

Molecular subtyping and external data sets

Gene signatures outlining six molecular subtypes of high-grade and advanced stage serous ovarian tumors as well as endometrioid, low-grade serous and borderline ovarian tumors (referred to as the "Tothill dataset") [18] were applied to the serous ovarian tumors in our cohort. Data were normalized and log2 transformed using the Gene Chip Operating Software Version 1.4 with Affymetrix default analysis settings. Probes with intensity values <4 and variance < 0.15 were excluded. The six ovarian cancer signatures contained in total 4,732 probes of which 4,099 probes, corresponding to 2,725 unique genes, with good quality were left after filtering away probes with bad quality. 1,295/2,725 (47.5%) of these genes were identified in our dataset and used for further analyses. To validate the classifier, the 1,295 genes present in our dataset were re-applied to the 285 ovarian tumors in the original cohort, thereby re-assigning subtypes to each tumor. A gene signature for intrinsic subtyping of breast cancer was also applied to our serous ovarian tumors [22]. Each tumor in our cohort was classified into the molecular subtypes of ovarian cancer (C1-C6) as well as the intrinsic subtypes of breast cancer (luminal A, luminal

Table 1. Clinicopathologic features of malignant and borderline ovarian tumors in the study cohort.

BC subtype	Normal	Normal	HER2	Normal	Basal	Luminal A	Basal	Luminal A	Basal	HER2	Basal	Normal	Basal	HER2	Basal	Basal	Normal	Basal	Basal	Basal	HER2	Basal	Basal	HER2	Luminal B	Basal	Basal	Normal						
C- signature	Cl	Cl	S	υ	2	CS	C4	C4	CS	Cl	C4	CS	٥	C1	ט	7	C4	C3	CI	C2	٥	Cl	٥	0	D	C2	C4	2	2	C4	S	C4	22	υ
Survival (years)	3	-	9	2	4	2	5	Alive	1,5	▽	2	1,5	-	2	-	2	Alive	▽	1	4	2	-	Alive	Alive	2	2	Alive	Alive	Alive	Alive	Alive	Alive	æ	Alive
Tissue type	Ovary	Omentum	Ovary	Ovary	N/A	N/A	N/A	Ħ	Ovary	N/A	Cystic fluid	Omentum	Omentum	N/A	E	N/A	Omentum	Ovary	Omentum	N/A	Ovary	Ovary	Omentum	N/A	Ovary	Ovary	N/A	Ovary	N/A	N/A	Ovary	Ovary	Ovary	Omentum
Age at diagnosis (years)	69	82	69	09	71	50	71	70	61	91	82	79	55	80	67	53	61	87	59	62	65	69	70	72	65	63	40	42	67	69	64	89	81	79
Grade	3	3	2	2	3	3	3	3	3	3	3	2	2	1	3	2	3	-	3	3	2	2	3	3	3	2	-	-	3	2	3	3	3	2
Stage	JII)III	C) C))	HIA	И))III)IIC	IIC	Σ	IIIC	IIC	IV1	IV3	IV3	IV3	IIIC)III	IIIB	⊞	IIC	IIIC	IIIB	IIC	IIC1	IIB	JIIC	-	JIIC	JIIC)
Feature	×	M	Σ	M	Σ	M	Σ	Σ	Σ	M	Σ	M	Σ	M	Σ	M	M	M	M	×	Σ	M	Σ	M	M	M	Σ	M	Σ	Σ	×	Σ	Σ	Σ
므	38	70	84	8	106	118	125	137	153	159	186	190	192	207	219	225	526	232	251	273	275	279	284	293	297	305	306	307	311	314	330	344	393	397

Table 1. Cont.

Р	Feature	Stage	Grade	Age at diagnosis (years)	Tissue type	Survival (years)	C- signature	BC subtype
420	×	IC1	-		Ovary	Alive	C4	Luminal B
438	Σ	⊇	8	53	Ovary	Alive	22	Basal
16	Во	⊻	N/A	40	Ovary	Alive	G	Luminal A
48	Во	⊻	N/A	51	Ovary	Alive	ß	Luminal A
98	Во	⊻	N/A	45	N/A	Alive	G	Luminal A
377	Во	17	N/A	29	Ovary	Alive	ß	Normal
385	Во	IC1	N/A	09	Ovary	Alive	C3	Luminal A
Feature	: M = Malignant, Bo	= Borderline;	Tissue type: tissu	Feature: $M = Malignant$, $Bo = Borderline$; Tissue type: tissue used for RNA extraction, FT = Fallopian Tube, $N/A = Pature$	· Unknown; <i>Survival</i> : Diseas	on, FT = Fallopian Tube, N/A = Unknown; Survival: Disease specific survival, Alive = alive at start of study; C-signature: corresponding ovarian molecular	t of study; C-signature: corre	sponding ovarian molecular

18]; BC subtype: corresponding intrinsic breast cancer subtype

B, basal-like, normal-like, and HER2 enriched) using nearest centroid classification. The methodology for nearest centroid classification is outlined by Johansson et al. [32]. Validation was performed by classifying the 285 ovarian tumors in the Tothill dataset into the intrinsic breast cancer subtypes. Furthermore, seven gene expression modules representing key biological processes in breast cancer (AURKA/proliferation, CASP3/ apoptosis, ERBb2/HER2 signaling, ESR1/ER signaling, STAT1/immune response, PLAU/tumor invasion and metastasis, VEGF/angiogenesis; referred to as the "Desmedt modules") were applied to the serous ovarian tumors in our dataset as well as the Tothill dataset [33], and their relationship to the previously described intrinsic breast cancer subtypes was investigated. These modules, derived from 917 breast cancers in publicly available datasets and characterized by computed module scores, comprise in total 889 genes.

Mutation analysis

KRAS mutation analysis was performed using the Roche cobas K-RAS Mutation Kit (product number 05852170190) (Roche, Pleasanton, CA), a CE-IVD real-time melting curve KRAS mutations assay, detecting mutations in codons 12, 13 and 61 of the KRAS oncogene. BRAF mutation analysis was performed using the Roche cobas BRAF V600 mutational analysis (product number 05985595190), which evaluates the BRAF V600 site in exon 15 and detects wildtype or mutated V600. The analyses were performed according to the manufacturer's instructions, and the assays were run on the z480 Lightcycler (Roche). The mutation analyses were performed at the Department of Pathology, Clinical Research Centre, Hvidovre Hospital, Denmark.

Statistical methods

Mann-Whitney U-test and Pearson correlation were used for comparison between expression profiles of the different tumor subsets using the MeV 4.6.02 software. Correlations between different subtype classifications were assessed using Fisher's exact test and between module scores using Mann-Whitney U-test and Kruskal Wallis test in SPSS (IBM SPSS Statistics 19). P-values < 0.05 were considered statistically significant.

Results

Comparison of malignant, benign and borderline serous ovarian cancers

Unsupervised hierarchical clustering of benign and malignant tumors based on the 20% of the probes that showed the most variability revealed two distinct clusters, one containing only malignant tumors and one containing all benign and four malignant tumors (Figure S1). All but one of the biological replicates clustered together pair-wise. The clusters remained stable after removal of the biological replicates, suggesting stable transcriptional differences between the clusters (Figure S2). Of note, of the four malignant tumors in the benign cluster one was grade 1, two were grade 2 and one was grade 3, but no significant differences regarding stage or mean age at diagnosis were seen between the malignant tumors in the two clusters.

Next, a SAM analysis was performed to explore transcriptional differences between benign and malignant tumors, revealing 5,944 significantly differentially expressed genes (FDR <0.01), of which 2,984 were upregulated and 2,960 were downregulated among the malignant tumors (figure 1, Table S1).

Cell cycle kinases (e.g. CDC2, CDC5, CDC7 and CDC20) as well as AURKA and S100A9, which can all broadly be linked to cell cycle regulation and mitosis, were upregulated in the

malignant tumors. Consistent with this, gene ontology analyses revealed a significant upregulation of cell cycle associated biological processes (ToppGene, FDR<0.05; table 2).

To investigate whether borderline tumors are more closely related to benign or malignant tumors, an unsupervised clustering based on the 20% most varying probes across the dataset, and a hierarchical clustering supervised by the 5,944 significantly differentially expressed genes between benign and malignant tumors, were performed on the whole dataset. The analyses resulted in two distinct clusters, one malignant and one benign. The borderline tumors were split between the two main clusters, implying heterogeneity within this group (Figure S3).

We next applied the gene signatures described by Tothill *et al.* [18] to study the representation of molecular ovarian subtypes ("C-signatures") in our dataset. Using nearest centroid classification, a specific ovarian cancer C-signature was assigned to each tumor in our cohort. 52/59 (88%) of the tumors had a correlation coefficient ≥ 0.2 . The centroid classifications revealed considerable heterogeneity across the tumors (p<0.001; table 3). These differences prompted us to investigate each signature individually, and significant correlations between the malignant tumors in our cohort and the C1, C2 and C4 signatures (p=0.020), and between the benign and borderline tumors in our cohort and the C3 signature (p<0.001) were revealed (Table S2). 251/285 tumors in the original Tothill cohort had an assigned C-signature, and 239/251 (95.2%) were correctly re-assigned to their respective C-signatures, thereby validating the classification method (Table S7).

Exploring similarities between ovarian and breast cancer

To investigate potential similarities between ovarian cancer and the widely acknowledged intrinsic subtypes of breast cancer, beyond the similarities between high-grade serous ovarian cancer and basal-like breast cancer that have been reported, we applied the signatures representing the intrinsic breast cancer subtypes to our cohort [19,22]. 40/59 (68%) of the tumors had a correlation coefficient ≥0.2. Classification of the intrinsic breast cancer subtypes was applied to the ovarian tumors in our cohort, revealing considerable heterogeneity (p<0.001; table 4). Significant correlations between the malignant ovarian tumors and the basal-like breast cancer subtype (p<0.001), and between the nonmalignant (benign and borderline) ovarian tumors and the normal-like breast cancer subtype were found (p<0.001) (Table S3). The borderline tumors in our cohort, all of which were most highly correlated to the ovarian cancer C3 signature, also had highest correlation to the luminal A breast cancer subtype (p< 0.001) (Table S4), thus extending the links between the two tumor

Validation of the intrinsic breast cancer subtypes in ovarian cancer

Next, potential correlations between ovarian cancer C-signatures and the intrinsic breast cancer subtypes were explored. When the C-signatures for the tumors in our cohort were correlated to the intrinsic breast cancer subtypes for the same tumors, a significant heterogeneity within the tumor cohort was observed (p<0.001; figure 2). The ovarian cancer C2 and C4 signatures correlated significantly with the basal-like breast cancer subtype (p=0.019 and p=0.001, respectively) (Table S5), while the C3 signature correlated to the normal-like breast cancer subtype (p<0.001) (Table S6). These observations suggest commonalities between the transcriptionally based molecular classifiers of ovarian and breast cancer (Figure S4). The results were verified by classifying the ovarian tumors in the Tothill dataset [18] into the intrinsic breast cancer subtypes; similar links between the C2




Figure 1. SAM analysis. Supervised hierarchical clustering of malignant (n=37) and benign (n=17) serous ovarian tumors (FDR < 1%). Red represents relative upregulation and green represents relative downregulation.

doi:10.1371/journal.pone.0107643.g001

signature and the basal-like subtype, and the C3 signature and normal-like subtype were observed (p<0.001; Table S8). Notably, 16/26 patients (61.5%) whose tumors resembled the basal-like and HER2 enriched intrinsic breast cancer subtypes died from their disease within four years from diagnosis, compared to only 3/9 patients (33.3%) whose tumors displayed gene expression profiles corresponding to the luminal A and luminal B intrinsic breast cancer subtypes.

To further explore potential connections between the molecular subtypes of ovarian and breast cancer, we applied the gene expression modules representing key biological features of breast cancer described by Desmedt et al. to our tumors [33]. As for the ovarian cancer C-signatures and the intrinsic breast cancer subtypes, considerable heterogeneity within the tumor cohort was found. The malignant tumors displayed a significantly higher module score than the benign and borderline tumors for the AURKA/proliferation, STAT1/immune response, CASP3/apoptosis, VEGF/angiogenesis and ERBb2/HER2 signaling modules. The borderline and benign tumors on the other hand correlated to the ESR1/ER signaling module (Figure S5). The somewhat surprising absence of a correlation between the malignant tumors and the PLAU/tumor invasion and metastasis module led us to investigate the correlation between the Desmedt modules and the ovarian C-signatures. Again, significant differences within the cohort were observed, with highly significant correlations between the C1 signature and PLAU/invasion

Table 2. The 10 most significantly enriched biological processes in the malignant ovarian tumors in the study cohort [31].

	Biological process*	p-value	Genes from input ^a	Genes in annotation ^b
1.	mitotic cell cycle	3.823E ⁻²⁴	213	874
2.	cell cycle process	1.544E ⁻²³	265	1192
3.	antigen processing and presentation of exogenous peptide antigen	2.768E ⁻¹⁹	67	171
4.	antigen processing and presentation of exogenous peptide antigen via MHC class I	3.943E ⁻¹⁹	43	80
5.	cellular response to stress	6.028E ⁻¹⁹	279	1370
б.	symbiosis, encompassing mutualism through parasitism	1.886E ⁻¹⁸	175	741
' .	interspecies interaction between organisms	1.886E ⁻¹⁸	175	741
8.	antigen processing and presentation of exogenous antigen	3.374E ⁻¹⁸	67	178
9.	mitotic cell cycel phase transition	5.793E ⁻¹⁸	121	445
10.	cell cycle phase transition	6.556E ⁻¹⁸	122	451

^{*}FDR<0.05 and ≥3 recognized genes/biological function were required to consider a gene ontology (GO) process significant. 731 significant. GO processes were identified.

module (p<0.001), the C2 signature and STAT1/immune response module (p<0.001), and the C4 signature and VEGF/angiogenesis module (p=0.001). The C5 signature showed a trend towards correlation to the AURKA/proliferation module compared to non-C5 tumors, but did not reach statistical significance (p=0.176). The C3 signature correlated significantly to the ESR1/ER module (p<0.001, figure 3). Finally, we also applied the Desmedt modules to the tumors in the Tothill dataset, and could verify the correlations described between the ovarian cancer C-signatures and the functional breast cancer derived gene expression modules (Figure S6). In addition, in this larger cohort, the C5/AURKA correlation was also found to be significant (p<0.001).

Mutation analysis

Mutation analyses revealed four tumors with *KRAS* mutations and two tumors with *BRAF* mutations in our cohort. Four of these mutations were present among the borderline tumors, all of which

corresponded to the C3 ovarian cancer signature and the luminal A breast cancer subtype, respectively. Two malignant tumors, both grade 1, harbored mutations in *KRAS* and corresponded to the C3/normal-like subtypes and C4/luminal B subtypes, respectively (table 5).

Discussion

Serous carcinomas account for about 50% of the malignant epithelial ovarian tumors and thereby constitute the predominant histologic subtype. Type 1 tumors, i.e. low-grade serous carcinomas, along with low-grade endometrioid, mucinous and clear cell tumors, are thought to develop step-wise from benign cystadenomas/adenofibromas via borderline tumors and are typically slowly proliferating and frequently harbor mutations in *KRAS*, *BRAF* and *PTEN*. In contrast, type 2 tumors (high-grade serous and high-grade endometrioid ovarian tumors, carcinosarcomas and undifferentiated carcinomas) are suggested to develop from

Table 3. Ovarian cancer subtypes.

	C-signatur	C-signature C-signature								
	C1	C2	СЗ	C4	C5	Total				
Malignant	12	8	3	9	5	37				
(% within group)	(32.4)	(21.6)	(8.1)	(24.3)	(13.5)	(100.0)				
Borderline	0	0	5	0	0	5				
(% within group)	(0.0)	(0.0)	(100.0)	(0.0)	(0.0)	(100.0)				
Benign	1	0	15	0	1	17				
(% within group)	(5.9)	(0.0)	(88.2)	(0.0)	(5.9)	(100.0)				
p<0.001										

Serous ovarian tumors in the study cohort with corresponding ovarian cancer subtypes ("C-signatures") [18]. The rows outline the tumor types with the representation in each subtype in percent within parentheses. The p-value is calculated using Fisher's exact test. doi:10.1371/journal.pone.0107643.t003

^aNumber of genes in the study cohort correlating to the GO process.

^bNumber of genes in the GO process.

doi:10.1371/journal.pone.0107643.t002

Table 4. Intrinsic breast cancer subtypes.

	Intrinsic breast cancer subtype										
	Luminal A	Luminal B	Basal-like	Normal-like	Her2	Total					
Malignant	2	3	21	6	5	37					
(% within group)	(5.4)	(8.1)	(56.8)	(16.2)	(13.5)	(100.0)					
Borderline	4	0	0	1	0	5					
(% within group)	(80.0)	(0.0)	(0.09	(20.0)	(0.0)	(100.0)					
Benign	1	0	0	16	0	17					
(% within group)	(5.9)	(0.0)	(0.0)	(94.1)	(0.0)	(100.0)					
p<0.001											

Serous ovarian tumors in the study cohort with corresponding intrinsic breast cancer subtypes [22]. The rows outline the tumor types with the representation in each subtype in percent within parentheses. The p-value is calculated using Fisher's exact test. doi:10.1371/journal.pone.0107643.t004

precursor lesions in the fallopian tube [24] and are characterized by rapid progression and frequent TP53 mutations [10,34–36]. Molecular subtyping of ovarian cancer is being increasingly recognized, with e.g. the six transcriptionally based ovarian C-signatures proposed by Tothill *et al.* being both descriptive (with good correlations to clinical factors) and predictive of outcome [18].

Molecular subtypes of breast cancer have been established and linked to clinical behavior and treatment response [20,21]. Although about 60% of all ovarian tumors display high expression of estrogen receptors (ER) [37], features shared between high-grade serous ovarian cancer and basal-like breast cancer (the majority of which are "triple negative", i.e. lack expression of estrogen and progesterone receptors and HER2 amplification) have recently been reported. Both tumor groups display frequent TP53 mutations and genomic instability and are clinically aggressive. Also, BRCA mutations are more frequent in high-grade serous ovarian cancers and in basal-like breast cancers in the case of BRCA1 [19]. The diagnostic and therapeutic potential in clarifying ovarian cancer heterogeneity and identifying mechanisms shared between ovarian cancer and breast cancer constituted the basis of our study.





Figure 2. Correlations between ovarian and breast cancer molecular subtypes. Correlations between specific ovarian cancer C-signatures [18] and the intrinsic breast cancer subtypes [22] in the serous ovarian tumors in the study cohort. Tumors within each ovarian cancer C-signature are shown along the X axis, and the colored bars represent the percentage (on the Y axis) of each intrinsic breast cancer subtype within the respective C-signatures. doi:10.1371/journal.pone.0107643.q002

SAM analysis between the malignant and benign ovarian tumors in our cohort revealed enrichment of cell cycle associated processes among the malignant tumors, in line with malignant tumors being highly proliferative *per se*. Although very few borderline tumors were included in our cohort, the fact that they were divided between the benign and the malignant tumor clusters, regardless of whether the clustering was supervised by significantly differentially expressed genes between benign and malignant tumors, or unsupervised, is in line with other studies indicating that borderline tumors constitute a very heterogeneous group with both benign and malignant features [38].

Classification of the ovarian tumors in our cohort using the Csignatures demonstrated the presence of all but the C6 signature. Since the C6 signature is characterized mainly by low-grade endometrioid tumors, which were not present in our cohort, this finding supports the ability of gene signatures to capture histological differences, and indicates that the C-signatures are stable and widely applicable across datasets and microarray platforms. The malignant tumors correlated significantly to the C1, C2 and C4 signatures, and as anticipated from the recent data reported from the TCGA, classification into the intrinsic breast cancer subtypes also revealed a significant correlation between malignant ovarian tumors and the basal-like breast cancer subtype [19]. This link was further supported by the finding that the ovarian tumors classified as basal-like in our cohort in turn correlated to the C2 and C4 signatures. A majority of both highgrade serous ovarian and basal-like breast cancers express high levels of proliferation genes [16,19]. Proliferation, among other biological processes, is captured by the Desmedt modules, and we could correlate the C-signatures to these modules and verify the finding in an independent dataset. The C2 "high immune signature" correlated significantly to the STAT1/immune response module, for example, and the C1 "high stromal response signature" to the PLAU/invasion and metastasis module. Taken together, the breast cancer derived Desmedt modules capture the nature of the C-signatures as outlined by both Tothill et al. and the TCGA well [16,18] and provide further biological information regarding the differences in phenotype between the subgroups. Moreover, the statistical correlations shown here further support the link between the subtypes of serous ovarian and breast cancer.

The C3 signature in the original study encompassed borderline (low malignant potential, LMP) tumors and, as expected, the benign and borderline tumors in our cohort correlated significantly to the C3 signature. This signature is characterized by a relative overexpression of genes in the MAPK/ERK pathway, in



modules by Desmedt *et al* [33]. Log2 mRNA values are presented on the Y axis. p-values for the highlighted boxes vs. the rest in each plot are calculated using the Mann-Whitney U Test. doi:10.1371/journal.pone.0107643.g003

Table 5. KRAS and BRAF mutations.

ld	Feature	C-signature	BC subtype	Mutation
420	М	C4	Luminal B	KRAS
232	M	C3	Normal	KRAS
86	Во	C3	Luminal A	KRAS
385	Во	C3	Luminal A	KRAS
16	Во	C3	Luminal A	BRAF
48	Во	C3	Luminal A	BRAF

Distribution of KRAS and BRAF mutations and their correlations to ovarian cancer C-signatures and intrinsic breast cancer subtypes. Feature: M = Malignant, Bo = Borderline; C-signature: corresponding molecular ovarian cancer subtype [18]; BC-subtype: corresponding intrinsic breast cancer subtype

doi:10.1371/journal.pone.0107643.t005

line with the fact that type 1 ovarian tumors often harbor mutations in KRAS and BRAF. Four of five borderline tumors in our cohort displayed KRAS or BRAF mutations, and the two malignant tumors harboring mutations in KRAS in turn correlated to the C3 and C4 ovarian signatures. Interestingly, a significant correlation between the few borderline tumors and the luminal A breast cancer subtype was found. Luminal breast tumors frequently display mutations in the MAPK/ERK pathway, thereby resembling the ovarian type 1 tumors (and the ovarian C3 signature). This is further supported by the independent finding of a significant correlation between the ovarian C3 signature and the ESR1/ER signaling breast cancer module by Desmedt et al. [33]. Hence, although the borderline tumors in our cohort were interspersed between the malignant and benign neighbors in the clustering analyses, upon comparison with the ovarian cancer Csignatures and the intrinsic breast cancer subtypes, they showed obvious similarities with benign and low-grade malignant tumors as anticipated based on the prototypic type 1 tumors they are described as. Furthermore, the luminal A and B (estrogen receptor positive) breast cancer subtypes differ in transcriptional profiles, mutation spectra and overall survival [19]; this is in line with the clinical spectrum observed in low-grade serous ovarian cancer, with low-grade, early-stage tumors showing a favorable prognosis, while low-grade, advanced stage tumors tend to respond poorly to chemotherapy. Despite the generally high expression of ER in ovarian cancer, the response to both tamoxifen and letrozole has been limited [39,40]. The different isoforms of ER seem to vary with the malignant potential, with the beta isoform (ERβ) reported to be less expressed in malignant ovarian tumors compared to borderline tumors and benign ovaries, but whether ER β or ER α influence outcome is not clear [41,42]. In contrast, ERa is a favorable prognostic factor in breast cancer [43]. A recent study by the Ovarian Tumor Tissue Analysis consortium (OTTA) focused on expression of $ER\alpha$ and the progesterone receptor (PR), and in a large series only strong expression of PR, but not ERa, was correlated to increased survival in high-grade serous ovarian cancer. No significant correlations between $\text{ER}\alpha$ or PR expression and survival were found in multivariate analyses of low-grade serous tumors [44]. The vast majority of the low-grade tumors were however ER and/or PR positive, but the few that were negative did not have a significantly different outcome despite the fact that the majority of the low-grade tumors were stage III–IV. Likewise, ER α is reported to be overexpressed in serous borderline tumors [41]. Taking these results into consideration in light of our findings of a correlation between the ovarian C3 signature and the luminal A breast cancer subtype, it would be interesting to study

the response to and the potential effect of endocrine treatment specifically in advanced type I ovarian tumors.

Conclusions

The findings in this study support that transcriptional signatures indeed capture the biology of transforming events and oncogenic mutations and also support similarities between molecular subtypes of ovarian and breast cancer beyond high-grade serous ovarian cancer and basal-like breast cancer. Though limited series are sensitive to overfitting, importantly, our findings were stable and reproducible in a large independent cohort. The similarities between molecular subtypes of ovarian and breast cancer may be of potential interest for further studies regarding targeted therapies and the use of chemotherapeutic agents in ovarian cancer, as well as biomarker studies. While the proposed similarities between lowgrade serous and borderline ovarian (type 1) tumors and luminal breast cancers may in part be attributable to similarities in proliferation rates compared to high-grade ovarian (type 2) and basal-like breast cancers, other biological similarities, such as potential endocrine responsiveness, are thought-provoking and merit further investigation.

Supporting Information

Figure S1 Hierarchical clustering. Unsupervised clustering of malignant and benign ovarian tumors using the 20% most varying probes and including biological replicates. n = 66 tumors. (TIF)

Figure S2 Hierarchical clustering. Unsupervised clustering of malignant and benign ovarian tumors using the 20% most varying probes. Clustering performed without biological replicates. n = 54 tumors. (TIF)

Figure S3 Hierarchial clustering. Supervised clustering of malignant, borderline and benign tumors based on significant probes from supervised analysis of malignant and benign tumors. Clustering performed without biological replicates. n = 59 tumors. (TIF)

Figure S4 Hierarchical clustering. The serous ovarian tumors in the study cohort with corresponding tumor features and assigned ovarian cancer C-signatures and intrinsic breast cancer subtypes.

(TIF)

Figure S5 Functional gene expression modules. Boxplots representing the correlations between serous ovarian malignant,

borderline and benign tumors in the study cohort and the respective gene expression modules by Desmedt et al. (Desmedt et al., Clin Cancer Res 2008). Log2 mRNA values are presented on the Y axes. p-values are calculated using the Kruskal-Wallis test. (TIF)

Figure S6 Functional gene expression modules. Boxplots representing the correlations between the C1-C6 ovarian cancer signatures in an independent, publicly available dataset and the respective breast cancer gene expression modules by Desmedt et al. (Desmedt et al., Clin Cancer Res 2008). Log2 mRNA values are presented on the Y axes. p-values for the highlighted boxes vs. the rest in each plot are calculated using the Mann-Whitney U Test. (TIF)

Table S1 Deregulated genes. All significantly deregulated genes (n = 5,944) between malignant and benign ovarian tumors in the study cohort. (XLSX)

Table S2 Molecular subtypes. Significant correlations between malignant, borderline and benign ovarian tumors in the study cohort (n = 59) and the ovarian cancer C-signatures. p-values for each part of the table (separated with double lines) are calculated using Fisher's exact test. (XLSX)

Table S3 Molecular subtypes. Significant correlations between malignant, borderline and benign ovarian tumors in the study cohort (n = 59) and the basal-like and normal-like breast cancer subtypes. p-values for each part of the table (separated with double lines) are calculated using Fisher's exact test. (XLSX)

Table S4 Molecular subtypes. Significant correlations between malignant, borderline and benign ovarian tumors in the study cohort (n = 59) and the luminal A breast cancer subtype. The p-value is calculated using Fisher's exact test. (XLSX)

Table S5 Correlations between molecular subtypes. Significant correlations between and between assigned C-signatures and the basal-like breast cancer subtype. p-values for each

References

- Kurman RJ, le Shih M (2010) The origin and pathogenesis of epithelial ovarian cancer: a proposed unifying theory. Am J Surg Pathol 34: 433-443.
- Piek JM, van Diest PJ, Zweemer RP, Jansen JW, Poort-Keesom RJ, et al. (2001) Dysplastic changes in prophylactically removed Fallopian tubes of women
- predisposed to developing ovarian cancer. J Pathol 195: 451–456. Kindelberger DW, Lee Y, Miron A, Hirsch MS, Feltmate C, et al. (2007) Intraepithelial carcinoma of the fimbria and pelvic serous carcinoma: Evidence for a causal relationship. Am J Surg Pathol 31: 161-169.
- 4. Heintz AP, Odicino F, Maisonneuve P, Quinn MA, Benedet JL, et al. (2006) Carcinoma of the ovary. FIGO 26th Annual Report on the Results of Treatment in Gynecological Cancer. Int J Gynaecol Obstet 95 Suppl 1: S161-192.
- Bookman MA, Brady MF, McGuire WP, Harper PG, Alberts DS, et al. (2009) Evaluation of new platinum-based treatment regimens in advanced-stage ovarian cancer: a Phase III Trial of the Gynecologic Cancer Intergroup. J Clin Oncol 27: 1419-1425
- Winter-Roach BA, Kitchener HC, Dickinson HO (2009) Adjuvant (post-surgery) chemotherapy for early stage epithelial ovarian cancer. Cochrane Database Syst
- Perren TJ, Swart AM, Pfisterer J, Ledermann JA, Pujade-Lauraine E, et al. (2011) A phase 3 trial of bevacizumab in ovarian cancer. N Engl I Med 365: 2484-2496
- 8. McCluggage WG (2011) Morphological subtypes of ovarian carcinoma: a review with emphasis on new developments and pathogenesis. Pathology 43: 420-432.
- Saad AF, Hu W, Sood AK (2010) Microenvironment and pathogenesis of epithelial ovarian cancer. Horm Cancer 1: 277-290.

part of the table (separated with double lines) are calculated using Fisher's exact test.

(XLSX)

Table S6 Correlations between molecular subtypes. Significant correlations between and between assigned C-signatures and the normal-like breast cancer subtype. The p-value is calculated using Fisher's exact test. (XLSX)

Table S7 Validation of the centroid classifier. Cross table comparing the original k.means groups (rows) for the tumors in an independent, publicly available dataset, consisting of malignant and borderline tumors, with the centroid classification for the same tumors (columns). The numbers represent number of tumors. Correlations between the C-signature classifications are highlighted in bold (diagonal). 251 tumors had an assigned k.means group. 239 of these (95.2%) were correctly re-assigned using nearest centroid classification.

(XLSX)

Table S8 Correlations between molecular subtypes. Correlations between ovarian cancer C-signatures and intrinsic breast cancer subtypes in an independent, publicly available dataset consisting of malignant and borderline ovarian tumors (n = 285). The C2 signature correlated significantly to the basallike breast cancer subtype and the C3 signature to the normal-like breast cancer subtype. p-values (*) are calculated using Fisher's exact test.

(XLSX)

Acknowledgments

We would like to acknowledge the patients who have generously contributed to the Skåne University Hospital ovarian tumor biobank.

Author Contributions

Conceived and designed the experiments: JMJ SM MN IH MJ IJ MDV. Performed the experiments: JMJ IJ MDV SK MJ AM JHB. Analyzed the data: JMJ IJ MDV SK AM SM MJ MN IH JHB PK. Contributed reagents/materials/analysis tools: JMJ SM PK AM MN. Contributed to the writing of the manuscript: JMJ IJ MDV SK MJ JHB PK AM SM MN

- 10. Prat J (2012) Ovarian carcinomas: five distinct diseases with different origins, genetic alterations, and clinicopathological features. Virchows Arch 460: 237-
- 11. Kobel M, Kalloger SE, Boyd N, McKinney S, Mehl E, et al. (2008) Ovarian carcinoma subtypes are different diseases: implications for biomarker studies. PLoS Med 5: e232.
- 12. Bonome T, Lee JY, Park DC, Radonovich M, Pise-Masison C, et al. (2005) Expression profiling of serous low malignant potential, low-grade, and highgrade tumors of the ovary. Cancer Res 65: 10602-10612.
- 13. Meinhold-Heerlein I, Bauerschlag D, Hilpert F, Dimitrov P, Sapinoso LM, et al. (2005) Molecular and prognostic distinction between serous ovarian carcinomas of varying grade and malignant potential. Oncogene 24: 1053-1065.
- 14. Marquez RT, Baggerly KA, Patterson AP, Liu J, Broaddus R, et al. (2005) Patterns of gene expression in different histotypes of epithelial ovarian cancer correlate with those in normal fallopian tube, endometrium, and colon. Clin Cancer Res 11: 6116-6126.
- 15. Marchini S, Mariani P, Chiorino G, Marrazzo E, Bonomi R, et al. (2008) Analysis of gene expression in early-stage ovarian cancer. Clin Cancer Res 14: 7850-7860.
- 16. The Cancer Genome Atlas Network (2011) Integrated genomic analyses of ovarian carcinoma. Nature 474: 609-615.
- 17. Yoshihara K, Tsunoda T, Shigemizu D, Fujiwara H, Hatae M, et al. (2012) High-risk ovarian cancer based on 126-gene expression signature is uniquely characterized by downregulation of antigen presentation pathway. Clin Cancer Res 18: 1374-1385.

- Tothill RW, Tinker AV, George J, Brown R, Fox SB, et al. (2008) Novel molecular subtypes of serous and endometrioid ovarian cancer linked to clinical outcome. Clin Cancer Res 14: 5198–5208.
- The Cancer Genome Atlas Network (2012) Comprehensive molecular portraits of human breast tumours, Nature 490: 61–70.
- Perou CM, Sorlie T, Eisen MB, van de Rijn M, Jeffrey SS, et al. (2000) Molecular portraits of human breast tumours. Nature 406: 747–752.
- Sorlie T, Perou CM, Tibshirani R, Aas T, Geisler S, et al. (2001) Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications. Proc Natl Acad Sci U S A 98: 10869–10874.
- Hu Z, Fan C, Oh DS, Marron JS, He X, et al. (2006) The molecular portraits of breast tumors are conserved across microarray platforms. BMC Genomics 7: 96.
- Anglesio MS, Arnold JM, George J, Tinker AV, Tothill R, et al. (2008) Mutation
 of ERBB2 provides a novel alternative mechanism for the ubiquitous activation
 of RAS-MAPK in ovarian serous low malignant potential tumors. Mol Cancer
 Res 6: 1678–1690.
- Shih Ie M, Kurman RJ (2004) Ovarian tumorigenesis: a proposed model based on morphological and molecular genetic analysis. Am J Pathol 164: 1511–1518.
- Silverberg SG (2000) Histopathologic grading of ovarian carcinoma: a review and proposal. Int J Gynecol Pathol 19: 7–15.
- Tavassoli FA, Devilee P (2003) Pathology and Genetics of Tumours of the Breast and Female Genital Organs; Tavassoli Fattaneh A DP, editor. Lyon: IARC Press.
- Edgar R, Domrachev M, Lash AE (2002) Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. Nucleic Acids Res 30: 207–210.
- Saeed AI, Sharov V, White J, Li J, Liang W, et al. (2003) TM4: a free, opensource system for microarray data management and analysis. Biotechniques 34: 374–378.
- Tusher VG, Tibshirani R, Chu G (2001) Significance analysis of microarrays applied to the ionizing radiation response. Proc Natl Acad Sci U S A 98: 5116– 5121.
- Eden E, Navon R, Steinfeld I, Lipson D, Yakhini Z (2009) GOrilla: a tool for discovery and visualization of enriched GO terms in ranked gene lists. BMC Bioinformatics 10: 48.
- Chen J, Aronow BJ, Jegga AG (2009) Disease candidate gene identification and prioritization using protein interaction networks. BMC Bioinformatics 10: 73.
- Johansson I, Nilsson C, Berglund P, Lauss M, Ringner M, et al. (2012) Gene expression profiling of primary male breast cancers reveals two unique

- subgroups and identifies N-acetyltransferase-1 (NAT1) as a novel prognostic biomarker. Breast Cancer Res $14\colon R31$.
- Desmedt C, Haibe-Kains B, Wirapati P, Buyse M, Larsimont D, et al. (2008) Biological processes associated with breast cancer clinical outcome depend on the molecular subtypes. Clin Cancer Res 14: 5158–5165.
- Hauptmann S, Dietel M (2001) Serous tumors of low malignant potential of the ovary-molecular pathology: part 2. Virchows Arch 438: 539–551.
- 35. Li J, Fadare O, Xiang L, Kong B, Zheng W (2012) Ovarian serous carcinoma: recent concepts on its origin and carcinogenesis. J Hematol Oncol 5: 8.
- Berns EM, Bowtell DD (2012) The changing view of high-grade serous ovarian cancer. Cancer Res 72: 2701–2704.
- Perez-Gracia JL, Carrasco EM (2002) Tamoxifen therapy for ovarian cancer in the adjuvant and advanced settings: systematic review of the literature and implications for future research. Gynecol Oncol 84: 201–209.
- Curry EW, Stronach EA, Rama NR, Wang YY, Gabra H, et al. (2013) Molecular subtypes of serous borderline ovarian tumor show distinct expression patterns of benign tumor and malignant tumor-associated signatures. Mod Pathol.
- Smyth JF, Gourley C, Walker G, MacKean MJ, Stevenson A, et al. (2007)
 Antiestrogen therapy is active in selected ovarian cancer cases: the use of letrozole in estrogen receptor-positive patients. Clin Cancer Res 13: 3617–3622.
- Hatch KD, Beecham JB, Blessing JA, Creasman WT (1991) Responsiveness of patients with advanced ovarian carcinoma to tamoxifen. A Gynecologic Oncology Group study of second-line therapy in 105 patients. Cancer 68: 269–271.
- Chan KK, Wei N, Liu SS, Xiao-Yun L, Cheung AN, et al. (2008) Estrogen receptor subtypes in ovarian cancer: a clinical correlation. Obstet Gynecol 111: 144–151.
- Burges A, Bruning A, Dannenmann C, Blankenstein T, Jeschke U, et al. (2010) Prognostic significance of estrogen receptor alpha and beta expression in human serous carcinomas of the ovary. Arch Gynecol Obstet 281: 511–517.
- 43. Early Breast Cancer Trialists' Collaborative Group (2005) Effects of chemotherapy and hormonal therapy for early breast cancer on recurrence and 15-year survival: an overview of the randomised trials. Lancet 365: 1687–1717.
- Sieh W, Kobel M, Longacre TA, Bowtell DD, deFazio A, et al. (2013)
 Hormone-receptor expression and ovarian cancer survival: an Ovarian Tumor Tissue Analysis consortium study. Lancet Oncol 14: 853–862.