



LUND UNIVERSITY

MAPK signaling pathway regulates cerebrovascular receptor expression in human cerebral arteries

Ansar, Saema; Eftekhari, Sajedah; Waldsee, Roya; Nilsson, Elisabeth; Nilsson, Ola; Säveland, Hans; Edvinsson, Lars

Published in:
BMC Neuroscience

DOI:
[10.1186/1471-2202-14-12](https://doi.org/10.1186/1471-2202-14-12)

2013

[Link to publication](#)

Citation for published version (APA):

Ansar, S., Eftekhari, S., Waldsee, R., Nilsson, E., Nilsson, O., Säveland, H., & Edvinsson, L. (2013). MAPK signaling pathway regulates cerebrovascular receptor expression in human cerebral arteries. *BMC Neuroscience*, 14, Article 12. <https://doi.org/10.1186/1471-2202-14-12>

Total number of authors:
7

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

RESEARCH ARTICLE

Open Access

MAPK signaling pathway regulates cerebrovascular receptor expression in human cerebral arteries

Saema Ansar^{1*}, Sajedah Eftekhari¹, Roya Waldsee¹, Elisabeth Nilsson¹, Ola Nilsson², Hans Sävemark² and Lars Edvinsson¹

Abstract

Background: Cerebral ischemia results in enhanced expression of contractile cerebrovascular receptors, such as endothelin type B (ET_B), 5-hydroxytryptamine type 1B (5-HT_{1B}), angiotensin II type 1 (AT₁) and thromboxane (TP) receptors in the cerebral arteries within the ischemic area. The receptor upregulation occurs via activation of the mitogen-activated protein kinases (MAPK) pathway. Previous studies have shown that inhibitors of the MAPK pathway diminished the ischemic area and contractile cerebrovascular receptors after experimental cerebral ischemia. The aim of this study was to examine if the upregulation of contractile cerebrovascular receptors after 48 h of organ culture of human cerebral arteries involves MAPK pathways and if it can be prevented by a MEK1/2 inhibitor. Human cerebral arteries were obtained from patients undergoing intracranial tumor surgery. The vessels were divided into ring segments and incubated for 48 h in the presence or absence of the specific MEK1/2 inhibitor U0126. The vessels were then examined by using in vitro pharmacological methods and protein immunohistochemistry.

Results: After organ culture of the cerebral arteries the contractile responses to endothelin (ET)-1, angiotensin (Ang) II and thromboxane (TP) were enhanced in comparison with fresh human arteries. However, 5-carboxamidotryptamine (5-CT) induced decreased contractile responses after organ culture as compared to fresh arteries. Incubation with U0126 diminished the maximum contraction elicited by application of ET-1, Ang II and U46619 in human cerebral arteries. In addition, the MEK1/2 inhibitor decreased the contractile response to 5-CT. Immunohistochemistry revealed that organ culture resulted in increased expression of endothelin ET_A, endothelin ET_B, angiotensin AT₂, 5-hydroxytryptamine 5-HT_{1B} and thromboxane A₂ receptors, and elevated levels of activated pERK1/2, all localized to the smooth muscle cells of the cerebral arteries. Co-incubation with U0126 normalized these proteins.

Conclusion: The study demonstrated that there is a clear association between human cerebrovascular receptor upregulation via transcription involving activation of the MAPK pathway after organ culture. Inhibition of the MAPK pathways attenuated the vasoconstriction mediated by ET, AT and TP receptors in human cerebral arteries and the enhanced expression of their receptors. The results indicate that MAPK inhibition might be a novel target for treatment of cerebrovascular disorders.

Keywords: Human cerebral arteries, Endothelin receptors, Angiotensin receptors, 5-hydroxytryptamine receptors, Thromboxane receptors, Mitogen activated protein kinase

* Correspondence: Saema.Ansar@med.lu.se

¹Department of Clinical Sciences, Division of Experimental Vascular Research, Lund University, Lund, Sweden

Full list of author information is available at the end of the article

Background

Large efforts have during the last few decades been made to understand the intracellular mechanisms involved in ischemia-induced cerebral damage and to develop drugs that protect the brain from damage once a stroke has occurred. However, despite extensive research into genetics and molecular biology associated with cerebral ischemia, few acute therapies have proven effective in the clinic [1]. Investigations have revealed that cerebral ischemia is accompanied by modifications in the expression of genes regulating receptor expressions in cerebrovascular smooth muscle cell (SMC)s associated with the cerebral ischemia [2]. Thus, experimental and clinical studies of cerebral ischemia have reported increased levels of the potent vasoconstrictor substances endothelin (ET) [3,4], 5-hydroxytryptamine (5-HT) [5,6], angiotensin (Ang II) [7] and thromboxane (TXA₂) [8,9]. ET-1, 5-HT, Ang II and TXA₂ are all potent vasoconstrictors of cerebral arteries that mediate effects through the family of G-protein coupled receptors (GPCRs) [10,11]; endothelin A (ET_A), endothelin B (ET_B) [11,12], 5-HT receptors [5], the angiotensin II type 1 (AT₁) and type 2 (AT₂) receptors [13,14] and the thromboxane receptor (TP) [15].

Cerebral ischemia is multifactorial, involves a number of neuronal and glial mechanisms; however, several cerebrovascular receptors are in addition involved in the pathophysiology of cerebral ischemia. There is upregulation (enhanced expression) of contractile ET_B, 5-HT_{1B}, AT₁ and TP receptors in major cerebral arteries from experimental focal and global ischemia, via enhanced transcription and translation [16-24]. This upregulation of cerebrovascular receptors leads to enhanced vasoconstriction and correlates with reduction in regional cerebral blood flow (rCBF) and degree of neurology deficit [21]. Blockade of the individual subtypes of receptors involved might prevent or reduce the cerebral ischemia to a certain degree; we hypothesize that treatment aimed at a common signaling pathway would be more beneficial by avoiding the administration of several antagonists with circulatory consequences.

The mitogen-activated protein kinase (MAPK) pathways are implicated in neuronal death and survival after stroke. A time study of the alteration in cerebrovascular MAPKs after experimental subarachnoid hemorrhage (SAH) revealed that there was early (within minutes) and sustained activation of the specific extracellular signal-regulated kinases (ERK)1/2 pathway, while the p38 and JNK pathways were activated first at 48 hours [25]. The ERK1/2 pathway can be inhibited at various points upstream such as at ras-raf-MEK1/2; inhibition of this pathway with a specific MEK1/2 (the MAPKK of ERK1/2) inhibitor abolished the receptor upregulation as well as preventing the CBF reduction and diminishes the infarct [16,26,27]. ERK1/2 belongs

to the family of MAPK and is phosphorylated and thereby activated by the MAP kinase/ERK kinase (MEK)1/2. Several studies have shown an involvement of the MEK/ERK1/2 signalling pathway in cerebral ischemia [26,28].

Organ culture is an *in vitro* method for investigating cellular mechanisms involved in upregulation of vasoconstrictile G-protein coupled receptors. Organ culture is not a model for stroke, however, changes in vasoconstrictor responses after *in vitro* organ culture show a remarkable similarity to changes observed in animal models of ischemic and hemorrhagic stroke. Thus, there is an upregulation of contractile G-protein receptors after SAH [21] and focal ischemia [22,23] which also is observed in organ culture [29]. This make the organ culture model an appropriate model for investigating the pharmacological characteristics and underlying molecular and cellular mechanism involved in the upregulation of vasoconstrictile G-protein coupled receptors. The upregulation of contractile receptors in the SMCs are prevented with MAPK inhibitor both in organ culture [29] and experimental stroke [27,30].

In the design of future cerebrovascular therapeutics it is important that the intracellular mechanisms are characterised in human subjects. Here we hypothesize that there is an upregulation of contractile cerebrovascular receptors after 48 h of organ culture in human cerebral arteries and that this upregulation occurs via the MAPK ERK1/2 pathway and can be inhibited by the MEK1/2 inhibitor U0126.

Methods

All procedures were carried out strictly within national laws and guidelines and approved by the Ethical Committee at the University of Lund (LU-818-01) and has been performed in accordance with the Declaration of Helsinki. A consent was obtained from the participants prior to surgery.

Tissue collection and organ culture procedure

Cortical arteries were obtained from patients undergoing neurological surgery for brain tumors. The arteries obtained were physiological arteries with surrounded tumor tissue, the arteries were carefully dissected free of connective tissue leaving the vessel with intact intima, media and adventitia. The arteries were immediately immersed in cold sterile Dulbecco's modified Eagle's medium (DMEM, Gibco, Invitrogen, Carlsbad, CA, USA) and transported to the laboratory. The arteries were cut into 1-mm long ring segments for *in vitro* pharmacological experiments and 3-mm for immunohistochemistry. The outer diameters were between 300 and 800 μ m.

Organ culture

The arterial segments were cultured for 48 hours at 37°C in humidified 5% CO₂ and air in Dulbecco's modified Eagle's medium supplemented with penicillin (100 U/ml),

streptomycin (100 µg/ml) and amphotericin B (25 µg/ml). The method of blood vessel culture has been described previously [31]. The segments were cultured in the absence or presence of the MEK1/2 inhibitors U0126 (5 µM). The selection of the inhibitor U0126 was based on previous detailed work on isolated arteries in organ culture, where U0126 was demonstrated to be the best of all available MEK1/2 inhibitors to inhibit the GPCRs and MAPK pathway [29,32].

In vitro pharmacology myograph experiments


For contractile experiments a sensitive myograph was used for recording the isometric tension in isolated cerebral arteries [33,34]. The vessels were cut into 1 mm long cylindrical segments and mounted on two 40 µm in diameter stainless steel wires in a Myograph (Danish Myo Technology A/S, Denmark). One wire was connected to a force displacement transducer attached to an analogue-digital converter unit (ADInstruments, Oxford, UK). The other wire was connected to a micrometer screw, allowing fine adjustments of vascular tone by varying the distance between the wires. Measurements were recorded on a computer by use of a PowerLab unit (ADInstruments). The segments were immersed in a temperature controlled buffer solution (37°C) of the following composition (mM) NaCl 119, NaHCO₃ 15, KCl 4.6, MgCl₂ 1.2, NaH₂PO₄ 1.2, CaCl₂ 1.5 and glucose 5.5. The buffer was continuously aerated with oxygen enriched with 5% CO₂ resulting in a pH of 7.4. Initially, the vessel segments were normalized and set to an initial resting tone of 2 mN that is the tone that it would have if relaxed and under a transmural pressure of 100 mmHg. The vessels were allowed to stabilize at this tone for 1 hour. The contractile capacity was determined by exposing the vessels to an isotonic solution containing 63.5 mM of K⁺, obtained by partial change of NaCl for KCl in the above buffer. The contraction induced by K⁺ was used as reference for the contractile capacity [34]. Only vessels responding by contraction of at least 2 mN to potassium were included in the study.

Concentration-response curves were obtained by cumulative application of 5-carboxamidotryptamine (5-CT; specific 5-HT₁ receptor agonist (Sigma, St. Louis, USA)) in the concentration range 10⁻¹² to 10⁻⁵ M, ET-1 (Endothelin ET_A and ET_B receptor agonist (AnaSpec, San Jose, USA)) in the concentration range 10⁻¹⁴ to 10⁻⁷ M, U46619 (Thromboxane A₂ receptor agonist (Sigma, St. Louis, USA)) in the concentration range 10⁻¹² to 10⁻⁶ M and Ang II (Angiotensin AT₁ and AT_{II} receptor agonist (Sigma, St. Louis, USA)) in the concentration range 10⁻¹² to 10⁻⁶ M.

Immunohistochemistry

For immunofluorescence the cerebral artery segments were embedded in Tissue TEK (Gibo, Invitrogen A/S, Taastrup,

Denmark), frozen at -80°C and subsequently sectioned into 10 µm thick slices. Cryostat sections were fixed for 10 minutes in ice-cold acetone (-20°C) and thereafter rehydrated in phosphate buffered-saline (PBS, pH 7.2) containing 0.25% Triton X-100 (PBST), for 3×5 minutes. The sections were then permeabilized and blocked for 1 h in blocking solution containing PBS and 5% normal donkey serum and then incubated over night at 4°C with either of the following primary antibodies; rabbit anti ET_A (1:50, Santa Cruz Biotechnology, CA, U.S.A., sc-33535), rabbit anti ET_B (1:200, Abcam, Cambridge, UK, ab1921), rabbit anti AT₁ (1:100, Santa Cruz Biotechnology, sc-1173), AT₂ (1:100, Santa Cruz Biotechnology, sc-9040), 5-HT_{1B} (1:100, Santa Cruz Biotechnology, sc-1460), rabbit TP-receptor (1:100, Cayman Chemical company, Michigan, U.S.A., 10004452) and rabbit anti phospho-ERK p44/42 MAPK (1:50, Cell Signaling Technology, Beverly, CA, U.S.A., #4376). The primary antibodies were diluted in PBST, 1% bovine serum albumin (BSA) and 3% normal donkey serum. On the second day sections were rinsed in PBST for 3×15 minutes and incubated with the secondary antibody (1 h, room-temperature). The secondary antibody used was CyTM2 conjugated donkey anti rabbit (1:200, Jackson ImmunoResearch, West Grove, PA, U.S.A., 711-165-152) diluted in PBST and 1% BSA. The sections were washed subsequently with PBST and mounted with Crystal mounting medium (Sigma, St. Louis, MO, U.S.A). Immunoreactivity was visualized with an Olympus Microscope (BX 60, Japan) at the appropriate wavelength. Negative controls for all antibodies were made by omitting primary antibodies. In all cases, no specific staining was found; only auto-fluorescence in lamina elastica interna was seen (Figure 1). To evaluate the auto-fluorescence in lamina elastica interna, controls were made with only primary antibodies.



Calculations and statistics

Data are expressed as mean \pm standard error of the mean (s.e.m.), and n refers to the number of patients. Statistical analyses were performed with Kruskal-Wallis non-parametric test with Dunn's post-hoc test, where $P < 0.05$ was considered significant.

In vitro pharmacology

Contractile responses in each segment are expressed as percentage of the 63.5 mM K^+ induced contraction. E_{max} value represents the maximum contractile response elicited by an agonist and the pEC_{50} the negative logarithm of the drug concentration that elicited half the maximum response.

Immunohistochemistry

Measurements were made in order to quantify the immunoreactivity of the protein expressions in the different groups. The evaluation of receptors and pERK expression was performed by measurements of the fluorescence intensity using the software image J <http://rsb.info.nih.gov/ij/>. The immunoreactivity of the individual receptors was visualized with the same microscope settings during the same day for all groups. The intensity measurements were performed in a blinded manner. The fluorescence intensity was measured in four given areas in the smooth muscle layer of each sample (always clockwise). Values were given by dividing the fluorescence intensity to measured area of each sample. These values are presented as percentage fluorescence in the cultured groups compared to the fresh groups (henceforth only mentioned as control), where the fresh (control) group is set to 100%.

Results

Functional in vitro pharmacology

K^+ -induced contraction, E_{max} and pEC_{50} values for respective group are presented in Table 1. Of all vessels tested 72% responded to a K^+ -induced contraction.

ET-1 receptor

Contractile response to ET-1

In cultured arteries ET-1 yielded contractions with an E_{max} of $143 \pm 22\%$ and a pEC_{50} of 8.74 ± 0.25 . These values were significantly higher than those observed in control arterial segments, in which an E_{max} of $107 \pm 12\%$ was observed. This is in accordance with previous results, which show a similar upregulation in human cerebral arteries after organ culture [35]. The presence of U0126 during the organ culture produced a significantly attenuated ET-1 induced response, with an E_{max} of $57 \pm 8\%$ compared to the cultured arteries (Figure 2A, Table 1).

Protein expression examined by immunohistochemistry

The ET_A receptor protein was increased after organ culture ($137 \pm 3\%$) as compared to control ($100 \pm 4\%$).

Table 1 Contractile effects of ET-1, U46619, Ang II and 5-CT in human cerebral arteries

	N	K^+ mean \pm s.e.m	E_{max} (%) \pm s.e.m	$pEC_{50} \pm$ s.e.m
ET-1				
Control	6	6.76 ± 1.32	107 ± 12^a	7.20 ± 0.89
Organ Culture	6	4.37 ± 1.21	$143 \pm 22^{a,b}$	8.74 ± 0.25
Organ Culture + U0126	7	3.53 ± 0.90	57 ± 8^b	8.49 ± 0.10
U46619				
Control	7	8.26 ± 1.98	102 ± 15^a	7.35 ± 0.74
Organ Culture	7	5.10 ± 1.60	$141 \pm 11^{a,b}$	7.18 ± 0.18
Organ Culture + U0126	4	3.65 ± 1.89	76 ± 40^b	6.64 ± 0.10
Ang II				
Control	6	1.67 ± 0.54	12 ± 2^a	
Organ Culture	5	4.44 ± 1.92	$43 \pm 12^{a,b}$	9.15 ± 0.82
Organ Culture + U0126	4	4.46 ± 2.41	1 ± 1^b	9.05 ± 1.13
5-CT				
Control	6	6.76 ± 1.36	50 ± 8	7.45 ± 0.66
Organ Culture	4	3.90 ± 1.59	17 ± 5^b	7.35 ± 1.21
Organ Culture + U0126	7	3.53 ± 0.90	2 ± 1^b	7.38 ± 0.73

Responses were characterized by E_{max} values, expressed as percent of 63 mM K^+ -induced contraction, and pEC_{50} values (negative logarithm of the molar concentration that produces half maximum contraction). Values are represented as mean \pm s.e.m and n represents number of patients.

^a = significant difference between organ culture and control groups,
^b = significant difference between organ culture and organ culture treated with U0126.

Incubation with U0126 prevented the increased expression of ET_A ($107 \pm 4\%$) receptor protein on the smooth muscle cells (Figure 2B and 2D, Table 2). In addition, the ET_B receptor protein was expressed in the smooth muscle cells and this signal was increased in organ culture ($128 \pm 3\%$) as compared to control arteries ($100 \pm 6\%$). Treatment with the MEK1/2 inhibitor U0126 prevented the upregulation of ET_B ($80 \pm 4\%$), receptor protein levels in the smooth muscle cell layer as compared to the organ culture (Figure 2C, 2D Table 2).

Thromboxane receptor

Contractile response

In cultured arteries U46619 yielded contractions with an E_{max} of $141 \pm 11\%$. This value was significantly higher than those observed in control arterial segments, in which an E_{max} of $102 \pm 15\%$ was observed (Figure 3A). The presence of the MEK1/2 inhibitor U0126 during the organ culture produced a significantly attenuated U46619 contractile response, compared to the cultured arteries. There was no significant difference in the E_{max} between control arteries and cultured arteries treated with U0126 (Figure 3A, Table 1).




Figure 2 (See legend on next page.)

(See figure on previous page.)

Figure 2 A. Concentration response curves elicited by cumulative application of ET-1 in human cerebral arteries incubated for 48 h with or without the MEK1/2 inhibitor U0126 (5 μ M) and control human arteries. The responses to ET-1 are clearly increased in the incubated arteries as compared to control arteries. The enhanced contraction observed after organ culture was inhibited in arteries cultured with U0126. Data are expressed as mean \pm s.e.m and n refer to the number of patients used. **B and C** Sections from the human cerebral artery showing ET_A and ET_B receptor immunoreactivity in the smooth muscle cell layer. **D** Expression of ET_A and ET_B protein levels in human cerebral arteries incubated for 48 h with or without the MEK1/2 inhibitor U0126 (5 μ M) and control human arteries. Data are expressed as percentage of control and given as mean \pm s.e.m. * P <0.05, ** P <0.01.

Protein expression examined by immunohistochemistry

The TP receptor protein was expressed in the smooth muscle cells and this signal was slightly increased in organ culture ($119 \pm 5\%$) as compared to control arteries ($100 \pm 3\%$). Treatment with the MEK1/2 inhibitor U0126 prevented the upregulation of TP ($106 \pm 3\%$), receptor protein levels in the smooth muscle cell layer as compared to the organ culture, however not significantly (Figure 3B-C and Table 2).

Angiotensin receptor

Contractile response to Ang II

In cultured arteries Ang II induced a concentration-dependent contraction with an E_{\max} of $43 \pm 15\%$ and a pEC_{50} of 9.15 ± 1.65 . These values were significantly higher than those observed in control arterial segments, in which an E_{\max} of $12 \pm 2\%$ was observed (Figure 4A, Table 1). The presence of U0126 during the organ culture produced a significantly attenuated Ang II induced response, compared to the cultured arteries. Interestingly there was no significant difference in the contractile response between control arteries and cultured arteries treated with U0126 (Figure 4A, Table 1).

Table 2 Receptor protein levels in human cerebral arteries

	Control	Organ. Culture	Organ Culture + treatment with U0126
ET _B (%) \pm s.e.m	100 ± 6 a	128 ± 3 ^{a, b}	80 ± 4 ^b
ET _A (%) \pm s.e.m	100 ± 4 a	137 ± 3 ^{a, b}	107 ± 4 ^b
TP (%) \pm s.e.m	100 ± 3	119 ± 5	106 ± 3
AT ₁ (%) \pm s.e.m	100 ± 2 a	62 ± 4 ^a	82 ± 7
AT ₂ (%) \pm s.e.m	100 ± 2 a	120 ± 2 ^a	111 ± 1
5-HT _{1B} (%) \pm s.e.m	100 ± 6 a	128 ± 3 ^{a, b}	84 ± 4 ^b
pERK1/2 (%) \pm s.e.m	100 ± 2 a	128 ± 2 ^{a, b}	86 ± 2 ^b

Activation of the different protein levels measured with immunohistochemistry in human arteries. Values are expressed as percentage of control and given as mean \pm s.e.m. ^a = significant difference between organ culture and control groups, ^b = significant difference between organ culture and organ culture treated with U0126.

In the presence of the AT₂ receptor antagonist PD12319 there was a diminished contraction after organ culture compared to control arteries, suggesting that the AT₂ receptors are responsible for the upregulated responses induced by organ culture (data not shown).

Protein expression examined with immunohistochemistry

Immunohistochemistry showed a decrease in AT₁ receptor protein in the smooth muscle cells after organ culture ($62 \pm 4\%$) as compared to control ($100 \pm 2\%$). Treatment with the MEK1/2 inhibitor U0126 prevented the down regulation of AT₁ ($82 \pm 7\%$), receptor protein levels in the smooth muscle cell layer as compared to the organ culture (Figure 4B and 4D). The AT₂ receptor protein was increased after organ culture ($119 \pm 2\%$) as compared to control ($100 \pm 2\%$). Incubation with U0126 prevented the increased expression of AT₂ ($111 \pm 3\%$) receptor protein on the smooth muscle cells, however not significantly (Figure 4C-D, Table 2).

5-HT_{1B} receptor

Contractile response

In cultured arteries 5-CT yielded significantly lower contractions than those observed in control arterial segments, presence of U0126 during the organ culture produced a significantly attenuated 5-CT induced response, compared to the cultured arteries (Figure 5A).

Protein expression examined by immunohistochemistry

The 5-HT_{1B} receptor protein was expressed in the smooth muscle cells and this signal was increased in organ culture ($128 \pm 3\%$) as compared to control ($100 \pm 6\%$). Treatment with the MEK1/2 inhibitor U0126 prevented the upregulation of 5-HT_{1B} ($84 \pm 4\%$), receptor protein levels in the smooth muscle cell layer as compared to the organ culture (Figure 5B-C, Table 2).

pERK1/2

Protein expression examined by immunohistochemistry

The pERK1/2 protein was expressed in the smooth muscle cells and this signal was increased in organ culture ($128 \pm 2\%$) as compared to control ($100 \pm 2\%$). Treatment with the MEK1/2 inhibitor U0126 prevented the upregulation of pERK1/2 ($86 \pm 2\%$), protein levels in




Figure 3 A. Concentration response curves elicited by cumulative application of U46619 in human cerebral arteries incubated for 48 h, human cerebral arteries incubated for 48 h with the MEK1/2 inhibitor U0126 (5 μM) and control human arteries. Data are expressed as mean ± s.e.m. **B.** Sections from the human cerebral artery showing TP receptor immunoreactivity in the smooth muscle cell layer. **C.** Expression of TP protein levels in human cerebral arteries incubated for 48 h with or without the MEK1/2 inhibitor U0126 (5 μM) and control human arteries. Data are expressed as percentage of fresh and given as mean ± s.e.m. * $P < 0.05$, ** $P < 0.01$.

the smooth muscle cell layer as compared to the organ culture (Figure 6A-B).

Discussion

This study demonstrates that there is a clear association between human cerebrovascular receptor upregulation via transcription involving activation of the MAPK pathway after organ culture. This was shown by the close association between pERK1/2 activation and the enhanced expression of the contractile receptors at both protein level and at a functional level.

The study design was based on previous work that has shown that organ culture induces upregulation of cerebrovascular receptors in a manner similar as that observed in experimental SAH and MCAO [36,37]. Organ culture is not a model for stroke, however, changes in vasoconstrictor responses after in vitro organ culture show a remarkable similarity to changes observed in animal models of ischemic and hemorrhagic stroke, which makes organ culture a useful in vitro method to study the pharmacological characteristics and underlying molecular and cellular mechanism of cerebrovascular receptor alterations.




Figure 4 **A.** Concentration response curves elicited by cumulative application of Ang II in human cerebral arteries incubated for 48 h, human cerebral arteries incubated for 48 h with the MEK1/2 inhibitor U0126 (5 μM) and control human arteries. Data are expressed as mean ± s.e.m. **B** and **C.** Sections from the human cerebral artery showing AT₁ and AT₂ receptor immunoreactivity in the smooth muscle cell layer. **D.** Expression of AT₁ and AT₂ protein levels in human cerebral arteries incubated for 48 h with or without the MEK1/2 inhibitor U0126 (5 μM) and control human arteries. Data are expressed as percentage of fresh and given as mean ± s.e.m. **P*<0.05, ** *P*<0.01.




Figure 5 A. Concentration response curves elicited by cumulative application of 5-CT in human cerebral arteries incubated for 48 h, human cerebral arteries incubated for 48 h with the MEK1/2 inhibitor U0126 (5 μM) and control human arteries. Data are expressed as mean ± s.e.m. **B.** Sections from the human cerebral artery showing 5-HT_{1B} receptor immunoreactivity in the smooth muscle cell layer. **C.** Expression of 5-HT_{1B} protein levels in human cerebral arteries incubated for 48 h with or without the MEK1/2 inhibitor U0126 (5 μM) and control human arteries. Data are expressed as percentage of fresh and given as mean ± s.e.m. **P*<0.05, ***P*<0.01.

In man cerebral vessels after a stroke contains elevated levels of several cerebrovascular receptor types ET_A, ET_B, AT₁, AT₂ and 5-HT_{1B} [38] and hence is in agreement with the experimental data obtained in animals. The present study revealed several novel and important findings: (i) While the contractile responses to ET-1 usually means activation of the ET_A receptors we found that there is an interaction between the ET_A and ET_B receptors in cerebral arteries following SAH both in experimental studies and in organ culture of human brain vessels [19,35]. The present immunohistochemistry provide experimental evidence that both ET_A and ET_B

receptors are upregulated after organ culture. (ii) The angiotensin II responses were increased. Contrary to animal data the responses were unaltered by specific angiotensin AT₁ blockers [22]. In animal models of stroke an upregulation of the contractile response to Ang II is observed and this contraction is mediated by the AT₁ receptor [22,36]. In our study in the human arteries the results showed that in presence of the AT₂ receptor antagonist PD12319 there was a diminished contraction after organ culture compared to control arteries, suggesting that the AT₂ receptors are responsible for the upregulated responses induced by organ culture. Since




Figure 6 A. Sections from the human cerebral artery showing pERK immunoreactivity in the smooth muscle cell layer. There are increased expressions of the ERK1/2 protein levels in the cultured arteries compared to the control segments. Treatment with U0126 prevented the increased protein expression in the smooth muscle cells. **B.** Expression of pERK1/2 protein levels in human cerebral arteries incubated for 48 h with or without the MEK1/2 inhibitor U0126 (5 μ M) and control human arteries. Data are expressed as percentage of fresh and given as mean \pm s.e.m. * P <0.05, ** P <0.01.

the immunocytochemistry revealed that it was only the AT_2 receptor protein that was elevated in the cerebral artery smooth muscle cells the evidence suggests that there is de novo upregulation of AT_2 receptors producing contraction in the human brain arteries after organ culture. (iii) The 5-HT responses were reduced by organ culture, a finding in concert with what we have seen in tMCAO using two different models as well as after 24 h of organ culture.

Although the decrease in 5-HT_{1B}-mediated contractile responses are less pronounced after organ culture than after experimental stroke [37]. In the subarachnoid hemorrhage model and global ischemia we have demonstrated an upregulation of the 5-HT_{1B} receptor [20,39]. Hence, whereas smooth muscle ET_B receptors are upregulated after all types of cerebral ischemia investigated so far (subarachnoid hemorrhage, global cerebral ischemia, and focal cerebral ischemia) as well as after organ culture, changes in the expression of the 5-HT_{1B} receptor appear to vary with the type of cerebral ischemia. The role of 5-HT and its receptor in ischemia are not clear; while some studies report a protective role for

5-HT receptor agonists, others show increased contractility and improvement with 5-HT receptor antagonist [40,41]. The results from the study of TP receptors revealed an increased response to the agonist but no significant upregulation of receptor protein. In experimental SAH a similar type of reaction appears [24]. The GPCR changes that we have observed in the human cerebral arteries after organ culture show a striking similarity to the changes observed in experimental cerebral ischemia and thus complements the picture of reactions.

Early during organ culture the raf-MEK-ERK pathway is activated and remains activated during the first two days of culture [32]. Other MAPK members such as p38 and JNK may also be activated but this seems to occur later during organ culture. Detailed study of major cerebral arteries and intracerebral microvessels were performed following experimental SAH [25]; the results clearly demonstrated that the MEK/ERK1/2 pathway was activated within minutes and remained activated until end of the 48 h period. On the other hand p38 and JNK reached significance only at 48 h. This is a model of SAH, however, a similar phenomenon was observed after

MCA occlusion for 2 h and then reperfusion for 48 h both in large cerebral arteries and in microvessels within the brain tissue [42]. In the present study we verified that organ culture results in enhanced expression of pERK1/2 in the smooth muscle cells. Co-administration of the specific MEK1/2 inhibitor U0126 abolished this, confirming that the MEK/ERK pathway is important.

In cultured human arteries, the specific blockade of the MAPK MEK1/2 activity abolished the vascular smooth muscle cell receptor upregulation. A number of mechanisms and receptors have been proposed to account for the late cerebral ischemia [43] but no drug exist with good effect. Clazosentan, an endothelin receptor antagonist, was in a recent clinical study shown to result in reduction in vasospasm as seen angiographically but the outcome was not altered [44]. This was taken as evidence that we may consider also other events than just arterial narrowing; early brain injury and cortical spreading depression. We propose that cerebrovascular receptor upregulation may be such a mechanism that could be of importance, the current study reveals a mechanism present in man that can be modified with inhibition of raf-MEK-ERK signaling.

Conclusion

In conclusion, we show that specific inhibition of the MAPK pathway using U0126 significantly attenuates the vasoconstriction mediated by ET, AT and TP receptors in human cerebral arteries and the enhanced expression of their receptors. The results indicate that MAPK inhibition might be a novel target for treatment of cerebrovascular disorders.

Authors' contributions

SA participated in the design of the study, guided the experimental procedures analyzed the data, and wrote the manuscript. SE performed the immunohistochemistry and prepared the images for it and reviewed the manuscript. RW performed the intensity measurements and reviewed the manuscript. EN carried out the organ culture and in vitro pharmacology myograph experiments. ON and HS performed the surgeries and reviewed the manuscript. LE conceived the study and participated in writing the manuscript. All authors read and approved the final manuscript.

Acknowledgement

This work was supported by the Swedish Research Council, the Heart and Lung foundation (Sweden) and the Royal Physiographic Society (Sweden).

Author details

¹Department of Clinical Sciences, Division of Experimental Vascular Research, Lund University, Lund, Sweden. ²Department of Neurosurgery, Lund University Hospital, Lund, Sweden.

Received: 29 June 2012 Accepted: 26 December 2012

Published: 23 January 2013

References

- O'Collins VE, Macleod MR, Donnan GA, Horky LL, van der Worp BH, Howells DW: **1,026 experimental treatments in acute stroke.** *Ann Neurol* 2006, **59**(3):467–477.
- Edvinsson L: **Cerebrovascular endothelin receptor upregulation in cerebral ischemia.** *Curr Vasc Pharmacol* 2009, **7**(1):26–33.
- Viossat I, Duverger D, Chapelat M, Pirotzky E, Chabrier PE, Braquet P: **Elevated tissue endothelin content during focal cerebral ischemia in the rat.** *J Cardiovasc Pharmacol* 1993, **22**(Suppl 8):S306–309.
- Lampl Y, Fleminger G, Gilad R, Galron R, Sarova-Pinhas I, Sokolovsky M: **Endothelin in cerebrospinal fluid and plasma of patients in the early stage of ischemic stroke.** *Stroke* 1997, **28**(10):1951–1955.
- Zervas NT: **Vasospasm: an update.** *Clin Neurosurg* 1979, **26**:643–656.
- Zervas NT, Lavyne MH, Negoro M: **Neurotransmitters and the normal and ischemic cerebral circulation.** *N Engl J Med* 1975, **293**(16):812–816.
- Saavedra JM, Benicky J, Zhou J: **Mechanisms of the Anti-Ischemic Effect of Angiotensin II AT(1) Receptor Antagonists in the Brain.** *Cell Mol Neurobiol* 2006, **26**(7–8):1099–1111.
- Koudstaal PJ, Ciabattini G, van Gijn J, Nieuwenhuis HK, de Groot PG, Sixma JJ, Patrono C: **Increased thromboxane biosynthesis in patients with acute cerebral ischemia.** *Stroke* 1993, **24**(2):219–223.
- Saloheimo P, Juvela S, Riutta A, Pyhtinen J, Hillbom M: **Thromboxane and prostacyclin biosynthesis in patients with acute spontaneous intracerebral hemorrhage.** *Thromb Res* 2005, **115**(5):367–373.
- Furuta H, Guo DF, Inagami T: **Molecular cloning and sequencing of the gene encoding human angiotensin II type 1 receptor.** *Biochem Biophys Res Commun* 1992, **183**(1):8–13.
- Masaki T, Vane JR, Vanhoutte PM: **International Union of Pharmacology nomenclature of endothelin receptors.** *Pharmacol Rev* 1994, **46**(2):137–142.
- Zimmermann M, Seifert V: **Endothelin and subarachnoid hemorrhage: an overview.** *Neurosurgery* 1998, **43**(4):863–875. discussion 875–866.
- Timmermans PB, Chiu AT, Herblin WF, Wong PC, Smith RD: **Angiotensin II receptor subtypes.** *Am J Hypertens* 1992, **5**(6 Pt 1):406–410.
- Chiu AT, Herblin WF, McCall DE, Ardecky RJ, Carini DJ, Duncia JV, Pease LJ, Wong PC, Wexler RR, Johnson AL, et al: **Identification of angiotensin II receptor subtypes.** *Biochem Biophys Res Commun* 1989, **165**(1):196–203.
- Coyle AT, Miggin SM, Kinsella BT: **Characterization of the 5' untranslated region of alpha and beta isoforms of the human thromboxane A2 receptor (TP). Differential promoter utilization by the TP isoforms.** *Eur J Biochem* 2002, **269**(16):4058–4073.
- Beg SA, Hansen-Schwartz JA, Vikman PJ, Xu CB, Edvinsson L: **ERK1/2 inhibition attenuates cerebral blood flow reduction and abolishes ET(B) and 5-HT(1B) receptor upregulation after subarachnoid hemorrhage in rat.** *J Cereb Blood Flow Metab* 2006, **26**(6):846–856.
- Beg SS, Hansen-Schwartz JA, Vikman PJ, Xu CB, Edvinsson L: **Protein kinase C inhibition prevents upregulation of vascular ET(B) and 5-HT(1B) receptors and reverses cerebral blood flow reduction after subarachnoid haemorrhage in rats.** *J Cereb Blood Flow Metab* 2007, **27**(1):21–32.
- Vikman P, Beg S, Khurana T, Hansen-Schwartz J, Edvinsson L: **Gene expression and molecular changes in cerebral arteries following subarachnoid hemorrhage in the rat.** *J Neurosurg* 2006, **105**(3):438–444.
- Hansen-Schwartz J, Hoel NL, Zhou M, Xu CB, Svendgaard NA, Edvinsson L: **Subarachnoid hemorrhage enhances endothelin receptor expression and function in rat cerebral arteries.** *Neurosurgery* 2003, **52**(5):1188–1194. 1194–1185.
- Hansen-Schwartz J, Hoel NL, Xu CB, Svendgaard NA, Edvinsson L: **Subarachnoid hemorrhage-induced upregulation of the 5-HT1B receptor in cerebral arteries in rats.** *J Neurosurg* 2003, **99**(1):115–120.
- Ansar S, Vikman P, Nielsen M, Edvinsson L: **Cerebrovascular ETB, 5-HT1B, and AT1 receptor upregulation correlates with reduction in regional CBF after subarachnoid hemorrhage.** *Am J Physiol Heart Circ Physiol* 2007, **293**(6):H3750–3758.
- Stenman E, Edvinsson L: **Cerebral ischemia enhances vascular angiotensin AT1 receptor-mediated contraction in rats.** *Stroke* 2004, **35**(4):970–974.
- Stenman E, Malmstro M, Uddman E, Gido G, Wieloch T, Edvinsson L: **Cerebral ischemia upregulates vascular endothelin ET(B) receptors in rat.** *Stroke* 2002, **33**(9):2311–2316.
- Ansar S, Larsen C, Maddahi A, Edvinsson L: **Subarachnoid hemorrhage induces enhanced expression of thromboxane A2 receptors in rat cerebral arteries.** *Brain Res* 2010, **1316**:163–172.
- Ansar S, Edvinsson L: **Subtype activation and interaction of protein kinase C and mitogen-activated protein kinase controlling receptor expression in cerebral arteries and microvessels after subarachnoid hemorrhage.** *Stroke* 2008, **39**(1):185–190.
- Namura S, Iihara K, Takami S, Nagata I, Kikuchi H, Matsushita K, Moskowitz MA, Bonventre JV, Alessandrini A: **Intravenous administration of MEK inhibitor U0126 affords brain protection against forebrain ischemia and focal cerebral ischemia.** *Proc Natl Acad Sci USA* 2001, **98**(20):11569–11574.

27. Henriksson M, Stenman E, Vikman P, Edvinsson L: **MEK1/2 inhibition attenuates vascular ETA and ETB receptor alterations after cerebral ischaemia.** *Exp Brain Res* 2007, **178**(4):470–476.
28. Irving EA, Bamford M: **Role of mitogen- and stress-activated kinases in ischemic injury.** *J Cereb Blood Flow Metab* 2002, **22**(6):631–647.
29. Sandhu H, Ansar S, Edvinsson L: **Comparison of MEK/ERK pathway inhibitors on the upregulation of vascular G-protein coupled receptors in rat cerebral arteries.** *Eur J Pharmacol* 2010, **644**(1–3):128–137.
30. Ansar S, Maddahi A, Edvinsson L: **Inhibition of cerebrovascular raf activation attenuates cerebral blood flow and prevents upregulation of contractile receptors after subarachnoid hemorrhage.** *BMC Neurosci* 2011, **12**:107.
31. Adner M, Cantera L, Ehler F, Nilsson L, Edvinsson L: **Plasticity of contractile endothelin-B receptors in human arteries after organ culture.** *Br J Pharmacol* 1996, **119**(6):1159–1166.
32. Henriksson M, Xu CB, Edvinsson L: **Importance of ERK1/2 in upregulation of endothelin type B receptors in cerebral arteries.** *Br J Pharmacol* 2004, **142**(7):1155–1161.
33. Mulvany MJ, Halpern W: **Contractile properties of small arterial resistance vessels in spontaneously hypertensive and normotensive rats.** *Circ Res* 1977, **41**(1):19–26.
34. Hogestatt ED, Andersson KE, Edvinsson L: **Mechanical properties of rat cerebral arteries as studied by a sensitive device for recording of mechanical activity in isolated small blood vessels.** *Acta Physiol Scand* 1983, **117**(1):49–61.
35. Hansen-Schwartz J, Nordstrom CH, Edvinsson L: **Human endothelin subtype A receptor enhancement during tissue culture via de novo transcription.** *Neurosurgery* 2002, **50**(1):127–133. discussion 133–125.
36. Edvinsson LI, Povlsen GK: **Vascular plasticity in cerebrovascular disorders.** *J Cereb Blood Flow Metab* 2011, **31**(7):1554–1571.
37. Povlsen GK, Waldsee R, Ahnstedt H, Kristiansen KA, Johansen FF, Edvinsson L: **In vivo experimental stroke and in vitro organ culture induce similar changes in vasoconstrictor receptors and intracellular calcium handling in rat cerebral arteries.** *Exp Brain Res* 2012, **219**(4):507–520.
38. Vikman P, Edvinsson L: **Gene expression profiling in the human middle cerebral artery after cerebral ischemia.** *Eur J Neurol* 2006, **13**(12):1324–1332.
39. Johansson S, Povlsen GK, Edvinsson L: **Expressional changes in cerebrovascular receptors after experimental transient forebrain ischemia.** *PLoS One* 2012, **7**(7):e41852.
40. Prehn JH, Welsch M, Backhaus C, Nüglisch J, Ausmeier F, Karkouti C, Kriegstein J: **Effects of serotonergic drugs in experimental brain ischemia: evidence for a protective role of serotonin in cerebral ischemia.** *Brain Res* 1993, **630**(1–2):10–20.
41. 3Guilbert F, Laine P, Dubreuil B, McCort G, O'Connor SE, Janiak P, Herbert JM: **Serotonin aggravates exercise-induced cardiac ischemia in the dog: effect of serotonin receptor antagonists.** *Eur J Pharmacol* 2004, **497**(1):55–63.
42. Maddahi A, Edvinsson L: **Enhanced expressions of microvascular smooth muscle receptors after focal cerebral ischemia occur via the MAPK MEK/ERK pathway.** *BMC Neurosci* 2008, **9**:85.
43. Zubkov AY, Nanda A, Zhang JH: **Signal transduction pathways in cerebral vasospasm.** *Pathophysiology* 2003, **9**(2):47–61.
44. Vajkoczy P, Meyer B, Weidauer S, Raabe A, Thome C, Ringel F, Breu V, Schmiedek P: **Clazosentan (AXV-034343), a selective endothelin A receptor antagonist, in the prevention of cerebral vasospasm following severe aneurysmal subarachnoid hemorrhage: results of a randomized, double-blind, placebo-controlled, multicenter phase IIa study.** *J Neurosurg* 2005, **103**(1):9–17.

doi:10.1186/1471-2202-14-12

Cite this article as: Ansar et al.: MAPK signaling pathway regulates cerebrovascular receptor expression in human cerebral arteries. *BMC Neuroscience* 2013 **14**:12.

Submit your next manuscript to BioMed Central and take full advantage of:

- **Convenient online submission**
- **Thorough peer review**
- **No space constraints or color figure charges**
- **Immediate publication on acceptance**
- **Inclusion in PubMed, CAS, Scopus and Google Scholar**
- **Research which is freely available for redistribution**

Submit your manuscript at
www.biomedcentral.com/submit

