



LUND UNIVERSITY

NOTCH1 mutations influence survival in chronic lymphocytic leukemia patients

Willander, Kerstin; Dutta, Ravi Kumar; Ungerback, Jonas; Gunnarsson, Rebeqa; Juliusson, Gunnar; Fredrikson, Mats; Linderholm, Mats; Soderkvist, Peter

Published in:
BMC Cancer

DOI:
[10.1186/1471-2407-13-274](https://doi.org/10.1186/1471-2407-13-274)

2013

[Link to publication](#)

Citation for published version (APA):

Willander, K., Dutta, R. K., Ungerback, J., Gunnarsson, R., Juliusson, G., Fredrikson, M., Linderholm, M., & Soderkvist, P. (2013). NOTCH1 mutations influence survival in chronic lymphocytic leukemia patients. *BMC Cancer*, 13, Article 274. <https://doi.org/10.1186/1471-2407-13-274>

Total number of authors:
8

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

RESEARCH ARTICLE

Open Access

NOTCH1 mutations influence survival in chronic lymphocytic leukemia patients

Kerstin Willander^{1*}, Ravi Kumar Dutta², Jonas Ungerback², Rebeqa Gunnarsson³, Gunnar Juliusson³, Mats Fredrikson², Mats Linderholm⁴ and Peter Söderkvist²

Abstract

Background: *NOTCH1* PEST domain mutations in chronic lymphocytic leukemia have recently been shown to be of prognostic relevance. Both *NOTCH1* and *NOTCH2* are constitutively activated in B-cell CLL but not expressed in normal B cells and may be involved in survival and resistance to apoptosis in CLL. We screened for mutations in different parts of both *NOTCH1* and *NOTCH2* genes and related the changes to survival and other known risk factors.

Methods: In a cohort of 209 CLL patients, we used single strand conformation analysis to determine which of the samples carrying the *NOTCH* mutations and direct dideoxy sequencing was used to determine the exact nucleotide changes. Kaplan-Meier curves and log rank test were used to determine overall survival for *NOTCH1* mutated cases and Cox regression analysis was used to calculate hazardous ratios.

Results: In the present study, we found *NOTCH1* PEST domain mutations in 6.7% of the cases. A shorter overall survival was found in patients with *NOTCH1* mutations compared to wildtype ($p = 0.049$). Further, we also examined the extracellular and the heterodimerisation domains of the *NOTCH1* gene and the PEST domain and heterodimerisation domain of the *NOTCH2* gene, but no mutations were found in these regions. *NOTCH1* mutations were most commonly observed in patients with unmutated IGHV gene (10/14), and associated with a more aggressive disease course. In addition, *NOTCH1* mutations were almost mutually exclusive with *TP53* mutations. In the combined group of *NOTCH1* (6.7%) or *TP53* (6.2%) mutations, a significant difference in overall survival compared to the wildtype *NOTCH1* and *TP53* was found ($p = 0.002$).

Conclusions: Both *NOTCH1* and *TP53* mutations seem to be independent predictive markers for worse outcome in CLL-patients and this study emphasizes the contention that *NOTCH1* mutations is a novel risk marker.

Keywords: Chronic lymphocytic leukemia, *NOTCH1* mutations, *TP53* mutations, Prognostic markers

Background

Chronic lymphocytic leukemia (CLL) is a heterogeneous disease with variable clinical course characterized by a monoclonal progressive accumulation of mature CD5+ B-lymphocytes avoiding apoptosis. Some patients with an indolent disease need no or little treatment while others have a more adverse disease at diagnosis. No common genetic lesion, which causes the disease, has been found [1], but recurrent mutations in CLL involve *TP53* and *ATM*, and novel mutations in the *NOTCH1*,

SF3B1, *MYD88*, *BIRC3* and *FBXW7* genes have been identified through next generation sequencing [2]. The CLL cases may be divided in two major groups regarding to mutated (M) or unmutated (UM) immunoglobulin heavy chain variable region gene (IGHV) where patients with an unmutated IGHV clone have a more adverse prognosis than patients with mutated IGHV gene [3,4]. By the means of FISH analysis, different chromosomal aberrations as deletion in 11q, 13q, 17p or trisomy 12 are found in about 80% of tumor cells in the CLL-patients [5].

Recently, *NOTCH1* mutations were found to be predictor of poor prognosis in CLL [6-11]. Furthermore a study of Rosati et al. [12] showed that *NOTCH1* and

* Correspondence: kerstin.willander@liu.se

¹Department of Clinical and Experimental Medicine, Department of Hematology, County Council of Östergötland, Linköping University, SE-58185 Linköping, Sweden

Full list of author information is available at the end of the article

Table 1 Primer sequences

Gene-exon-segment	Application	Forward primer	Reverse primer	Product size/bp	T _A /°C
Notch1- 26-a	PCR/SSCA	AGCCCCCTGTACGACCAGTA	CTTGCGCAGCTCCTCCTC	283	63.5
Notch1- 26-b	PCR/SSCA	ACACGGCCAGCAGATGAT	GAGAGTTGCGGGGATTGAC	231	57.1
Notch1- 27	PCR/SSCA	GTGGCGTCATGGGCCTCA	TAGCAACTGGCACAAACAGC	342	63.2
Notch1- 34a	PCR/SSCA	AACCACCTGCCTGGGATG	CGCATTGACCAATTCAACTG	232	57.1
Notch1- 34b	PCR/SSCA	GGGCCCTGAATTCCTGT	AGGCCCTGGTAGCTCATCAT	229	60
Notch1- 34c	PCR/SSCA	GCTGCACAGTAGCCTTGCT	CTGAGCTCACGCCAAGGT	224	58
Notch1- 34d	PCR/SSCA	ACATCCAGCAGCAGCAAAG	GTGGGACCAGCGAGGATG	222	58
Notch1- 34e	PCR/SSCA	CACTATTCTGCCCCAGGAGA	CAGTCGGAGACGTTGGAATG	234	58
Notch1- 34f	PCR/SSCA	ACAGCTACTCTCGCTGTG	AAGGCTTGGGAAAGGAAGC	248	58
Notch1-6	PCR/SSCA	GCAGCTGCCCGGGGCCGACA	TCAGGCTGGCCCATGTGA	330	62
Notch1-7	PCR/SSCA	ATGCCTGGCCAGGGGCCGT	TCGACTTCTCATCGTTCT	273	58
Notch1-8	PCR/SSCA	CCGATGGGGTGGTGTGCAGT	TGCCCAGCCTCGACTCGGTT	331	63
Notch1-11	PCR/SSCA	AGTCCTAAGTCTTCTGTGCC	AGGCCCGCCCTGCCCACT	325	65
Notch1-12	PCR/SSCA	AGGACTGACCGACACGTG	TCTGAGCACAGTGCAGTCA	183	53
Notch1-13	PCR/SSCA	TGGGCGCTGGGCCTCGGA	ACTGATGTGTCCCATGA	268	54
Gene-exon-segment	Application	Forward primer	Reverse primer	Product size/bp	T _A /°C
Notch2- 26-1	PCR/SSCA	TTCTCTGCTTCCCTTACCT	TTAATGCGCAGGTTGGTGT	250	54.1
Notch2- 26-2	PCR/SSCA	TGGTATTGATGCCACCTGAA	GCCTTGAAGTTCAGAAACCAA	240	54.1
Notch2- 27	PCR/SSCA	TACCCCATCTCTCCTCTC	AATTGTTCCCAATTGACA	250	55.2
Notch2- 34-1	PCR/SSCA	TCCCCTGTTGATTCCCTAGA	CACAATGTGGTGGTGGGATA	249	55.2
Notch2- 34-2	PCR/SSCA	GCACTGTGCTTCCCTCAGT	CTGCCCTTAGGGATGAGCTG	298	55.2
Notch2- 34-3	PCR/SSCA	ACCCATCCTGGCATAGCTC	TAGGCTGGGAGAATGGTCTG	287	55.2
Notch2- 34-4	PCR/SSCA	TTTGCCAGTGTGGCTTT	GGTGATGAACTTGACCACTG	249	57.1
Notch2- 34-5	PCR/SSCA	ACACCCAGTCACAGTGGTCA	TGTCTCTACTGGAGGTGGAC	242	61.6

NOTCH2, together with their ligands Jagged1 –and 2 are constitutively activated in B-CLL cells but not in normal B cells, suggesting that NOTCH signaling is involved in survival and resistance to apoptosis in CLL.

The NOTCH receptor is a membrane bound protein that consists of an extracellular, transmembrane and intracellular domain that can be released upon ligand interaction and transactivate target genes. The NOTCH signal pathway is activated by a ligand on a neighboring cell and plays an essential role in controlling proliferation, differentiation and survival. Following the receptor-ligand binding, the NOTCH receptor first undergoes a S2 proteolytic cleavage by ADAM proteinase in the extracellular domain, which then is followed by a S3 cleavage by a γ -secretase complex in the transmembrane domain releasing the intracellular NOTCH domain that translocates to the nucleus where it interacts with a transcription complex and acts as a transcriptional activator for multiple target genes [13]. The C-terminal part of the intracellular domain consists of a PEST region that is important for proteasomal degradation of the NOTCH receptor by binding to FBXW7, an E3 ubiquitin ligase, to limit duration of the NOTCH activity. A CT deletion in the C-terminal region results in removal of



the PEST domain, a truncated NOTCH protein, and impaired NOTCH degradation and constitutive transcriptional activation of NOTCH target genes in CLL [7,14].

In the present study we have screened for mutations in different parts of both the *NOTCH1* and *NOTCH2* gene in a cohort of 209 CLL-patients. There is a high structural similarity between *NOTCH1* and *NOTCH2* genes and recent *NOTCH2* gain-of-function mutations are found in B-cell lymphomas [15]. Further, as NOTCH2 is involved in overexpression of CD23, one of the hallmarks of CLL [16], it prompted us to screen both the *NOTCH1* and *NOTCH2* genes for genetic alterations. Mutations were only found in the PEST region in the *NOTCH1* gene in our cohort and emerged as an independent factor of poor overall survival and disease stage, in addition to *TP53* mutations and IGHV gene status.

Methods

Patients

In this study, peripheral blood from 209 CLL patients (145 men and 64 women) was collected between 1996 and 2006 at the Department of Hematology, Linköping University Hospital. Mononuclear cells were isolated by



Ficoll-Paque gradient centrifugation and genomic DNA was extracted by proteinase K digestion and stored frozen until used as earlier described [17]. The samples were collected either at the time of diagnosis or prior to the first treatment. For all patients, follow-up data were available, and for 106 live patients the median follow-up time was 6.8 years (range 1.6-14.9 years). The median age at diagnosis was 62.5 years (range 38.3-87.0 years). The immunophenotype and the Binet staging system were according to the IWCLL guidelines [18]. The

Table 2 Clinical and biological characteristics of the 209 CLL-patients

	NOTCH1						P value
	Total		Wildtype		Mutated		
	n	%	n	%	n	%	
Number of patients	209		195		14		
Sex							0.67
male	145	69	136	70	9	64	
female	64	31	59	30	5	36	
Binet stage							0.11
A	101	48	98	50	3	21	
B/C	91	44	82	42	9	64	
Not determined	17	8	15	8	2	14	
Treatment							0.22
yes	182	87	168	86	14	100	
no	20	10	20	10	0	0	
Not determined	7	3	7	4	0	0	
IGHV status							0.22
Mutated	64	31	61	31	3	21	
Unmutated	123	59	113	58	10	71	
Not determined	22	10	21	11	1	7	
TP 53 status							0.88
wild-type	196	94	183	94	13	93	
mutated	13	6	12	6	1	7	

immunoglobulin heavy chain variable region genes (IGHV) and *TP53* gene status were analysed and reported in an earlier study [17]. Informed consent was obtained from the patients and the study was approved by the regional ethical committee (Dnr 02-459) in Linköping and conducted in accordance with the ethical guidelines of the Helsinki Declaration.

NOTCH mutation status detection

The *NOTCH1* mutations status were analyzed for the extracellular region (exon 6, 7, 8, 11, 12 and 13), the heterodimerisation domain (exon 26, 27) and the PEST region (exon 34) and the *NOTCH2* was only analyzed for mutations in the heterodimerisation and the PEST domains by PCR amplification followed by single strand conformation analysis (SSCA) according to the original protocol [19] and direct dideoxy sequencing to determine the exact nucleotide change and compared to corresponding *NOTCH1* and *NOTCH2* germline sequence (NM_017617.3 and NM_024408.3 respectively). Primer sequences are shown in Table 1.

TP53 and IGHV gene status detection

TP53 gene mutation analysis was performed for exons 5-8 (the DNA binding domains) by the PCR-single strand conformation analysis (SSCA) technique and samples displaying mobility shifts were sequenced with the dideoxy termination method to confirm the nucleotide changes.

The IGHV gene mutational status was performed by PCR amplification on genomic DNA by using specific VH/JH primers [20], followed by DNA sequencing of both forward and reverse strands. To determine the IGHV gene identity the sequences were aligned by using the IMGT/V-QUEST database (<http://imgt.org>), $\geq 98\%$ identity to the corresponding germline sequence was considered as an unmutated IGHV gene.

Table 3 Analysis for overall survival

	N	HR	95% CI	P	HR ¹	95%CI	P
Wildtype <i>NOTCH1</i> and <i>TP53</i>	183	1			1		
Mutated <i>NOTCH1</i> or <i>TP53</i>	26	2.27	1.32-3.91	0.003	2.16	1.25-3.72	0.006
Wildtype <i>NOTCH1</i>	195	1			1		
Mutated <i>NOTCH1</i>	14	2.04	0.98-4.25	0.056	1.80	0.86-3.76	0.12
Wildtype <i>TP53</i>	196	1			1		
Mutated <i>TP53</i>	13	2.54	1.17-5.54	0.019	2.54	1.17-5.53	0.002

HR, hazard ratio; CI, confidence interval.

¹Adjusted for age and sex.

Statistical analysis

Kaplan-Meier curves were used to show the overall survival and the log-rank test was used to compare the survival between the groups. To calculate hazard ratios (HR) the Cox proportional hazard model (Cox-regression) was used. For all statistical analyses Stata v12.1 was used (StataCorp LP, College Station, TX, USA). P-values less than 0.05 were considered significant. Overall survival was measured from date of diagnosis until the last follow-up or death.

Results and discussion


NOTCH1 heterozygous mutations in the PEST domain occurred in a frequency of 14 out of 209 patients (6.7%) in our study. Thirteen of the mutations correspond to a 2-bp frameshift deletion, c.7541_7542delCT and one is a novel GT deletion at c.6988_6989delGT, both generating frameshift mutations, with subsequent stopcodon and truncated proteins eliminating the PEST domain. The Kaplan-Meier curve for the CLL-patients with *NOTCH1* mutations revealed a shorter overall survival (OS) compared to *NOTCH1* wildtype patients ($p = 0.049$) (Figure 1). Clinical and biological characteristics of the CLL patients in relation to the *NOTCH1* status are summarized in Table 2. Our study showed a similarity to other CLL studies, which have reported *NOTCH1* PEST domain mutations in 4.7% - 12.2% [6-11].

T cell acute lymphoblastic leukemia (T-ALL) display high frequency of activating *NOTCH1* mutations including the extracellular heterodimerisation domain, in addition to elimination C-terminal PEST region mutations

[21]. Studies in head and neck cancer have also identified mutations in the extracellular epidermal growth factor repeat domain in the *NOTCH1* gene [22]. These studies prompted us to screen for mutations in those parts of the *NOTCH1* gene, but no mutations could be detected. *NOTCH2* has a role in marginal zone B cell fate decision, similar to the critical role for *NOTCH1* in determining the T-cell fate [13]. *NOTCH2* mutations are found in the C-terminal part close to the PEST region in splenic marginal zone lymphoma [23,24], but we did not find any mutations in the PEST region or heterodimerisation domain of the *NOTCH2* gene in our cohort.

Mutations in the *TP53* gene are associated with poor prognosis in CLL [25]. Mutations in the *NOTCH1* gene are almost mutually exclusive with mutations in the *TP53* gene with or without 17p or 11q deletions, only one sample harbored mutations in both *NOTCH1* and *TP53*. Combined, *NOTCH1* (6.7%) and *TP53* (6.2%) mutations represent 12.9% of the patients in this cohort and indicated a significant worse survival as compared to wildtype *NOTCH1* and *TP53* (Log rank analysis, $p = 0.002$) (Figure 2). *NOTCH1* mutations may appear together with trisomy 12 [26,27], and in our cohort, trisomy 12 was found in 15/152 patients by BAC (bacterial artificial chromosome) microarray analysis and two of these carried *NOTCH1* mutation; this association was not significant ($p = 0.56$). By univariate analysis, the HR for death increased to 2.27 (1.32-3.91; 95% confidence interval) for tumors mutated in *NOTCH1* or *TP53* compared to *NOTCH1* and *TP53* wildtype tumors ($p = 0.003$) (Table 3). At the molecular level there seems to be an intriguing and complex link between p53 and *NOTCH1*. P53 induce *NOTCH1* expression and seems to initiate an anti-apoptotic feedback mechanism with subsequent increased cell survival that may limit p53 promoting therapy with e.g. nutlins [28,29]. *NOTCH* signaling blockade by γ -secretase inhibitors to stimulate apoptosis may be considered to be of therapeutic value at least for wt p53 CLL patients [28].

Among CLL patients with a mutated *NOTCH1* gene 10/14 (71%) had an unmutated IGHV gene in contrast to 113/195 (58%) with a wildtype *NOTCH1* gene, a difference that did not reach statistical significance ($p = 0.22$)



(Figure 3 and Table 2). CLL with *NOTCH1* mutations seemed to be more progressive, with a high frequency of unmutated IGHV gene and advanced Binet stages, indicating a more aggressive disease course.

Five patients in this cohort had *NOTCH1* mutation at the time of diagnosis. For these patients the median time to first treatment was 101 days (range 21 to 145 days). For the whole group the median time from diagnosis to the first treatment was 438 days (range 0–6021 days). Further and expected, all patients with *NOTCH1* mutations identified at diagnosis had the more advanced Binet stages B and C, a tendency that due to few observations did not reach significance ($p = 0.11$) (Table 2). The frequency of *NOTCH1* mutations is also reported to be significantly higher in Richter syndrome, i.e., a progression of CLL into diffuse large lymphoma with often dismal outcome [8,30], however our cohort contained no information on the prevalence of Richter syndrome.

It is now recommended to perform *TP53* mutation analysis in patients with CLL as *TP53* mutations occur in about 5% of cases in absence of 17p deletion and represent an independent prognostic factor associated with worse outcome [31]. CLL patients with 17p deletion and/or *TP53* mutations are strongly associated with refractory disease, and also activated *NOTCH1* mutations were recently suggested to cause refractoriness to fludarabine [8,9,32].

Conclusions

Our study confirms other recent reports that *NOTCH1* mutation eliminating the PEST domain, has a prognostic value as a novel risk marker in CLL similar to *TP53* mutations. Thus both *NOTCH1* and *TP53* mutation may be an indication for earlier and more active treatment or as an indicator for transplantation therapy.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

KW collected data, performed experiments, analyzed and interpreted data, wrote the manuscript; RKD performed experiments, analyzed and interpreted data; JU performed experiments, analyzed and interpreted data; RG performed experiments and interpreted data; GJ interpreted data; MF performed statistical analysis; ML designed experiments; PS designed experiments, wrote the manuscript. All authors were involved in writing the manuscript. All authors read and approved the final manuscript.

Acknowledgements

This study is supported by FORSS grant #235261.

Author details

¹Department of Clinical and Experimental Medicine, Department of Hematology, County Council of Östergötland, Linköping University, SE-58185 Linköping, Sweden. ²Department of Clinical and Experimental Medicine, Linköping University, Linköping, Sweden. ³Department of Laboratory Medicine, Stem Cell Center, Hematology and Transplantation, Lund University, Lund, Sweden. ⁴Department of palliative care, Stockholms Sjukhem, Stockholm, Sweden.

Received: 21 December 2012 Accepted: 29 May 2013

Published: 4 June 2013

References

- Chiorazzi N, Rai KR, Ferranini M: Chronic lymphocytic leukemia. *N Engl J Med* 2005, **352**:805–815.
- Wang L, Lawrence MS, Wan Y, Stojanov P, Sougnez C, Stevenson K, Werner L, Sivachenko A, DeLuca DS, Zhang L, Zhang W, Zhang W, Vartanov AR, Fernandes SM, Goldstein NR, Folco EG, Cibulskis K, Tesar B, Sievers QL, Shefler E, Gabriel S, Hacohen N, Reed R, Meyerson M, Golub TR, Lander ES, Neuberger D, Brown JR, Getz G, Wu CJ: **SF3B1 and other novel cancer genes in chronic lymphocytic leukemia.** *N Engl J Med* 2011, **365**:2497–2506.
- Damle RN, Wasil T, Fais F, Ghiotto F, Valetto A, Allen SL, Buchbinder A, Budman D, Dittmar K, Kolitz J, Lichtman SM, Schulman P, Vinciguerra VP, Rai KR, Ferranini M, Chiorazzi N: **IgV gene mutation status and CD 38 expression as novel prognostic indicators in chronic lymphocytic leukemia.** *Blood* 1999, **94**:1840–1847.
- Hamblin TJ, Davis Z, Gardiner A, Oscier DG, Stevenson FK: **Unmutated IgV (H) genes are associated with a more aggressive form of chronic lymphocytic leukemia.** *Blood* 1999, **94**:1848–1854.
- Döhner H, Stielgenbauer S, Benner A, Leupolt E, Kröber A, Bullinger L, Döhner K, Bentz M, Lichter P: **Genomic aberrations and survival in chronic lymphocytic leukemia.** *N Engl J Med* 2000, **343**:1910–1916.
- Sportoletti P, Baldoni S, Cavalli L, Del Papa B, Bonifacio E, Ciunelli R, Bell AS, Di Tommaso A, Rosati E, Crescenzi B, Mecucci C, Screpanti I, Marconi P, Martelli MF, Di Ianni M, Falzetti F: **Notch1 PEST domain mutation is an adverse prognostic factor in B-CLL.** *British Journal of Haematology* 2010, **151**:404–406.
- Puente XS, Pinyol M, Quesada V, Conde L, Ordóñez GR, Villamor N, Escaramis G, Jares P, Beà S, González-Díaz M, Bassaganyas L, Baumann T, Juan M, López-Guerra M, Colomer D, Tubío JM, López C, Navarro A, Tornador C, Aymerich M, Rozman M, Hernández JM, Puente DA, Freije JM, Velasco G, Gutiérrez-Fernández A, Costa D, Carrió A, Gujjarro S, Enjuanes A, et al: **Whole genome sequencing identifies recurrent mutations in chronic lymphocytic leukemia.** *Nature* 2011, **475**:101–105.
- Fabbri G, Rasi S, Rossi D, Trifonov V, Khiabanian H, Ma J, Grunn A, Fangazio M, Capello D, Monti S, Cresta S, Gargiulo E, Forconi F, Guarini A, Arcaini L, Paulli M, Laurenti L, Larocca LM, Marasca R, Gattei V, Oscier D, Berton F, Mullighan CG, Foà R, Pasqualucci L, Rabadan R, Dalla-Favera R, Gaidano G: **Analysis of the chronic lymphocytic leukemia coding genome: role of NOTCH1 mutational activation.** *J Exp Med* 2011, **208**:1389–1401.
- Rossi D, Rasi S, Fabbri G, Spina V, Fangazio M, Forconi F, Marasca R, Laurenti L, Bruscaggini A, Cerri M, Monti S, Cresta S, Famà R, De Paoli L, Bulian P, Gattei V, Guarini A, Deaglio S, Capello D, Rabadan R, Pasqualucci L, Dalla-Favera R, Foà R, Gaidano G: **Mutations of Notch1 are an independent predictor of survival in chronic lymphocytic leukemia.** *Blood* 2012, **119**:521–529.
- Shedden K, Li Y, Quillet P, Malek SN: **Characteristics of chronic lymphocytic leukemia with somatically acquired mutations in exon 34.** *Leukemia* 2011, **26**:1108–1110.
- Mansouri L, Cahill N, Gunnarsson R, Smedby KE, Tjønnefjord E, Hjalgrim H, Juliusson G, Geisler C, Rosenquist R: **NOTCH1 and SF3B1 mutations can be added to the hierarchical prognostic classification in chronic lymphocytic leukemia.** *Leukemia* 2013, **27**:512–514.
- Rosati E, Sabatini R, Rampino G, Tabilio A, Di Ianni M, Fettucciari K, Bartoli A, Coaccioli S, Screpanti I, Marconi P: **Constitutively activated Notch signaling is involved in survival and apoptosis resistance of B-CLL cells.** *Blood* 2009, **113**:856–865.
- Radtke F, Wilson A, Mancini SJC, Robson MacDonald H: **Notch regulation of lymphocyte development and function.** *Nature Immunology* 2004, **5**:247–253.
- Palomero T, Lim WK, Odom DT, Sulis ML, Real PJ, Margolin A, Barnes KC, O'Neil J, Neuberger D, Weng AP, Aster JC, Sigaux F, Soulier J, Look AT, Young RA, Califano A, Ferrando AA: **Notch1 directly regulates c-MYC and activates a feed-forward-loop transcriptional network promoting leukemic cell growth.** *PNAS* 2006, **103**:18261–18266.
- Lee SY, Kumano K, Nakazaki K, Sanada M, Matsumoto A, Yamamoto G, Nannya Y, Suzuki R, Ota S, Ota Y, Izutsu K, Sakata-Yanagimoto M, Hangaishi A, Yagita H, Fukayama M, Seto M, Kurokawa M, Ogawa S, Chiba S: **Gain-of-function mutations and copy number increases of Notch2 diffuse large B-cell lymphoma.** *Cancer Science* 2009, **100**:920–926.

16. Hubmann R, Schwarzmeier JD, Sheta M, Hilgarth M, Duechler M, Dettke M, Berger R: **Notch2 is involved in overexpression of CD23 in B-cell chronic lymphocytic leukemia.** *Blood* 2002, **99**:3742–3747.
17. Willander K, Ungerback J, Karlsson K, Fredrikson M, Söderkvist P, Linderholm M: **MDM2 SNP309 promoter polymorphism, an independent prognostic factor in chronic lymphocytic leukemia.** *European Journal of Haematology* 2010, **85**:251–256.
18. Hallek M, Cheson BD, Catovsky D, Caligaris-Cappio F, Dighiero G, Döhner H, Hillmen P, Keating MJ, Montserrat E, Rai KR, Kipps TJ: **Guidelines for diagnosis and treatment of chronic lymphocytic leukemia: a report from an international workshop on chronic lymphocytic leukemia updating the national cancer institute working group 1996 guidelines.** *Blood* 2008, **111**:5446–5456.
19. Orita M, Suzuki Y, Sekiya T, Hayashi K: **Rapid and sensitive detection of point mutations and DNA polymorphism using the polymerase chain reaction.** *Genomics* 1989, **5**:874–879.
20. van Dongen JJ, Langerak AW, Brüggemann M, Evans PA, Hummel M, Lavender FL, Delabesse E, Davi F, Schuurink E, García-Sanz R, van Krieken JH, Droese J, González D, Bastard C, White HE, Spaargaren M, González M, Parreira A, Smith JL, Morgan GJ, Kneba M, Macintyre EA: **Design and standardization of PCR primers and protocols for detection of clonal immunoglobulin and T-cell receptor gene recombinations in suspect lymphoproliferations: report of BIOMED-2 Concerted Action BMH4-CT98-3936.** *Leukemia* 2003, **17**:2257–2317.
21. Weng AP, Ferrando AA, Lee W, Morris JP 4th, Silverman LB, Sanchez-Irizarry C, Blacklow SC, Look AT, Aster JC: **Activating mutations of NOTCH1 in human T cell acute lymphoblastic leukemia.** *Science* 2004, **306**:269–271.
22. Stransky N, Egloff AM, Tward AD, Kostic AD, Cibulskis K, Sivachenko A, Kryukov GV, Lawrence MS, Sougnez C, McKenna A, Shefler E, Ramos AH, Stojanov P, Carter SL, Voet D, Cortés ML, Auclair D, Berger MF, Saksena G, Guiducci C, Onofrio RC, Parkin M, Romkes M, Weissfeld JL, Seethala RR, Wang L, Rangel-Escareño C, Fernandez-Lopez JC, Hidalgo-Miranda A, Melendez-Zajgla J, et al: **The mutational landscape of head and neck squamous cell carcinoma.** *Science* 2011, **333**:1157–1160.
23. Rossi D, Trifonov V, Fangazio M, Bruscaggin A, Rasi S, Spina V, Monti S, Vaisitti T, Arruga F, Famà R, Ciardullo C, Greco M, Cresta S, Piranda D, Holmes A, Fabbri G, Messina M, Rinaldi A, Wang J, Agostinelli C, Piccaluga PP, Lucioni M, Tabbò F, Serra R, Franceschetti S, Deambrogi C, Daniele G, Gattei V, Marasca R, et al: **The coding genome of splenic marginal zone lymphoma: activation of NOTCH2 and other pathways regulating marginal zone development.** *J Exp Med* 2012, **209**:1537–1551.
24. Kiel MJ, Velusamy T, Betz BL, Zhao L, Weigelin HG, Chiang MY, Huebner-Chan DR, Bailey NG, Yang DT, Bhagat G, Miranda RN, Bahler DW, Medeiros LJ, Lim MS, Elenitoba-Johnson KS: **Whole-genome sequencing identifies recurrent somatic NOTCH2 mutations in splenic marginal zone lymphoma.** *J Exp Med* 2012, **209**:1553–1565.
25. Zenz T, Kröber A, Scherer K, Häbe S, Bühler A, Benner A, Denzel T, Winkler D, Edelmann J, Schwänen C, Döhner H, Stilgenbauer S: **Monoallelic TP53 inactivation is associated with poor prognosis in chronic lymphocytic leukemia: results from a detailed genetic characterization with long-term follow-up.** *Blood* 2008, **112**:3322–3329.
26. Baletti V, Bottoni A, Palamarchuk A, Alder H, Rassenti LZ, Kipps TJ: **Notch1 mutations in CLL associated with trisomy 12.** *Blood* 2012, **119**:329–331.
27. Del Giudice I, Rossi D, Chiaretti S, Marinelli M, Tavaloro S, Gabrielli S, Lauretti L, Marasca R, Rasi S, Fangazio M, Guarini A, Gaidano G, Foà R: **Notch1 mutations in +12 chronic lymphocytic leukemia (CLL) confer and unfavorable prognosis, induce a distinctive transcriptional profiling and refine the intermediate prognosis of +12 CLL.** *Haematologica* 2012, **97**:437–441.
28. Secchiero P, Melloni E, di Iasio MG, Tiribelli M, Rimondi E, Corallini F, Gattei V, Zauli G: **Nutlin-3 up-regulates the expression of Notch1 in both myeloid and lymphoid leukemic cells, as part of negative feedback antiapoptotic mechanism.** *Blood* 2009, **113**:4300–4308.
29. Paolo Dotto G: **Crosstalk of Notch with p53 and p63 in cancer growth control.** *Nature Rev. Cancer* 2009, **9**:587–595.
30. Rossi D, Rasi S, Spina V, Fangazio M, Monti S, Greco M, Ciardullo C, Famà R, Cresta S, Bruscaggin A, Lauretti L, Martini M, Musto P, Forconi F, Marasca R, Larocca LM, Foà R, Gaidano G: **Different impact of NOTCH1 and SF3B1 mutations on the risk of chronic lymphocytic leukemia transformation to Richter syndrome.** *British Journal of Haematology* 2012, **158**:426–229.
31. Pospisilova S, Gonzalez D, Malcikova J, Trbusek M, Rossi D, Kater AP, Cymbalista F, Eichhorst B, Hallek M, Döhner H, Hillmen P, van Oers M, Gribben J, Ghia P, Montserrat E, Stilgenbauer S, Zenz T: **ERIC recommendations on TP53 mutation analysis in chronic lymphocytic leukemia.** *Leukemia* 2012, **26**:1458–1461.
32. Zenz T, Gribben JG, Hallek M, Döhner H, Keating MJ, Stilgenbauer S: **Risk categories and refractory CLL in era of chemoimmunotherapy.** *Blood* 2012, **119**:4101–4107.

doi:10.1186/1471-2407-13-274

Cite this article as: Willander et al.: **NOTCH1 mutations influence survival in chronic lymphocytic leukemia patients.** *BMC Cancer* 2013 **13**:274.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

