



# LUND UNIVERSITY

## Neurogenesis from Neural Stem Cells, Ependymal Cells and Fibroblasts

Devaraju, Karthikeyan

2014

[Link to publication](#)

*Citation for published version (APA):*

Devaraju, K. (2014). *Neurogenesis from Neural Stem Cells, Ependymal Cells and Fibroblasts*. [Doctoral Thesis (compilation), Neurology, Lund]. Neurology, Lund.

*Total number of authors:*

1

### General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00

*Academic Dissertation*

# **Neurogenesis from Neural Stem Cells, Ependymal Cells and Fibroblasts**

**Karthikeyan Devaraju**

2014

With the approval of the Faculty of Medicine, Lund University, Sweden,  
this thesis will be defended on  
March 21, 2014 at 09.00 in  
Segerfalksalen, Wallenberg Neurocentrum,  
Lund, Sweden.

Faculty opponent

**Urban Lendahl**

Professor, Vice-Chairman  
Department of Cell and Molecular Biology  
Karolinska institute, Stockholm, Sweden



**LUND**  
UNIVERSITY

Organization LUND UNIVERSITY Department of Clinical Sciences Division of Neurology, Lund University Lund, Sweden	Document name DOCTORAL DISSERTATION	
	Date of issue March 21, 2014	
	Sponsoring organization	
Author(s) Karthikeyan Devaraju		
Title and subtitle Neurogenesis from neural stem cells, ependymal cells and fibroblasts		
Abstract <p>Stroke is a major cause of death and disability around the world. Stroke leads to loss of neurons and also other cells in the brain due to lack of blood supply. Currently no therapies are available to treat stroke-related disability. It has been shown that stroke leads to increased neurogenesis, birth of new neurons, within the brain. This increased neurogenesis is not sufficient to restore lost function. There is a need to develop therapies for neuronal replacement by improving neurogenesis within the brain and / or transplanting neurons. Cortical strokes lead to more disability after stroke as compared to those affecting the striatum, and whether cortical neurogenesis occurs after stroke is controversial. Cell transplantation may be the key to cortical repair after stroke.</p> <p>Reports have identified positive but very few negative regulators of neurogenesis after stroke, and suggested that ependymal cells can also contribute to stroke-induced neurogenesis. Transplantation of neurons generated from different sources such as fetal brain, embryonic stem cells and induced pluripotent stem cells are associated with ethical issues and carry the risk of immune rejection and tumorigenicity. Direct conversion of patient's own skin cells to neurons could overcome these problems and potentially restore function after transplantation in stroke-damaged brain.</p> <p>In this thesis we have used transgenic models, viral vectors, electroporation-mediated gene delivery and overexpression of transcription factors to demonstrate neurogenesis from neural stem cells, ependymal cells in the lateral ventricular wall and fibroblasts.</p> <p>We show that Lnk, a known inhibitor of hematopoietic stem cell self-renewal, is also expressed in the brain. Overexpression or removal of Lnk expression leads to decreased or increased neurogenesis in vitro respectively. When brain is damaged by stroke there is increased proliferation of neural stem cells in animals without Lnk expression. This was not observed in status epilepticus, a severe form of epilepsy. We determined that upregulation of STAT1/3 after stroke leads to increased Lnk expression. Subsequently Lnk inhibits cellular response to increased IGF1 stimulation after stroke, by decreasing Akt phosphorylation. We have identified LNK signaling as a novel mechanism of influencing neurogenic response to stroke.</p> <p>We next determined if ependymal cells in lateral ventricular wall of adult rat brain contribute to neurogenesis after stroke. We identified FoxJ1 as a marker of ependymal cells in rats similar to mice, and used FoxJ1 promoter in piggyBac system to genetically label these cells with fluorescent reporter proteins GFP or RFP by electroporation. Tracing the lineage of the labeled cells in intact and stroke-damaged brain, we identified that FoxJ1 expressing cells contribute to olfactory bulb neurogenesis while the striatal neurogenic response was not significant. Thus, FoxJ1 expressing cells probably have only a minor role in repair after stroke.</p> <p>We then tested whether human fetal lung fibroblasts could be directly converted to cortical neurons. We overexpressed sets of transcription factors that are known to be involved in cortical neuron development. We found that overexpression of different sets of these factors in fibroblasts converted them to cortical-like neurons. These neurons expressed markers of cortical neurons and were functional by electrophysiology.</p> <p>In summary, these results raise the possibility that inhibition of Lnk, a negative regulator of neurogenesis from the brain's own neural stem cells, and intracortical transplantation of cortical neurons directly converted from fibroblasts could be developed into novel therapeutic strategies for stroke in the future.</p>		
Key words: Adult neurogenesis, stroke, neural stem cells, Lnk, ependymal cells, FoxJ1, cortical reprogramming		
Classification system and/or index terms (if any):		
Supplementary bibliographical information:	Language	
ISSN and key title: 1652-8220	ISBN 978-91-87651-56-4	
Recipient's notes	Number of pages 116	Price
	Security classification	

Distribution by (name and address)

I, the undersigned, being the copyright owner of the abstract of the above-mentioned dissertation, hereby grant to all reference sources permission to publish and disseminate the abstract of the above-mentioned dissertation.

Signature \_\_\_\_\_

Date 2014-02-10

# Neurogenesis from Neural Stem Cells, Ependymal Cells and Fibroblasts

**Karthikeyan Devaraju**

Laboratory of Stem Cells and Restorative Neurology  
Division of Neurology  
Department of Clinical Sciences  
Lund Stem Cell Center  
Lund University  
Lund, Sweden



**LUND**  
UNIVERSITY

Academic Dissertation

2014

## Cover image

The image shows the delaminated and migrating ependymal cells from the lateral ventricular wall in adult rat brain, within the stroke-damaged striatum. The cells express FoxJ1 indicated by the presence of green and red fluorescent proteins while they do not immunostain for BrdU (blue).

Copyright © Karthikeyan Devaraju and respective publishers

ISBN 978-91-87651-56-4

ISSN 1652-8220

Lund University, Faculty of Medicine Doctoral Dissertation Series 2014:32

Printed in Sweden by Media-Tryck, Lund University

Lund 2014

*To my wonderful family*

*"In a day, when you don't come across any problems –  
you can be sure that you are travelling in a wrong path"*

*- Swami Vivekananda*



# TABLE OF CONTENTS

ORIGINAL PAPERS AND MANUSCRIPT.....	9
SUMMARY.....	11
SVENSKSAMMANFATTNING.....	12
தமிழ் முன்னுரை.....	15
ABBREVIATIONS.....	17
INTRODUCTION.....	19
Neurogenesis.....	19
Adult Neurogenesis.....	19
<i>Subventricular zone</i> .....	19
<i>Subgranular zone</i> .....	21
Stroke-induced neurogenesis.....	21
<i>Lnk in Neurogenesis</i> .....	22
<i>Ependymal cells and FoxJ1</i> .....	22
<i>Cortical neuron replacement</i> .....	23
Cortical neurons.....	24
Direct conversion to functional neurons.....	25
AIMS OF THE THESIS.....	27
RESULTS AND DISCUSSION.....	29
Paper I.....	29
<i>Lnk expression in brain and influence on neural stem cells</i> .....	29
<i>Lnk negatively regulates stroke-induced proliferation of NSPCs</i> .....	29
<i>Upstream regulators of Lnk and downstream effectors of Lnk after stroke</i> .....	30
<i>Discussion</i> .....	31
Paper II.....	32
<i>Specific labeling of LVW cells in adult rat brain by electroporation</i> .....	32
<i>FoxJ1+ cells in adult rat brain contribute to olfactory bulb neurogenesis</i> .....	32
<i>Stroke-induced striatal neurogenesis from FoxJ1+ cells</i> .....	34
<i>Discussion</i> .....	34
Paper III.....	35
<i>Conversion of human fibroblasts to functional pyramidal-shaped neuron-like cells</i> .....	36
<i>Layer- and region-specific functional cortical neurons</i> .....	36
<i>Discussion</i> .....	37
CONCLUDING REMARKS AND PROSPECTS.....	39
EXPERIMENTAL PROCEDURES.....	41
Animals.....	41
Experimental models.....	41
Human tissue.....	42
Dissection and cell culture.....	42
Plasmid constructs.....	43
Viral vectors.....	44
Neuronal reprogramming.....	45



Cell sorting and flow cytometry.....	45
RNA extraction and Quantitative PCR.....	46
Transcription factor binding site prediction.....	46
Chromatin immunoprecipitation PCR.....	47
Enzyme linked immunosorbent assay.....	47
BrdU administration.....	47
Immunostaining.....	48
Microscopical analysis.....	49
Electrophysiological analysis.....	50
Statistical analysis.....	51
ACKNOWLEDGEMENTS.....	55
REFERENCES.....	57
APPENDIX.....	67

## ORIGINAL PAPERS AND MANUSCRIPTS

This thesis is based on the following papers and manuscripts

- I. Henrik Ahlenius, **Karthikeyan Devaraju**, Emanuela Monni, Koichi Oki, Somsak Wattananit, Vladimer Darsalia, Robert E. Iosif, Olof Torper, James C. Wood, Sebastian Braun, Lucas Jagemann, Ulrike A. Nuber, Elisabet Englund, Sten-Erik W. Jacobsen, Olle Lindvall and Zaal Kokaia. Adaptor Protein LNK Is a Negative Regulator of Brain Neural Stem Cell Proliferation after Stroke.  
*The Journal of Neuroscience* 2012, 32(15): 5151-5164
- II. **Karthikeyan Devaraju**, Fanie Barnabé-Heider, Zaal Koakaia and Olle Lindvall. FoxJ1-expressing cells contribute to neurogenesis in forebrain of adult rats: Evidence from *in vivo* electroporation combined with piggyBac transposon.  
*Experimental Cell Research* 2013, 319(18): 2790-2800.
- III. **Karthikeyan Devaraju**, Giedre Miskinyte, Emanuela Monni, Marita Grønning Madsen, Daniel Tornero Prieto, Niels-Bjarne Woods, Olle Lindvall and Zaal Kokaia. Direct conversion of human fibroblasts to functional cortical neurons.  
*Manuscript*.



## SUMMARY

Stroke is a major cause of death and disability around the world. Stroke leads to loss of neurons and also other cells in the brain due to lack of blood supply. Currently no therapies are available to treat stroke-related disability. It has been shown that stroke leads to increased neurogenesis, birth of new neurons, within the brain. This increased neurogenesis is not sufficient to restore lost function. There is a need to develop therapies for neuronal replacement by improving neurogenesis within the brain and / or transplanting neurons. Cortical strokes lead to more disability after stroke as compared to those affecting the striatum, and whether cortical neurogenesis occurs after stroke is controversial. Cell transplantation may be the key to cortical repair after stroke.

Reports have identified positive but very few negative regulators of neurogenesis after stroke, and suggested that ependymal cells can also contribute to stroke-induced neurogenesis. Transplantation of neurons generated from different sources such as fetal brain, embryonic stem cells and induced pluripotent stem cells are associated with ethical issues and carry the risk of immune rejection and tumorigenicity. Direct conversion of patient's own skin cells to neurons could overcome these problems and potentially restore function after transplantation in stroke-damaged brain.

In this thesis we have used transgenic models, viral vectors, electroporation-mediated gene delivery and overexpression of transcription factors to demonstrate neurogenesis from neural stem cells, ependymal cells in the lateral ventricular wall and fibroblasts.

We show that Lnk, a known inhibitor of hematopoietic stem cell self-renewal, is also expressed in the brain. Overexpression or removal of Lnk expression leads to decreased or increased neurogenesis *in vitro* respectively. When brain is damaged by stroke there is increased proliferation of neural stem cells in animals without Lnk expression. This was not observed in status epilepticus, a severe form of epilepsy. We determined that upregulation of Stat1/3 after stroke leads to increased Lnk expression. Subsequently Lnk inhibits cellular response to increased IGF1 stimulation after stroke, by decreasing Akt phosphorylation. We have identified Lnk signaling as a novel mechanism of influencing neurogenic response to stroke.

We next determined if ependymal cells in lateral ventricular wall of adult rat brain contribute to neurogenesis after stroke. We identified FoxJ1 as a marker of ependymal cells in rats similar to mice, and used FoxJ1 promoter in piggyBac system to genetically label these cells with fluorescent reporter proteins GFP or RFP by electroporation. Tracing the lineage of the labeled cells in intact and stroke-damaged brain, we identified that FoxJ1 expressing cells contribute to olfactory bulb neurogenesis while the striatal neurogenic response was not significant. Thus, FoxJ1 expressing cells probably have only a minor role in repair after stroke.

We then tested whether human fetal lung fibroblasts could be directly converted to cortical neurons. We overexpressed sets of transcription factors that are known to be involved in cortical neuron development. We found that overexpression of different sets of these factors in fibroblasts converted them to cortical-like neurons. These neurons expressed markers of cortical neurons and were functional by electrophysiology.

In summary, these results raise the possibility that inhibition of Lnk, a negative regulator of neurogenesis from the brain's own neural stem cells, and intracortical transplantation of cortical neurons directly converted from fibroblasts could be developed into novel therapeutic strategies for stroke in the future.

## SVENSK SAMMANFATTNING

Stroke är en ledande orsak till dödlighet och handikapp i hela världen. Stroke leder till förlust av nervceller och även andra celler i hjärnan på grund av bristande blodtillförsel. För tillfället finns få möjligheter att behandla funktionshinder relaterade till stroke. Det har visats att stroke leder till ökad neurogenes, nybildning av nervceller i hjärnan. Denna ökade neurogenes är dock förmodligen inte tillräcklig för att återställa förlorad funktion. Därför finns ett behov av att utveckla nya terapier för att ersätta förlorade nervceller genom att förbättra neurogenes i hjärnan och/eller transplantatio av nervceller. Stroke som drabbar kortex leder till svårare handikapp jämfört med de som påverkar striatum, och hurvida kortikal neurogenes sker efter stroke är kontroversiellt. Därför kan celltransplantation kan vara nyckeln till reparation av kortex efter stroke.

Tidigare studier har identifierat många positiva men mycket få negativa regulatorer av nybildning av nervceller efter stroke. Det har också föreslagits att endymala celler kan bidra till stroke-inducerad neurogenes.

Nervceller till transplantation kan genereras från olika källor såsom fostrets hjärna, embryonala stamceller och inducerade pluripotenta stamceller. Dessa är dock förknippade med etiskt svåra frågor och har risker associerade med avstötning och uppkomst av cancer. Direkt omvandling av patientens egna hudceller till nervceller, och använda dessa till transplantation, skulle kunna övervinna dessa problem och potentiellt återställa funktionen i stroke-skadad hjärna.

I denna avhandling har vi använt transgena modeller, virala vektorer, elektroporation-medierad genleverans och överuttryck av transkriptionsfaktorer för att påvisa neurogenes från neurala stamceller, endymalceller och fibroblaster.

Vi visar att Lnk, en känd hämmare av hematopoietiska stamcellers självförnyelse, också uttrycks i hjärnan. Överuttryck eller avlägsnande av Lnk leder till minskad eller ökad neurogenes respektive in vitro. Efter stroke ses en ökad proliferation av neurala stamceller hos djur utan Lnk uttryck. Detta observerades dock inte i djur med status eplipeticus, en allvarlig form av epilepsi som också är associerad med ökad neurogenes. Vi visade att uppreglering av STAT1/3 efter stroke leder till ökat Lnk uttryck. Varpå Lnk hämmar neurala stamcellers svar på ökad IGF1 stimulering efter stroke, genom att minska AKT fosforylering. Vi har därigenom identifierat Lnk signalering som en ny mekanism för att påverka neurogenes efter stroke.

Vi undersökte sedan hurvida endymalceller i ventrikelväggen på vuxna råttor bidrar till neurogenes efter stroke. Vi identifierade FoxJ1 som en markör av endymala celler i råttor. Vi använde sedan FoxJ1 promotorn och piggyBac systemet för att genetiskt märka dessa celler med fluorescerande reporterproteiner, GFP eller RFP, genom elektroporering. Genom att följa de märkta cellerna i intakt och strokeskadad hjärna, identifierade vi att FoxJ1 uttryckande celler bidrar till neurogenes i luktbulberna men väldigt lite i striatum. Således har FoxJ1 uttryckande celler förmodligen bara en mindre roll i reparation av striatum efter stroke.

Vi testade sedan om humana fetala lungfibroblaster kan direkt omvandlas till kortikala nervceller. Vi överuttryckte olika kombinationer av transkriptionsfaktorer som är kända för att vara inblandade i utvecklingen av kortikala nervceller. Vi upptäckte att överuttryck av olika uppsättningar av dessa faktorer i fibroblaster konverterade dem till kortikal-liknande nervceller. Dessa nervceller uttryckte markörer för kortikala nervceller och visades vara funktionella med hjälp av elektrofysiologi.

Sammanfattningsvis, visar dessa resultat, att hämning av Lnk, en negativ regulator av neurogenes från hjärnans egna neurala stamceller, och intrakortikal transplantation av kortikala nervceller direkt omvandlade från fibroblaster kan utvecklas till nya terapeutiska strategier för stroke i framtiden.

# தமிழ் முன்னுரை

உலகமெங்கிலும் இறப்புக்கும் இயலாமைக்கும் பக்கவாதம் ஒரு தலையாய காரணமாகவுள்ளது. மூளைக்கு செல்லவேண்டிய குருதியளவில் பற்றாக்குறை ஏற்படுமேயானால் மூளையிலுள்ள நரம்பணுக்கள் உட்பட இதர சில அணுக்கள் சிதைவுறும். இச்சிதைவின் காரணமாக பக்கவாதம் ஏற்படுகிறது. பக்கவாதத்தினால் பாதிக்கப்படுபவர்கள் பேச்சாற்றல் நினைவாற்றல் இழப்பதுமட்டுமன்றி உடல் ஊனமுற்றும் போகிறார்கள். குறிப்பாக புரணிமூளை பக்கவாதத்தால் பாதிக்கப்படுபவர்கள் பேரியலாமைக்குள்ளாகிறார்கள். இவ்வாறான ஆற்றலின்மைக்கும் உடலூனத்திற்கும் தற்பொழுதுள்ள சிகிச்சை முறைகள் முழுமையான தீர்வளிப்பதில்லையென ஆய்வுகள் குறிப்பிடுகின்றன.

பக்கவாதத்தினால் இழந்த ஆற்றலையும் செயல்பாட்டையும் திரும்பப்பெற மூளையில் நரம்பாக்கம் போதிய அளவில் நடைபெற வேண்டும். மூளையில் நரம்பாக்கமொரு தொடர் நிகழ்வு ஆனால் புரணிமூளையில் நரம்பாக்கம் நடைபெறுவதில்லை. பக்கவாதம் ஏற்பட்டபின்னர் நரம்பாக்கம் அதிகரிக்கின்றதெனினும் அது இருகிழமைகள் மட்டுமே நீடிக்கின்றது. அதன் பின்னர் நரம்பாக்கம் குறையத்துவங்குகிறது. ஆதலால் இந்த நரம்பாக்க அதிகரிப்பு, இழந்த செயல்பாட்டை மீட்டுப்பெற போதுவதில்லை. எனவே பக்கவாதத்தை குணப்படுத்த நரம்பாக்கத்தை அதிகரிக்கும் வகையிலான நரம்பணு மாற்றறுவை சிகிச்சைகளை உருவாக்க வேண்டும்.

பக்கவாதம் ஏற்பட்ட பின்னர் நரம்பாக்கத்தின் எதிர்மறை-கட்டுப்பாட்டு காரணிகள் நரம்பாக்கத்தை குறைக்கின்றன. மேலும் மூளையில் உள்ள எப்பன்டைமல் செல்கள் பக்கவாதம் ஏற்பட்ட பிறகு நரம்பாக்கத்திற்கு பங்களிக்கின்றன. கருஉயிரணு மற்றும் கருஆற்றல் மிக்க உயிரணு மாற்றறுவை சிகிச்சைகளில் நெறிமுறை சிக்கல்கள், நோய் எதிர்ப்பு நிராகரிப்பு மற்றும் புற்று நோய் ஏற்படும் ஆபத்து உள்ளன. நோயாளியின் சொந்த தோல் உயிரணுக்களை நேரடியாக நரம்பணுக்களாக மாற்றினால் சிகிச்சைக்கு பின்னர் மூளையின் செயல்பாட்டை மீட்க முடியும் மற்றும் மேற்கண்ட சிக்கல்கள் ஏற்படாது.

இந்த ஆய்வறிக்கையில் மரபணு விலங்குகள், வைரஸ் விநியோக-கலங்கள், மின்சாரம் மூலம் மரபணு செலுத்தி மற்றும் மரபணு வெளிப்பாட்டு காரணிகளை பயன்படுத்தி கீழ்கண்ட ஆய்வுகளை மேற்கொண்டோம்.

Lnk, மூளையில் வெளிப்படுத்தப்படுகிறது என்றும், அதன் வெளிப்பாடு பக்கவாதம் ஏற்பட்டால் அதிகரிக்கின்றது. இது IGF1-AKT பாதையை கட்டுப்



படுத்தி பக்கவாதத்தினால் அதிககரிக்கும் நரம்பாக்கத்தை குறைக்கின்றது.

எலியின் மூளையில் எப்பன்டைமல் உயிரணுக்களை FoxJ1 அடையாளம் காட்டுகின்றது. பிக்கிபேக் மரபணுவை மின்சாரம் மூலம் செலுத்தி எப்பன்டைமல் உயிரணுக்களை குறித்தோம். இதன் மூலம் மூளையின் நுகர்தல் தண்டு பகுதியில் நரம்பாக்கத்திற்கு பங்கேற்-கின்றன என கண்டுபிடித்தோம். பக்கவாதத்திற்கு இவ்வுயிரணுக்களின் பங்களிப்பு குறைவு என கண்டறிந்தோம்.

மனித கருவிலிருந்து எடுக்கப்பட்ட நுரையீரல் உயிரணுக்களை நேரடியாக புறணிமூளை நரம்பணுக்களாக மாற்ற முயன்றோம். குறிப்பிட்ட மரபணு வெளிப்பாட்டு காரணிகளை பயன்படுத்தி நுரையீரல் உயிரணுக்களை புறணிமூளை நரம்பணுக்கள் போன்ற உயிரணுக்களாக மாற்றியுள்ளோம்.

ஆகவே, எதிர்காலத்தில் பக்கவாத சிகிச்சைகளுக்கு இவ்வாய்வுகள் வழிவகுக்கும்.

## ABBREVIATIONS

ABM	Ascl1, Brn2 and Myt1l
ANOVA	Analysis of variance
AMPA	$\alpha$ -Amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid
BrdU	Bromo deoxy-Uridine
CA	Constitutively Active
CCA	Common Carotid Artery
ChIP	Chromatin Immuno Precipitation
DCX	Doublecortin
DN	Dominant Negative
DNA	Deoxyribo Nucleic Acid
dpi	days post induction
EPO	Erythropoietin
EGF	Epidermal Growth Factor
FACS	Fluorescent Activated Cell Sorting
FBS	Fetal Bovine Serum
FGF	Fibroblast Growth Factor
GABA	$\gamma$ Amino Butyric Acid
GFAP	Glial Fibrillary Acidic Protein
GFP	Green Fluorescent Protein
HEFL2	Human Fetal Lung Fibroblasts
ICA	Internal carotid artery
iCtx	induced Cortical neurons
iDA	induced Dopaminergic neurons
IGF1	Insulin-like Growth Factor 1
iMN	induced Motor Neurons
iN	induced Neurons
i.p.	Intraperitoneal
LVW	Lateral Ventricular Wall
MCAO	Middle Cerebral Artery Occlusion
MCA	Middle Cerebral Artery
NeuN	Neuronal Nuclei / Fox3a
NMDA	N-methyl-D-aspartic acid
NSPC	Neural stem and progenitor cells
OB	Olfactory Bulb
PBS	Phosphate Buffered Saline
PCR	Polymerase Chain Reaction
PFA	paraformaldehyde
PH	Pleckstrin Homology
PTX	Picrotoxin
qPCR	quantitative Polymerase Chain Reaction
RFP	Red Fluorescent Protein

RNA	Ribo Nucleic Acid
RT	Room Temperature
RT-PCR	Reverse transcriptase Polymerase Chain Reaction
rtPA	recombinant tissue Plasminogen Activator
SE	Status Epilepticus
SEM	Standard Error of Mean
SGZ	Subgranular zone
SH2	Src Homolog 2
SVZ	Subventricular zone
TAP	Transit Amplifying Progenitor cells
TEA	Tetraethylammonium
TTX	Tetrodotoxin
TUNEL	Terminal deoxynucleotidyl transferase dUTP nick end labeling
Wt	Wild type

# INTRODUCTION

## Neurogenesis


Neurogenesis, birth of neurons, starts from the embryonic stage and continues in adult brain declining with age until death. The precursor of all neurons and macroglia in the central nervous system are the neuroepithelial cells that give rise to radial glial cells. The radial glial cells line the ventricle and extend processes to the pial surface. The radial glial cells expand by symmetric division and then give rise to neural progenitors by asymmetric cell division (Huttner and Brand, 1997). Thus they maintain a pool of radial glial cells while also giving rise to neural progenitors (Gotz and Huttner, 2005). The regional specification of neural precursors into forebrain, midbrain and hindbrain is orchestrated by morphogen gradients and specific gene expression patterns (Sansom and Livesey, 2009; Shimamura and Rubenstein, 1997; Tiberi et al., 2012). The neuronal diversity is determined by specific gene expression patterns. The temporal and spatial expression of genes determines the fate of the neurons, different regions and functions of the brain (Greig, Woodworth et al. 2013).

## Adult Neurogenesis

Adult neurogenesis is confined to two regions of the brain, the subventricular zone (SVZ) and the subgranular zone (SGZ). Initial reports of neurogenesis in adult brain were made with findings from H<sup>3</sup>-Thymidine (Altman and Das, 1965) in hippocampus of adult rats. Later, several reports have clearly shown that neurogenesis from these regions has functional significance (Abrous et al., 2005; Schinder and Gage, 2004). The demonstration that cells grow and self renew in the form of spheres when isolated from SVZ and SGZ has been a hallmark of the functional assay for neural stem cells (Reynolds and Weiss, 1992). Sox2, BLBP, GFAP, CD133, Nestin and Vimentin are markers of neural stem cells, but these are also expressed in other cell types in the brain. So far, there is no unique marker to identify these cells. Partly the problem has been the slow division of cells to give rise to progenitors that rapidly proliferate. These proliferating cells have been identified by BrdU pulse chasing experiments, markers of cell cycle such as p-H3 and Ki-67, and retroviral labeling.

### *Subventricular zone*

The SVZ, present around the lateral ventricles of adult brain, gives rise to new olfactory bulb (OB) neurons. The neural stem cells, probably single ciliated ventricle-contacting astrocytes, reside within the niche consisting of the multiciliated ependymal cells that line the ventricles, proliferating progenitors, migrating neuroblasts and endothelial cells. In mice, the neurogenic SVZ consists of the lateral ventricular wall (LVW) lined by the ependymal cells and ventricle-contacting astrocytes that are arranged in pinwheel architecture (Mirzadeh et al., 2008). The ventricle-contacting astrocytes, express markers of neural stem cells GFAP, Nestin and Vimentin, and are also referred to as type B cells. The B cells give rise to proliferating type C cells that express Mash1, Dlx1 and give rise to doublecortin (DCX) expressing neuroblasts,



*Figure 1 Subventricular zone niche.*

The subventricular zone consists of the multiciliated ependymal cells separating the cerebrospinal fluid containing lateral ventricles from the parenchyma. The ependymal cell layer also have ventricle contacting astrocytes (type B cells), which have a single cilium. These astrocytes give rise to progenitor cells (type C cells) that proliferate and give rise to migrating neuroblasts (type A cells). These cells are closely associated with blood vessels. Modified from Riquelme et al.,(2008).

referred to as Type A cells. The type A cells migrate along the rostral migratory stream as chains of neuroblasts and end up in the OB becoming interneurons (Alvarez-Buylla et al., 2002; Doetsch, 2003a, b; Doetsch et al., 1999). Depending upon the site of origin within the SVZ they become granule cell or the periglomerular interneurons. They express markers such as calretinin and calbindin while a few of them also express tyrosine hydroxylase and Tbr1/2 (Brill et al., 2009; Ihrie and Álvarez-Buylla, 2011; Kohwi et al., 2005; Kohwi et al., 2007). These new neurons are constantly produced to replace the dying neurons in the granule cell layer and the periglomerular region. Species difference between mice, rats and humans have been observed. In rats the type C cells have not been identified (Danilov et al., 2009). The

organization of the SVZ niche in humans is different with a hypocellular gap between the LVW lined by ependymal cells and the astrocytes that give rise to progenitors (Quiñones-Hinojosa et al., 2006). The type C cells have not been identified and there is absence of chain migration of neuroblasts.

### *Subgranular zone*

The SGZ of the dentate gyrus in the hippocampus is the other neurogenic region in the adult brain. The GFAP+ neural stem cells give rise to proliferating intermediate progenitors. These intermediate progenitors give rise to DCX+ neuroblasts that integrate into the granule cell layer of the dentate gyrus. These immature cells integrate into the network and mature by establishing connections with the adjacent granule cells and interneurons. They send axonal projections to the CA3 area of hippocampus (Abrous et al., 2005; Zhao et al., 2008). The addition of these new neurons plays a major role in learning and memory function (Deng et al., 2010; Gage, 2002). SGZ neurogenesis is increased after brain damage such as stroke and epilepsy (Benzon et al., 1997; Parent et al., 1997).

## **Stroke-induced neurogenesis**

Stroke is a major cerebrovascular disease that leads to disability and mortality. The most common cause of stroke is ischemic stroke, caused by blockage of arteries that supply the brain. Currently there are no therapies for stroke except recombinant tissue plasminogen activator (rtPA) that is used to lyse the clot causing the stroke (Frey, 2005). But, rtPA has a therapeutic window of 3 – 6 hours only in selected patients to be effective (Wardlaw et al., 2012). There is a need to develop therapies towards functional recovery of stroke-affected brain.

Experimentally occluding the blood vessels supplying the brain physically or chemically induces stroke in animal models. The most commonly used model to mimic clinical condition is intraluminal occlusion of the middle cerebral artery (MCAO) (Durukan and Tatlisumak, 2007; Iosif et al., 2008; Kokaia et al., 1998). Depending on the duration of occlusion the damage extends from striatum only to involving both the cortex and striatum. Occlusion of the distal branch of the MCA induces cortical lesion. Stroke has been shown to induce increased neurogenesis in the SVZ and migration of these neuroblasts towards the peri-infarct region in rodents and humans (Arvidsson et al., 2002; Jin et al., 2006; Martí-Fàbregas et al., 2010; Parent et al., 2002; Thored et al., 2006). The increased proliferation and migration of these neuroblasts is influenced by many factors including IGF1 (Thored et al., 2009), glial cell-derived neurotrophic factor (Kobayashi et al., 2006), vascular endothelial growth factor (Wang et al., 2007), erythropoietin (Tsai et al., 2006) and notch signaling (Androutsellis-Theotokis et al., 2006; Carlen et al., 2009). The migrating neuroblasts integrate into the stroke-damaged brain and become mature neurons expressing markers such as NeuN (Arvidsson et al., 2002; Parent et al., 2002). Only a few of these neuroblasts survive and become functional neurons that also project to distant targets (Arvidsson et al., 2002; Sun et al., 2012b). It has been shown that neurogenic response is necessary for functional recovery and reducing the stroke-induced

tissue loss (Sun et al., 2012a). The increased neurogenic response lasts for two weeks and returns to baseline levels (Thored et al., 2006). Only a few of the migrating neuroblasts survive and mature as neurons (Arvidsson et al., 2002; Parent et al., 2002). Due to this functional recovery is never complete due to the less number of neurons replacing the lost neurons. Thus there is a need to enhance the generation, survival, migration, differentiation and functional integration of these newborn neurons to restore function after stroke. Most importantly the restoration of the phenotype of the lost neurons is critical to restore function. Even a minor improvement in function will enhance the recovery in stroke patients.

### *Lnk in Neurogenesis*

The increased neurogenic response after stroke persists for two weeks and returns to basal levels (Arvidsson et al., 2002; Parent et al., 2002; Thored et al., 2006). There is still striatal neurogenesis and migration towards the stroke-damaged brain for few months after stroke. Most of the studies have shown factors that enhance neurogenesis, but factors that reduce the neurogenic response also need attention. It has been shown that TNF- $\alpha$  is an inhibitor of neurogenesis after stroke, but there might be other factors inhibiting neurogenesis (Iosif et al., 2008). Interestingly, in hematopoietic system, Lnk has been shown to inhibit proliferation and expansion of hematopoietic stem cells. Lack of Lnk has been shown to increase the number of hematopoietic stem cells and also B-cell genesis (Buza-Vidas et al., 2006; Takaki et al., 2000). It has been implicated in regulating neovascularization by inhibiting endothelial progenitor proliferation (Kamei et al., 2010; Kwon et al., 2009). The role of Lnk in brain has not been elucidated and its expression in brain is unknown.

Lnk, also known as SH2B3, is an adapter protein belonging to the SH2B adaptor family of proteins. These proteins are involved in signaling by growth factor and cytokine receptors. SH2B3 is conserved across humans, chimpanzee, rhesus monkey, dog, cow, mouse, chicken and zebrafish. It contains Src Homology 2 (SH2) and pleckstrin homology (PH) domains along with a phenylalanine zipper. The SH2 domain helps to bind to phosphorylated tyrosine residues on other proteins and gives rise to their involvement in signal transduction of receptor tyrosine kinase pathways. The PH domain allows them to bind to phosphatidylinositols, protein kinase C and  $\beta\gamma$  subunits of G-proteins, which are involved in intracellular signaling. The phenylalanine zipper allows the proteins to dimerize. Lnk is involved in c-Kit and erythropoietin (EPO) signaling pathways (Tong et al., 2005; Velazquez et al., 2002), which have been shown to influence NSPC proliferation, survival, migration and differentiation (Sun et al., 2004; Tsai et al., 2006). Whether Lnk is expressed in the brain and if expressed, what is its role in neurogenesis in intact and stroke-damaged brain is not known. Loss or gain of function studies with Lnk would help us to identify whether Lnk is a modulator of stroke-induced neurogenesis.

### *Ependymal cells and FoxJ1*

The ependymal cells are multiciliated and are an important component of the SVZ niche. They help to maintain the cerebrospinal fluid flow in the ventricles and also maintain a gradient of factors such as BMP and Noggin that allows the migration of neuroblasts to the olfactory

bulb (Sawamoto et al., 2006). Disruption of ependymal cells and the layer integrity leads to hydrocephalus and impaired neurogenesis (Jimenez et al., 2009). Ependymal cells have also been shown to be neural stem cells (Chiasson et al., 1999; Johansson et al., 1999) but their contribution to neurogenesis has been controversial. Studies have shown that ependymal cells also give rise to OB and striatal neurogenesis after stroke in a notch dependent manner (Carlen et al., 2009). In mice the ependymal cell derived cells become gliogenic differentiating to astrocytes or oligodendrocytes. In rats displaced ependymal cells have been observed after stroke but not much is known about their fate (Danilov et al., 2012). Interestingly in intact human brain displaced ependymal cells are observed in the astrocytic ribbon layer of the SVZ niche (Quiñones-Hinojosa et al., 2006). Thus it is possible that these cells can improve neurogenic response and survival of the migrating neuroblasts into the peri-infarct region. Ependymal cells express multiple markers such as Vimentin, S100 $\beta$ ,  $\beta$ -Catenin clusters, CD24, CD133 and LRP2. But these markers are not unique to ependymal cells as these are also expressed by astrocytes and neural stem cells. To study these ependymal cells genetically a unique marker is essential.

FoxJ1 has been shown as a marker of ependymal cells in mice and is necessary for development of ependymal cells from radial glia (Jacquet et al., 2009). FoxJ1 is a marker of ciliated cells in the body (Ostrowski et al., 2003) and is conserved across many species including chimpanzee, Rhesus monkey, dog, cow, mouse, chicken, and zebrafish. It belongs to the forkhead family of proteins and regulates the transcription of genes involved in motile ciliary assembly. Studies have shown that prenatal and early postnatal OB neurogenesis can happen from FoxJ1+ cells (Jacquet et al., 2011)2011. Using the FoxJ1-cre mice it has been shown that the FoxJ1+ ependymal cells contribute to striatal neurogenesis after stroke (Carlen et al., 2009). The stroke lesion size and neurogenic response are robust in rats compared to mice models. Since lesion size determines neurogenic response, the variability of lesion size in mice leads to varied neurogenic response within the same experimental setup. Thus it is even more important to study the neurogenic response and mechanisms in rats. The lack of transgenic rats is a huge drawback and hence there is a necessity to use both mice and rats to study stroke therapies. Whether FoxJ1 is expressed in rats is not known and if expressed whether it could be used to study ependymal cells after stroke needs to be studied. Genetic labeling experiments using FoxJ1 could help us to study rat ependymal cell response after stroke.

### *Cortical neuron replacement*

Cortical damage causes major disability compared to striatal damage following stroke in humans (Delavaran et al., 2013). Thus restoration of cortical function is important to functional recovery. There have been conflicting reports about cortical neurogenesis in stroke-damaged brain. The presence of neuroblasts in stroke damaged cortex was not observed in some reports (Arvidsson et al., 2002; Parent et al., 2002). On the other hand, there are reports showing that SVZ derived neuroblasts migrate to the stroke-damaged cortex (Jiang et al., 2001; Kreuzberg et al., 2010; Ziv et al., 2007). The subpial cortical layer I progenitor cells were shown to proliferate and give rise to GABAergic neurons after stroke (Ohira et al., 2010). These variations can be



attributed to factors including stroke model, infusion of growth factors and extent of lesion. Given the complexity of the cortex with multiple neuronal phenotypes and connections this endogenous neurogenic response may not be sufficient for neuronal replacement after stroke. Replacement strategies by cell transplantation remain the viable option for replacing the lost cortical neurons. Fetal brain tissue transplantation in humans restores function in Parkinson's disease (Lindvall et al., 1990). Transplantation of cells derived from embryonic stem cells (ESC) (Espuny-Camacho et al., 2013) show that they follow development patterns and establish proper connections in mouse brain. Transplantation of cortically fated induced pluripotent stem cells (iPSC) have been shown to restore function after stroke (Torneró et al., 2013). Fetal brain tissue derived cells have ethical problems and are a limited source of transplantable cells. ESC and iPSC carry the risk of tumorigenicity from undifferentiated pluripotent cells. Direct conversion of fibroblasts to neurons is a viable alternative to the ESC and iPSC derived neural transplants. Similar to iPSC technology (Takahashi and Yamanaka, 2006), fibroblasts can be converted to functional neurons without going through pluripotent state, by using a defined set of transcription factors (Vierbuchen et al., 2010). The greatest challenge is to derive the different subtypes of neurons present in the six-layered cortex.

## Cortical neurons

The cortex is divided into six distinct layers, layers I - VI, dorsoventrally in rodents and superoinferiorly in humans, based on the morphology, function and gene expression pattern. The cortex is further divided into distinct functional areas, each having different combinations of transcription factor expression. The functional or regional specification within the cortex is determined by gene expression patterns and adhesion molecules expressed (Greig et al., 2013; Srinivasan et al., 2012). For example *Bhlhb5* and *Lmo4* transcription factor expression along with *cadherin-8* (*Cdh8*) determine the boundary between sensory and motor areas (Azim et al., 2009; Cederquist et al., 2013; Greig et al., 2013). The projection neurons in the different layers project to the thalamus (layer VI), hindbrain and spinal cord (layer V) and to inter- or intra-hemispheric projections (layer IV). Functional recovery in damaged cortex requires restoration of the different layers and their connections in a given region of the brain. This requires that the cells transplanted have the necessary molecular and phenotypic signatures as observed in cortical neurons. Understanding the cortical development and determining the key molecules involved in this process could help us to develop transplantable cells. Cortical development could be divided into three phases, 1) the development of specific layers, 2) the development of different functional regions within the cortex and 3) the molecular mechanisms of how these neurons find their targets and maintain these connections.

The first step in developing layer-specific neurons would be the identification of key molecules that determine cortical neuron specificity. *Fezf2* (Arlotta et al., 2005; Rouaux and Arlotta, 2010), *Ctip2* (Chen et al., 2008), *Satb2* (Alcamo et al., 2008), *Tbr1* (Hevner et al., 2001) and *Cux2* (Franco et al., 2012; Molyneaux et al., 2009) have been identified as critical molecules for cortical layer specification. Whether the expression of these molecules in fibroblasts will convert them to cortical neurons is not yet explored.




Figure 2. Laminar organization and functional areas of the rodent cortex.

The representative figure illustrates the complexity of the cortex in an adult rodent brain. The different functionally distinct areas, namely auditory (A), motor (M), somatosensory (S) and visual (V) areas in the cortex are depicted on the left hemisphere along the rostro-caudal axis of the brain. The coronal section on the right hand side shows the laminar organization of the cortex into six layers along with their known projections. The different neuron phenotypes mostly found in these layers along with their characteristic molecular markers are depicted. The other neuronal subtypes and cells are not shown. Adapted from Greig et al. (2013) and Molyneaux et al. (2007).

## Direct conversion to functional neurons

Direct conversion of somatic cells to neurons, called induced neurons (iN), has been shown in mouse and human cells (Marro et al., 2011; Pang et al., 2011; Pfisterer et al., 2011b; Vierbuchen et al., 2010). The overexpression of proneural genes such as *Ascl1* or *Ngn2* along with other transcription factors (Ladewig et al., 2012; Vierbuchen et al., 2010) in somatic cells gives rise to iN cells. Subtype-specific neurons such as dopaminergic neurons (iDA) (Caiazzo et al., 2011; Liu et al., 2012; Pfisterer et al., 2011a) and spinal motor neurons (iMN) (Son et al., 2011) have been demonstrated. These cells were derived by expression of proneural genes along with subtype-specific transcription factors. Transplantation studies have shown that these neurons survive in the host brain (Torper et al., 2013) and reverse the functional deficits in a mouse model of Parkinson's disease (Kim et al., 2011). Since this is an emerging technology, their safety and efficacy to restore function in brain damage needs to be analyzed in detail. Even though these studies demonstrated the feasibility of deriving mixed population of neurons (Vierbuchen et al., 2010) or subtype-specific neurons, the number of neurons derived or neuronal conversion

efficiency from fibroblasts is less. To be useful for disease modeling and clinical transplantation studies, more neuronal cells have to be derived. Using synergistic SMAD (Mothers against decapentaplegic homolog) inhibition, the conversion efficiency has been enhanced manifold (Ladewig et al., 2012). Thus, a highly enriched population of neurons can be derived from somatic cells in mice and humans. The possibility of reprogramming fibroblasts to cortical neurons would open up options for transplantation studies in stroke. Whether combining the proneural factors with cortical layer specific transcription factors would give rise to cortical neurons remains to be explored.

## AIMS OF THE THESIS

The main objective of the thesis has been to explore different approaches for neuronal replacement in the stroke-damaged brain. The main strategies were to investigate the neurogenic response from endogenous neural stem cells and ependymal cells after stroke, and direct conversion of fibroblasts to neurons. The specific aims of the thesis have been:

- I. To investigate the role of Lnk in adult neurogenesis from subventricular zone after stroke (*paper I*)
- II. To investigate the role of ependymal cells in lateral ventricular wall in adult neurogenesis after stroke (*paper II*)
- III. To derive cortical neurons by direct conversion of human fetal lung fibroblasts for transplantation after stroke (*paper III*)



## RESULTS AND DISCUSSION

### Adaptor protein Lnk is a negative regulator of brain neural stem cell proliferation after stroke (paper I)

Adult neurogenesis is modulated by pathological conditions such as stroke and epilepsy. SVZ neurogenesis increases for the first two weeks after stroke and plateaus later. The mechanisms behind the decrease in neurogenesis are not fully understood. Cytokines such as C-KIT and EPO are potent modulators of NSPC after stroke. Lnk, an adaptor protein, negatively regulates cytokine pathways including C-KIT and EPO. The role of Lnk in normal and pathological brain has not been elucidated.

#### *Lnk expression in brain and influence on neural stem cells*

Lnk belongs to adaptor protein family that has SH2 and PH domains with a C-terminal tyrosine phosphorylation site. The expression and role of Lnk in hematopoietic system has been delineated. So far it has been detected in embryonic rat cortex (Wang et al., 2011), but the presence and function in human or mouse SVZ has not been determined. We detected Lnk expression in mouse and human SVZ tissue and also in neurospheres derived from mouse SVZ and human fetal ganglionic eminences. The expression was observed in SOX2+, DCX+ and IBA1+ cells of SVZ but also in other cell types that did not express these markers. We then used the Lnk<sup>-/-</sup> mice and could not detect abnormalities in the gross structure and fine cytoarchitecture of brain. We grew neurospheres from SVZ of Lnk<sup>-/-</sup> mice and observed that over subsequent passages, the numbers of proliferating cells were higher compared to Wt mice. Data from BrdU flow cytometry showed that Lnk<sup>-/-</sup> cells proliferate faster by shortening G0/G1 phase. We then overexpressed Lnk in neurospheres from Wt mice and found the size of the Lnk-overexpressing neurospheres was dramatically reduced along with more TUNEL+ cells. These results clearly showed that Lnk is an inhibitor of NSPC proliferation.

#### *Lnk negatively regulates stroke-induced proliferation of NSPCs*

Having found that Lnk inhibits NSPC proliferation *in vitro*, we then determined if Lnk inhibits NSPC proliferation *in vivo*. We used BrdU pulse chase analysis coupled with p-H3 immunofluorescence in Lnk<sup>-/-</sup> and Wt mice. We could not detect any differences in the p-H3+ and BrdU+ population in SVZ of these mice, indicating that Lnk has no influence on NSPC proliferation in SVZ of intact mice. Having shown that stroke induces transient increase in NSPC proliferation in SVZ (Arvidsson et al., 2002; Thored et al., 2006), we decided to check the role of LNK in stroke-induced neurogenesis. We compared the NSPC proliferation and survival of newly formed cells 7 days after stroke in Lnk<sup>-/-</sup> and Wt mice by p-H3 and BrdU staining. We found p-H3+ and BrdU+ cells were increased in Lnk<sup>-/-</sup> SVZ compared to Wt mice. We also found significant increase in Sox2+/BrdU+ cells and an increase in DCX+/BrdU+ cells, while no increased proliferation of CD31+ endothelial cells, Iba1+ microglial or FoxJ1+ cells were observed. The stroke lesion size was not different between the Lnk<sup>-/-</sup> and Wt mice, since lesion size also determines neurogenic response. We then determined if the increased NSPC

proliferation is confined to SVZ or also to SGZ. SVZ neurogenesis is also increased after status epilepticus (SE), a severe model of epilepsy that mimics the condition seen in humans. We induced SE in *Lnk*<sup>-/-</sup> and Wt mice and analyzed the SVZ for NSPC proliferation. We did not detect any differences in SVZ cell proliferation between *Lnk*<sup>-/-</sup> and Wt mice. We then compared the SGZ cell proliferation in the stroke and SE lesioned brains of *Lnk*<sup>-/-</sup> and Wt mice. We found increased cell proliferation in SGZ of *Lnk*<sup>-/-</sup> mice but no differences were observed between stroke and SE lesions. We did not find any differences in the numbers of Iba1+ microglia in the intact, stroke or SE lesioned brains of *Lnk*<sup>-/-</sup> and Wt mice. This ruled out the possibility of *Lnk* suppressing NSPC proliferation by microglial activation.

### *Upstream regulators of Lnk and downstream effectors of Lnk after stroke*

To determine the influence of stroke on *Lnk* upregulation, we checked the *Lnk* expression levels in intact, stroke- and SE-lesioned mice brain by qPCR. *Lnk* expression was upregulated after stroke while downregulated after SE. We analysed the *Lnk* promoter region for transcription factor binding motifs and found SP1, E2F1, Stat1 and Stat3 binding sites. We detected by qPCR that Stat1 and Stat3 were upregulated after stroke, indicating that Stat1 or Stat3 could influence *Lnk* upregulation after stroke. We transfected CA-Stat1 and DN-Stat1 plasmids to Wt SVZ derived neurospheres and found increased *Lnk* expression in CA-Stat1 transfected cells. By ChIP analysis with Stat1 antibody, we found that Stat1 was specifically binding to first exon of *Lnk*. Stat1 and Stat3 work in synergy, hence we cloned the *Lnk* genomic sequence including the Stat1 and Stat3 binding region to a luciferase reporter plasmid. We also mutated the Stat1 and Stat3 binding sites of the above-mentioned *Lnk* sequence. These plasmids were transiently expressed in 3T3 and Stat1-deficient U3A cells and also separately co-transfected CA-Stat1 and CA-Stat3 plasmids. In the presence of CA-Stat1 and CA-Stat3 plasmids, we found increased luciferase activity in *Lnk* sequence while the mutated *Lnk* sequence had decreased luciferase activity. These findings showed that Stat1 and Stat3 regulate increased *Lnk* expression after stroke.

To identify mechanisms through which *Lnk* might suppress stroke-induced NSPC proliferation, we first investigated using qPCR the expression of known and potential *Lnk* targets in SVZ tissue from intact mice and at 1 week after stroke or SE. Gene expression of Epo, Fgfr1, Fgfr2, Egfr, Fgf2 and Igf1 were upregulated only after stroke and not in SE. Based on the specific alterations of gene expression induced by stroke, we hypothesized that *Lnk* may act on the EPO, EGF, FGF and IGF1 signaling pathways.

*Lnk* is known to inhibit the phosphorylation cascade in receptor tyrosine kinase signaling by direct interaction with receptors (Takaki et al., 2000; Tong et al., 2005) or with downstream signaling (Bersenev et al., 2008). One common signaling pathway of EGF, FGF, and EPO is the phosphorylation of ERK1/2, while IGF1 signaling leads to Akt phosphorylation by PI3K. We analyzed the degree of phosphorylation of ERK1/2 and Akt in neurosphere cells from SVZ of *Lnk*<sup>-/-</sup> and Wt mice in response to EGF/FGF, IGF1, and EPO stimulation. We found marked increase of AKT phosphorylation in *Lnk*<sup>-/-</sup> neurospheres stimulated with IGF-1 compared with Wt controls. We then transfected CA-Akt or DN-Akt plasmids into the lateral ventricle of Wt mice by *in vivo* electroporation. 3 days later, animals were injected with BrdU, and SVZ

proliferation was assessed using BrdU and p-H3 immunohistochemistry. Consistent with a regulatory role of AKT on NSPC proliferation, we found that CA-Akt increased while DN-Akt decreased the number of BrdU+ and p-H3+ cells in the transfected SVZ.

The Lnk-induced changes in growth factor signaling could lead to alterations in gene expression and subsequently in cell proliferation. To explore this further, we analyzed the expression of several genes that have been implicated in NSPC proliferation and maintenance in the Lnk-overexpressing neurosphere cells. We found a significant reduction in Hes5 and Gli1 gene expression.

Together, our findings provide evidence that Lnk inhibits IGF1 signaling by attenuating phosphorylation of AKT, possibly also EGF/FGF signaling via ERK1/2, and decreases gene expression of Hes5 and Gli1, resulting in reduction of stroke-induced NSPC proliferation.

## *Discussion*

We have shown here that Lnk is expressed in adult brain and has a functional role in regulating NSPC proliferation after stroke. Interestingly, NSPC proliferation is regulated in stroke and not in SE. Previous reports in hematopoietic system have shown that Lnk is a negative regulator of HSC self-renewal and proliferation (Buza-Vidas et al., 2006; Seita et al., 2007), and B-cell progenitor proliferation (Takaki et al., 2003). It is also involved in regulation of endothelial cell proliferation (Kamei et al., 2010; Kwon et al., 2009). Mutations in Lnk gene have been observed in myeloproliferative neoplasms (Oh et al., 2010). We did not find increased endothelial cell proliferation or vascularization after stroke in Lnk<sup>-/-</sup> mice. We found no evidence for altered microglial activation in SVZ of Lnk<sup>-/-</sup> mice after stroke, while in absence of immune cells or microglia, Lnk overexpression in neurospheres decreased NSPC proliferation. Therefore, observed effect of increased NSPC proliferation in Lnk<sup>-/-</sup> mice cannot be attributed to effect of other cell types. Since Lnk is expressed in NSPCs, LNK influences NSPC proliferation cell autonomously.

We found that Lnk does not influence NSPC proliferation in intact brain of Wt or Lnk<sup>-/-</sup> mice in vivo. We observed increased proliferation of neurospheres from Lnk<sup>-/-</sup> mice compared to Wt mice. This could be due to the abundance of growth factors in the culture conditions similar to what is observed after stroke. We found that Lnk expression increased after stroke and decreased after SE, indicating that it is regulated in brain after pathological insults. We also found increased levels of Stat1 and Stat3 after stroke indicating that they upregulate Lnk after stroke. STAT1 (Takagi et al., 2002) and STAT3 upregulation after stroke is mediated by cytokine release after insult (Liesz et al., 2011; Yamashita et al., 2005). Cytokines such as IL-6 and IFN  $\alpha$  and  $\gamma$  are known to activate STAT family members (Takeda and Akira, 2000). Thus STAT1 / 3 might mediate Lnk upregulation after stroke.

The increased levels of FGF, EGF, EPO and IGF1 after stroke but not after SE might be responsible for increased neurogenesis in Lnk<sup>-/-</sup> mice after stroke. After stroke IGF-1 levels are increased (Thored et al., 2009) and mediate NSPC proliferation (Yan et al., 2006). LNK mediates receptor tyrosine kinase signaling by inhibiting or decreasing phosphorylation of



downstream kinases (Bersenev et al., 2008; Takaki et al., 2000; Tong et al., 2005). We found increased levels of AKT phosphorylation after IGF-1 stimulation and ERK1/2 phosphorylation after EGF and FGF stimulation in Lnk<sup>-/-</sup> neurospheres. Similarly, we found increased and decreased NSPC proliferation in SVZ after CA-Akt and DN-Akt electroporation, respectively. Thus, LNK might inhibit ERK1/2 and AKT phosphorylation in response to growth factor signaling. In hematopoietic stem cells and endothelial precursors LNK inhibits AKT and ERK1/2 phosphorylation (Seita et al., 2007; Tong et al., 2005; Wan et al., 2006). The targets of AKT and ERK1/2 are Hes and Gli1 proteins, involved in NSPC proliferation (Cayuso et al., 2006; Gregory et al., 2010; Stecca and Ruiz i Altaba, 2009), which we also observed to be downregulated with Lnk overexpression. Thus Lnk inhibits NSPC proliferation by inhibiting the IGF1-AKT pathway.

## **FoxJ1-expressing cells contribute to neurogenesis in forebrain of adult rats: Evidence from in vivo electroporation combined with piggyBac transposon (paper II)**


The SVZ neurogenic niche consists of multiciliated ependymal cells and ventricle contacting astrocytes with single cilium along with the vasculature (Mirzadeh et al., 2008). These single ciliated astrocytes are putative neural stem cells that give rise to the neural progenitors. The ependymal cells considered to be post-mitotic, have been shown to be neurogenic especially after stroke (Carlen et al., 2009). But the fate of these ependymal cell derived neuroblasts has not been determined. Whether the ependymal cells in adult rat brain contribute to stroke induced neurogenesis is not known. In mice FoxJ1 has been shown to be a marker of ependymal cells. Whether FoxJ1 is expressed in rat brain and if they contribute to neurogenesis is not yet known. Genetic labeling studies of FoxJ1-expressing cells in adult rat brain would help us to answer these questions.

### *Specific labeling of LVW cells in adult rat brain by electroporation*

The LVW consists of the ependymal cell and ventricle-contacting astrocytes. Labeling of LVW cells has relied upon viral vectors (Johansson et al., 1999) but these vectors label the entire ventricular wall. Electroporation has been shown to effectively label the LVW in adult mouse brain (Barnabe-Heider et al., 2008; Carlen et al., 2009). We attempted to label the LVW in adult rat brain using electroporation. We observed GFP+ cells in the LVW of adult rat brain and found the method to be safe without causing any damage to the neurogenic niche and neurogenesis. After optimizing the electroporation parameters, we observed efficient labeling of the cells in LVW with 175 V/cm pulse of 50ms duration at 1Hz frequency.

### *FoxJ1+ cells in the adult rat brain contribute to olfactory bulb neurogenesis*

FoxJ1 is a key transcription factor in functionally motile cilia development (Ostrowski et al., 2003). It is important for the development of ependymal cells and a subset of astrocytes from radial glia (Jacquet et al., 2009) 2009. FoxJ1 has been used as a specific marker of ependymal cells in the adult mouse brain (Carlen et al., 2009; Jacquet et al., 2009). We detected the expression



*Figure 3: Electroporation in lateral ventricular wall of adult rodent brain.*

Electroporation is a method of delivering DNA to the target cells by applying an electric potential. The plasmid is first injected into the lateral ventricle of brain by stereotaxy. Then the electrodes with conductive gel is applied on the head parallel to the injected site and electroporated. DNA is taken up by cells along the side of cathode due to the transient permeability caused by electroporation. The lateral ventricular wall cells are later analyzed for plasmid expression.

of FoxJ1 in the LVW of adult rat brain by using FoxJ1 antibody. Next, a Cre-expressing plasmid under FoxJ1 promoter, along with a floxed reporter construct was electroporated. Based on the Cre recombinase system, we identified that FoxJ1 is expressed in ependymal cells and also in a subset of astrocytes lining the LVW.

Having established the expression of FoxJ1 in adult rat brain, we needed to establish a stable genetic reporter for lineage tracing of FoxJ1+ cells. We used the *piggyBac* transposon system (Cadinanos and Bradley, 2007) to genetically label the FoxJ1+ cells and its progeny. The *piggyBac* transposon, with FoxJ1 promoter based donor (pPB-FoxJ1-eGFP) and helper (pFoxJ1-mPB) plasmids, was electroporated to the LVW of adult rat brain. We also used donor plasmid with RFP reporter (pPB-UbC-DsRed2) along with pFoxJ1-mPB for lineage tracing. We observed expression of both GFP+ and RFP+ cells in the LVW. Most of the GFP+ and RFP+ cells were FoxJ1+, while some of the RFP+ cells were DCX+ and GFAP+. We also found RFP+ and GFP+ cells in the OB of the electroporated brains and most of the cells were NeuN+. We did not observe any disturbances in the LVW and shows that OB neurons would have originated from the LVW cells. These results clearly showed that the *piggyBac* system is ideal for lineage tracing in adult rat brain. Most importantly we found that FoxJ1+ cells in the LVW of adult rat brain contribute to OB neurogenesis.

### *Stroke-induced striatal neurogenesis from FoxJ1+ cells*

Stroke induced striatal neurogenesis involves recruitment of migrating neuroblasts from the RMS to the stroke-damaged striatum. Since we found FoxJ1+ cells contributed to OB neurogenesis, we electroporated the pFoxJ1-mPB along with pPB-FoxJ1-eGFP and pPB-UbC-DsRed2. One week later, we induced stroke by 30 minutes and 2 hour MCAO models. One week following MCAO, we administered BrdU for seven days to detect proliferating LVW cells. We observed an increased neurogenic response after stroke, but very few GFP+/RFP+ cells were observed in the striatum. We observed both GFP+/RFP+ and GFP-/RFP+ cells with some BrdU+ among the GFP-/RFP+ cells. We could not determine the phenotype of these migrating cells. The OB neurogenesis from labeled cells was intact and was similar when compared to the intact brain.

Taken together our findings show that FoxJ1 expressing cells in the LVW of adult rat brain contribute to olfactory bulb neurogenesis in intact and stroke-damaged brain, while some ependymal cells delaminate and migrate into the stroke-damaged striatum.

### *Discussion*

We have used genetic lineage tracing with electroporation to show that FoxJ1 expressing cells in LVW contribute to neurogenesis in adult rat forebrain.

We show that electroporation is a safe and valuable method to label LVW of adult rat brain similar to what is described mice (Barnabe-Heider et al., 2008). No evidence for damage to the SVZ, rat brain or the rats in general from electroporation was found. Markers used to identify ependymal cells S100 $\beta$ , Vimentin, CD24 and CD133 are also expressed by astrocytes and NSPCs (Järlestedt et al., 2010; Pfenninger et al., 2007; Pruszek et al., 2009). FoxJ1 has been used to identify ependymal cells in mice (Carlen et al., 2009; Jacquet et al., 2011; Jacquet et al., 2009). We also determined by immunostainings and using FoxJ1-Cre (Carlen et al., 2009) and Cre-reporter plasmid electroporation that the ependymal cells express FoxJ1. A small population of astrocytes expressing FoxJ1 is consistent with similar findings in FoxJ1EGFP

mice (Jacquet et al., 2009).

We found long-term expression, 12 weeks in our study, of reporter proteins and also in the progeny of FoxJ1+ cells. We show the first evidence for *in vivo* labeling in adult brain using *piggyBac* transposon as an effective genome-targeting tool. *piggyBac* is known to stably express transgenes in human and rodent genomes (Cadinanos and Bradley, 2007; Ding et al., 2005; Jang and Behringer, 2007; Saridey et al., 2009; VandenDriessche et al., 2009; Wilson et al., 2007). With *piggyBac* transposon labeling, we were able to detect OB neurons arising from FoxJ1 expressing cells. This has not been detected so far in the rat brain. Previously, FoxJ1+ ependymal cells and their progeny in perinatal period were shown to be necessary for adult OB neurogenesis (Jacquet et al., 2011) 2011. FoxJ1+ cells have been shown to give rise to neurospheres (Jacquet et al., 2011) 2011 and these could have arisen from FoxJ1+ astrocytes. Since ependymal cells are post-mitotic, it is probable that the OB neurons observed here are from the FoxJ1+/GFAP+ cells. We observed that GFP expression under FoxJ1 promoter was weak in labeled neuroblasts and OB neurons, while RFP expression under FoxJ1 promoter was strong in all labeled populations. These results clearly showed that for effective labeling transposase expression should be under cell specific promoter.

Ependymal cells have previously been shown to give rise to neuroblasts (Carlen et al., 2009) and transform into radial glia after stroke (Zhang et al., 2007). They have also been shown to generate progeny by asymmetric division after 6-hydroxydopamine lesion (Gleason et al., 2008). We performed MCAO one week after electroporation with the GFP and RFP donor plasmids. GFP+/RFP+/BrdU- cells were found in the stroke-damaged striatum 6 weeks after stroke. We also found GFP-/RFP+/BrdU+ neuroblasts-like cells in the striatum but their fate could not be determined. This clearly shows that these cells originate from FoxJ1 expressing cells, while loss of GFP expression shows that FoxJ1 is downregulated after giving rise to progeny. We also observed RFP+/NeuN+ and RFP+/BrdU+ OB neurons after stroke. The GFP+/RFP+/BrdU- cells could be similar to the displaced ependymal cells found in striatum after stroke in rats (Danilov et al., 2012). Thus we show that FoxJ1 expressing cells contribute to OB neurogenesis in intact and stroke-damaged brain, but contribution to striatal repair after stroke in these experimental conditions is limited.

### **Direct conversion of fibroblasts to functional cortical neurons (paper III)**

Reprogramming of mature somatic cells to pluripotent stem cells (iPSC) (Takahashi and Yamanaka, 2006) and neurons (iN) (Vierbuchen et al., 2010), have opened up possibilities for patient specific transplantation to diseased brain. Most importantly these are also powerful tools to create human disease models *in vitro*. Many diseases such as neonatal hypoxia, head trauma, stroke and Alzheimer's disease affect the cortex and cause cortical dysfunction. Cortical neuronal transplants would be useful to restore function in diseased brain. Cells produced by direct conversion probably are not tumorigenic and do not require long-term immunosuppression after transplantation. Thus, cortical neurons from fibroblasts (iCtx) would be an invaluable source for intracerebral transplantation.

## *Conversion of human fibroblasts to functional pyramidal-shaped neuron-like cells*

Cortical projection neurons are glutamatergic and exhibit pyramidal morphology. We decided to use the transcription factors that are critical for development of projection neurons with glutamatergic phenotype. Brn2 (B), Zic1 (Z), Ctip2 (C), Fezf2 (F) and Tbr1 are critical for the formation of cortical projection neurons from layers II - VI. The combination of the transcription factors Ascl1 (A), Brn2 and Myt1l (ABM) has been shown to reprogram mouse fibroblasts to Tbr1+ neurons (Vierbuchen et al., 2010), a marker of layer VI cortical neurons. Brn2 and Zic1, part of transcription factor combinations (BMZ) for iN reprogramming (Vierbuchen et al., 2010) are expressed in layers II - V (Molyneaux et al., 2007). We initially tested whether the combinations ABM or BMZ would give rise to layer V and layer VI neurons alone or in combination with Fezf2 and/or Ctip2. However, we observed increased cell death and hardly any cells survived at 25 days after doxycycline addition.

We continued to search for combinations of transcription factors that could give rise to surviving neurons with pyramidal morphology. Factors critical for glutamatergic phenotype and NeuroD1 (Nd) have been shown in reprogramming studies to be more effective in converting fibroblasts to neurons (Pang et al., 2011). Brn2 and Myt1l have been shown to produce iN cells without Ascl1 in combination with other transcription factors (Vierbuchen et al., 2010). Hence, we included Brn2 and Myt1l along with cortical transcription factors to test if they would give rise to cortical layer-specific neurons. We observed surviving neurons and the combinations were analyzed for neuronal conversion efficiency and cortical projection neuron morphology. qPCR analysis for expression of cortical layer markers revealed that compared to other combinations, maximal expression was observed for Satb2, Cux2, Fezf2 and Tbr1. Combinations with NeuroD1 had more GAD1 expression compared to vGlut1 and vGlut2 expression. Whole-cell patch-clamp recordings revealed that cells from these combinations were able to generate action potentials. Based on the electrophysiological and morphological analysis along with the expression of layer specific markers, we identified BMNgn2, MNgn2 and BMF as the most promising combinations for inducing cortical layer-specific iCtx cells.

## *Layer- and region-specific functional cortical neurons*

We evaluated three combinations in more detail for their ability to generate iCtx cells expressing cortical layer- and region-specific markers. Cultured cells were transduced with SynI-GFP to distinguish neurons i.e. iCtx cells, from fibroblasts. Cells were FACS sorted for GFP expression and qPCR was performed in BioMark HD with 48.48 chip. First combination of transcription factors expressed Bhlhb5, Emx2 and Satb2 maximally while Tbr2, Pax6, Cux2, Ctip2 and Tbr1 were expressed at lower levels. Second combination expressed weakly Emx2, Tbr2, Bhlhb5, Satb2 and Ctip2. Interestingly, third combination expressed Tbr2, Satb2 and Ctip2 at very low levels. We could not detect expression of Fezf2 in any of the conditions. Electrophysiological recordings revealed that most of the iCtx cells from the three combinations exhibited the characteristics of mature neurons. We also found that BMF or BMNgn2 combinations were more responsive to Glutamate and GABA application compared to MNgn2. Taken together, these findings show that the neurons derived from human fetal lung fibroblasts by direct con-

version exhibit the characteristics of mature neurons and express functional glutamate and GABA receptors. By immunostaining we detected that cells reprogrammed with BMNgn2 and BMF were immunopositive for *Satb2* and *Ctip2* respectively.

Taken together, our findings show that cortical transcription factor combinations can induce cortical projection neuron phenotype from human fetal lung fibroblasts that is layer- and region-specific.

## *Discussion*

We have shown that expression of specific combinations of transcription factor can lead to specific cortical projection neuron subtypes. The cerebral cortex has six functionally distinct layers (layers I to VI) with combination of transcription factor expression. The initial iN study (Vierbuchen et al., 2010) showed that majority of mouse iN cells were *Tbr1*<sup>+</sup> and glutamatergic. *Tbr1* is a marker of layer VI neurons (Hevner et al., 2001) and is also expressed by olfactory bulb interneurons (Brill et al., 2009) and cerebellar neurons (Fink et al., 2006). Evaluation of other layer specific markers such as *Sox5* (layer VI), *Fezf2* (layer V), *Ctip2* (layer V), *Satb2* (layer IV) and *Cux2* (layer II – III) (Greig et al., 2013; Molyneaux et al., 2007) is necessary to confirm if these iN cells are cortical in nature. We did not have surviving neurons with ABM combination to evaluate their cortical phenotype, which is in agreement with other reports (Pang et al., 2011; Qiang et al., 2011).

The iCtx cells expressed layer specific genes *Satb2*, *Cux2*, *Tbr1* and *Fezf2*. Interestingly, the iCtx cells also expressed markers of cortical neuron development, *Emx2*, *Pax6* and *Tbr2*. These molecules are necessary for rostro-caudal patterning in the mouse brain (Bishop et al., 2000; Gulisano et al., 1996; Hamasaki et al., 2004; Muzio and Mallamaci, 2003). They also expressed *Bhlhb5* critical for sensory area development in the brain (Joshi et al., 2008). During cortical development, layer VI neurons are formed first followed by layers V, IV, III and II (Greig et al., 2013). It is probable that these fibroblast-derived cells also exhibit patterns observed in cortical development. We have previously shown that iPSC-derived cortically fated neurons express *Tbr1* *in vitro* and also *in vivo* after transplantation (Tornerio et al., 2013). Studies on human ESC-derived cortical neurons (Espuny-Camacho et al., 2013) have also reported that differentiation follows the cortical development pattern. Thus, a complex phenotype such as cortical neurons could be derived by forced expression of sets of transcription factors.



## CONCLUDING REMARKS AND PROSPECTS

Stroke is a major disabling disease and till date there are no therapies to enhance the functional recovery. Rehabilitation helps in improving the plastic response within the brain after stroke, but is not sufficient to improve the recovery in many stroke patients. Enhancing endogenous neurogenesis and neuronal replacement by cell transplantation could potentially promote functional recovery after stroke. In this thesis, we have analyzed three different approaches for neuronal replacement after stroke. We have identified a negative regulator of neurogenesis, a new subset of cells that contribute to adult OB neurogenesis and transcription factor combinations that can convert fibroblasts to cortical neurons.

Lnk is an inhibitory regulator of NSPC proliferation after stroke in rodents. Since we observed Lnk expression in human SVZ, it is probable that Lnk also suppresses increased neurogenesis after stroke in humans. Targeting Lnk could potentially be developed into a therapeutic strategy for stroke by increasing neurogenesis. Given its expression in other cells such as microglia, neuroblasts and endothelial cells in the brain, it is probable that Lnk acts upon these cells too. Even though we show that Lnk inhibits post-stroke SVZ neurogenesis by inhibiting the IGF1 – AKT pathway, it is probable that other pathways may also be involved. Recovery in the post-stroke brain involves functional and structural reorganization such as plasticity of surviving neurons, reorganization of vasculature in the peri-infarct region and SVZ, and activation of the immune system. Lnk might also influence these mechanisms. It remains to be explored in more detail how manipulation of Lnk can be used to stimulate post-stroke recovery.

We show that FoxJ1 expressing cells contribute to olfactory bulb neurogenesis in the adult rat brain. FoxJ1 is predominantly a marker of ependymal cells and we also found a subset of astrocytes that express FoxJ1. Rats are useful animal models for many neurodegenerative diseases and electroporation enabled us to trace the lineage of the LVW cells in intact and stroke-damaged brain, thereby overcoming the lack of genetic models. This opens up the possibility to do genetic labeling or ablation studies in rat LVW that was not possible before. We show that the OB neurons arise from the FoxJ1+ cells, but whether these FoxJ1+ cells are a finite source of these OB neurons is not known. It remains to be established whether these OB neurons arise from the FoxJ1+/GFAP+ subset. The functional significance of neurogenesis from FoxJ1+ cells in intact brain is not known. We found that ependymal cells delaminated from the LVW and migrated into the striatum after stroke. How these cells delaminate and what factors stimulate their migration is not known. The contribution to striatal neurogenesis from the FoxJ1+ cells is minor, whether it could be enhanced for functional recovery remains to be studied.

Cortical strokes cause more disability than other types of stroke and we tried to formulate a replacement strategy using patient's own somatic cells. Cortical neurons expressing Satb2 and Ctip2 were derived from human fetal lung fibroblasts by using defined sets of transcription factors. These cells also express markers for cortical layer and area development. It remains to be explored if these cells would integrate into the intact cortex and also reconstruct the stroke-damaged cortex. Improvement of conversion efficiency of these neurons to achieve



the numbers necessary for transplantation is needed. The peri-infarct region is characterized by glial scar and increased gliogenesis. Whether these astrocytes can be converted into cortical neurons *in situ* by gene delivery, leading to improved functional recovery, also needs to be explored.

To conclude, the results in this thesis raise the possibility that inhibition of Lnk, identified here as a negative regulator of neurogenesis from the brain's own neural stem cells, and intracortical transplantation of fibroblast-derived cortical neurons could be developed into novel therapeutic strategies for stroke in the future.

# EXPERIMENTAL PROCEDURES

## Animals

All experimental procedures were approved by the Malmö-Lund Ethical Committee for the use of laboratory animals and were conducted in accordance with European Union directive on the subject of animal rights. Animals were housed under 12 h light/12 h dark cycle with unlimited access to food and water. Wild type and transgenic mice (*paper I*) as well as adult male Wistar rats (*paper II*) were used for the thesis.

## Experimental models

### *Stroke*

Mice and rats were subjected to experimental stroke by intraluminal filament MCAO model. In mice (*paper I*), under isoflurane anesthesia, the left external carotid was ligated and temporary sutures were placed around the common and internal carotid arteries. An 8–0 monofilament (Alcon) coated with silicone was advanced through the internal carotid artery until it blocked the blood flow in the MCA. For reperfusion, mice were re-anesthetized after 40 min of occlusion and the filament was removed (Iosif et al., 2008). Rats were subjected to experimental stroke one week after electroporation (*paper II*), on the electroporated side (ipsilateral side). Briefly, under isoflurane anesthesia, the right common carotid artery (CCA) and its proximal branches were isolated. The CCA and external carotid artery (ECA) were ligated, while the internal carotid artery (ICA) was temporarily occluded using a metal microvessel clip. A nylon monofilament was inserted and advanced through the CCA and ICA until resistance was felt, past the origin of middle cerebral artery (MCA). The nylon filament was carefully removed 30 min or 2 h after the start of occlusion, and ECA was ligated permanently (Kokaia et al., 1998). Both the mice and rats were allowed to awake in between occlusion and reperfusion.

### *Status Epilepticus*

Mice were subjected to experimental epilepsy by induction of SE (*paper I*), a severe form of epilepsy observed in humans. Mice were anesthetized and implanted with a stimulating/recording electrode (Plastics One) unilaterally into ventral hippocampal CA1–CA3 region. Ten days later, mice were subjected to electrically induced SE. Animals received 1 h of suprathreshold stimulation consisting of 10 s trains of 1ms biphasic square wave pulses at a frequency of 50 Hz. Stimulation was interrupted for 1 min every 10 min to allow for electroencephalographic (EEG) recording and measurement of afterdischarges (MacLab). After ending the stimulation, all mice exhibited self-sustained, continuous ictal activity in EEG, and associated motor behavioral convulsions. Epileptic activity was arrested with pentobarbital at 2 h after stimulation offset (Iosif et al., 2006).

## *In vivo electroporation*

Electroporation is a safe method of transfecting plasmids to cells. It is used extensively *in vitro* for transfecting cells in culture and *in utero* for transfecting developing embryo. Recently it has been shown to be effective in transfecting the cells lining the lateral ventricular wall (LVW) of adult brain (Barnabe-Heider et al., 2008). The mechanism behind electroporation is widely accepted to involve transient pore formation in the cell membrane when electric potential is applied across the cell (Neumann et al., 1999). Since the electric current travels from negative electrode to positive electrode, the plasmid DNA molecule, being negatively charged, travels along with the electrical current through the pores in the cell membrane towards the positive electrode. Since the electric potential is applied for a fraction of a second and pores are transient, the plasmid DNA is retained inside the cells. For *in vivo* transfection of LVW and subventricular zone (SVZ) cells, CUY21 EDIT square wave electroporator (NEPA Gene, Japan) with platinum electrodes was used. Plasmids were injected stereotactically into the lateral ventricle of adult mice or rats under isoflurane anesthesia. Immediately after injection, electrodes were applied with electro-conductive gel, the cathode (positive electrode) on the same side as the intraventricular injection (ipsilateral side) and anode (negative electrode) on the opposite side (contralateral side) to complete the circuit. Three to five electrical pulses (100-200 V/cm) of 50 ms duration with 1Hz frequency were applied. Electrodes were then taken off followed by removal of the injection needle. The wound was closed with Histoacryl (Braun, Germany) and animals were allowed to recover. Grooming behavior, food and water intake and signs of burns or scars at the site of electrode application were monitored (*paper I and II*).

## **Human tissue**

Human tissue was obtained from Skåne University Hospitals at Lund and Malmö according to the guidelines approved by the Lund-Malmö Ethical Committee. Dead aborted human fetuses 6- to 9-weeks post-conception were obtained from Skåne University hospital, Lund, with informed consent. Ganglionic eminences (GE) (*paper I*) and cortex (*paper III*) were sub-dissected from brain while the lung tissue (*paper III*) was microdissected, under a stereomicroscope (Leica, Germany) in ice-cold hibernation medium. Anonymized adult human brain samples of SVZ from routine clinical diagnostic autopsy for neuropathological investigations were obtained in accordance with national regulations.

## **Dissection and cell culture**

### *Neurospheres*

Neurospheres were generated from SVZ of intact 8- to 12-week-old mice (*paper I*). The SVZ was microdissected and single cell suspension was made by enzymatic digestion of the tissue. Human fetal GE was subdissected from the fetal brain, mechanically dissociated and

cultured. For neurosphere formation, cells were grown at clonal density (10 cells/ $\mu$ l) in DMEM/F12 with N2, 2  $\mu$ g/ml heparin, 10 ng/ml leukemia inhibitory factor (Sigma-Aldrich), and 20 ng/ml epidermal growth factor (EGF) and basic fibroblast growth factor (bFGF). Differentiation of neurospheres was performed by plating on poly-D-lysine (PDL)/laminin coated coverslips and culturing in medium without growth factors but with N2 and 1% fetal bovine serum (FBS).

### *Human fetal lung fibroblasts*

Human fetal lung fibroblasts (henceforth referred to as HEFL2) were isolated from aborted human fetuses 7-9 week post-conception (*paper III*). After removal of the central nervous system and spinal ganglia, the trachea was exposed and resected at the bifurcation of the mainstem bronchi. The sub-dissected pulmonary tissue was digested with 0.25% trypsin-EDTA (Sigma-Aldrich) at 37°C for 10 minutes and manually triturated to reach a single cell suspension. Cells were plated onto 0.1% gelatin (Sigma-Aldrich) in DMEM 4.5g/L glucose supplemented with 2mM glutamax and 10% FBS (all from Life technologies) and were passaged after reaching confluence with 0.25% trypsin-EDTA. The HEFL2 cell line was used for neuronal differentiation protocols after passage 3 to avoid contamination from neural tissue and other cell types (Pfisterer et al., 2011a; Vierbuchen et al., 2010).

### *Astrocytes*

Fetal cortical-derived astrocytes obtained from dead aborted fetuses were plated onto PDL/Laminin (both 10 $\mu$ g/mL; Sigma-Aldrich) coated glass coverslips at a density of 20,000 cells / cm<sup>2</sup> and maintained in DMEM:F12 (1:1 ratio) supplemented with 10% FBS until co-culturing (*paper III*).

## **Plasmid constructs**

Plasmids were constructed by standard molecular methods, extracted with phenol-chloroform and resuspended in 10mM Tris chloride for all experiments. The promoter sequences for cloning were amplified by PCR from genomic DNA (*paper I*) or from transgenic targeting plasmid construct (*paper II*). The Lnk promoter sequence was mutated with Quickchange Lightning Kit (Agilent) (*paper I*).

### *piggyBac transposon*

*piggyBac* transposon was used for genetic labeling of the LVW cells (*paper III*) (Cadinanos and Bradley, 2007). Transposons are mobile DNA elements, remnants of viral genome. They can integrate or excise genetic sequences within inverted terminal repeats into / from the host genome. The mobility of transposons is conferred by the transposase enzyme that transposes the genetic sequences, within inverted terminal repeats of a helper plasmid or genetic sequence, into the genome. The transposase enzyme can also cleave out the inserted sequence thus having the ability to safely insert and excise a given sequence from the genome (Elick et al., 1996; Elick et al., 1997). This enables genetic labeling of target cells that could be

removed later. Transposons are found in all organisms and have the ability to insert even 100 kb sequence into the host genome. piggyBac system was identified in grasshoppers and later shown to be effective in targeting mammalian genomes including human genome (Jang and Behringer, 2007; Li et al., 2011; Lu et al., 2009; Wilson et al., 2007). The advantages of this system over other transposon system are its highly efficient genomic targeting and that there is no footprint upon excision in the genome (Wu et al., 2006). We used the helper plasmid mPB with donor plasmid pPB-UbC-eGFP (UbC: Ubiquitin-C promoter; kind gift of Dr. A. Bradley, The Wellcome Trust Sanger Institute, UK). The FoxJ1 promoter from pTg-FoxJ1-CreGFP was amplified by PCR and cloned to generate the pPB-FoxJ1-eGFP and pFoxJ1-mPB donor and helper plasmids, respectively. pPB-UbC-DsRed2 was generated from pPB-UbC-eGFP by replacing the eGFP with DsRed2 from pIRES2-DsRed2 (Clontech). The plasmids pFoxJ1-mPB with pPB-UbC-DsRed2 or pPB-FoxJ1-eGFP were used in 1:1 ratio for fate mapping experiments in intact animals. The plasmids pFoxJ1-mPB with pPB-UbC-DsRed2 and pPB-FoxJ1-eGFP were electroporated in 1:1:1 ratio, (total DNA concentration 36 µg) for fate mapping experiments following MCAO.

## Viral vectors

All viral vectors were handled in a class III biosafety laboratory and a multiplicity of infection (MOI) of 2 for each vector was used for all viral transductions. All the vectors were packaged with the VSVG capsid and produced at the Viral Vector Core, Lund Stem Cell Center, Lund University.

### *Retroviral vectors*

Murine Stem Cell Virus (MSCV) encoding IRES-GFP (MIG) or IRES-GFP-Lnk (MIG-Lnk), kindly provided by Drs. J. Lodish and W. Tong, were used to transduce neurospheres (*paper I*). Transduced cells were allowed to grow for an additional 5 days in vitro before cell sorting.

### *Lentiviral vectors*

The lentiviral vectors encoding the murine transcription factors Ascl1, Brn2 and Myt1l immediately downstream of the tetracycline response element (TRE) in FUW-TRE--WPRE and pLD-puro-2A-M2rtTA-TcVA encoding the tetracycline activator expression were obtained from Addgene (*paper III*). The human consensus coding sequences (CCDS) of cortical transcriptino factors were codon optimized for human / mouse codon usage and synthesized commercially (Genscript, CA, USA). The synthesized sequences were cloned into pBOB-TRE--WPRE, immediately downstream of the TRE. pHG-hSynl-GFP, a kind gift of Dr. Cecilia Lundberg, was used to transduce reprogrammed neurons for further analysis (*paper III*).

## Neuronal reprogramming

The human fetal lung fibroblasts cells were transduced with pLD-puro-2A-M2rtTA-TcVA

(Mak et al., 2010) (paper III). The transduced cells were selected with puromycin (1 µg/ml; Life Technologies) for rtTA expression and grown to 90% confluency before passage. Upon passage, cells were transduced with the respective pool of transcription factor expressing lentiviral vectors in microcentrifuge tubes and plated 1 h later along with the viral particles. The cells were plated to 6 well plates with coverslips, T75 flasks or T175 flasks and 48 h after transduction the medium was changed to DMEM / 10% FBS with Doxycycline (2.5 µg/ml; Sigma-Aldrich) to induce expression of transcription factors. The expression of the pool of transcription factors would induce cortical neurons from fibroblasts (iCtx). Three days after doxycycline was added, the medium was changed to neuronal induction medium (iCtx medium: Neurobasal medium, 2% B27 without vitamin A, 0.5mM Glutamine and 10U/ml Pen/Strep; all from Life Technologies). The medium was changed once every 3 days. Laminin (1µg/ml; Life Technologies) was added to the culture medium every week to improve neuronal cell survival. Cells were cultured in iCtx medium until 25 - 38 days after doxycycline addition (dpi). Cells were transduced with pHG-hSyn1-GFP lentiviral vector one week before electrophysiology analysis and cell sorting. Cells grown on gelatin-coated coverslips were used for electrophysiology and immunocytochemistry.

## Cell Sorting and flow cytometry

### *Fluorescence-activated cell sorting*

Neurospheres were passaged and SVZ tissue dissociated by enzymatic digestion (*paper I*) while the iCtx cells were trypsinized (*paper III*) to make a single cell suspension. Cells were washed and resuspended in PBS containing 30mM glucose and 5% FBS (*paper I*) or L-15 medium containing 0.1% IgG-free BSA (Jackson ImmunoResearch), 10U/ml DNaseI (Sigma-Aldrich) and 1:10 ratio Citrate-Phosphate-Dextrose (Sigma-Aldrich) (FACS buffer; *paper III*). iCtx cells were incubated at room temperature with APC conjugated CD44 antibody (BD Biosciences)(*paper III*). Suspensions were passed through a 70µm filter and cells were sorted using fluorescence-activated cell sorting (FACS) Aria (BD Biosciences). The sorted cells were directly sorted to medium or lysis buffer for cell culture or RNA extraction, respectively.

### *Flow cytometry*

WT and Lnk<sup>-/-</sup> neurospheres were pulsed with BrdU and cells harvested at 3, 6, and 10 h (*paper I*). The harvested cells were labeled with FITC conjugated BrdU antibody and propidium iodide. Cells were analyzed for BrdU incorporation and cell cycle with the data acquired on an LSR II flow cytometer, with Diva software version 6.0 (Becton Dickinson). The cytometry data was further analyzed on FlowJo software version 9.4.9 (TreeStar).

### *Magnetic activated cell sorting*

One week after transduction with Syn1-GFP lentiviral vector, iCtx cells were trypsinized

and made into single cell suspension in FACS buffer (*paper III*). The cells were incubated with the human anti-fibroblast antibody labeled with magnetic beads (Miltenyi Biotec, Germany) according to manufacturer's instructions and the cell suspension was passed through a separation column held within a magnet, by gravity. The column was removed from the magnet and remaining cells eluted out by gravity with FACS buffer. The sorted cells were used for co-culture experiments with human astrocytes.

## RNA extraction and Quantitative PCR

RNA was isolated from sub-dissected SVZ primary tissue, expanded neurospheres, (*paper I*), fluorescence-activated sorted cells (*paper I and III*) or cultured cells (*paper III*). Total RNA was isolated using RNAeasy kit (Qiagen) with DNase treatment according to manufacturer's instructions. RNA was reverse transcribed using oligoDT primers and superscript-II (Invitrogen) (*paper I*) or iScript advanced cDNA synthesis kit (Bio-Rad)(*paper III*). For some experiments (*paper III*) the cDNA was pre-amplified with TaqMan PreAmp Master mix (Applied Biosystems) according to manufacturer instructions. PCR was performed with cDNA using the primers listed in Table 1 for some of the experiments (*paper I*). Quantitative-PCR (Q-PCR) was performed with TaqMan gene expression assays (Applied Biosystems) on iQ5 thermal cycler (Bio-Rad) (*paper I and III*), or with 48.48 microfluidic chips BioMark HD (Fluidigm, CA) (*paper III*). cDNA input was normalized to GAPDH (*paper I*) or GAPDH,  $\beta$ Actin, UBC, TBP, HPRT and YWHKAZ (*paper III*). Relative gene expression was calculated using the  $\Delta\Delta C_t$  method (*paper I*).  $\Delta C_t$  values were log transformed to base of 2 ( $\log_2 2^{-\Delta C_t}$ ) and then z-scores, calculated by subtracting the  $\Delta C_t$  of a sample with the mean  $\Delta C_t$  value of all samples for a given gene and dividing it by the corresponding standard deviation, were calculated. The z-score values or  $\log_2 2^{-\Delta C_t}$  were visualized as a heat-map using Gene-E software (<http://www.broadinstitute.org/cancer/software/GENE-E/index.html>). Hierarchical clustering was performed for initial set of qPCR results in Gene-E by average linkage of one minus Pearson's correlation values to identify the clustering of transcription factor combinations with respect to cortical markers (*paper III*). The list of TaqMan assays used in this thesis are listed in Table 2.

## Transcription factor binding site prediction

DNA sequences for SH2B3 (LNK) (*paper I*) corresponding to conserved regions between the human, mouse, rat, horse, and dog species upstream of the 5' UTR and including a part of the 5' UTR were retrieved using the VISTA Browser (<http://pipeline.lbl.gov>) (Mayor et al., 2000) and the UCSC Genome Browser (<http://genome.ucsc.edu/>) (Kent et al., 2002). These regions corresponded to the following positions in the human genome: Human SH2B3 (March 2006) chr12: 110319473–110319590, chr12: 110320884 –110320980, chr12: 110325532–110325770, and chr12: 110327105–110328342. We applied MotifScanner 3.1.1 (<http://med.kuleuven.be/lcb/toucan/help/WebServices/motifscanner.htm>) (Aerts et al., 2005) for binding site prediction using matrix files from TRANSFAC and murine and human conserved noncoding sequences as background model (*paper I*).

## Chromatin immunoprecipitation PCR

Neurospheres were fixed with 1% formaldehyde for 10 min at RT, and the process stopped by the addition of glycine at a final concentration of 0.0125 M for 5 min. Cells were washed and resuspended in 1 ml lysis buffer with protease inhibitor mixture and 1 mM serine protease inhibitor, phenylmethylsulfonyl fluoride (PMSF; Roche). Sonication was performed with a Branson 450 probe, yielding DNA fragments with a bulk size of 50–300bp. The debris-cleared lysate was frozen at -80°C. Magnetic Dynabeads (Invitrogen) were incubated (1:2) with blocking buffer (Lonza) containing 1 mg/ml herring sperm DNA (Promega) and 1 mg/ml bovine serum albumin (Sigma-Aldrich) overnight at 4°C on a rotator. For each reaction, beads in 100 µl of the slurry-blocking buffer mix were separated with a DynaMag magnet (Invitrogen) and incubated for 20 min at room with rabbit anti-STAT1, normal rabbit IgG (both from Cell Signaling Technology), or H<sub>2</sub>O, diluted in 200 µl Binding & Washing Buffer (Invitrogen), and washed with Binding & Washing Buffer. For immunoprecipitation, 100 µl cell lysate was diluted with 900 µl of radioimmunoprecipitation assay buffer containing PMSF and protease inhibitor, and incubated with the prepared beads for 2 h. Beads were washed three times with Washing buffer. Cross-linking was reversed by resuspending beads in 250 µl digestion buffer containing 10mM Tris, 10 mM EDTA and 1% SDS, addition of proteinase K to a concentration of 50µg/ml, and incubation for 2 h at 68°C. 10 µl of raw lysate were diluted in 240 µl digestion buffer and treated equally to the ChIP samples, served as input control. Finally, the material was phenol-chloroform extracted and ethanol precipitated. DNA was resuspended in 50 µl of nuclease-free water and 2 µl was used as a template for PCR (*paper I*).

## Enzyme Linked Immunosorbent Assay (ELISA)

Neurosphere cells were grown adherent in 96-well plates in expansion medium until semiconfluent, when growth factors were withdrawn overnight. Cells were stimulated with 10 ng/ml EGF and 10 ng/ml bFGF, 50 ng/ml IGF1, or 5 U/ml EPO for 10 min. Levels of total and phosphorylated ERK1/2 and AKT were analyzed using FACE in-cell Western phospho ELISAs (Active Motif) (*paper I*).

## BrdU administration

BrdU (10 µM) was used for cell cycle analysis by flow cytometry (*paper I*). BrdU (50 mg/kg, i.p.), dissolved in PBS, was given four times with a 2 h interval to intact mice and at 7 d after SE, and animals were killed 2 h thereafter (*paper I*). Mice subjected to stroke received BrdU once daily for 7 d after the insult and were killed the following day (*paper I*). For BrdU label retention studies, mice received two daily BrdU injections during day 9 and 10 after stroke and were killed 8 weeks later (*paper I*). Rats received BrdU (50 mg/kg, i.p.) one week after stroke, twice daily for seven days and were killed 4 weeks after the last injection (*paper II*).



## Immunostaining

### *Immunocytochemistry in vitro*

Neurospheres or cells attached to coverslips or slides were fixed with 4% paraformaldehyde (PFA), pre-incubated in potassium PBS (KPBS) with 0.025% Triton X-100 and 5% serum for 1 h, and incubated with primary antibodies overnight at 4°C (Table 3). After washing, cells were incubated with respective secondary antibodies conjugated to Cy2 / Alexa488, Cy3 and Cy5 / Alexa 657 fluorophores. Biotinylated secondary antibodies were incubated with fluorophore conjugated streptavidin. The cells that had been labeled with biocytin from electrophysiology experiments (*paper III*) were stained with Streptavidin conjugated Alexa488 or Alexa647 along with the secondary antibodies. After rinsing thrice with PBS, cell nuclei were stained with Hoechst 33342 for 10 min in room temperature. Coverslips were mounted on glass slides or slides were coverslipped with PVA-Dabco after rinsing with distilled water (*papers I and III*).

### *Immunohistochemistry in vivo*

Animals were perfused transcardially and brains were removed, post-fixed overnight in PFA, and cryopreserved in 20% sucrose solution. Free-floating frozen sections (30 µm thick) were cut on a sliding microtome (Leica, Germany) in the coronal plane. Sections were pre-incubated in 0.25% Triton X-100 in phosphate-buffered saline (PBS) containing 5% serum for 1 h. Sections were then incubated with primary antibody overnight at 4°C (Table 3). Sections were washed with PBS and then incubated for 2 h with secondary antibodies conjugated to Cy2 / Alexa488, Cy3 and Cy5 / Alexa 657 fluorophores at room temperature (RT). Sequential staining was carried out if two of the primary antibodies were made in the same species. For BrdU staining, DNA was denatured in 1N HCl for 30 min at 65°C (*paper I*) or in 2N HCl at RT for 1 or 2 h (*paper I and II respectively*), washed three times with PBS, and incubated overnight with rat anti-BrdU (1:100; OBT0300, AbD Serotec, Germany). Biotin-conjugated donkey anti-rat (1:200; Jackson ImmunoResearch) secondary antibody was incubated for 2 h at RT followed by incubation with Streptavidin-conjugated Alexa-647 for 2 h at RT (*paper III*). The sections were washed with PBS after secondary antibody incubation and stained with Hoechst 33342 for 10 minutes at RT. The sections were washed three times, mounted onto gelatin coated slides, air-dried and coverslipped with PVA-DABCO.

### *TUNEL staining*

TUNEL staining was performed using the in situ cell death detection kit (Roche) according to manufacturer's instructions. The sections were rinsed three times in ice-cold PBS and then in ice-cold permeabilization solution (0.1% Triton X-100 in 0.1% Sodium Citrate) for 2 min. Sections were rinsed three times in PBS and incubated in TUNEL enzyme and label mix for 1 h at 37 °C. After rinsing once in PBS, nuclei were stained with Hoechst 33342 (10 µg/ml). The sections were then rinsed twice in PBS, air-dried and coverslipped with PVA-DABCO mounting medium (*paper I and II*).

## Microscopical analysis

Immunofluorescence was analyzed in Olympus BX61 (Olympus, Germany) epifluorescence microscope, images were obtained with CellSens Dimension software (Olympus, Germany). Cells were counted in four evenly spaced coronal sections that contained the SVZ (*paper I*) or LVW (*paper II*), or eight evenly spaced olfactory bulb sections (*paper II*). Confocal images were obtained for co-localization of markers on Leica TCS-SP2 (*paper I and II*) or Zeiss LSM 780 (*paper III*). Cell counts on MCAO lesioned animals were also performed on confocal images (*paper II*).

## Conversion efficiency

*In vitro* quantification was done in at least 20 regions of interest from three different coverslips per condition in an epifluorescence microscope (*paper III*). The total number of MAP2/ $\beta$ III Tubulin positive cells with neuronal morphology was quantified 25-30 days after infection. Transduction efficiency was calculated as previously described (Pang et al., 2011; Vierbuchen et al., 2010). Briefly, average number of neuronal cells present in 20 randomly selected 20x visual fields was estimated. The area of 20x visual field was then used to determine MAP2+/ $\beta$ III Tubulin+ cell density in the entire dish. This number was divided by the number of plated cells during transduction to get the percentage of starting population of cells that acquired neuronal-like characteristics (*paper III*).

## Cell soma size

The cell soma size was measured from 60 cells randomly chosen from 3 coverslips for each condition in CellSens imaging software (Olympus, Japan). The images were obtained on using 20X objective and the outline of the cell body was delineated manually within the software. The area covered by the outline corresponding to the cell soma size was calculated by the software and used for analyzing the average cell soma size in the given transcription factor combinations (*paper III*).

## Pyramidal morphology index

The pyramidal morphology index (PMI) was defined as the ratio between the width of the largest process and the total number of processes crossing a sampling circle of fixed diameter ( $\varnothing 25 \mu\text{m}$ ) (Hand et al., 2005) in CellSens imaging software (Olympus, Japan). To determine the PMI, at least 60 cells per condition were randomly chosen. The number of processes crossing the sampling circle was counted and the width of the widest neurite was measured. The index was calculated for the multipolar cells that had more than two processes. These cells were used for analyzing the percentage of cells that were multipolar in a given condition (*paper III*).

## Immunoreactivity assay

Assessment of MAP2/ $\beta$ III Tubulin immunoreactivity was done using cellSens Dimension (Olympus, Japan) imaging software, to determine the neurite density. 20 images of randomly chosen regions in three representative coverslips were acquired. Using a 10X objective, 15

fields were chosen randomly in 3 coverslips for each condition. In each coverslip, areas of MAP2/ $\beta$ III Tubulin immunoreactivity were identified using defined representative ranges of threshold for specific signal. Using these defined parameters, the images of each area were analyzed by software, which calculated the total area covered by the specific immunopositive signal ( $X$  in  $\mu\text{m}^2$ ). The number of MAP2/ $\beta$ III Tubulin cells ( $y$ ) that had Hoechst 33342 positive nuclei in each field was counted. This number was then used to calculate the immunoreactivity area per cell ( $X/y$ ,  $\mu\text{m}^2$  / cell) to get a measure of the neurite density. The cell density in a given coverslip was calculated using  $y/X$  (cells /  $\mu\text{m}^2$ ) (*paper III*).

## Electrophysiological analysis

Whole-cell patch-clamp recordings were performed with EPC10 double patch clamp amplifier (HEKA Elektronik, Germany) using PatchMaster software (HEKA Elektronik, Germany) for data acquisition. Cells grown on coverslips were transferred to the recording chamber mounted on Olympus BX62 upright microscope. The cells for recording were identified using a 40X water immersion lens with infrared differential interference contrast optics, while cells labeled with SynI-GFP virus were visualized with fluorescence optics. The coverslip was constantly perfused (1ml/min) with carbogenated artificial cerebral spinal fluid (aCSF, in mM : 119 NaCl, 2.5 KCl, 1.3  $\text{MgSO}_4$ , 2.5  $\text{CaCl}_2$ , 26  $\text{NaHCO}_3$ , 1.25  $\text{NaH}_2\text{PO}_4$ , and 25 glucose, pH 7.2-7.4, 295-300 mOsm) at 34°C. Recording pipettes were filled with intracellular solution containing (in mM): 122.5 KGLu, 12.5 KCl, 10.0 HEPES, 8.0 NaCl, 2.0 MgATP, and 0.3  $\text{Na}_2\text{GTP}$  for recordings of intrinsic properties, 135 CsGlu, 10 HEPES, 10 NaCl, 1  $\text{MgCl}_2$ , 2 MgATP, and 0.4  $\text{Na}_2\text{GTP}$  for recordings of evoked excitatory post synaptic currents (EPSC), and 135 CsCl, 10 HEPES, 10 NaCl, 2 MgATP, and 0.3 NaGTP for recordings of evoked inhibitory post synaptic currents (IPSC). Intracellular solutions had a pH of 7.2-7.4, an osmolarity of 285-295 mOsm and had a resistance of 2.5-9.5 M $\Omega$ . Biocytin (2-4 mg/ml) was added to the internal solution prior to recording for post-hoc identification of the recorded cell. Voltage values were not corrected for the liquid junction potential, which were 13.82 mV, 15.55 mV and 5.10 mV for KGLu-, CsGlu- and CsCl-based internal solutions, respectively. Voltage- and current-clamp recordings were used for the electrophysiological characterization. Sodium and potassium currents were evoked by a series of 200 ms long voltage steps (from -70 mV to +40 mV in 10 mV steps) and their sensitivity to respectively, 1  $\mu\text{M}$  TTX, (Tocris, UK) and 2-10 mM TEA (Tocris) were determined. A series of current steps (0-200 pA in 10 pA steps) lasting 500 ms were performed from a membrane potential of around -70mV (current was injected when needed to keep the membrane potential around -70mV) to determine the cells ability to generate action potentials. EPSCs and IPSCs were evoked by puff application (0.5-0.75 bar) of 100 mM glutamate or 100 mM GABA lasting 0.5-1 s using a pneumatic drug ejection system (PDES-02DE-2, NPI electronic, Germany). AMPA and NMDA receptors were blocked by 5  $\mu\text{M}$  NBQX and 50  $\mu\text{M}$  D-APV, respectively. GABA receptors were blocked by 0.1 mM PTX. Data was analyzed offline with FitMaster and IgorPro (*paper III*).

## **Statistical analysis**

Statistical analysis and graphs were made in Prism software (GraphPad, CA, USA). Comparisons were performed using paired or unpaired Student's t-test, ratio paired Student's t-test, two-way ANOVA followed by Scheffé's post hoc test, Tukey's multiple comparisons test or Uncorrected Fisher's LSD, Kruskal-Wallis test followed by Dunn's multiple comparison test and Wilcoxon matched-pairs signed rank test where appropriate. Differences are considered significant with  $P < 0.05$ .

**Table 1.** RT-PCR primer sequences

<b>Gene</b>	<b>Forward primer</b>	<b>Reverse primer</b>
<b>Mouse</b>		
Lnk	ATGCCTGACAACCTCTACAC	ATTCACAGTCTGCCTCTCT
$\beta$ -actin	ACACTGTGCCCATCTACGAG	CTAGAAGCACTTGC GG TGCA
<b>Human</b>		
Lnk	CCAGGAGAAGCACCTTGTT	CGGACAGACATCATCTTTGAGG
$\beta$ -actin	AATCTGGCACCCACCTTCTACA	CGACGTAGCACGCTTCTCCTTA

Sequences are in 5' to 3' orientation

**Table 2.** List of Taqman Gene Expression Assays

Paper I		Paper III	
Gene	Assay Id	Gene	Assay Id
bFGF	Mm01289199_m1	ACTB	Hs99999903_m1
$\beta$ -Catenin	Mm00483039_m1	BCL11B	Hs00256257_m1
Bmi1	Mm03053308_g1	(CTIP2)	
C-KIT	Mm00445212_m1	BHLHE22	Hs01084964_s1
E2F1	Mm00432941_g1	(BHLHB5)	
EGF	Mm01316968_m1	COL1A1	Hs00164004_m1
EGFR	Mm00433023_m1	CUX2	Hs00390035_m1
Epo	Mm01202754_g1	EMX2	Hs00244574_m1
EpoR	Mm00438760_m1	EOMES	Hs00172872_m1
FGFR1	Mm01219185_g1	(TBR2)	
FGFR2	Mm00438941_m1	FEZF2	Hs00215962_m1
GAPDH	Mm99999915_g1	FOXP1	Hs01850784_s1
Gli1	Mm00494646_g1	GAD1	Hs01065893_m1
Hes1	Mm01342805_m1	GAPDH	Hs03929097_g1
Hes5	Mm00439311_g1	HPRT1	Hs01003267_m1
IGF1	Mm00439560_m1	LMO4	Hs01086790_m1
IGF1R	Mm00802841_m1	MAPT	Hs00902194_m1
Lnk	Mm00493156_m1	OTX1	Hs00951099_m1
SCF	Mm00442972_m1	PAX6	Hs00240871_m1
SP1	Mm00489039_m1	SATB2	Hs00392652_m1
STAT1	Mm00439531_m1	SLC17A7	Hs00220404_m1
STAT3	Mm01219775_m1	SLC17A6	Hs00220439_m1
TGF $\alpha$	Mm00446231_m1	SOX5	Hs00753050_s1
TLX	Mm00455855_m1	TBP	Hs00427621_m1
		TBR1	Hs00232429_m1
		UBC	Hs00824723_m1
		YWHAZ	Hs03044281_g1

**Table 3.** List of primary antibodies

Antigen	Made in	Provider	Dilution	Marker of
BrdU*	Rat	AbD Serotec	1:100	Nuclei of dividing cells
$\beta$ -Catenin <sup>#</sup>	Rabbit	Millipore	1:200	
$\beta$ III tubulin	Rabbit	Covance	1:2000	Immature and mature neurons
	Mouse	Covance	1:500	
CD31	Rat	BD Bioscience	1:200	Endothelial cells
Collagen I	Rabbit	Abcam	1:50	Fibroblasts
Ctip2	Rat	Abcam	1:2000	Subcortical projection neurons
DCX	Goat	Santa Cruz Biotechnology	1:400	Migrating neuroblasts
FoxJ1	Mouse	eBioscience	1:1000	Ciliated cells (ependymal cells in this thesis)
GFAP	Rabbit	Zymed	1:400	Astrocytes
	Mouse	Sigma	1:400	
	Mouse	DAKO	1:500	
GFP	Chicken	Abcam	1:3000	
			( <i>in vitro</i> )	
			1:5000 ( <i>in vivo</i> )	
Iba1	Rabbit	Wako	1:1000	Microglia
Ki-67	Rabbit	Leica	1:200	Mitotic nuclei
Lnk	Goat	Santa Cruz Biotechnology	1:100	
MAP2a/b	Mouse	Sigma-Aldrich	1:500	Mature neurons
Nestin	Mouse	Millipore	1:200	Neural stem cells
NeuN	Mouse	Millipore	1:400	Neuronal nuclei
pH3	Rabbit	Millipore	1:400	Mitotic nuclei
RFP	Rabbit	Abcam	1:1000	
Satb2	Mouse	Abcam	1:20	Cortico-cortical projection neurons
S100 $\beta$	Rabbit	Millipore	1:200	Ependymal cells and a subtype of astrocytes
Sox2	Rabbit	Millipore	1:500	Neural stem cells
	Mouse	R&D systems	1:50	
vGlut 1/2	Rabbit	Synaptic Systems	1:200	Glutamatergic neurons
Vimentin	Mouse	DAKO	1:200	Ependymal cells

\* Antigen retrieval with 1 or 2 N HCl is required

<sup>#</sup> Antigen retrieval with Citrate buffer is required

## REFERENCES

- Abrous, D.N., Koehl, M., and Le Moal, M. (2005). Adult Neurogenesis: From Precursors to Network and Physiology. *Physiol Rev* 85, 523-569.
- Aerts, S., Van Loo, P., Thijs, G., Mayer, H., de Martin, R., Moreau, Y., and De Moor, B. (2005). TOUCAN 2: the all-inclusive open source workbench for regulatory sequence analysis. *Nucleic Acids Research* 33, W393-W396.
- Alcamo, E.A., Chirivella, L., Dautzenberg, M., Dobрева, G., Fariñas, I., Grosschedl, R., and McConnell, S.K. (2008). *Satb2* Regulates Callosal Projection Neuron Identity in the Developing Cerebral Cortex. *Neuron* 57, 364-377.
- Altman, J., and Das, G.D. (1965). Autoradiographic and histological evidence of postnatal hippocampal neurogenesis in rats. *The Journal of Comparative Neurology* 124, 319-335.
- Alvarez-Buylla, A., Seri, B., and Doetsch, F. (2002). Identification of neural stem cells in the adult vertebrate brain. *Brain Res Bull* 57, 751-758.
- Androutsellis-Theotokis, A., Leker, R.R., Soldner, F., Hoepfner, D.J., Ravin, R., Poser, S.W., Rueger, M.A., Bae, S.-K., Kittappa, R., and McKay, R.D.G. (2006). Notch signalling regulates stem cell numbers in vitro and in vivo. *Nature* 442, 823-826.
- Arlotta, P., Molyneaux, B.J., Chen, J., Inoue, J., Kominami, R., and Macklis, J.D. (2005). Neuronal Subtype-Specific Genes that Control Corticospinal Motor Neuron Development In Vivo. *Neuron* 45, 207-221.
- Arvidsson, A., Collin, T., Kirik, D., Kokaia, Z., and Lindvall, O. (2002). Neuronal replacement from endogenous precursors in the adult brain after stroke. *Nat Med* 8, 963-970.
- Azim, E., Shnider, S.J., Cederquist, G.Y., Sohur, U.S., and Macklis, J.D. (2009). *Lmo4* and *Clim1* progressively delineate cortical projection neuron subtypes during development. *Cereb Cortex* 19 Suppl 1, i62-69.
- Barnabe-Heider, F., Meletis, K., Eriksson, M., Bergmann, O., Sabelstrom, H., Harvey, M.A., Mikkers, H., and Frisen, J. (2008). Genetic manipulation of adult mouse neurogenic niches by in vivo electroporation. *Nat Meth* 5, 189-196.
- Bengzon, J., Kokaia, Z., Elmér, E., Nanobashvili, A., Kokaia, M., and Lindvall, O. (1997). Apoptosis and proliferation of dentate gyrus neurons after single and intermittent limbic seizures. *Proceedings of the National Academy of Sciences* 94, 10432-10437.
- Bersenev, A., Wu, C., Balcerak, J., and Tong, W. (2008). *Lnk* controls mouse hematopoietic stem cell self-renewal and quiescence through direct interactions with JAK2. *The Journal of Clinical Investigation* 118, 2832-2844.
- Bishop, K.M., Goudreau, G., and O'Leary, D.D.M. (2000). Regulation of Area Identity in the Mammalian Neocortex by *Emx2* and *Pax6*. *Science* 288, 344-349.
- Brill, M.S., Ninkovic, J., Winpenny, E., Hodge, R.D., Ozen, I., Yang, R., Lepier, A., Gascon, S., Erdelyi, F., Szabo, G., et al. (2009). Adult generation of glutamatergic olfactory bulb interneurons. *Nat Neurosci* 12, 1524-1533.
- Buza-Vidas, N., Antonchuk, J., Qian, H., Månsson, R., Luc, S., Zandi, S., Anderson, K., Takaki, S., Nygren, J.M., Jensen, C.T., et al. (2006). Cytokines regulate postnatal hematopoietic stem cell expansion: opposing roles of thrombopoietin and LNK. *Genes & Development* 20, 2018-2023.
- Cadinanos, J., and Bradley, A. (2007). Generation of an inducible and optimized piggyBac transposon system. *Nucleic Acids Res* 35, e87--e87-.
- Caiazzo, M., Dell'Anno, M.T., Dvoretzkova, E., Lazarevic, D., Taverna, S., Leo, D., Sotnikova, T.D., Menegon,



- A., Roncaglia, P., Colciago, G., et al. (2011). Direct generation of functional dopaminergic neurons from mouse and human fibroblasts. *Nature* 476, 224-227.
- Carlen, M., Meletis, K., Goritz, C., Darsalia, V., Evergren, E., Tanigaki, K., Amendola, M., Barnabe-Heider, F., Yeung, M.S., Naldini, L., et al. (2009). Forebrain ependymal cells are Notch-dependent and generate neuroblasts and astrocytes after stroke. *Nat Neurosci* 12, 259-267.
- Cayuso, J., Ulloa, F., Cox, B., Briscoe, J., and Martí, E. (2006). The Sonic hedgehog pathway independently controls the patterning, proliferation and survival of neuroepithelial cells by regulating Gli activity. *Development* 133, 517-528.
- Cederquist, G.Y., Azim, E., Shnider, S.J., Padmanabhan, H., and Macklis, J.D. (2013). Lmo4 Establishes Rostral Motor Cortex Projection Neuron Subtype Diversity. *The Journal of Neuroscience* 33, 6321-6332.
- Chen, B., Wang, S.S., Hattox, A.M., Rayburn, H., Nelson, S.B., and McConnell, S.K. (2008). The Fezf2-Ctip2 genetic pathway regulates the fate choice of subcortical projection neurons in the developing cerebral cortex. *Proceedings of the National Academy of Sciences* 105, 11382-11387.
- Chiasson, B.J., Tropepe, V., Morshead, C.M., and van der Kooy, D. (1999). Adult Mammalian Forebrain Ependymal and Subependymal Cells Demonstrate Proliferative Potential, but only Subependymal Cells Have Neural Stem Cell Characteristics. *J Neurosci* 19, 4462-4471.
- Danilov, A.I., Gomes-Leal, W., Ahlenius, H., Kokaia, Z., Carlemalm, E., and Lindvall, O. (2009). Ultrastructural and antigenic properties of neural stem cells and their progeny in adult rat subventricular zone. *Glia* 57, 136-152.
- Danilov, A.I., Kokaia, Z., and Lindvall, O. (2012). Ectopic ependymal cells in striatum accompany neurogenesis in a rat model of stroke. *Neuroscience* 214, 159-170.
- Delavaran, H., Sjunnesson, H., Arvidsson, A., Lindvall, O., Norrving, B., van Westen, D., Kokaia, Z., and Lindgren, A. (2013). Proximity of brain infarcts to regions of endogenous neurogenesis and involvement of striatum in ischaemic stroke. *European Journal of Neurology* 20, 473-479.
- Deng, W., Aimone, J.B., and Gage, F.H. (2010). New neurons and new memories: how does adult hippocampal neurogenesis affect learning and memory? *Nat Rev Neurosci* 11, 339-350.
- Ding, S., Wu, X., Li, G., Han, M., Zhuang, Y., and Xu, T. (2005). Efficient transposition of the piggyBac (PB) transposon in mammalian cells and mice. *Cell* 122, 473-483.
- Doetsch, F. (2003a). The glial identity of neural stem cells. *Nat Neurosci* 6, 1127-1134.
- Doetsch, F. (2003b). A niche for adult neural stem cells. *Curr Opin Genet Dev* 13, 543-550.
- Doetsch, F., Caille, I., Lim, D., Garcia-Verdugo, J., and Alvarez-Buylla, A. (1999). Subventricular zone astrocytes are neural stem cells in the adult mammalian brain. *Cell* 97, 703-716.
- Durukan, A., and Tatlisumak, T. (2007). Acute ischemic stroke: Overview of major experimental rodent models, pathophysiology, and therapy of focal cerebral ischemia. *Pharmacology Biochemistry and Behavior* 87, 179-197.
- Elick, T.A., Bauser, C.A., and Fraser, M.J. (1996). Excision of the piggyBac transposable element in vitro is a precise event that is enhanced by the expression of its encoded transposase. *Genetica* 98, 33-41.
- Elick, T.A., Lobo, N., and Fraser Jr, M.J. (1997). Analysis of the cis-acting DNA elements required for piggyBac transposable element excision. *Mol Gen Genet* 255, 605-610.
- Espuny-Camacho, I., Michelsen, Kimmo A., Gall, D., Linaro, D., Hasche, A., Bonnefont, J., Bali, C., Orduz, D., Bilheu, A., Herpoel, A., et al. (2013). Pyramidal Neurons Derived from Human Pluripotent Stem Cells Integrate Efficiently into Mouse Brain Circuits In Vivo. *Neuron* 77, 440-456.

- Fink, A.J., Englund, C., Daza, R.A.M., Pham, D., Lau, C., Nivison, M., Kowalczyk, T., and Hevner, R.F. (2006). Development of the Deep Cerebellar Nuclei: Transcription Factors and Cell Migration from the Rhombic Lip. *The Journal of Neuroscience* 26, 3066-3076.
- Franco, S.J., Gil-Sanz, C., Martinez-Garay, I., Espinosa, A., Harkins-Perry, S.R., Ramos, C., and Müller, U. (2012). Fate-Restricted Neural Progenitors in the Mammalian Cerebral Cortex. *Science* 337, 746-749.
- Frey, J.L. (2005). Recombinant Tissue Plasminogen Activator (rtPA) for Stroke: The Perspective at 8 Years. *The Neurologist* 11, 123-133.
- Gage, F.H. (2002). Neurogenesis in the Adult Brain. *Journal of Neuroscience* 22, 612-613.
- Gleason, D., Fallon, J.H., Guerra, M., Liu, J.C., and Bryant, P.J. (2008). Ependymal stem cells divide asymmetrically and transfer progeny into the subventricular zone when activated by injury. *Neuroscience* 156, 81-88.
- Gotz, M., and Huttner, W.B. (2005). The cell biology of neurogenesis. *Nat Rev Mol Cell Biol* 6, 777-788.
- Gregory, G.D., Miccio, A., Bersenev, A., Wang, Y., Hong, W., Zhang, Z., Poncz, M., Tong, W., and Blobel, G.A. (2010). FOG1 requires NuRD to promote hematopoiesis and maintain lineage fidelity within the megakaryocytic-erythroid compartment. *Blood* 115, 2156-2166.
- Greig, L.C., Woodworth, M.B., Galazo, M.J., Padmanabhan, H., and Macklis, J.D. (2013). Molecular logic of neocortical projection neuron specification, development and diversity. *Nat Rev Neurosci* 14, 755-769.
- Gulisano, M., Broccoli, V., Pardini, C., and Boncinelli, E. (1996). Emx1 and Emx2 Show Different Patterns of Expression During Proliferation and Differentiation of the Developing Cerebral Cortex in the Mouse. *European Journal of Neuroscience* 8, 1037-1050.
- Hamasaki, T., Leingärtner, A., Ringstedt, T., and O'Leary, D.D.M. (2004). EMX2 Regulates Sizes and Positioning of the Primary Sensory and Motor Areas in Neocortex by Direct Specification of Cortical Progenitors. *Neuron* 43, 359-372.
- Hand, R., Bortone, D., Mattar, P., Nguyen, L., Heng, J.I.-T., Guerrier, S., Boutt, E., Peters, E., Barnes, A.P., Parras, C., et al. (2005). Phosphorylation of Neurogenin2 Specifies the Migration Properties and the Dendritic Morphology of Pyramidal Neurons in the Neocortex. *Neuron* 48, 45-62.
- Hevner, R.F., Shi, L., Justice, N., Hsueh, Y.-P., Sheng, M., Smiga, S., Bulfone, A., Goffinet, A.M., Campagnoni, A.T., and Rubenstein, J.L.R. (2001). Tbr1 Regulates Differentiation of the Preplate and Layer 6. *Neuron* 29, 353-366.
- Huttner, W.B., and Brand, M. (1997). Asymmetric division and polarity of neuroepithelial cells. *Current Opinion in Neurobiology* 7, 29-39.
- Ihrie, Rebecca A., and Álvarez-Buylla, A. (2011). Lake-Front Property: A Unique Germinal Niche by the Lateral Ventricles of the Adult Brain. *Neuron* 70, 674-686.
- Iosif, R.E., Ahlenius, H., Ekdahl, C.T., Darsalia, V., Thored, P., Jovinge, S., Kokaia, Z., and Lindvall, O. (2008). Suppression of stroke-induced progenitor proliferation in adult subventricular zone by tumor necrosis factor receptor 1. *J Cereb Blood Flow Metab* 28, 1574-1587.
- Iosif, R.E., Ekdahl, C.T., Ahlenius, H., Pronk, C.J.H., Bonde, S., Kokaia, Z., Jacobsen, S.-E.W., and Lindvall, O. (2006). Tumor Necrosis Factor Receptor 1 Is a Negative Regulator of Progenitor Proliferation in Adult Hippocampal Neurogenesis. *The Journal of Neuroscience* 26, 9703-9712.
- Jacquet, B.V., Muthusamy, N., Sommerville, L.J., Xiao, G., Liang, H., Zhang, Y., Holtzman, M.J., and Ghashghaei, H.T. (2011). Specification of a Foxj1-Dependent Lineage in the Forebrain Is Required for Embryonic-to-Postnatal Transition of Neurogenesis in the Olfactory Bulb. *The Journal of Neuroscience* 31, 9368-9382.

- Jacquet, B.V., Salinas-Mondragon, R., Liang, H., Therit, B., Buie, J.D., Dykstra, M., Campbell, K., Ostrowski, L.E., Brody, S.L., and Ghashghaei, H.T. (2009). FoxJ1-dependent gene expression is required for differentiation of radial glia into ependymal cells and a subset of astrocytes in the postnatal brain. *Development* 136, 4021-4031.
- Jang, C.-W., and Behringer, R.R. (2007). Transposon-Mediated Transgenesis in Rats. *Cold Spring Harb Protoc* 2007, pdb.prot4866–pdb.prot4866-.
- Järlestedt, K., Rousset, C.I., Faiz, M., Wilhelmsson, U., Ståhlberg, A., Sourkova, H., Pekna, M., Mallard, C., Hagberg, H., and Pekny, M. (2010). Attenuation of Reactive Gliosis Does Not Affect Infarct Volume in Neonatal Hypoxic-Ischemic Brain Injury in Mice. *PLoS ONE* 5, e10397.
- Jiang, W., Gu, W., Brännström, T., Rosqvist, R., and Wester, P. (2001). Cortical Neurogenesis in Adult Rats After Transient Middle Cerebral Artery Occlusion. *Stroke* 32, 1201-1207.
- Jimenez, A.J., Garcia-Verdugo, J.M., Gonzalez, C.A., Batiz, L.F., Rodriguez-Perez, L.M., Paez, P., Soriano-Navarro, M., Roales-Bujan, R., Rivera, P., Rodriguez, S., et al. (2009). Disruption of the neurogenic niche in the subventricular zone of postnatal hydrocephalic hyh mice. *Journal of neuropathology and experimental neurology* 68, 1006-1020.
- Jin, K., Wang, X., Xie, L., Mao, X.O., Zhu, W., Wang, Y., Shen, J., Mao, Y., Banwait, S., and Greenberg, D.A. (2006). Evidence for stroke-induced neurogenesis in the human brain. *Proceedings of the National Academy of Sciences* 103, 13198-13202.
- Johansson, C.B., Momma, S., Clarke, D.L., Risling, M., Lendahl, U., and Frisén, J. (1999). Identification of a Neural Stem Cell in the Adult Mammalian Central Nervous System. *Cell* 96, 25-34.
- Joshi, P.S., Molyneaux, B.J., Feng, L., Xie, X., Macklis, J.D., and Gan, L. (2008). Bhlhb5 Regulates the Postmitotic Acquisition of Area Identities in Layers II-V of the Developing Neocortex. *Neuron* 60, 258-272.
- Kamei, N., Kwon, S.-M., Alev, C., Ishikawa, M., Yokoyama, A., Nakanishi, K., Yamada, K., Horii, M., Nishimura, H., Takaki, S., et al. (2010). Lnk Deletion Reinforces the Function of Bone Marrow Progenitors in Promoting Neovascularization and Astroglialosis Following Spinal Cord Injury. *STEM CELLS* 28, 365-375.
- Kent, W.J., Sugnet, C.W., Furey, T.S., Roskin, K.M., Pringle, T.H., Zahler, A.M., Haussler, and David (2002). The Human Genome Browser at UCSC. *Genome Research* 12, 996-1006.
- Kim, J., Su, S.C., Wang, H., Cheng, A.W., Cassady, J.P., Lodato, M.A., Lengner, C.J., Chung, C.-Y., Dawlaty, M.M., Tsai, L.-H., et al. (2011). Functional Integration of Dopaminergic Neurons Directly Converted from Mouse Fibroblasts. *Cell Stem Cell* 9, 413-419.
- Kobayashi, T., Ahlenius, H., Thored, P., Kobayashi, R., Kokaia, Z., and Lindvall, O. (2006). Intracerebral Infusion of Glial Cell Line-Derived Neurotrophic Factor Promotes Striatal Neurogenesis After Stroke in Adult Rats. *Stroke* 37, 2361-2367.
- Kohwi, M., Osumi, N., Rubenstein, J.L.R., and Alvarez-Buylla, A. (2005). Pax6 Is Required for Making Specific Subpopulations of Granule and Periglomerular Neurons in the Olfactory Bulb. *J Neurosci* 25, 6997-7003.
- Kohwi, M., Petryniak, M.A., Long, J.E., Ekker, M., Obata, K., Yanagawa, Y., Rubenstein, J.L.R., and Alvarez-Buylla, A. (2007). A Subpopulation of Olfactory Bulb GABAergic Interneurons Is Derived from Emx1- and Dlx5/6-Expressing Progenitors. *J Neurosci* 27, 6878-6891.
- Kokaia, Z., Andsberg, G., Martinez-Serrano, A., and Lindvall, O. (1998). Focal cerebral ischemia in rats induces expression of p75 neurotrophin receptor in resistant striatal cholinergic neurons. *Neuroscience* 84, 1113-1125.

- Kreuzberg, M., Kanov, E., Timofeev, O., Schwaninger, M., Monyer, H., and Khodosevich, K. (2010). Increased subventricular zone-derived cortical neurogenesis after ischemic lesion. *Experimental Neurology* 226, 90-99.
- Kwon, S.-M., Suzuki, T., Kawamoto, A., Ii, M., Eguchi, M., Akimaru, H., Wada, M., Matsumoto, T., Masuda, H., Nakagawa, Y., et al. (2009). Pivotal Role of Lnk Adaptor Protein in Endothelial Progenitor Cell Biology for Vascular Regeneration. *Circulation Research* 104, 969-977.
- Ladewig, J., Mertens, J., Kesavan, J., Doerr, J., Poppe, D., Glaue, F., Herms, S., Wernet, P., Kogler, G., Muller, F.J., et al. (2012). Small molecules enable highly efficient neuronal conversion of human fibroblasts. *Nat Methods* 9, 575-578.
- Li, M.A., Turner, D.J., Ning, Z., Yusa, K., Liang, Q., Eckert, S., Rad, L., Fitzgerald, T.W., Craig, N.L., and Bradley, A. (2011). Mobilization of giant piggyBac transposons in the mouse genome. *Nucleic Acids Res* 39, e148.
- Liesz, A., Zhou, W., Mrcskó, É., Karcher, S., Bauer, H., Schwarting, S., Sun, L., Bruder, D., Stegemann, S., Cerwenka, A., et al. (2011). Inhibition of lymphocyte trafficking shields the brain against deleterious neuroinflammation after stroke. *Brain* 134, 704-720.
- Lindvall, O., Brundin, P., Widner, H., Rehnström, S., Gustavii, B., Frackowiak, R., Leenders, K., Sawle, G., Rothwell, J., Marsden, C., et al. (1990). Grafts of fetal dopamine neurons survive and improve motor function in Parkinson's disease. *Science* 247, 574-577.
- Liu, X., Li, F., Stubblefield, E.A., Blanchard, B., Richards, T.L., Larson, G.A., He, Y., Huang, Q., Tan, A.-C., Zhang, D., et al. (2012). Direct reprogramming of human fibroblasts into dopaminergic neuron-like cells. *Cell Res* 22, 321-332.
- Lu, Y., Lin, C., and Wang, X. (2009). PiggyBac transgenic strategies in the developing chicken spinal cord. *Nucleic Acids Res* 37, e141--e141-.
- Mak, A.B., Ni, Z., Hewel, J.A., Chen, G.I., Zhong, G., Karamboulas, K., Blakely, K., Smiley, S., Marcon, E., Roudeva, D., et al. (2010). A lentiviral-based functional proteomics approach identifies chromatin remodelling complexes important for the induction of pluripotency. *Molecular & Cellular Proteomics* 9, 811-823.
- Marro, S., Pang, Zhiping P., Yang, N., Tsai, M.-C., Qu, K., Chang, Howard Y., Südhof, Thomas C., and Wernig, M. (2011). Direct Lineage Conversion of Terminally Differentiated Hepatocytes to Functional Neurons. *Cell Stem Cell* 9, 374-382.
- Martí-Fàbregas, J., Romaguera-Ros, M., Gómez-Pinedo, U., Martínez-Ramírez, S., Jiménez-Xarrié, E., Marín, R., Martí-Vilalta, J.-L., and García-Verdugo, J.-M. (2010). Proliferation in the human ipsilateral subventricular zone after ischemic stroke. *Neurology* 74, 357-365.
- Mayor, C., Brudno, M., Schwartz, J.R., Poliakov, A., Rubin, E.M., Frazer, K.A., Pachter, L.S., and Dubchak, I. (2000). VISTA : visualizing global DNA sequence alignments of arbitrary length. *Bioinformatics* 16, 1046-1047.
- Mirzadeh, Z., Merkle, F.T., Soriano-Navarro, M., Garcia-Verdugo, J.M., and Alvarez-Buylla, A. (2008). Neural stem cells confer unique pinwheel architecture to the ventricular surface in neurogenic regions of the adult brain. *Cell Stem Cell* 3, 265-278.
- Molyneaux, B.J., Arlotta, P., Fame, R.M., MacDonald, J.L., MacQuarrie, K.L., and Macklis, J.D. (2009). Novel Subtype-Specific Genes Identify Distinct Subpopulations of Callosal Projection Neurons. *The Journal of Neuroscience* 29, 12343-12354.
- Molyneaux, B.J., Arlotta, P., Menezes, J.R.L., and Macklis, J.D. (2007). Neuronal subtype specification in the cerebral cortex. *Nat Rev Neurosci* 8, 427-437.

- Muzio, L., and Mallamaci, A. (2003). Emx1, Emx2 and Pax6 in Specification, Regionalization and Realization of the Cerebral Cortex. *Cerebral Cortex* 13, 641-647.
- Neumann, E., Kakorin, S., and Toensing, K. (1999). Fundamentals of electroporative delivery of drugs and genes. *Bioelectrochemistry and Bioenergetics* 48, 3-16.
- Oh, S.T., Simonds, E.F., Jones, C., Hale, M.B., Goltsev, Y., Gibbs, K.D., Merker, J.D., Zehnder, J.L., Nolan, G.P., and Gotlib, J. (2010). Novel mutations in the inhibitory adaptor protein LNK drive JAK-STAT signaling in patients with myeloproliferative neoplasms. *Blood* 116, 988-992.
- Ohira, K., Furuta, T., Hioki, H., Nakamura, K.C., Kuramoto, E., Tanaka, Y., Funatsu, N., Shimizu, K., Oishi, T., Hayashi, M., et al. (2010). Ischemia-induced neurogenesis of neocortical layer 1 progenitor cells. *Nat Neurosci* 13, 173-179.
- Ostrowski, L.E., Hutchins, J.R., Zakel, K., and O'Neal, W.K. (2003). Targeting Expression of a Transgene to the Airway Surface Epithelium Using a Ciliated Cell-Specific Promoter. *Mol Ther* 8, 637-645.
- Pang, Z.P., Yang, N., Vierbuchen, T., Ostermeier, A., Fuentes, D.R., Yang, T.Q., Citri, A., Sebastiano, V., Marro, S., Sudhof, T.C., et al. (2011). Induction of human neuronal cells by defined transcription factors. *Nature* 476, 220-223.
- Parent, J.M., Vexler, Z.S., Gong, C., Derugin, N., and Ferriero, D.M. (2002). Rat forebrain neurogenesis and striatal neuron replacement after focal stroke. *Annals of Neurology* 52, 802-813.
- Parent, J.M., Yu, T.W., Leibowitz, R.T., Geschwind, D.H., Sloviter, R.S., and Lowenstein, D.H. (1997). Dentate Granule Cell Neurogenesis Is Increased by Seizures and Contributes to Aberrant Network Reorganization in the Adult Rat Hippocampus. *The Journal of Neuroscience* 17, 3727-3738.
- Pfenninger, C.V., Roschupkina, T., Hertwig, F., Kottwitz, D., Englund, E., Bengzon, J., Jacobsen, S.E., and Nuber, U.A. (2007). CD133 Is Not Present on Neurogenic Astrocytes in the Adult Subventricular Zone, but on Embryonic Neural Stem Cells, Ependymal Cells, and Glioblastoma Cells. *Cancer Res* 67, 5727-5736.
- Pfisterer, U., Kirkeby, A., Torper, O., Wood, J., Nelander, J., Dufour, A., Björklund, A., Lindvall, O., Jakobsson, J., and Parmar, M. (2011a). Direct conversion of human fibroblasts to dopaminergic neurons. *Proceedings of the National Academy of Sciences* 108, 10343-10348.
- Pfisterer, U., Wood, J., Nihlberg, K., Hallgren, O., Bjermer, L., Westergren-Thorsson, G., Lindvall, O., and Parmar, M. (2011b). Efficient induction of functional neurons from adult human fibroblasts. *Cell Cycle* 10, 3311-3316.
- Pruszkowski, J., Ludwig, W., Blak, A., Alavian, K., and Isacson, O. (2009). CD15, CD24, and CD29 Define a Surface Biomarker Code for Neural Lineage Differentiation of Stem Cells. *STEM CELLS* 27, 2928-2940.
- Qiang, L., Fujita, R., Yamashita, T., Angulo, S., Rhinn, H., Rhee, D., Doege, C., Chau, L., Aubry, L., Vanti, W.B., et al. (2011). Directed Conversion of Alzheimer's Disease Patient Skin Fibroblasts into Functional Neurons. *Cell* 146, 359-371.
- Quiñones-Hinojosa, A., Sanai, N., Soriano-Navarro, M., Gonzalez-Perez, O., Mirzadeh, Z., Gil-Perotin, S., Romero-Rodriguez, R., Berger, M.S., Garcia-Verdugo, J.M., and Alvarez-Buylla, A. (2006). Cellular composition and cytoarchitecture of the adult human subventricular zone: A niche of neural stem cells. *The Journal of Comparative Neurology* 494, 415-434.
- Reynolds, B., and Weiss, S. (1992). Generation of neurons and astrocytes from isolated cells of the adult mammalian central nervous system. *Science* 255, 1707-1710.
- Riquelme, Patricio A, Elodie Drapeau, and Fiona Doetsch. "Brain Micro-Ecologies: Neural Stem Cell Niches in the Adult Mammalian Brain." *Philosophical Transactions of the Royal Society B: Biological Sciences* 363, no. 1489 (2008): 123-137.

- Rouaux, C., and Arlotta, P. (2010). Fezf2 directs the differentiation of corticofugal neurons from striatal progenitors in vivo. *Nat Neurosci* 13, 1345-1347.
- Sansom, S.N., and Livesey, F.J. (2009). Gradients in the Brain: The Control of the Development of Form and Function in the Cerebral Cortex. *Cold Spring Harbor Perspectives in Biology* 1.
- Saridey, S.K., Liu, L., Doherty, J.E., Kaja, A., Galvan, D.L., Fletcher, B.S., and Wilson, M.H. (2009). PiggyBac Transposon-based Inducible Gene Expression In Vivo After Somatic Cell Gene Transfer. *Mol Ther* 17, 2115-2120.
- Sawamoto, K., Wichterle, H., Gonzalez-Perez, O., Cholfin, J.A., Yamada, M., Spassky, N., Murcia, N.S., Garcia-Verdugo, J.M., Marin, O., Rubenstein, J.L.R., et al. (2006). New Neurons Follow the Flow of Cerebrospinal Fluid in the Adult Brain. *Science* 311, 629-632.
- Schinder, A.F., and Gage, F.H. (2004). A Hypothesis About the Role of Adult Neurogenesis in Hippocampal Function. *Physiology* 19, 253-261.
- Seita, J., Ema, H., Oeohara, J., Yamazaki, S., Tadokoro, Y., Yamasaki, A., Eto, K., Takaki, S., Takatsu, K., and Nakauchi, H. (2007). Lnk negatively regulates self-renewal of hematopoietic stem cells by modifying thrombopoietin-mediated signal transduction. *Proceedings of the National Academy of Sciences* 104, 2349-2354.
- Shimamura, K., and Rubenstein, J.L. (1997). Inductive interactions direct early regionalization of the mouse forebrain. *Development* 124, 2709-2718.
- Son, E.Y., Ichida, J.K., Wainger, B.J., Toma, J.S., Rafuse, V.F., Woolf, C.J., and Eggan, K. (2011). Conversion of Mouse and Human Fibroblasts into Functional Spinal Motor Neurons. *Cell Stem Cell* 9, 205-218.
- Srinivasan, K., Leone, D.P., Bateson, R.K., Dobрева, G., Kohwi, Y., Kohwi-Shigematsu, T., Grosschedl, R., and McConnell, S.K. (2012). A network of genetic repression and derepression specifies projection fates in the developing neocortex. *Proceedings of the National Academy of Sciences* 109, 19071-19078.
- Stecca, B., and Ruiz i Altaba, A. (2009). A GLI1-p53 inhibitory loop controls neural stem cell and tumour cell numbers. *The EMBO Journal* 28, 663-676.
- Sun, F., Wang, X., Mao, X., Xie, L., and Jin, K. (2012a). Ablation of Neurogenesis Attenuates Recovery of Motor Function after Focal Cerebral Ischemia in Middle-Aged Mice. *PLoS ONE* 7, e46326.
- Sun, L., Lee, J., and Fine, H.A. (2004). Neuronally expressed stem cell factor induces neural stem cell migration to areas of brain injury. *The Journal of Clinical Investigation* 113, 1364-1374.
- Sun, X., Zhang, Q.-W., Xu, M., Guo, J.-J., Shen, S.-W., Wang, Y.-Q., and Sun, F.-Y. (2012b). New striatal neurons form projections to substantia nigra in adult rat brain after stroke. *Neurobiology of Disease* 45, 601-609.
- Takagi, Y., Harada, J., Chiarugi, A., and Moskowitz, M.A. (2002). STAT1 is activated in neurons after ischemia and contributes to ischemic brain injury. *J Cerebr Blood F Met* 22, 1311-1318.
- Takahashi, K., and Yamanaka, S. (2006). Induction of Pluripotent Stem Cells from Mouse Embryonic and Adult Fibroblast Cultures by Defined Factors. *Cell* 126, 663-676.
- Takaki, S., Sauer, K., Iritani, B.M., Chien, S., Ebihara, Y., Tsuji, K.-i., Takatsu, K., and Perlmutter, R.M. (2000). Control of B Cell Production by the Adaptor Protein Lnk: Definition of a Conserved Family of Signal-Modulating Proteins. *Immunity* 13, 599-609.
- Takaki, S., Tezuka, Y., Sauer, K., Kubo, C., Kwon, S.-M., Armstead, E., Nakao, K., Katsuki, M., Perlmutter, R.M., and Takatsu, K. (2003). Impaired Lymphopoiesis and Altered B Cell Subpopulations in Mice Overexpressing Lnk Adaptor Protein. *The Journal of Immunology* 170, 703-710.
- Takeda, K., and Akira, S. (2000). STAT family of transcription factors in cytokine-mediated biological responses. *Cytokine & Growth Factor Reviews* 11, 199-207.

- Thored, P., Arvidsson, A., Cacci, E., Ahlenius, H., Kallur, T., Darsalia, V., Ekdahl, C.T., Kokaia, Z., and Lindvall, O. (2006). Persistent Production of Neurons from Adult Brain Stem Cells During Recovery after Stroke. *STEM CELLS* 24, 739-747.
- Thored, P., Heldmann, U., Gomes-Leal, W., Gisler, R., Darsalia, V., Taneera, J., Nygren, J.M., Jacobsen, S.-E.W., Ekdahl, C.T., Kokaia, Z., et al. (2009). Long-term accumulation of microglia with proneurogenic phenotype concomitant with persistent neurogenesis in adult subventricular zone after stroke. *Glia* 57, 835-849.
- Tiberi, L., Vanderhaeghen, P., and van den Aemele, J. (2012). Cortical neurogenesis and morphogens: diversity of cues, sources and functions. *Current Opinion in Cell Biology* 24, 269-276.
- Tong, W., Zhang, J., and Lodish, H.F. (2005). Lnk inhibits erythropoiesis and Epo-dependent JAK2 activation and downstream signaling pathways. *Blood* 105, 4604-4612.
- Tornero, D., Wattananit, S., Grønning Madsen, M., Koch, P., Wood, J., Tatarishvili, J., Mine, Y., Ge, R., Monni, E., Devaraju, K., et al. (2013). Human induced pluripotent stem cell-derived cortical neurons integrate in stroke-injured cortex and improve functional recovery. *Brain* 136, 3561-3577.
- Torper, O., Pfisterer, U., Wolf, D.A., Pereira, M., Lau, S., Jakobsson, J., Björklund, A., Grealish, S., and Parmar, M. (2013). Generation of induced neurons via direct conversion in vivo. *Proceedings of the National Academy of Sciences* 110, 7038-7043.
- Tsai, P.T., Ohab, J.J., Kertesz, N., Groszer, M., Matter, C., Gao, J., Liu, X., Wu, H., and Carmichael, S.T. (2006). A Critical Role of Erythropoietin Receptor in Neurogenesis and Post-Stroke Recovery. *The Journal of Neuroscience* 26, 1269-1274.
- VandenDriessche, T., Ivics, Z., Izsvak, Z., and Chuah, M.K.L. (2009). Emerging potential of transposons for gene therapy and generation of induced pluripotent stem cells. *Blood* 114, 1461-1468.
- Velazquez, L., Cheng, A.M., Fleming, H.E., Furlonger, C., Vesely, S., Bernstein, A., Paige, C.J., and Pawson, T. (2002). Cytokine Signaling and Hematopoietic Homeostasis Are Disrupted in Lnk-deficient Mice. *The Journal of Experimental Medicine* 195, 1599-1611.
- Vierbuchen, T., Ostermeier, A., Pang, Z.P., Kokubu, Y., Südhof, T.C., and Wernig, M. (2010). Direct conversion of fibroblasts to functional neurons by defined factors. *Nature* 463, 1035-1041.
- Wan, M., Li, Y., Xue, H., Li, Q., and Li, J. (2006). TNF- $\alpha$  Induces Lnk Expression Through PI3K-Dependent Signaling Pathway in Human Umbilical Vein Endothelial Cells. *Journal of Surgical Research* 136, 53-57.
- Wang, T.-C., Chiu, H., Chang, Y.-J., Hsu, T.-Y., Chiu, I.-M., and Chen, L. (2011). The Adaptor Protein SH2B3 (Lnk) Negatively Regulates Neurite Outgrowth of PC12 Cells and Cortical Neurons. *PLoS ONE* 6, e26433.
- Wang, Y., Jin, K., Mao, X.O., Xie, L., Banwait, S., Marti, H.H., and Greenberg, D.A. (2007). VEGF-overexpressing transgenic mice show enhanced post-ischemic neurogenesis and neuromigration. *Journal of Neuroscience Research* 85, 740-747.
- Wardlaw, J.M., Murray, V., Berge, E., del Zoppo, G., Sandercock, P., Lindley, R.L., and Cohen, G. (2012). Recombinant tissue plasminogen activator for acute ischaemic stroke: an updated systematic review and meta-analysis. *The Lancet* 379, 2364-2372.
- Wilson, M.H., Coates, C.J., and George, A.L., Jr. (2007). PiggyBac transposon-mediated gene transfer in human cells. *Mol Ther* 15, 139-145.
- Wu, S.C., Meir, Y.J., Coates, C.J., Handler, A.M., Pelczar, P., and Moisyadi, S. (2006). piggyBac is a flexible and highly active transposon as compared to sleeping beauty, Tol2, and Mos1 in mammalian cells. *Proc Natl Acad Sci USA* 103, 15008-15013.



- Yamashita, T., Sawamoto, K., Suzuki, S., Suzuki, N., Adachi, K., Kawase, T., Mihara, M., Ohsugi, Y., Abe, K., and Okano, H. (2005). Blockade of interleukin-6 signaling aggravates ischemic cerebral damage in mice: possible involvement of Stat3 activation in the protection of neurons. *Journal of Neurochemistry* 94, 459-468.
- Yan, Y.-P., Sailor, K.A., Vemuganti, R., and Dempsey, R.J. (2006). Insulin-like growth factor-1 is an endogenous mediator of focal ischemia-induced neural progenitor proliferation. *European Journal of Neuroscience* 24, 45-54.
- Zhang, R.L., Zhang, Z.G., Wang, Y., LeTourneau, Y., Liu, X.S., Zhang, X., Gregg, S.R., Wang, L., and Chopp, M. (2007). Stroke induces ependymal cell transformation into radial glia in the subventricular zone of the adult rodent brain. *J Cereb Blood Flow Metab* 27, 1201-1212.
- Zhao, C., Deng, W., and Gage, F.H. (2008). Mechanisms and Functional Implications of Adult Neurogenesis. *Cell* 132, 645-660.
- Ziv, Y., Finkelstein, A., Geffen, Y., Kipnis, J., Smirnov, I., Shpilman, S., Vertkin, I., Kimron, M., Lange, A., Hecht, T., et al. (2007). A Novel Immune-Based Therapy for Stroke Induces Neuroprotection and Supports Neurogenesis. *Stroke* 38, 774-782.