



# LUND UNIVERSITY

## The role of N-cadherin and S1P/S1P1 in pancreas development

Johansson, Jenny

2010

[Link to publication](#)

*Citation for published version (APA):*

Johansson, J. (2010). *The role of N-cadherin and S1P/S1P1 in pancreas development*. [Doctoral Thesis (compilation), Stem Cell Center]. Lund University: Faculty of Medicine.

*Total number of authors:*

1

### General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00

# The role of N-cadherin and S1P/S1P<sub>1</sub> in pancreas development

**Jenny Johansson**

Stem Cell and Pancreas Developmental Biology  
Department of Laboratory Medicine Lund  
Lund Strategic Research Center for Stem Cell Biology and Cell Therapy  
Faculty of Medicine, Lund University  
Sweden

With the approval of the Lund University Faculty of Medicine, this thesis  
will be defended on June 10<sup>th</sup> 2010 at 13.15 in the Segerfalk lecture hall,  
BMC A10, Lund

**Supervisor:**

Henrik Semb

**Faculty Opponent:**

Mikael Nilsson, Ph.D.


Department of Medical Chemistry and Cell Biology  
Göteborg University, Sweden

Organization LUND UNIVERSITY  Stem Cell and Pancreas Developmental Biology Department of Laboratory Medicine Lund	Document name DOCTORAL DISSERTATION	
	Date of issue 2010-06-10	
	Sponsoring organization	
Author(s) Jenny Johansson		
Title and subtitle The role of N-cadherin and S1P/S1P1 in pancreas development		
Abstract <b>ABSTRACT</b> Organogenesis of epithelial organs requires interaction between epithelial and mesenchymal tissues. During pancreas development, mesenchyme-derived structures induce specification, growth and differentiation of pancreatic epithelial cells. For instance, notochord, endothelial cells, and pancreatic mesenchyme are essential for the development of the pancreas. N-cadherin deficient-embryos suffer from dorsal pancreas agenesis due to apoptosis of the mesenchyme surrounding the pancreatic epithelium. N-cadherin-deficient embryos expressing N-cadherin or E-cadherin ectopically in the heart rescue dorsal pancreas formation, indicating that the phenotype is secondary to cardio-vascular defects. These results suggest that circulating factors are important for dorsal pancreas development. Here we identify a sphingolipid, sphingosine-1-phosphate, to be important for dorsal pancreas formation. S1P acts on receptors (S1P1, S1P2, and S1P3) expressed in the pancreatic mesenchyme, inducing mesenchymal growth. Studies on the S1P1 receptor knockout showed a reduced size of the dorsal pancreatic bud due to decreased proliferation of Pdx1+ cells. Other epithelial organs revealed growth defects. The lung epithelium had fewer branches and the stomach was smaller. This suggests that endothelial cells are essential for early development of the pancreatic epithelium. Since N-cadherin deficient embryos die at embryonic day E10, due to the cardio-vascular phenotype, we used a conditional N-cadherin deficient-mouse which was intercrossed with a Pdx-1-Cre driven transgenic line to generate a pancreas specific knockout. Our results indicate that N-cadherin neither is important for pancreas morphogenesis nor endocrine specification. This could be due to variable recombination efficiency of the Pdx-1-Cre line during development. However, mutant $\beta$ -cells have fewer mature insulin secreting granules in the N-cadherin deleted islets indicating that N-cadherin is required for $\beta$ -cell granule turnover.		
Key words: N-cadherin, S1P, S1P <sub>1</sub> , pancreas, cell adhesion, beta cell, insulin, diabetes		
Classification system and/or index terms (if any):		
Supplementary bibliographical information:		Language English
ISSN and key title: 1652-8220		ISBN 978-91-86443-82-5
Recipient's notes	Number of pages	Price
	Security classification	

Distribution by (name and address)

I, the undersigned, being the copyright owner of the abstract of the above-mentioned dissertation, hereby grant to all reference sources permission to publish and disseminate the abstract of the above-mentioned dissertation.

Signature



Date

4/5-10

# The role of N-cadherin and S1P/S1P<sub>1</sub> in pancreas development

**Jenny Johansson**

Stem Cell and Pancreas Developmental Biology  
Department of Laboratory Medicine Lund  
Lund Strategic Research Center for Stem Cell Biology and Cell Therapy  
Faculty of Medicine, Lund University  
Sweden



**LUND**  
UNIVERSITY

ISSN 1652-8220

ISBN 978-91-86443-82-5

Lund University, Faculty of Medicine Doctoral Dissertation Series 2010:66

©2010 Jenny Johansson

Printed by Media-Tryck, Lund, Sweden

Till minne av farmor, Elin Johansson

# TABLE OF CONTENTS

<b>LIST OF PAPERS.....</b>	<b>10</b>
<b>ABBREVIATIONS.....</b>	<b>11</b>
<b>ABSTRACT.....</b>	<b>14</b>
<b>POPULÄRVETENSKAPLIG SAMMANFATTNING.....</b>	<b>15</b>
<b>INTRODUCTION.....</b>	<b>17</b>
<b>CELL ADHESION AND CADHERINS.....</b>	<b>17</b>
CELL ADHESION.....	17
CELL ADHESION MOLECULES (CAMs).....	17
THE CADHERIN SUPERFAMILY.....	18
CADHERINS IN CELLULAR JUNCTIONS.....	19
N-CADHERIN.....	19
<b>EPITHELIAL ORGAN DEVELOPMENT.....</b>	<b>21</b>
<b>MORPHOGENESIS.....</b>	<b>22</b>
CELLULAR POLARIZATION.....	22
CELL MIGRATION.....	23
CELL SORTING.....	24
CELL SURVIVAL.....	24
<b>DIFFERENTIATION.....</b>	<b>25</b>
<b>PANCREAS DEVELOPMENT.....</b>	<b>26</b>

ANATOMY AND FUNCTION.....	26
INITIATION OF THE PANCREATIC PROGRAM.....	27
PANCREATIC MORPHOGENESIS.....	28
SPECIFICATION AND DIFFERENTIATION OF THE PANCREAS.....	29
EXOCRINE LINEAGE.....	29
DUCT LINEAGE.....	29
ENDOCRINE LINEAGE.....	30
CELL ADHESION MOLECULES IN PANCREAS DEVELOPMENT.....	30
BLOODVESSELS IN PANCREAS DEVELOPMENT.....	31
<b>SPHINGOSINE-1-PHOSPHATE.....</b>	<b>32</b>
<b>SPHINGOSINE-1-PHOSPHATE RECEPTOR-1.....</b>	<b>32</b>
<b>INSULIN CELL MATURATION AND GLUCOSE HOMEOSTASIS.....</b>	<b>33</b>
<b>DIABETES.....</b>	<b>35</b>
<b>TECHNICAL CONSIDERATIONS.....</b>	<b>36</b>
THE GENE-TRAP SYSTEM.....	36
THE <i>Cre/loxP</i> -SYSTEM.....	36
RECOMBINATION VARIABILITY'S IN <i>Cre</i> TRANSGENES.....	36
<b>AIM OF THE THESIS.....</b>	<b>37</b>
<b>PAPERS IN SUMMARY.....</b>	<b>38</b>
PAPER I.....	38



INTRODUCTION.....38

RESULTS.....38

Cardiac-rescued *N-cadherin*-deficient mice develop a dorsal pancreas due to an intact circulatory system

S1P rescues the formation of the dorsal pancreas in *N-cadherin*-deficient pancreatic explants

S1P receptors in pancreas

S1P regulates dorsal pancreatic mesenchymal cell proliferation

SUMMARY.....39

PAPER II.....40

INTRODUCTION.....40

RESULTS.....40

Growth and branching morphogenesis of the pancreatic epithelium is compromised in *S1P<sub>1</sub>*-deficient embryos

*S1P<sub>1</sub>* is required for proliferation of Pdx1<sup>+</sup> pancreatic progenitors

*S1P<sub>1</sub>*-deficiency results in defective pancreatic morphogenesis in vitro

Blood vessel ablation does not mimic the *S1P<sub>1</sub>* phenotype

SUMMARY.....41

PAPER III.....42

INTRODUCTION.....42

RESULTS.....42

N-cadherin expression during pancreas development

*Pdx1-Cre*-mediated ablation of *N-cadherin*

Pancreatic morphogenesis and endocrine specification is not affected in conditional *N-cadherin*-knockout mice

N-cadherin controls insulin granule turnover

N-cadherin regulates insulin secretion

SUMMARY.....	44
<b>CONCLUDING REMARKS.....</b>	<b>44</b>
PAPER I.....	44
PAPER II.....	44
PAPER III.....	45
<b>FUTURE PERSPECTIVES.....</b>	<b>45</b>
PAPER I AND II.....	45
PAPER III.....	46
<b>ACKNOWLEDGEMENT.....</b>	<b>47</b>
<b>REFERENCES.....</b>	<b>50</b>
<b>PAPER I-III.....</b>	<b>61</b>

## LIST OF PAPERS

This thesis was based on the following papers, which will be referred to by their roman numerals (I-III):

**I. Vascular function and sphingosine-1-phosphate regulate development of the dorsal pancreatic mesenchyme.**

Edsbagge J\*, Johansson JK\*, Esni F, Luo Y, Radice GL, Semb H.

*Development. 2005 Mar;132(5):1085-92.*

\*Contributed equally to this work

**II. S1P<sub>1</sub> signaling control endoderm development.**

Fredrik Wolfhagen Sand, Christina Lorén\*, Jenny K. Johansson\*, Josefina Edsbagge, Anders Ståhlberg, Judith Magenheim, Ohad Ilovich, Eyal Mishani, Yuval Dor, Richard L. Proia, Ulf Ahlgren, and Henrik Semb.

Manuscript, \*Contributed equally to this work

**III. N-cadherin is dispensable for pancreas development but required for  $\beta$ -cell granule turnover.**

Jenny K Johansson, Ulrikke Voss, Gokul Kesavan, Igor Kostetskii, Nils Wierup, Glenn L Radice, Henrik Semb.

*Accepted in Genesis April 2010*

## ABBREVIATIONS

ADP	Adenosine di phosphate
Arx	Aristaless-related homeobox gene
ATP	Adenosine tri phosphate
B-Geo	$\beta$ -galactosidase-Neo <sup>R</sup> fusion
BMP	Bone morphogenetic protein
BrdU	Bromodeoxyuridine
CAM	Cell adhesion molecule
cAMP	Cyclic adenosine monophosphate
Cdc	Cell division cycle
cKO	Conditional knockout
Cpa1	Carboxypeptidase A1
Cre	Cyclization recombination
DE-cadherin	Drosophila E-cadherin
Dlg	Disc large
DNA	Deoxyribonucleic acid
E	Embryonic day
EC	Extracellular cadherins
E-cadherin	Epithelial cadherin
ECM	Extra cellular matrix
EGF	Epidermal growth factor
EMT	Epithelial to mesenchymal transition
EP	EPLIN
FGF	Fibroblast growth factor

GLP1	Glucagon-like-peptide 1
GLUT2	Glucose transporter 2
HAV	Histidine, Alanine, Valine
HDL	High density lipoprotein
hESC	Human embryonic stem cells
Hlxb9	Homeo box HB9
Hnf	Hepatocyte nuclear factor
Ig	Immunoglobulin
Ihh	Indian hedgehog
Isl1	Islet 1
Lgl	Lethal giant larvae
loxP	Locus of X-over P1
N-cadherin	Neural cadherin
Ngn3	Neurogenin-3
OPT	Optical Projection Tomography
Par	Partitioning defective
PALS1	Protein linked to protein associated with Lin-7
PATJ	Pals-associated tight junction protein
Pax	Paired box gene
P-cadherin	Placental cadherin
PDGF	Platelet-derived growth factor
Pdx1	Pancreatic and duodenal homeobox 1
Ptf1a	Pancreas specific transcription factor 1a
Rac1	RAS-related C3 botulinum substrate 1
R-cadherin	Retinal cadherin
RER	Rough endoplasmic reticulum

Rho-GTPase	Ras homology family of GTPase
RRP	Readily releasable pool
RT-PCR	Reverse transcription polymerase chain reaction
Shh	Sonic hedgehog
S1P	Sphingosine-1-phosphate
S1P <sub>1-5</sub>	Sphingosine-1-phosphate receptor 1-5
Sox2	(Sex determining region Y)-box 2
Sox9	(Sex determining region Y)-box 9
Stat3	Signal transducer and activator of transcription-3
TGF	Transforming growth factor
TGN	Trans-golgi network
VE-cadherin	Vascular endothelial cadherin
VEGF	Vascular endothelial growth factor

## ABSTRACT

Organogenesis of epithelial organs requires interaction between epithelial and mesenchymal tissues. During pancreas development, mesenchyme-derived structures induce specification, growth and differentiation of pancreatic epithelial cells. For instance, notochord, endothelial cells, and pancreatic mesenchyme are essential for the development of the pancreas. *N-cadherin* deficient-embryos suffer from dorsal pancreas agenesis due to apoptosis of the mesenchyme surrounding the pancreatic epithelium. *N-cadherin*-deficient embryos expressing N-cadherin or E-cadherin ectopically in the heart rescue dorsal pancreas formation, indicating that the phenotype is secondary to cardiovascular defects. These results suggest that circulating factors are important for dorsal pancreas development. Here we identify a sphingolipid, sphingosine-1-phosphate, to be important for dorsal pancreas formation. S1P acts on receptors (S1P<sub>1</sub>, S1P<sub>2</sub>, and S1P<sub>3</sub>) expressed in the pancreatic mesenchyme, inducing mesenchymal growth.

Studies on the S1P<sub>1</sub> receptor knockout showed a reduced size of the dorsal pancreatic bud due to decreased proliferation of Pdx1<sup>+</sup> cells. Other epithelial organs revealed growth defects. The lung epithelium had fewer branches and the stomach was smaller. This suggests that endothelial cells are essential for early development of the pancreatic epithelium.

Since N-cadherin deficient embryos die at embryonic day E10, due to the cardiovascular phenotype, we used a conditional *N-cadherin* deficient-mouse which was intercrossed with a *Pdx-1-Cre* driven transgenic line to generate a pancreas specific knockout. Our results indicate that N-cadherin neither is important for pancreas morphogenesis nor endocrine specification. This could be due to variable recombination efficiency of the *Pdx-1-Cre* line during development. However, mutant  $\beta$ -cells have fewer mature insulin secreting granules in the *N-cadherin* deleted islets indicating that N-cadherin is required for  $\beta$ -cell granule turnover.

## POPULÄRVETENSKAPLIG SAMMANFATTNING

$\beta$ -celler producerar insulin som frisätts när blodsockret höjs, t.ex. efter en måltid. Patienter med diabetes saknar de insulinproducerande  $\beta$ -cellerna eller har nedsatt känslighet för insulin. För de patienter som saknar insulin kan det vara svårt att reglera blodsockernivåerna i blodet trots dagliga injektioner med insulin, varför transplanterbara insulinproducerande  $\beta$ -celler skulle kunna vara av stor nytta för dessa patienter. Hittills har försök gjorts med att injicera hormonproducerande celler från donatorer till patienter. Dessa patienter har klarat sig utan insulin i några år men har sedan behövt återgå till insulininjektioner. Ett annat problem med denna behandling är bristen på donatorer. För att komma runt detta görs försök med att framställa insulinproducerande  $\beta$ -celler från humana embryonala stamceller, dvs de stamceller som ger upphov till ett människo-foster. För att driva de humana embryonala stamcellerna till att bli insulinproducerande  $\beta$ -celler behövs forskning om hur dessa celler bildas normalt i bukspottkörtelns utveckling.

Tidigt i utvecklingen bildar bukspottkörteln två utbuktningar, så kallade buddar, från tarmen. Dessa två buddar växer sedan samman och bildar bukspottkörteln som består av exokrin och endokrin vävnad samt kanaler. Den exokrina vävnaden bildar de enzymer som används vid matspjälkningen. Dessa enzymer transporteras från de exokrina cellerna via kanalerna till tarmen. Den endokrina vävnaden bildar hormonproducerande celler. Dessa celler utsöndrar hormoner till blodet, t.ex. insulin.

I den här avhandlingen har vi fokuserat på blodkärl och proteinet N-cadherins inverkan på bukspottkörtelns utveckling. I det första arbetet har vi använt en musmodell där endast en av de två buddarna bildas. Denna musmodell saknar N-cadherin i embryots alla celler. Detta leder till att embryot dör tidigt under utveckling på grund av avsaknad av blodcirkulation. Forskare har visat att signalsubstanser ifrån blodcirkulationen är viktigt för bukspottkörtelns utveckling. Här har vi visat att plasma ifrån blodet är kapabel till att bilda den budd som saknas i musmodellen. Vi har även identifierat en lipid, sphingosine-



1-phosphate (S1P), som är nödvändig för bildandet av budden.

I det andra arbetet har vi använt oss av en genmodifierad mus som saknar receptorn S1P<sub>1</sub> och har en dramatisk kärlddefekt. I denna modell bildas båda buddarna men de är mindre än buddarna hos kontrollmössen, även storleken på mage, lever och lungor är mindre.

Syftet med det tredje arbetet var att studera hur N-cadherin reglerar bukspottkörtelns utveckling. För att studera hur N-cadherin reglerar bukspottkörtelns utveckling använde vi oss av en musmodell som saknar N-cadherin endast i bukspottkörteln. Resultaten från denna studie i embryon visade att N-cadherin inte har en funktion i utvecklingen av bukspottkörteln eller i bildandet av de hormonproducerande cellerna. N-cadherin har dock visat sig ha en betydelse för utsöndrandet av insulin i vuxna möss. Detta beror på att mindre insulin bildas.


Sammanfattningsvis visar våra resultat att bukspottkörteln behöver signaler ifrån blodet, i detta fall sphingosine-1-phosphate, för att initiera bildandet av bukspottkörteln, samt att N-cadherin är viktigt för insulinfrisättningen.

# INTRODUCTION

## CELL ADHESION AND CADHERINS

### CELL ADHESION

Cell adhesion mediates adhesive contacts between cells or helps cells to adhere to the extra cellular matrix (ECM) (Figure 1). Cell adhesion molecules (CAMs) prevent the tissue from dissociating. This was first studied in the 1950's, when Townes and Holfreter dissociated amphibian embryos into single cells and mixed them. Strikingly, upon reaggregation the cells sorted out to form their particular germ-layer [1]. From this experiment the researchers formed the idea of cell adhesion molecules expressed on the cell-surface. Today cell adhesion is implicated in signal transduction, morphogenesis, and in diseases such as cancer.



**Figure 1. Adherens junction.** Cadherins form homophilic interactions with their extracellular part. The intracellular part binds to a catenin complex. The catenin complex binds to EPLIN. EPLIN is required for linkage between cadherins and F-actin, and for maintaining the circumferential actin belt. Abbreviations: B,  $\beta$ -Catenin; a,  $\alpha$ -Catenin; EP, EPLIN. The picture is modified from Nishimura et al 2009.

### CELL ADHESION MOLECULES (CAMs)


CAMs are important during both the development of new tissue and the controlled growth and turnover of adult tissue. CAMs are cell surface glycoproteins that display two different kinds of adhesive mechanisms,  $\text{Ca}^{2+}$ -dependent and non- $\text{Ca}^{2+}$ -dependent [2]. Based on their structures, CAMs are classified into

four major families; cadherins, selectins, integrins and the immunoglobulin (Ig) superfamily [3]. Cadherins and selectins belong to the  $\text{Ca}^{2+}$ -dependent group, while integrins and immunoglobins belong to  $\text{Ca}^{2+}$ -independent group. Cadherins and immunoglobins promote cell-cell adhesion, whereas integrins mediate cell-matrix interactions [4, 5]. Selectins are involved in inflammatory responses.

### THE CADHERIN SUPERFAMILY

The cadherin family of proteins can be divided into several groups. These subgroups include classical cadherins type I and type II, fat-like cadherins, seven-pass transmembrane cadherins, DCad102F-like cadherins, Desmosomal cadherins, and Protocadherins [6]. The classification of cadherins into different groups is based on the number of the extracellular cadherins repeats (EC). Each repeat consists of specific residues that bind calcium. Calcium provides structural rigidity of the extracellular domain, and is essential for the cadherins' adhesive function and protection against protease digestion (Figure 2) [7].

Classical cadherins are divided into type I and II based on the presence or absence of HAV (histidine, alanine, valine) tri-peptide within the most extracellular N-terminal repeat EC1. The EC1 domain promotes homophilic adhesive interactions of cadherins on opposing cells. The classical cadherins have five EC domains, a single-pass transmembrane part, and an intracellular part binding to the catenin complex [8].



**Figure 2. Structure of classical cadherins.** Classical cadherins have an extracellular part containing five extracellular domains, a single transmembrane part, and an intracellular part. The intracellular part binds to the catenin complex and mediates the binding

between cadherins and F-actin. Abbreviations: N, N-terminal; EC, extracellular domain; B,  $\beta$ -Catenin; a,  $\alpha$ -Catenin; EP, EPLIN. Picture modified from Ivanov et al 2001.

## **CADHERINS IN CELLULAR JUNCTIONS**

Cells are connected with multiple types of junctions; tight junctions, adherens junctions, gap junctions, and desmosomes. The cell-cell junctions in epithelial cells contain tight and adherens junctions as well as desmosomes. These three junctional components are clustered together at the apical-lateral cell-cell contacts, forming the apical “junctional complex” [9]. The adherens junctions and desmosomes located in the junctional complex are termed zonal adherens (Figure 1).

The adhesive function of cadherins requires catenins ( $\alpha$ -,  $\beta$ -,  $\gamma$ -catenin, and p120) that bind the cadherins to the actin fibers. The cytoplasmic part of classical cadherins binds to p120 [10] and  $\beta$ -catenin. P120 regulates cytoskeleton dynamics through direct modulation of the Rho GTPases [11] by inhibiting RhoA and activating Rac1 and Cdc42 [12-14].  $\beta$ -catenin binds directly to  $\alpha$ -catenin, which in turn, binds and bundles actin filaments and interact with other actin partners. Studies performed by Drees and colleagues and Yamada and colleagues showed that the cadherin- $\beta$ -catenin- $\alpha$ -catenin complex associating with F-actin can not bind directly to F-actin. Only free  $\alpha$ -catenin could link directly to F-actin. Later it has been shown that a molecule called EPLIN mediate the linkage between the cadherin- $\beta$ -catenin- $\alpha$ -catenin complex and the F-actin [15, 16] (Figure1).

## **N-CADHERIN**

N-cadherin was first identified as a cell adhesion molecule expressed in neural tissue [17], but was later found to be expressed in various non-neural tissues, such as the cardiac muscle [18], testis [19], kidney [20], and liver [21].

During embryogenesis, the expression pattern of N-cadherin changes as the cells undergo different morphological events such as gastrulation, neurogenesis, cardiogenesis, and somitogenesis [22, 23]. During gastrulation, when the cells delaminate from the primitive streak to form mesoderm, they switch from E-cadherin to N-cadherin expression. During neurogenesis the neural plate switches from E-cadherin to N-cadherin expression. After the neural tube closure, E-cadherin is lost and N-cadherin proteins are the major cadherin.

However, N-cadherin is not expressed in the neural crest cells where the neural tube closes [24]. In heart development, the mesoderm is divided into somatic mesoderm (dorsal) and the splanchnic mesoderm (ventral). N-cadherin expression is downregulated in the somatic mesoderm, but continues to be expressed in the splanchnic mesoderm [25]. When the somites are forming, N-cadherin is redistributed to the luminal side of the epithelium.


N-cadherin mutants die at E10. They express several developmental abnormalities such as malformed somites and yolk sac, undulated neural tube, dorsal pancreatic bud agenesis and severe cardiovascular defect, which is the cause of death [26, 27]. Due to the fact that N-cadherin is widely expressed in the embryo it is difficult to ascertain which phenotypes those are primary or secondary to the mutation. Therefore, the cardiac-mediated adhesion was restored by expressing either N-cadherin or E-cadherin in the heart of mutant embryos. These embryos survived one to two days longer than mutant *N-cadherin* embryos. N-cadherin and E-cadherin can restore cardiovascular phenotype and N-cadherin mediated cell-cell adhesion is required for cell survival [28].

As shown in rescue experiments, cadherins can compensate for each other. For instance, cadherin-11 is expressed in somites and may substitute for the loss of N-cadherin. There is also data showing that cadherin-11 can substitute for N-cadherin in skeletal muscle differentiation [26]. In embryonic stem cells E-cadherin and N-cadherin have been shown to compensate for each other [29].

N-cadherin is involved in many processes such as cell sorting, cell migration, and cell aggregation. In endothelial cells, N-cadherin regulates proliferation and motility by controlling VE-cadherin expression at the cell membrane [30]. In the cerebral cortex N-cadherin has been shown to be important for normal architecture of the neuroepithelial or radial glial cells and ablation of *N-cadherin* randomizes the internal structure of the cortex [31]. In endothelial cells, N-cadherin is required for pericyte recruitment [32].

## EPITHELIAL ORGAN DEVELOPMENT

Endoderm has two functions; 1) to induce the formation of several mesodermal organs (notochord, heart, and blood vessels), and 2) to construct the lining of the digestive tube and respiratory tube. The digestive tube forms the pharynx, esophagus, stomach, small intestine and colon, while the respiratory tube forms the lungs (Figure 3). Caudal to the stomach liver, gallbladder, and pancreas are formed. The digestive and respiratory tubes are surrounded by mesenchyme that induces different endodermal structures along the anterior-posterior axis. Initially the expression of specific transcription factors is established to define the areas along the endodermal tube. These areas will become specific organs (e.g. Pdx1 expression) that define the future pancreatic and duodenal region. The endodermal epithelium then differentiates into the organ specific cell types. Concomitantly with specification and differentiation the tissues undergo morphogenesis. During this process the cells create cell-cell contacts and polarity, the tissues migrate and branches, and epithelial-mesenchymal interactions occurs. Coordination between these multiple processes results in the development of epithelial organs [33].



**Figure 3. Anterior/posterior patterning of the digestive tract.** Anterior foregut gives rise to esophagus, trachea, lung, and forestomach expressing Sox2. Posterior foregut gives rise to antral stomach, dorsal bud, ventral bud and liver expressing Pdx1. Midgut and hindgut develop small and large intestine expressing Cdx2 and Cdx1. Abbreviations: lu, lung; eso, esophagus; tra, trachea; fs, forestomach; as, antral stomach; dp, dorsal pancreas; liv, liver; vp, ventral pancreas; si, small intestine; li, large intestine. Picture modified from Guney et al 2009.


## **MORPHOGENESIS**

Morphogenesis establishes appropriate environment for cells to interact and respond to signals. During epithelial organ development several cellular processes such as initiation and expansion of cell polarization, coordinated migration of cells towards particular chemo attractants, segregation of cells into specific groups, and cell survival occur. These morphogenetic processes can be triggered by signals like fibroblast growth factors (FGFs) secreted from the mesenchyme or by signaling pathways such as notch. For example, FGF has been shown to be important for lung development in mouse, while Notch has been proven to be required for tooth morphogenesis [34, 35].

### **CELLULAR POLARIZATION**

Epithelial cells have three distinct surfaces, the apical surface facing the lumen, the lateral surface facing neighboring cells and the basal surface facing the ECM. Epithelial cells are polarized and their cell polarity is regulated by intrinsic mechanisms, such as coordinated targeting of vesicles with apical and basolateral proteins to distinct surfaces. There are three groups of proteins responsible for forming and maintaining baso-apical polarity: (1) the Par complex (Par proteins and atypical protein kinase C); (2) the Scribble (Scrib) complex (Scrib, Disc large (Dlg), and Lethal giant larvae (Lgl)); (3) the Crumbs (Crb) complex (Crb, PALS1, and PATJ) [36] (Figure 4). Extrinsic factors regulate polarity by forming junctional complexes to create barriers between apical-basal surfaces and cell contacts to the ECM. This influences the vesicular transport as well as transcription of genes necessary for establishing polarity.

Adherens junction proteins like cadherins are shown to play an important role during cell polarization. Disruption of adherens junctions can alter cell polarization. This was investigated by McNeill and colleagues by transfecting non-polar L-cells with E-cadherin. The cells expressing E-cadherin redistributed  $\text{Na}^+$ ,  $\text{K}^+$  ATPase from a diffuse pattern to the regions of cell-cell contacts. This indicates a direct role of cell adhesion molecules as inducers of cell polarity [37].



**Figure 4. Cell polarization by polarity complexes.** The apical side express Crumbs complex, while zonal adherens express Par complex and the baso-lateral domain express Scribble complex.

## CELL MIGRATION

Cell migration is essential for normal development but also for responses to tissue damage and infections. Cell migration also occurs in diseases; cancer, atherosclerosis, and rheumatoid arthritis. Thus, preventing the migration of specific cell types could inhibit disease progression [38].

Migrating cells undergo changes in the cytoskeleton, cell-substrate adhesions and the extracellular matrix. Many cell types migrate as single cells, including leukocytes, and neuronal cells. Epithelial and endothelial cells migrate as sheets or group of cells. Cells moving as sheets or groups are connected via cell-cell adhesion to neighboring cells that generate strong forces, whereas single cells rely on cell-substrate interactions providing less force.

Migrating cells are guided via different chemo attractant ligands, most commonly chemokines or growth factors. For instance, angiogenic sprouting of blood vessels is dependent on vascular epidermal growth factor A (VEGF-A) for migration [39]. Cells migrate towards an area expressing higher levels of VEGF-A and will continue migrating until the sprouts migrate into a region of local hypoxia. Local hypoxia downregulate VEGF-A and cause the cells to stop migrating [40]. PDGFA is important for migration in mesodermal cells. Loss of PDGFA results in randomized migration [41]. Epithelial cells undergoing epithelial to mesenchymal transition (EMT), which is triggered by several signaling pathways; bone morphogenetic protein (BMP), FGF, and Wnt, is usually associated with the enhanced migratory behaviour. FGF induces EMT by down regulating E-cadherin and represses polarization via the transcription factor Snail.



Cadherins are not only important for EMT, but also to promote cell-cell adhesion. When cells migrate cadherin-mediated cell-cell interactions have to break and reform. DE-cadherin has been shown to be important in border cell migration in *Drosophila*. Cadherin 11 regulates filipodia and lamellipodia formation via small RhoGTPases [42]. Lamellipodia formation is dependent on fibronectin-integrin interactions. The connection between the ECM and the integrins is also essential for cell-adhesion, development of directed protrusions, and cell polarity-processes highly relevant to migratory cells.

## **CELL SORTING**

Cell sorting is a process where differentiated cells separate from one another and form clusters that constitute organized tissues. This process is mediated by classical cadherins. Cadherins are expressed to a variable degree in distinct tissues during development. For example, E-cadherin is expressed in all epithelial cells, N-cadherin in neural tissue and muscles, R-cadherin in forebrain and bone, and VE-cadherin in endothelial cells and so on. The specific expression pattern of cadherins combined with homophilic adhesive properties, may facilitate sorting of specific cell types in tissues. Experiments performed by mixing cadherin-negative L-cells transfected with E-cadherin and L-cells transfected with P-cadherin, showed that the cells were able to sort out and cluster [43]. Even mixtures of cells expressing the same cadherin but with different levels of adhesion sorted from one another [44].

## **CELL SURVIVAL**

Cellular interactions with neighboring cells induce a variety of signaling events, including survival and differentiation. Studies in different genetic models where cadherins have been deleted reveal an essential role for these molecules in cell survival. For instance, studies on R-cadherin, N-cadherin, and E-cadherin show that cadherins are important for cell survival.

Ablation of *R-cadherin* in the ureteric bud epithelium show an altered morphology and branching behavior, including increased apoptosis [45]. It is not known if the decreased cell survival is directly due to the deletion of *R-cadherin* or if it is secondary due to the changes in morphology of the epithelium.

*N-cadherin* deficient embryos exhibit a decrease in cell survival. The deficient embryos have increased apoptosis within the collapsing neural folds and somites [26]. Since *N-cadherin* deficient embryos have a cardio-vascular phenotype, the researchers performed a rescue experiment expressing N-cadherin or E-cadherin in the heart. The embryos survived longer but were still smaller and exhibited increased apoptosis, demonstrating that N-cadherin is directly involved in cell survival [28, 46].

E-cadherin is also known to mediate cell survival. Arulanandam and colleagues have shown that E-cadherin promotes cell survival via *Rac1* and *Cdc42* by activating the signal transducer and activator of transcription-3 (Stat3). *Stat3* increases both the expression levels and activates *Rac1* and *Cdc42*. Moreover, inhibitors of *Rac1* and *Cdc42* block E-cadherin-mediated *Stat3* activation, indicating that *Rac1* and *Cdc42*, are the mediators of the cadherin signal to *Stat3*.

ES-cells lacking E-cadherin undergo a dramatic induction of apoptosis. The survival signals may be mediated by an increase of *Rac1* and *Cdc42* activity, leading to *Stat3* stimulation. This could explain why Stat3 inhibition in cells expressing cell-cell adhesion can induce apoptosis [47].

## DIFFERENTIATION

The generation of cellular diversity is called differentiation. Differentiation depends on many signaling pathways, e.g. Wnt, Hedgehog, and Notch. These signaling pathways activate various transcription factors, which in turn initiate cell differentiation, for example, suppression of Wnt signaling in the anterior endoderm is required for both liver and pancreas development [48]. The mesodermal tissues surrounding the posterior foregut provide secreted signals that promote liver and pancreas. In the ventral posterior foregut, BMP secreted by the septum transversum mesenchyme and FGF1 and FGF2 produced by the cardiac mesoderm promote liver development while concomitantly suppressing the pancreatic differentiation program [46, 49-54].

Initially, transcription factors pattern the endoderm along the anterior to posterior axis. For example, *Sox2* is expressed in the anterior domain of the endoderm that will give rise to the esophagus and stomach; *Pdx1* expression is found in the antral stomach, presumptive pancreas, common bile duct, and rostral duodenum [55]; *Cdx2* is expressed in the entire postgastric epithelium

in the regions which will form intestine [56] (Figure 3). Initially, the presumptive pancreatic domain is marked by overlapping expression of Pdx1, Ptf1a, and Hlxb9/Hb9 [55, 57-59].


## **PANCREAS DEVELOPMENT**

### **ANATOMY AND FUNCTION**

The pancreas is an endoderm derived organ consisting of two major cell types: exocrine and endocrine cells (Figure 5). The exocrine part of the pancreas consists of acinar cells and ductal cells forming a highly branched duct system. The acinar cells produce digestive enzymes, such as proteases, lipases, and nucleases. These enzymes are secreted into the ductal system and transported to the intestine by the ampulla of Vater. The enzymes produced by the exocrine tissue are necessary for digesting food.

Approximately 98% of the pancreas consists of exocrine tissue; the other 2% are endocrine tissue [54]. The endocrine cells form clusters called islets of Langerhans. The islets are divided into five cell types;  $\alpha$ -,  $\beta$ -,  $\epsilon$ -,  $\delta$ -, and pancreatic-polypeptide cells. These five cell types produce the hormones; glucagon, insulin, ghrelin, somatostatin, and pancreatic polypeptide respectively. The core of the islets contains mainly  $\beta$ -cells; which are surrounded by  $\alpha$ -,  $\epsilon$ -,  $\delta$ -, and PP-cells. Insulin-producing  $\beta$ -cells make up for ~60-80% of the islet, while 15-20% are glucagon-producing  $\alpha$ -cells; 5-10% are somatostatin-producing  $\delta$ -cells, <2% are pancreatic-polypeptide-producing cells, and <1% is ghrelin-producing  $\epsilon$ -cells [60].

Endocrine cells are important for controlling blood glucose homeostasis. After food intake, blood sugar levels increases, and insulin is released from  $\beta$ -cells. This is the signal for liver, muscle, and fat to store glucose and thereby control glucose homeostasis. In contrast, glucagon secretion is stimulated at low blood-sugar levels. The release of glucagon primes  $\beta$ -cells to secrete more insulin when the glucose levels rises. Somatostatin and pancreatic polypeptide apply inhibitory effects on both endocrine and exocrine secretion [60].




**Figure 5. Anatomy of mouse pancreas.** Pancreas develops posterior to the stomach and is connected to the gut via Ampulla of water. The pancreas consists of exocrine and endocrine tissues. The exocrine part is composed of acinar cells and ducts. The endocrine part forms islets of Langerhans.

### INITIATION OF THE PANCREATIC PROGRAM

The pancreas develops from the posterior foregut endoderm expressing *Pdx1*/*lpf1*. At E9.5 in the mouse, the dorsal and ventral buds of the pancreas begin to evaginate. All cell lineages in the pancreas originate from *Pdx1*<sup>+</sup> cells [61]. Deletion of *Pdx1* results in arrested pancreas development at E10.5. Before the dorsal pancreas starts budding the endoderm is in close proximity to the notochord (Figure 6). Factors from the notochord, such as activin- $\beta$ B and FGF2, prevent the expression of the hedgehog genes. If the hedgehog genes, sonic hedgehog (*Shh*) and Indian hedgehog (*Ihh*) are not suppressed, the intestinal differentiation program will be initiated at the expense of dorsal pancreas development [62]. The ventral pancreas is not in contact with the notochord indicating that another pathway inhibits hedgehog genes in the ventral pancreas [63]. Patterning of the ventral bud relies on signals from the overlying cardiac mesenchyme and the lateral plate mesoderm.

When the dorsal pancreas starts to protrude, the notochord is no longer in close proximity to the dorsal pancreas. Instead the dorsal aortas fuse and become in direct contact with the dorsal pancreatic endoderm. Signals from the dorsal aorta have been shown to be important for differentiation and growth of the

pancreatic endoderm [64], partly by inducing expression of *Ptf1a* [65]. *Ptf1a* is expressed in the pancreas from E9.5 and *Ptf1a*<sup>+</sup> cells give rise to all endoderm derived cell lineages of the pancreas. *Ptf1a* knockouts fail to form a ventral pancreas and formation of the dorsal pancreas does not proceed beyond the initial budding.



**Figure 6. Early pancreas development.** The developing gut endoderm is in contact with the notochord at E8.5. One day later the dorsal aortas have fused and are in close vicinity to the dorsal pancreatic endoderm. At E10 the mesenchyme is surrounding the dorsal pancreatic bud. Abbreviations: da, dorsal aorta; m, mesenchyme; dp, dorsal pancreas; vp, ventral pancreas. Picture modified from Slack, J 1995.

### PANCREATIC MORPHOGENESIS

At E10 the pancreatic epithelium is surrounded by mesenchymal cells that control its growth and differentiation [66, 67]. Several mesenchymal factors are essential for proper pancreas growth. The transcription factor *Isl1* is expressed throughout the dorsal mesenchyme during bud formation. In *Isl1* mutants, the mesenchyme is absent, and there is an associated failure of exocrine cell differentiation in the dorsal but not ventral pancreas [68]. There is also a complete loss of differentiated islet cells. By adding wild type mesenchyme to pancreas *Isl1*<sup>-/-</sup> epithelium, the exocrine phenotype could be rescued. This shows that *Isl1* in the mesenchyme is important for dorsal exocrine pancreas and that *Isl1* in the endoderm is required for the generation of all endocrine cells.

## SPECIFICATION AND DIFFERENTIATION OF THE PANCREAS

Generation of a ductal tree reprints proliferation of the pancreatic epithelium and mesenchyme, branching morphogenesis, and fusion of the dorsal and ventral buds at E12.5. This branched epithelial organ contains the precursor cells for islets, acini, and ducts [69]. The tip cells in the ductal tree are marked by expression of Carboxypeptidase A1 (Cpa1), Pdx1, Ptf1a, and c-Myc [70]. At early stages, these cells contribute to all lineages by acting as multipotent progenitors. Later in development cells from the stalk in the ductal tree will give rise to endocrine- or ductal-cells, while the tip cells differentiate into acinar cells. The peak in exocrine and endocrine differentiation, begins at E13.5 in mouse, and is referred to as the secondary transition.

### EXOCRINE LINEAGE

Differentiation of the epithelium into exocrine tissue is mesenchyme dependent. Cultures with pancreas epithelium lacking mesenchyme promote endocrine cell differentiation resulting in impaired exocrine differentiation [71-73]. Mouse pancreatic epithelium exposed to the ligands (activin and transforming growth factor- $\beta$ 1 (TGF- $\beta$ 1) for TGF- $\beta$  signaling promotes development of endocrine cells, particular  $\beta$ -cells and PP-cells [74], and disrupts epithelial branching and acinar formation [75]. Furthermore, an antagonist of TGF- $\beta$  signaling (follistatin) promotes exocrine differentiation [73].


### DUCT LINEAGE

By E12.5 the specification of different lineages (exocrine, ductal, and endocrine) is initiated. The factors regulating ductal differentiation are not well understood [76]. However, several gene deletions have branching phenotypes. Inactivation of FGF10 or FGFR2IIIb leads to a reduced number of branches and decreased proliferation [77, 78]. The epidermal growth factor (EGF) family is also important for growth and branching [79]. Ablation of hepatocyte nuclear factor-6 (*Hnf-6*) demonstrate enlarged lumina and multiple cysts within the ducts as well as impaired formation of primary cilia and mislocation of the adherens junction proteins, including  $\beta$ -catenin [80]. Sox9 mutants also reveal cystic ducts and reduced exocrine differentiation [81]. Small Rho GTPases such as Cdc42 have also been demonstrated to be required for pancreatic duct differentiation. Ablation of *Cdc42* increased acinar differentiation at the expense of endocrine and duct lineage commitment [82].

## ENDOCRINE LINEAGES

All endocrine cells are derived from Neurogenin-3<sup>+</sup> cells (Ngn3). Ngn3<sup>+</sup> cells arise from Sox9 progenitor cells within the duct epithelium. These cells differentiate, delaminate and migrate to form the islets of Langerhans (Figure 7). Mutant *Ngn3* lack all endocrine cells and die shortly after birth due to hyperglycemia [83, 84]. Forced expression of Ngn3 at different time point indicates that timing is important for epithelial cells to respond to Ngn3 and differentiate into distinct endocrine cell lineages [85]. Maturation and maintenance of endocrine cells is also dependent on *Ngn3* [86].

Downstream targets of *Ngn3* are the *Pax* and *Nkx* genes, *NeuroD*, *Isl1*, and *Arx*. Together these genes maintain the balance between the different endocrine cells [87]. Mice lacking expression of *Isl1*, *Pax6*, *NeuroD*, *Nkx 2.2*, or *Nkx 6.1* exhibit defects in endocrine formation, while *Pax4* and *Arx* mutants show a shift in endocrine lineages.



**Figure 7. Endocrine differentiation and morphogenesis.** The ductal epithelium consists of endocrine progenitor cells expressing Ngn3. The Ngn3<sup>+</sup> cells delaminate from the ducts and cluster into islets of Langerhans. The islets migrate away from the ductal epithelium and are found within the exocrine tissue.

## CELL ADHESION MOLECULES IN PANCREAS DEVELOPMENT

Several cell adhesion molecules are expressed during pancreas development. For instance, E-cadherin is expressed in all pancreatic epithelial cells when the pancreas starts to form and continues to be expressed in exocrine, ductal, and endocrine lineage throughout development and adulthood. To investigate if cadherins mediate organization of endocrine cells into islets of Langerhans, a dominant negative mutant of mouse *E-cadherin* was expressed in insulin-

producing cells, resulting in displacement of both E-cadherin and N-cadherin in the  $\beta$ -cells. As a result, the architecture of the islets was disorganized.  $\beta$ -cells were found as individual cells within the tissue, whereas  $\alpha$ -cells aggregated into islets [88]. R-cadherin is also expressed in the majority of the cells in the pancreatic buds. In adults, R-cadherin is primarily expressed in the ductal system of the pancreas, on the apical side of the exocrine tissue, in intraductal endocrine cells, but low or no expression in endocrine cells [89, 90]. The *R-cadherin* mutant mice had no pancreatic phenotype (Dahl and Semb unpublished data). N-cadherin has been shown to be expressed in both dorsal and ventral bud at embryonic day E9.5 and being specifically expressed in the islets of adult pancreas [27, 91]. Ablation of *N-cadherin* resulted in a lack of the dorsal bud [27].

Another cell-cell adhesion molecule expressed in the pancreas is N-CAM. It is expressed both in pancreatic mesenchyme and epithelium. The expression becomes restricted to endocrine cells, and peripheral nerve endings and ganglia. *N-CAM* mutants develop a normal pancreas, consisting of all pancreatic cell types and islets of normal size and number, scattered within the exocrine tissue. However, lack of NCAM results in disrupted organization and morphology of the islets.  $\alpha$ -cells were distributed centrally in the islets and not in the periphery. This suggests that N-CAM regulates cell type segregation within islets. Islet cell polarity was also affected in *N-CAM* mutant mice. E-cadherin and N-cadherin were distributed subcellularly, whereas R-cadherin distribution was unaffected [92].

### **BLOOD VESSELS IN PANCREAS DEVELOPMENT**

During initiation of pancreas organogenesis the buds are in close vicinity to large vessels, the aorta dorsally and the vitelline veins ventrally. Several experiments have shown that endothelial signaling is important for pancreas development. Cleaver and colleagues removed the endothelial cell precursors of the dorsal aorta from frog embryos. Thus endocrine gene expression was inhibited, while the development of liver and neural tube proceeded normally [64, 93]. The result suggests that endothelial cells are required for endocrine development.

Recombination of prepatterned dorsal endoderm with dorsal aorta, notochord, and neural tube indicated that dorsal endoderm together with dorsal aorta can only initiate Pdx1 expression and insulin expression. Dorsal endoderm with



notochord and neural tube did not induce insulin. However, the notochord did induce Pdx1. This experiment showed that endothelial cells supply signals sufficient for insulin expression [64].

The third experiment showing that endothelial cells are important for insulin expression, was to analyze the role of VEGF-A. Overexpression of VEGF-A in the Pdx1-expressing domain resulted in increased vascularization. These endothelial cells instructed foregut cells to differentiate into pancreatic cells.

The results presented above all indicate that endothelial cells promote pancreas development. In vitro experiment on *Flk1*<sup>-/-</sup> embryos (VEGFR2) lacking endothelial cells induced Pdx1, indicating that aorta or endothelial cells are not required for initial pancreas development. The expression of Ptf1a was lost in the in the *Flk1*<sup>-/-</sup> embryos, indicating that endothelial cells are important for the appearance of the dorsal bud and to maintain Pdx1 expression [65].

## **SPHINGOSINE-1-PHOSPHATE**

Sphingosine-1-phosphate (S1P) is a metabolite formed by the phosphorylation of sphingosine by sphingosine kinase [94, 95]. S1P can be recycled back to sphingosine by S1P phosphatase or be degraded by an S1P lyase [96, 97, 98]. S1P is produced and secreted by red blood cells, platelets, monocytes, mast cells, and possibly by endothelial cells [99-101]. Since S1P is highly soluble in water it is bound to high density lipoprotein (HDL) or to albumin in plasma [102].

S1P triggers several activities in cells including proliferation, migration, cytoskeletal changes, adhesion molecule expression, and anti-apoptotic effects [103-107]. Many of these activities are produced via the interaction of S1P with G protein-coupled receptors [108].

## **SPHINGOSINE-1-PHOSPHATE RECEPTOR-1**

S1P bind to five different receptors, named S1P<sub>1-5</sub> [109]. The receptors bind to different G proteins. S1P<sub>1</sub> bind to G<sub>i/o</sub>, S1P<sub>2,3</sub> bind to G<sub>i/o</sub>, G<sub>q</sub>, G<sub>12/13</sub>, and S1P<sub>4,5</sub> bind to G<sub>i/o</sub>, G<sub>12/13</sub> and thereby mediate different signaling pathways. Many cell types express several S1P receptors. For example, endothelial cells express S1P<sub>1</sub> and S1P<sub>3</sub>, vascular smooth muscle cells express S1P<sub>2</sub> and S1P<sub>3</sub>, and cardiac muscles express S1P<sub>1</sub>, S1P<sub>2</sub>, and S1P<sub>3</sub> [109].

S1P<sub>1</sub> is highly expressed in the cardiovascular system during mouse development [110] and in the adult mouse; in humans S1P<sub>1</sub> is expressed at significant levels in brain, heart, lung, spleen, liver and low expression was detected in kidney, muscle and thymus [111-113]. Mice lacking *S1P<sub>1</sub>* die at E12.5-14.5 due to vascular maturation defects. These embryos develop a normal blood vessel network at E12.5, indicating that S1P<sub>1</sub> was not essential for endothelial cell differentiation, proliferation, migration and tube formation during vasculogenesis or for sprouting and branching of vessels during angiogenesis [110]. Instead, the association of vascular smooth muscle cells and pericytes was defected. The dorsal aorta was not covered by vascular smooth muscle cells. Cells were lining the ventral part of the aorta and not the dorsal. The defects in vascular smooth muscle cell coverage extended to arteries and capillaries in the brain and also to small vessels in the limbs. The lack of vascular smooth muscle cells resulted in a weakened vasculature with disrupted and leaky vessels. The defect seen in the *S1P<sub>1</sub>* mutant might be caused by impaired vascular smooth muscle cell and pericyte differentiation, proliferation and/or migration. For example, fibroblast cells from the *S1P<sub>1</sub>* mutant failed to migrate towards S1P in vitro [114].


## **INSULIN CELL MATURATION AND GLUCOSE HOMEOSTASIS**

After digestion, a variety of nutritional factors in circulation, including amino acids, fatty acids and glucose initiate release of insulin. A series of signaling cascades responsible for the suppression of hepatic glucose output, increased synthesis of glycogen and triglycerides, and stimulation of peripheral tissue uptake of glucose begin. In addition, hormones from the gut (incretins) are secreted, which increases insulin secretion from the pancreas in a glucose-dependent manner [115].

Insulin is synthesized as proinsulin on the rough endoplasmic reticulum (RER). It is converted into proinsulin and transferred to the Trans-Golgi Network (TGN), where it is packed into immature secretory granules. These vesicles fuse directly to the plasma membrane or enter the recycling endosome system. The immature granules undergo several maturation processes, such as acidification, conversion of proinsulin into insulin and C-peptide, and loss of the coat protein clathrin [116, 117].

There are two pools of insulin secretory granules, the readily releasable pool (RRP) that is responsible for the initial phase of insulin secretion and a second reserve pool that is responsible for a more prolonged insulin secretion phase [118-121]. The RRP is pre-docked to the membrane and is released within 1-5 min. During this first phase approximately 15 granules per minute are released. The second phase continues for 5-60 min and releases 5 granules per minute [119]. A mouse  $\beta$ -cell contains more than 10 000 secretory granules [122]. Only 1-5% of the secretory granules belong to the RRP.

High glucose in the blood is sensed by glucose transporter 2 (GLUT2) in the plasma membrane. Via glycolysis the ratio of ATP/ADP increases and closes the ATP-sensitive  $K^+$  channels. This leads to a depolarization that activates  $Ca^{2+}$  channels resulting in extracellular  $Ca^{2+}$  influx and fusion of insulin granules with the plasma membrane. At the same time GLP-1 binds to its receptor and activates cAMP that binds to EPAC. EPAC increases the numbers of insulin granules in the RRP at the plasma membrane (Figure 8). Together these processes results in insulin secretion [119].



**Figure 8. Insulin signaling.** Glucose increase ATP/ADP ratio leading to closing of  $K^+$  channels. Depolarization of the membrane, increased concentration of  $Ca^{2+}$  within the cell, and fusion of insulin granules with the plasma membrane increase cAMP. cAMP binds to EPAC to increase  $Ca^{2+}$  and thereby increasing the insulin granules within the readily releasable pool. Light blue dots represent insulin granules. Picture modified from Hou et al 2009.

## DIABETES

The most common forms of diabetes are type 1 and 2, however there are several different types of diabetes. Type 1 diabetes is an autoimmune disease resulting in destruction of  $\beta$ -cells by the immune system. Type 2 diabetes is caused by insulin resistance. The pancreas in type 2 diabetic patient has an increased  $\beta$ -cell mass to compensate for the elevated need for insulin. Eventually this compensatory mechanism fails and the patient develops diabetes.

Diabetes is a disease that is associated with major complications, including retinopathy, neuropathy, kidney failure and cardiovascular defects. The diabetic retinopathy causes blindness, while diabetic neuropathy causes symptoms of tingling, pain, numbness, or weakness in the feet and hands. Kidney failure as well as cardiovascular defects is also common causes of death in diabetes patients (World Health Organization; [www.who.int](http://www.who.int)).

The most common treatment for patients with type 1 diabetes is daily injections of recombinant insulin. This is a fairly efficient treatment but excursions in blood glucose levels may still be difficult to control. Thus alternative treatment options are currently explored. Type 1 diabetic patients have been injected with donor islets. This procedure was able to restore euglycemia [123]. However, two major obstacles are associated with this procedure: only 10% of the patients were insulin free after 5 years of transplantation, and secondly there are not enough donors.

Thus alternative sources for  $\beta$ -cells are needed. One possibility would be to differentiate human Embryonic Stem Cells (hESC) into transplantable  $\beta$ -cells. hESC have been proven to be able to differentiate into insulin producing  $\beta$ -cells [124] and in vitro studies have also succeeded in differentiating hESC into insulin producing  $\beta$ -cells [125-127]. Efficient and reproducible differentiation protocols will be needed to develop fully functional, transplantable  $\beta$ -cells.

## TECHNICAL CONSIDERATIONS

### THE GENE-TRAP SYSTEM

Gene-trap mutagenesis is a technique that randomly generates loss-of-function mutations and reports the expression of many mouse genes. A Gene-trap vector contains a splice acceptor site immediately upstream of a promoterless reporter. Since the Gene-trap has no promoter of its own, it is driven by an endogenous promoter close to the insertion site. The Gene-trap also contains a  $\beta$ -galactosidase-Neo<sup>R</sup>fusion ( $\beta$ -Geo) cassette in order to work as a reporter. A disadvantage using a Gene-trap vector is that, because the insertion occurs in an intron, alternative splicing can sometimes take place. This might lead to lower levels of the wild-type transcripts, resulting in hypomorphic alleles [128].

### THE *Cre/loxP*-SYSTEM

Traditional knockout mice may be hard to investigate since many mutants are lethal during embryogenesis. However, even if the mutants do survive until adulthood it might be hard to interpret the data. Since an observed phenotype might be due to a primary defect in another tissue. A solution to this problem is to use the *Cre-loxP* recombination system. The *Cre-loxP* system allows deletion of a gene and its expression in a particular tissue or cell type. *Cre* is a recombinase that recognizes 34 base-pair *loxP*-sequences and catalyses the recombination between two *loxP* sites by excising the intervening DNA [129]. A sequence that is flanked with two *loxP* sites is referred to as a floxed sequence. The floxed sequence will get excised from the genome in the presence of a *Cre*-recombinase.

### RECOMBINATION VARIABILITY'S IN *Cre* TRANSGENES

A known problem with *Cre* transgenic mouse strains is the variability of recombination. This could be due to several factors, including integration site. Integration of a transgene into a site with an open chromatin structure is expected to increase availability for recombination events. Increased variability of transgenic expression is observed with increased number of backcrosses [130]. Different backgrounds of the responder mice may also affect variability in recombination efficiency.

## AIM OF THE THESIS

The overall objective of this thesis was to understand the roles of S1P/S1P<sub>1</sub> and N-cadherin in pancreas development

*The specific aims were:*

- Paper I. To address whether the pancreatic phenotype in *N-cadherin* deficient mice is due to a cell-autonomous function of N-cadherin within the mesenchyme or if it is secondary to other defects, e.g. cardiac and/or vascular function.
- Paper II. To study the functional role of S1P signaling from endothelial cells in pancreas development using *S1P<sub>1</sub>* deficient embryos.
- Paper III. To investigate the consequences of *N-cadherin* ablation in the developing pancreas, specifically in morphogenesis and endocrine cell differentiation.

# PAPERS IN SUMMARY

## PAPER I

### INTRODUCTION

N-cadherin is required for the formation of the dorsal pancreas, by mediating dorsal pancreatic mesenchymal cell survival [27]. *N-cadherin* deficient mice die of cell adhesion defects in the heart at E9.5. These mice also exhibit defects in the development of the nervous system, somites and yolk sac. Importantly, when cadherin function was restored in the heart of *N-cadherin* mutant mice, the embryos survived until E10.5-11 due to rescued heart and vascular function. The aim of this investigation was to address whether the pancreatic phenotype in *N-cadherin*-deficient mice reflects a cell-autonomous function of N-cadherin within the pancreatic mesenchyme or if it is secondary to other defects, e.g. cardiac and/or vascular function.

### RESULTS

#### **Cardiac-rescued *N-cadherin*-deficient mice develop a dorsal pancreas due to an intact circulatory system**

Cardiac rescued *N-cadherin*-knockout mice were analyzed to assess if the lack of the dorsal pancreas in *N-cadherin*-deficient mice is due to a cell-autonomous function within the mesenchyme or if it secondarily caused by other defects (e.g. cardiac and/or vascular function). The most dramatic consequence of expressing N-cadherin within the heart in the *N-cadherin*-mutant mice was the restoration of an intact circulatory system. As a consequence of a functioning circulatory system the dorsal pancreas was formed. The lack of mesenchyme recruitment in the *N-cadherin*-deficient mice was not observed in the rescued *N-cadherin*-knockout mice. This indicates that N-cadherin does not play a cell-autonomous role in mesenchymal cell survival. To identify how the cardiac/vascular function affects pancreas development, explants from *N-cadherin*-deficient embryos were incubated with beads soaked in plasma. The dorsal pancreas was rescued. Endoderm with and without mesenchyme was also cultured with plasma to address whether plasma acts on mesenchyme or epithelium. Our results show that plasma acts primarily on mesenchyme.

### **S1P rescues the formation of the dorsal pancreas in *N-cadherin*-deficient pancreatic explants.**

The dorsal aorta is in close vicinity to the pancreatic endoderm and the vascular smooth muscle cells of the dorsal aorta are derived from mesenchymal cells. It has been demonstrated that vascular maturation is deficient in mice lacking the S1P receptor S1P<sub>1</sub> due to deficient vascular smooth muscle cell recruitment. S1P is a lipid present in the circulation. To test whether the plasma-induced rescue of the dorsal pancreas could be mediated by S1P, we incubated explants with beads soaked in S1P. The explants developed a dorsal pancreas and S1P's effect was specifically seen on the mesenchyme.

### **S1P receptors in pancreas**

S1P binds to receptors S1P<sub>1-5</sub> with high affinity. RT-PCR and in situ hybridization of these receptors show that S1P<sub>1</sub> is expressed in endothelial cells and S1P<sub>2-3</sub> are preferentially expressed in the mesenchyme, but that S1P<sub>4-5</sub> cannot be detected. S1P receptors are G-protein coupled and to block their activity we cultured explants with pertussis toxin that inactivates G<sub>i</sub>. Pertussis toxin blocked the S1P-mediated rescue of the dorsal pancreas as well as early pancreas development in wt explants.

### **S1P regulates dorsal pancreatic mesenchymal cell proliferation**

The mechanism by which S1P rescued the dorsal pancreas development may involve a direct effect on pancreatic mesenchyme. Mesenchymal development involves both cell migration and proliferation, and to investigate if S1P act on any of these processes, we cultured primary mesenchymal cells. BrdU-labeling experiments demonstrated that S1P increase the number of proliferating cells. Importantly, pertussis toxin inhibits this effect.

## **SUMMARY**

We showed that restoring cardiac and circulatory function in *N-cadherin*-deficient mice by cardiac-specific expression of N-cadherin, rescues formation of the dorsal pancreas. This indicates that the phenotype observed is secondary to defects related to cardiac/vascular function. Based on this observation, we demonstrated that plasma and S1P-mediated G-protein-coupled signaling rescues formation of the dorsal pancreas in vitro.



## PAPER II

### INTRODUCTION

During early pancreatic development major blood vessels are in close contact with the pancreatic endoderm. The dorsal bud forms close to the dorsal aorta, while the ventral buds emerge close to the vitelline veins. Endothelial signals have been shown to be important for initiation of the dorsal bud. However, endothelial signals are not required for initiation of the ventral bud.

Later in development the blood vessels and the pancreatic endoderm get separated by invading mesenchyme. Blood vessels are embedded within the mesenchyme and endoderm-endothelial interactions continue. Previously, we have shown that a functional vascular system is required for initiating dorsal pancreas formation. Signaling cues from the circulation, in particular the bioactive sphingolipid metabolite S1P is essential. *S1P<sub>1</sub>* deficient embryos were analyzed to investigate the role of S1P signaling in endothelial cells during pancreas development.

### RESULTS

#### **Growth and branching morphogenesis of the pancreatic epithelium is compromised in *S1P<sub>1</sub>*-deficient embryos**

To measure the effects of specific ablation of *S1P<sub>1</sub>* in the pancreas, the pancreatic volume was scanned with Optical Projection Tomography (OPT). Analysis showed that the pancreatic buds from *S1P<sub>1</sub>*-deficient embryos did not protrude as far as pancreatic buds from wild type embryos. The dorsal bud was significantly smaller in the *S1P<sub>1</sub>*-deficient embryos. The ventral bud was also smaller but not significantly smaller. There were no differences in the structure of the tubular network in the distal parts of the dorsal and ventral buds. However, the proximal parts were less branched in the *S1P<sub>1</sub>*-deficient embryos.

Other endoderm-derived organs, such as liver, ventricle, and lung, revealed a more general effect of *S1P* deletion on endodermal development, for example the lung had fewer branches, and liver and ventricle were smaller.

### **S1P<sub>1</sub> is required for proliferation of Pdx1<sup>+</sup> pancreatic progenitors**

Possible explanations for smaller dorsal and ventral buds in the *S1P<sub>1</sub>*-deficient embryos are less proliferating cells and/or more apoptosis. Wild type and mutant embryos were BrdU-pulse labelled and stained for Pdx1 and BrdU. There were significantly fewer proliferating Pdx1<sup>+</sup> cells in the *S1P<sub>1</sub>*-deficient embryos compared to littermate controls. To investigate if there were more apoptotic cells, we stained for Caspase 3. No significant change was observed. We also looked at endocrine differentiation but there were no significant change between *S1P<sub>1</sub>*-deficient and control embryos. Thus, S1P<sub>1</sub> is required for proliferation of multipotent Pdx1<sup>+</sup> pancreatic progenitor cells, but not for endocrine cell specification.

### **S1P<sub>1</sub> deficiency results in defective pancreatic morphogenesis in vitro**

To exclude that the phenotype observed is caused by the cardiovascular defect, in vitro explants studies were performed. All control explants developed normally, while the *S1P<sub>1</sub>*-deficient explants showed a rather diverse phenotype. The explants development spanned from severe growth defects to showing only size reduction and less branching.

### **Blood vessel ablation does not mimic the S1P<sub>1</sub> phenotype**

Our results indicate that endothelial cells, via S1P<sub>1</sub>, provide inductive cues that are necessary for pancreatic endoderm development. To test this, we ablated all blood vessels using quinolin-urea. After two days of incubation, starting at E11.5, there were no affect on the pancreatic endoderm, suggesting that expansion of progenitor cells beyond E11.5 does not require endothelial cells.

## **SUMMARY**

Developing organism are all dependent on blood vessels providing oxygen and nutrients as well as inductive cues that control cell proliferation and cell specification. It is known that endothelial cells control dorsal pancreas outgrowth and early endocrine cell specification. However, it is not understood how endothelial cells control these processes. Here, we identify S1P<sub>1</sub> as a new signaling pathway that is necessary for development of foregut derived organs, such as lung, liver, stomach and pancreas. Ablation of *S1P<sub>1</sub>* results in reduced size of the dorsal pancreatic bud due to decreased proliferation of Pdx1<sup>+</sup> cells.

## PAPER III

### INTRODUCTION

Previously we demonstrated that N-cadherin is expressed in the pancreatic epithelium at E9.5, but later becomes restricted to endocrine aggregates in mice. Furthermore, in the absence of N-cadherin the dorsal pancreatic bud fails to form. This is a secondary phenotype due to cardiac failure. The early lethality of *N-cadherin*-deficient embryos excludes a complete analysis of N-cadherin function in pancreas development. We generated a tissue specific knockout of *N-cadherin* in the early pancreatic epithelium to study the role of N-cadherin during pancreas formation and function.

### RESULTS

#### **N-cadherin expression during pancreas development**

At E10.5-13.5, N-cadherin is expressed throughout the Pdx1<sup>+</sup> pancreatic epithelium. Later, at E14.5 N-cadherin is expressed at low levels in Sox9<sup>+</sup> cells, acinar cells and at high levels in mature endocrine cells. At E15.5, N-cadherin expression becomes restricted to a subpopulation of Sox9<sup>+</sup> cells, and never in Ngn3<sup>+</sup> cells. In addition Isl1<sup>+</sup> endocrine progenitors (hormone negative) exhibit a mosaic expression of N-cadherin and all hormone producing cells express N-cadherin. At this point acinar cells do not express N-cadherin. Throughout development N-cadherin is also expressed in neurons and blood vessels. From E18.5 and onwards N-cadherin is restricted to hormone producing cells, neurons and blood vessels.

#### ***Pdx1*Cre-mediated ablation of *N-cadherin***

To study the function of N-cadherin during pancreas development *N-cadherin* was conditionally deleted in *Pdx1*<sup>+</sup> cells (cKO). To ensure that *N-cadherin* was ablated, N-cadherin expression was examined by immunofluorescence, and immunoblotting. At E13.5, the efficacy of *N-cadherin* ablation varied from <5% to almost complete ablation of *N-cadherin*. At E15.5, the expression varied between littermates, but N-cadherin expression was consistently maintained in more than 50% of the cells. From E18.5 and onwards, N-cadherin was no longer detectable in the pancreas of cKO individuals.

### **Pancreatic morphogenesis and endocrine specification is not affected in conditional *N-cadherin*-knockout mice.**

To determine if N-cadherin is important for cell lineage specification, expression of specific markers for acinar, ductal, and endocrine cells was analyzed. No developmental defects were observed within the exocrine and endocrine compartment. To investigate if N-cadherin is important for initiation and/or maintenance of islet cell polarity, the distribution of characteristic epithelial junctional, apical and lateral markers was analyzed. However, the normal allocation of these cell polarity markers indicates that islet cell contacts and polarity is not altered. To understand if microtubule dynamics are altered in islets,  $\alpha$ - and  $\beta$ -tubulin were analyzed. There was no difference observed between control and cKO. To investigate the role of N-cadherin in endocrine cell specification and islets formation we measured insulin area versus E-cadherin area. This experiment did not reveal any difference between control and cKO, suggesting that N-cadherin is not required for  $\beta$ -cell specification. To study if other hormone-producing cells were affected, the ratio of glucagon<sup>+</sup>, PP<sup>+</sup>, Somatostatin<sup>+</sup> cells versus insulin<sup>+</sup> cells, respectively, were estimated in adult mice. The ratio was not altered, suggesting that N-cadherin appears to be dispensable for endocrine development.

### **N-cadherin controls insulin granule turnover**

Transmission electron microscopy studies of islets showed a significant overall reduction (27%) of insulin secreting granules in mutant islets. 16% of the  $\beta$ -cells contained very few insulin granules which are a 67% reduction. Even if the  $\beta$ -cells with very few insulin granules were not included in the analysis, the difference was statistical significant. In contrast there is no difference in immature granules in the cKO, suggesting that the decrease in mature granules is not due to a change in biogenesis of granules.

### **N-cadherin regulates insulin secretion**

Insulin secretion was studied in response to low and high concentration of glucose. In response to low glucose insulin secretion was significantly reduced in cKO. At high glucose, insulin secretion was also reduced but the change was not significant.

## SUMMARY

By ablating *N-cadherin* specifically during pancreas development we show that N-cadherin is not essential for pancreas organogenesis. However, the late onset and high variability of recombination, suggest that a potential requirement of N-cadherin during pancreas development prior to E15.5 cannot be ruled out. *N-cadherin*-deficient endocrine cells aggregate into islets with normal morphology, indicating that N-cadherin is not required for islets morphogenesis. However, ultrastructural analysis of adult islets revealed a reduction in number of mature insulin secretory granules in *N-cadherin*-deficient  $\beta$ -cells. In conclusion our findings suggest that N-cadherin is dispensable to pancreas morphogenesis and cell fate specification, but is required for insulin secretory granules turnover and insulin secretion.

## CONCLUDING REMARKS

### PAPER I

N-cadherin does not act cell-autonomously on the mesenchyme surrounding the dorsal pancreas. When expressing cadherins in the heart the mesenchyme was recruited and the dorsal pancreas started to form. This data show that the agenesis seen in *N-cadherin* deficient mice is secondarily due to cardiac-vascular defects. The rescue experiment led us to believe that circulation is important for dorsal pancreas development. Therefore we soaked beads in plasma from E15.5 embryos and cultured explants from *N-cadherin*-deficient mice for two days. The addition of plasma was able to rescue the dorsal pancreas. S1P, a lipid produced and secreted by red blood cells, platelets, monocytes, mast cells, and possibly by endothelial cells, was also able to rescue dorsal pancreas formation. S1P secreted from the dorsal aorta binds to its receptors in the mesenchyme (S1P<sub>1-3</sub>) and induce mesenchymal proliferation, which induce dorsal pancreas formation.

### PAPER II

S1P<sub>1</sub> knockout have defects in endodermal organ formation, such as pancreas, stomach, liver and lung. In the pancreas this is due to less proliferation of the endoderm and not by increased cell death. To evaluate if the phenotype is due to cardiovascular defects, blood vessels were ablated from E11.5 wild type explants. The S1P<sub>1</sub> phenotype was not observed, suggesting that S1P<sub>1</sub> only

has an effect early in development and not during secondary transition. In conclusion we identified a pathway, by which endothelial cells, via S1P<sub>1</sub>, control early pancreas development.

### **PAPER III**

Ablation of *N-cadherin* specifically during pancreas development demonstrates that N-cadherin is not essential for pancreas organogenesis. However, due to the late onset and high variability of recombination, a role of N-cadherin during early pancreatic development can not be ruled out.

Concomitant with islet formation the efficiency of recombination was high. Normal architecture was seen in the *N-cadherin* deficient islets indicating that N-cadherin is not required for islet morphogenesis. However, ultrastructural analysis of adult islets revealed a reduction in the number of mature insulin secretory granules in *N-cadherin*-deficient  $\beta$ -cells. Since the number of immature insulin secretory granules was unaffected we conclude that N-cadherin is not required for granule biogenesis. Consistent with the low number of mature insulin secretory granules, *N-cadherin*-deficient islets displayed less insulin secretion when stimulated with low glucose. We speculate that this is due to fewer granules docked to the membrane and/or fewer granules in the ready releasable pool. Our findings suggest that N-cadherin is dispensable for pancreas morphogenesis and cell fate specification, but that N-cadherin is required for insulin secretory granules turnover and insulin secretion.

## **FUTURE PERSPECTIVES**

### **PAPER I AND II**

Understanding how S1P regulates growth in endoderm derived organs would be of great importance. Most likely the mechanism involves endothelial cells that secrete molecule/molecules that is/are important for endodermal growth.

It could also be of interest to identify other compounds from plasma that can induce dorsal pancreas development. Preliminary results show that lysophosphatidic acid (LPA) is one potential compound.

### PAPER III

To further analyze if N-cadherin plays a role in pancreas development, it is of importance to use a *Pdx1-Cre* transgenic mouse line with high recombination efficiency or other *Cre*-lines with an early onset of expression in the pancreas, such as *Ptf1a*. Thereby, the role of N-cadherin in morphogenesis can be studied in greater detail.

To analyze how N-cadherin is required for insulin granule turnover, different insulin secreting pathways need to be studied. It would be important to clarify if fewer granules are docked to the membrane or if there are fewer granules in the ready releasable pool. TEM studies could be used to address this issue. It could potentially be of interest to see if any of the components in the insulin secreting pathway is defected in the cKO. Insulin secretion studies on islet stimulated with different components involved in insulin signaling pathway would answer this. To analyze if there is less insulin content, perfusion studies on islets will be performed. By adding high percentage of triton, the membrane breaks and all insulin granules are secreted. Preliminary data show that the cKO has lower insulin content.

## ACKNOWLEDGEMENT

Vem hade kunnat tro att det skulle ta så här lång tid att doktorera, inte jag iallafall. Även om den här resan har varit väldigt lång har jag fått uppleva många fantastiska saker och lärt mig mycket på vägen. Jag vill härmed passa på att tacka de människor som på ett eller annat sätt bidragit till min avhandling:

Min handledare **Henrik Semb**. För att du har givit mig friheten att driva mina projekt självständigt, på gott och ont. Detta har gjort att jag utvecklas både som forskare och människa. Jag vill även tacka för att du har skapat en bra miljö att forska i, samt att du har det goda omdömet att rekrytera intressanta medarbetare som var och en på sitt sätt förgyller vardagen.

I would also like to express my gratitude to **Glenn Radice** for all help with the N-cadherin projects. It has been much appreciated.

Jag skulle också vilja tacka alla mina nuvarande kollegor. **Gokul** for being a great companion. I have really enjoyed working with you. **Fredrik** för att du är en god människa med ett stort hjärta. Tänk om du hade lagt all din energi på forskning istället för skitsnack, då hade du varit professor vid det här laget. **Jackie** för din omtanke och för din vilja att alltid hjälpa till om du kan. Hävdar fortfarande att det vore bättre om du tänkte innan du pratar. **Marie** för att du är så lätt att ha att göra med och för att du alltid ställer upp när jag behöver hjälp. **Anti** och **Karro** för att ni underlättar livet på labbet. Utan er hade det inte gått. **Ingar** för att du sköter om beställningar och gör det bästa fiket. **Zarah** för du är en god medarbetare med den rätta takten. **Elvira** för att du ger labbet en helt ny dimension. **Siqin** för många intressanta diskussioner som ofta får mig att skratta. **Pia** för dina nya initiativ på labbet och för att du verkar vara genuint intresserad av forskning. **Yvonne** för att du är down to earth. Uppskattas i denna dramadrottningarnas högborg. **Novo-tjejerna (Jenny, Nina och Katja)** för att ni kryddar till livet på labbet lite extra. Till de nya rekryterna **Hanna, Maria** och **Karen** vill jag bara säga: Livet på labbet kan liknas vid ett dagis för vuxna. Roligt, frustrerande, men samtidigt utvecklande.

Jag vill även tacka våra inneboende i labbet. **Isabella** för all hjälp med allt. Jag uppskattar det verkligen. **Magda** för att du precis som jag tycker om ordning och reda samt ett välstädat labb. **Jesper** för du är easy going.



Har man varit i ett labb så länge som jag har, har man också många före detta kollegor. Jag vill tacka alla som passerat dörrarna på BMC i Lund. **Thomas** det har varit fantastiskt roligt att ha haft dig som kollega under nästan hela resan. Du stod vid ditt ord länge men sköt ut dig i sista stund. Vet få människor som kan vara så entusiastiska som du. **Karin** min gamla student och sedermera arbetskamrat för ditt alldeles egna sätt att se på saker och ting. **Martina** för att du är den bästa rumskamrat man kan ha. **Camilla** för all hjälp genom åren med mössen. **Anders** för att du introducerade QPCRen i våra liv samt försökte få oss att tänka outside the box. **Sune** för den idéspruta du är. **Maria H** för att du höll koll på allt på jobbet. Vet få människor som besitter den förmågan. **Ingrid** för att du alltid bidrog med god stämning, inte minst på Ålagillet.

Jag vill även tacka mina gamla kollegor från tiden på MedKem i Göteborg. **Fina** för ett fantastiskt samarbete. Din sprudlande energi smittar av sig och gör att allt blir mycket roligare. **Gabriella** för din omtanke och fantastiska diskussioner om allt. Vet ingen som kan älta ett problem så som du. **Jocke** för lång och trogen tjänst. Ditt engagemang i saker du tror på är enormt. **Xiaojie** for being the strongest woman on earth. I admire you! **Gunilla** för att du är den bästa BMA man kan önska sig. Hade gärna velat ha med dig ner till Lund. **Nico** för att du är en oerhört fascinerande person. **Mikael** du bidrog kanske inte så mycket till forskningen men desto mer till en helt fantastiskt galen atmosfär att jobba i. **Anna** för att du är den mest ambitiösa person jag någonsin träffat.

Jag vill också passa på att tacka alla andra i min omgivning för en trevlig arbetsmiljö. Grupperna som var på MedKem när det begav sig. Medlemmarna av **CBz's** grupp, **Per Lindahls** grupp, **Sven Enerbäcks** grupp, **Pekna** och **Pekny**-gruppen. Ni gjorde MedKem till den bästa av arbetsplatser. I Lund vill jag passa på att tacka alla på **BMC B10**, och övriga grupper på BMC såsom **Bo Ahrens** grupp för all hjälp under åren, **Nils Wierups** grupp för hjälp med insulinsekretionsstudier samt **Udo Häckers** grupp, **Stefan Baumgartner** och **Edgar Peras** grupp för gemensamma journal clubs.

Till alla mina vänner (ingen nämnd, ingen glömd) vill jag bara säga att ni är de coolaste, smartaste och roligaste vännerna man kan ha. Jag är oerhört tacksam för att jag fortfarande får ha er som vänner, fast jag är så kiss kass värdelös på att höra av mig. Tänker på er ofta!

Jag vill även tacka min familj. Till mormor **Emma Matilda** för att du försöker förstå vad jag gör med de stackars mössen. Till min syster **Mia** för att du är den bästa syster man kan ha. Min svåger **Håkan** tackar jag för all uppmuntran under vägen. Min systerson **Plutten** för att du tillför så mycket glädje till familjen och för att du är nästan lika envis som jag. Till min **mamma** och **pappa** för all kärlek, stöd och uppmuntran som ni gett mig under årens lopp. Jag är oerhört glad för att ni har trott på mig och låtit mig göra som jag vill. Sist men inte minst det största av tack till min älskade **Mattias**. Tack för att du har stöttat mig under upp- och nedgångar under de här åren. Du gör livet så mycket roligare och jag är väldigt tacksam att få vara med dig.

## REFERENCES

1. Townes, P.L.a.H., J., *Directed movements and selective adhesion of embryonic amphibian cells*. J. Exp. Zool., 1955. **128**: p. 53-120.
2. Takeichi, M., *Functional correlation between cell adhesive properties and some cell surface proteins*. J Cell Biol, 1977. **75**(2 Pt 1): p. 464-74.
3. Kemler, R., *Classical cadherins*. Semin Cell Biol, 1992. **3**(3): p. 149-55.
4. Tuckwell, D.S. and M.J. Humphries, *Molecular and cellular biology of integrins*. Crit Rev Oncol Hematol, 1993. **15**(2): p. 149-71.
5. DeSimone, D.W., *Adhesion and matrix in vertebrate development*. Curr Opin Cell Biol, 1994. **6**(5): p. 747-51.
6. Tepass, U., et al., *Cadherins in embryonic and neural morphogenesis*. Nat Rev Mol Cell Biol, 2000. **1**(2): p. 91-100.
7. Ivanov, D.B., M.P. Philippova, and V.A. Tkachuk, *Structure and functions of classical cadherins*. Biochemistry (Mosc), 2001. **66**(10): p. 1174-86.
8. Hatta, K., et al., *Cloning and expression of cDNA encoding a neural calcium-dependent cell adhesion molecule: its identity in the cadherin gene family*. J Cell Biol, 1988. **106**(3): p. 873-81.
9. Farquhar, M.G. and G.E. Palade, *Junctional complexes in various epithelia*. J Cell Biol, 1963. **17**: p. 375-412.
10. Thoreson, M.A., et al., *Selective uncoupling of p120(ctn) from E-cadherin disrupts strong adhesion*. J Cell Biol, 2000. **148**(1): p. 189-202.
11. Anastasiadis, P.Z., *p120-ctn: A nexus for contextual signaling via Rho GTPases*. Biochim Biophys Acta, 2007. **1773**(1): p. 34-46.
12. Anastasiadis, P.Z., et al., *Inhibition of RhoA by p120 catenin*. Nat Cell Biol, 2000. **2**(9): p. 637-44.
13. Grosheva, I., et al., *p120 catenin affects cell motility via modulation of activity of Rho-family GTPases: a link between cell-cell contact formation and regulation of cell locomotion*. J Cell Sci, 2001. **114**(Pt 4): p. 695-707.
14. Noren, N.K., et al., *p120 catenin regulates the actin cytoskeleton via Rho family GTPases*. J Cell Biol, 2000. **150**(3): p. 567-80.

15. Abe, K. and M. Takeichi, *EPLIN mediates linkage of the cadherin catenin complex to F-actin and stabilizes the circumferential actin belt*. Proc Natl Acad Sci U S A, 2008. **105**(1): p. 13-9.
16. Maul, R.S., et al., *EPLIN regulates actin dynamics by cross-linking and stabilizing filaments*. J Cell Biol, 2003. **160**(3): p. 399-407.
17. Grunwald, G.B., R.S. Pratt, and J. Lilien, *Enzymic dissection of embryonic cell adhesive mechanisms. III. Immunological identification of a component of the calcium-dependent adhesive system of embryonic chick neural retina cells*. J Cell Sci, 1982. **55**: p. 69-83.
18. Volk, T. and B. Geiger, *A 135-kd membrane protein of intercellular adherens junctions*. EMBO J, 1984. **3**(10): p. 2249-60.
19. Andersson, A.M., K. Edvardsen, and N.E. Skakkebaek, *Expression and localization of N- and E-cadherin in the human testis and epididymis*. Int J Androl, 1994. **17**(4): p. 174-80.
20. Nouwen, E.J., et al., *Stage- and segment-specific expression of cell-adhesion molecules N-CAM, A-CAM, and L-CAM in the kidney*. Kidney Int, 1993. **44**(1): p. 147-58.
21. Nuruki, K., et al., *E-cadherin but not N-cadherin expression is correlated with the intracellular distribution of catenins in human hepatocellular carcinomas*. Oncol Rep, 1998. **5**(5): p. 1109-14.
22. Hatta, K. and M. Takeichi, *Expression of N-cadherin adhesion molecules associated with early morphogenetic events in chick development*. Nature, 1986. **320**(6061): p. 447-9.
23. Hatta, K., et al., *Spatial and temporal expression pattern of N-cadherin cell adhesion molecules correlated with morphogenetic processes of chicken embryos*. Dev Biol, 1987. **120**(1): p. 215-27.
24. Nakagawa, S. and M. Takeichi, *Neural crest cell-cell adhesion controlled by sequential and subpopulation-specific expression of novel cadherins*. Development, 1995. **121**(5): p. 1321-32.
25. Linask, K.K., *N-cadherin localization in early heart development and polar expression of Na<sup>+</sup>,K<sup>(+)</sup>-ATPase, and integrin during pericardial coelom formation and epithelialization of the differentiating myocardium*. Dev Biol, 1992. **151**(1): p. 213-24.
26. Radice, G.L., et al., *Developmental defects in mouse embryos lacking N-cadherin*. Dev Biol, 1997. **181**(1): p. 64-78.
27. Esni, F., et al., *Dorsal pancreas agenesis in N-cadherin- deficient mice*. Dev Biol, 2001. **238**(1): p. 202-12.

28. Luo, Y., et al., *Rescuing the N-cadherin knockout by cardiac-specific expression of N- or E-cadherin*. *Development*, 2001. **128**(4): p. 459-69.
29. Larue, L., et al., *A role for cadherins in tissue formation*. *Development*, 1996. **122**(10): p. 3185-94.
30. Luo, Y. and G.L. Radice, *N-cadherin acts upstream of VE-cadherin in controlling vascular morphogenesis*. *J Cell Biol*, 2005. **169**(1): p. 29-34.
31. Kadowaki, M., et al., *N-cadherin mediates cortical organization in the mouse brain*. *Dev Biol*, 2007. **304**(1): p. 22-33.
32. Tillet, E., et al., *N-cadherin deficiency impairs pericyte recruitment, and not endothelial differentiation or sprouting, in embryonic stem cell-derived angiogenesis*. *Exp Cell Res*, 2005. **310**(2): p. 392-400.
33. Gilbert, S.F., *Developmental Biology, Seventh Edition* 2003.
34. Metzger, R.J. and M.A. Krasnow, *Genetic control of branching morphogenesis*. *Science*, 1999. **284**(5420): p. 1635-9.
35. Mitsiadis, T.A., L. Regaudiat, and T. Gridley, *Role of the Notch signalling pathway in tooth morphogenesis*. *Arch Oral Biol*, 2005. **50**(2): p. 137-40.
36. Assemat, E., et al., *Polarity complex proteins*. *Biochim Biophys Acta*, 2008. **1778**(3): p. 614-30.
37. McNeill, H., et al., *Novel function of the cell adhesion molecule uvomorulin as an inducer of cell surface polarity*. *Cell*, 1990. **62**(2): p. 309-16.
38. Ridley, A.J., *Rho GTPases and cell migration*. *J Cell Sci*, 2001. **114**(Pt 15): p. 2713-22.
39. Gerhardt, H., et al., *VEGF guides angiogenic sprouting utilizing endothelial tip cell filopodia*. *J Cell Biol*, 2003. **161**(6): p. 1163-77.
40. Stone, J., et al., *Development of retinal vasculature is mediated by hypoxia-induced vascular endothelial growth factor (VEGF) expression by neuroglia*. *J Neurosci*, 1995. **15**(7 Pt 1): p. 4738-47.
41. Nagel, M., et al., *Guidance of mesoderm cell migration in the *Xenopus* gastrula requires PDGF signaling*. *Development*, 2004. **131**(11): p. 2727-36.
42. Jaffe, A.B. and A. Hall, *Rho GTPases: biochemistry and biology*. *Annu Rev Cell Dev Biol*, 2005. **21**: p. 247-69.

43. Nose, A., A. Nagafuchi, and M. Takeichi, *Expressed recombinant cadherins mediate cell sorting in model systems*. Cell, 1988. **54**(7): p. 993-1001.
44. Steinberg, M.S. and M. Takeichi, *Experimental specification of cell sorting, tissue spreading, and specific spatial patterning by quantitative differences in cadherin expression*. Proc Natl Acad Sci U S A, 1994. **91**(1): p. 206-9.
45. Dahl, U., et al., *Genetic dissection of cadherin function during nephrogenesis*. Mol Cell Biol, 2002. **22**(5): p. 1474-87.
46. Semb, H., *Cadherins in Development*. HEP, 2004. **165**: p. 53-68.
47. Arulanandam, R., et al., *Activated Rac1 requires gp130 for Stat3 activation, cell proliferation and migration*. Exp Cell Res. **316**(5): p. 875-86.
48. McLin, V.A., S.A. Rankin, and A.M. Zorn, *Repression of Wnt/beta-catenin signaling in the anterior endoderm is essential for liver and pancreas development*. Development, 2007. **134**(12): p. 2207-17.
49. Deutsch, G., et al., *A bipotential precursor population for pancreas and liver within the embryonic endoderm*. Development, 2001. **128**(6): p. 871-81.
50. Rossi, J.M., et al., *Distinct mesodermal signals, including BMPs from the septum transversum mesenchyme, are required in combination for hepatogenesis from the endoderm*. Genes Dev, 2001. **15**(15): p. 1998-2009.
51. Zhang, W., et al., *Regulation of Hex gene expression and initial stages of avian hepatogenesis by Bmp and Fgf signaling*. Dev Biol, 2004. **268**(2): p. 312-26.
52. Shin, D., et al., *Bmp and Fgf signaling are essential for liver specification in zebrafish*. Development, 2007. **134**(11): p. 2041-50.
53. Chung, W.S., C.H. Shin, and D.Y. Stainier, *Bmp2 signaling regulates the hepatic versus pancreatic fate decision*. Dev Cell, 2008. **15**(5): p. 738-48.
54. Guney, M.A. and M. Gannon, *Pancreas cell fate*. Birth Defects Res C Embryo Today, 2009. **87**(3): p. 232-48.
55. Offield, M.F., et al., *PDX-1 is required for pancreatic outgrowth and differentiation of the rostral duodenum*. Development, 1996. **122**(3): p. 983-95.

56. Silberg, D.G., et al., *Cdx1 and cdx2 expression during intestinal development*. Gastroenterology, 2000. **119**(4): p. 961-71.
57. Harrison, K.A., et al., *Pancreas dorsal lobe agenesis and abnormal islets of Langerhans in Hlxb9-deficient mice*. Nat Genet, 1999. **23**(1): p. 71-5.
58. Li, H., et al., *Selective agenesis of the dorsal pancreas in mice lacking homeobox gene Hlxb9*. Nat Genet, 1999. **23**(1): p. 67-70.
59. Kawaguchi, Y., et al., *The role of the transcriptional regulator Ptf1a in converting intestinal to pancreatic progenitors*. Nat Genet, 2002. **32**(1): p. 128-34.
60. Edlund, H., *Pancreatic organogenesis--developmental mechanisms and implications for therapy*. Nat Rev Genet, 2002. **3**(7): p. 524-32.
61. Ohlsson, H., K. Karlsson, and T. Edlund, *IPF1, a homeodomain-containing transactivator of the insulin gene*. EMBO J, 1993. **12**(11): p. 4251-9.
62. Apelqvist, A., U. Ahlgren, and H. Edlund, *Sonic hedgehog directs specialised mesoderm differentiation in the intestine and pancreas*. Curr Biol, 1997. **7**(10): p. 801-4.
63. Kim, S.K., M. Hebrok, and D.A. Melton, *Notochord to endoderm signaling is required for pancreas development*. Development, 1997. **124**(21): p. 4243-52.
64. Lammert, E., O. Cleaver, and D. Melton, *Induction of pancreatic differentiation by signals from blood vessels*. Science, 2001. **294**(5542): p. 564-7.
65. Yoshitomi, H. and K.S. Zaret, *Endothelial cell interactions initiate dorsal pancreas development by selectively inducing the transcription factor Ptf1a*. Development, 2004. **131**(4): p. 807-17.
66. Golosow, N. and C. Grobstein, *Epitheliomesenchymal interaction in pancreatic morphogenesis*. Dev Biol, 1962. **4**: p. 242-55.
67. Wessells, N.K., *Differentiation of epidermis and epidermal derivatives*. N Engl J Med, 1967. **277**(1): p. 21-33.
68. Ahlgren, U., et al., *Independent requirement for ISL1 in formation of pancreatic mesenchyme and islet cells*. Nature, 1997. **385**(6613): p. 257-60.
69. Kim, S.K. and R.J. MacDonald, *Signaling and transcriptional control of pancreatic organogenesis*. Curr Opin Genet Dev, 2002. **12**(5): p. 540-7.



70. Zhou, Q., et al., *A multipotent progenitor domain guides pancreatic organogenesis*. Dev Cell, 2007. **13**(1): p. 103-14.
71. Rutter, W.J., et al., *An analysis of pancreatic development: role of mesenchymal factor and other extracellular factors*. Symp Soc Dev Biol, 1978(35): p. 205-27.
72. Gittes, G.K., et al., *Lineage-specific morphogenesis in the developing pancreas: role of mesenchymal factors*. Development, 1996. **122**(2): p. 439-47.
73. Miralles, F., et al., *TGF-beta plays a key role in morphogenesis of the pancreatic islets of Langerhans by controlling the activity of the matrix metalloproteinase MMP-2*. J Cell Biol, 1998. **143**(3): p. 827-36.
74. Sanvito, F., et al., *TGF-beta 1 influences the relative development of the exocrine and endocrine pancreas in vitro*. Development, 1994. **120**(12): p. 3451-62.
75. Ritvos, O., et al., *Activin disrupts epithelial branching morphogenesis in developing glandular organs of the mouse*. Mech Dev, 1995. **50**(2-3): p. 229-45.
76. Grapin-Botton, A., *Ductal cells of the pancreas*. Int J Biochem Cell Biol, 2005. **37**(3): p. 504-10.
77. Bhushan, A., et al., *Fgf10 is essential for maintaining the proliferative capacity of epithelial progenitor cells during early pancreatic organogenesis*. Development, 2001. **128**(24): p. 5109-17.
78. Pulkkinen, M.A., et al., *The IIIb isoform of fibroblast growth factor receptor 2 is required for proper growth and branching of pancreatic ductal epithelium but not for differentiation of exocrine or endocrine cells*. Mech Dev, 2003. **120**(2): p. 167-75.
79. Cras-Meneur, C., et al., *Epidermal growth factor increases undifferentiated pancreatic embryonic cells in vitro: a balance between proliferation and differentiation*. Diabetes, 2001. **50**(7): p. 1571-9.
80. Pierreux, C.E., et al., *The transcription factor hepatocyte nuclear factor-6 controls the development of pancreatic ducts in the mouse*. Gastroenterology, 2006. **130**(2): p. 532-41.
81. Seymour, P.A., et al., *SOX9 is required for maintenance of the pancreatic progenitor cell pool*. Proc Natl Acad Sci U S A, 2007. **104**(6): p. 1865-70.
82. Kesavan, G., et al., *Cdc42-mediated tubulogenesis controls cell specification*. Cell, 2009. **139**(4): p. 791-801.



83. Gradwohl, G., et al., *neurogenin3 is required for the development of the four endocrine cell lineages of the pancreas*. Proc Natl Acad Sci U S A, 2000. **97**(4): p. 1607-11.
84. Gu, G., J. Dubauskaite, and D.A. Melton, *Direct evidence for the pancreatic lineage: NGN3+ cells are islet progenitors and are distinct from duct progenitors*. Development, 2002. **129**(10): p. 2447-57.
85. Johansson, K.A., et al., *Temporal control of neurogenin3 activity in pancreas progenitors reveals competence windows for the generation of different endocrine cell types*. Dev Cell, 2007. **12**(3): p. 457-65.
86. Wang, S., et al., *Sustained Neurog3 expression in hormone-expressing islet cells is required for endocrine maturation and function*. Proc Natl Acad Sci U S A, 2009. **106**(24): p. 9715-20.
87. Wilson, M.E., D. Scheel, and M.S. German, *Gene expression cascades in pancreatic development*. Mech Dev, 2003. **120**(1): p. 65-80.
88. Dahl, U., A. Sjodin, and H. Semb, *Cadherins regulate aggregation of pancreatic beta-cells in vivo*. Development, 1996. **122**(9): p. 2895-902.
89. Hutton, J.C., et al., *Molecular cloning of mouse pancreatic islet R-cadherin: differential expression in endocrine and exocrine tissue*. Mol Endocrinol, 1993. **7**(9): p. 1151-60.
90. Sjodin, A., U. Dahl, and H. Semb, *Mouse R-cadherin: expression during the organogenesis of pancreas and gastrointestinal tract*. Exp Cell Res, 1995. **221**(2): p. 413-25.
91. Moller, C.J., et al., *Differential expression of neural cell adhesion molecule and cadherins in pancreatic islets, glucagonomas, and insulinomas*. Mol Endocrinol, 1992. **6**(8): p. 1332-42.
92. Esni, F., et al., *Neural cell adhesion molecule (N-CAM) is required for cell type segregation and normal ultrastructure in pancreatic islets*. J Cell Biol, 1999. **144**(2): p. 325-37.
93. Cleaver, O. and P.A. Krieg, *VEGF mediates angioblast migration during development of the dorsal aorta in Xenopus*. Development, 1998. **125**(19): p. 3905-14.
94. Olivera, A. and S. Spiegel, *Sphingosine kinase. Assay and product analysis*. Methods Mol Biol, 1998. **105**: p. 233-42.
95. Kohama, T., et al., *Molecular cloning and functional characterization of murine sphingosine kinase*. J Biol Chem, 1998. **273**(37): p. 23722-8.

96. van Veldhoven, P.P. and G.P. Mannaerts, *Sphingosine-phosphate lyase*. *Adv Lipid Res*, 1993. **26**: p. 69-98.
97. Saba, J.D., et al., *The BST1 gene of Saccharomyces cerevisiae is the sphingosine-1-phosphate lyase*. *J Biol Chem*, 1997. **272**(42): p. 26087-90.
98. Zhou, J. and J.D. Saba, *Identification of the first mammalian sphingosine phosphate lyase gene and its functional expression in yeast*. *Biochem Biophys Res Commun*, 1998. **242**(3): p. 502-7.
99. Yatomi, Y., et al., *Quantitative measurement of sphingosine 1-phosphate in biological samples by acylation with radioactive acetic anhydride*. *Anal Biochem*, 1995. **230**(2): p. 315-20.
100. Venkataraman, K., et al., *Vascular endothelium as a contributor of plasma sphingosine 1-phosphate*. *Circ Res*, 2008. **102**(6): p. 669-76.
101. Pappu, R., et al., *Promotion of lymphocyte egress into blood and lymph by distinct sources of sphingosine-1-phosphate*. *Science*, 2007. **316**(5822): p. 295-8.
102. Murata, N., et al., *Interaction of sphingosine 1-phosphate with plasma components, including lipoproteins, regulates the lipid receptor-mediated actions*. *Biochem J*, 2000. **352 Pt 3**: p. 809-15.
103. Goetzl, E.J. and S. An, *A subfamily of G protein-coupled cellular receptors for lysophospholipids and lysosphingolipids*. *Adv Exp Med Biol*, 1999. **469**: p. 259-64.
104. Goetzl, E.J., Y. Kong, and B. Mei, *Lysophosphatidic acid and sphingosine 1-phosphate protection of T cells from apoptosis in association with suppression of Bax*. *J Immunol*, 1999. **162**(4): p. 2049-56.
105. Edsall, L.C., et al., *Sphingosine kinase expression regulates apoptosis and caspase activation in PC12 cells*. *J Neurochem*, 2001. **76**(5): p. 1573-84.
106. Hobson, J.P., et al., *Role of the sphingosine-1-phosphate receptor EDG-1 in PDGF-induced cell motility*. *Science*, 2001. **291**(5509): p. 1800-3.
107. Paik, J.H., et al., *Sphingosine 1-phosphate-induced endothelial cell migration requires the expression of EDG-1 and EDG-3 receptors and Rho-dependent activation of alpha vbeta3- and beta1-containing integrins*. *J Biol Chem*, 2001. **276**(15): p. 11830-7.
108. Proia, R.R., et al., *The effect of endothelial cell overexpression of plasminogen activator inhibitor-1 on smooth muscle cell migration*. *J Vasc Surg*, 2002. **36**(1): p. 164-71.

109. Rosen, H., et al., *Sphingosine 1-phosphate receptor signaling*. Annu Rev Biochem, 2009. **78**: p. 743-68.
110. Liu, Y., et al., *Edg-1, the G protein-coupled receptor for sphingosine-1-phosphate, is essential for vascular maturation*. J Clin Invest, 2000. **106**(8): p. 951-61.
111. Liu, C.H. and T. Hla, *The mouse gene for the inducible G-protein-coupled receptor edg-1*. Genomics, 1997. **43**(1): p. 15-24.
112. Zhang, G., et al., *Comparative analysis of three murine G-protein coupled receptors activated by sphingosine-1-phosphate*. Gene, 1999. **227**(1): p. 89-99.
113. An, S., E.J. Goetzl, and H. Lee, *Signaling mechanisms and molecular characteristics of G protein-coupled receptors for lysophosphatidic acid and sphingosine 1-phosphate*. J Cell Biochem Suppl, 1998. **30-31**: p. 147-57.
114. Allende, M.L. and R.L. Proia, *Sphingosine-1-phosphate receptors and the development of the vascular system*. Biochim Biophys Acta, 2002. **1582**(1-3): p. 222-7.
115. Hou, J.C., L. Min, and J.E. Pessin, *Insulin granule biogenesis, trafficking and exocytosis*. Vitam Horm, 2009. **80**: p. 473-506.
116. Orci, L., *Macro- and micro-domains in the endocrine pancreas*. Diabetes, 1982. **31**(6 Pt 1): p. 538-65.
117. Orci, L., et al., *Clathrin-immunoreactive sites in the Golgi apparatus are concentrated at the trans pole in polypeptide hormone-secreting cells*. Proc Natl Acad Sci U S A, 1985. **82**(16): p. 5385-9.
118. Bratanova-Tochkova, T.K., et al., *Triggering and augmentation mechanisms, granule pools, and biphasic insulin secretion*. Diabetes, 2002. **51 Suppl 1**: p. S83-90.
119. Rorsman, P. and E. Renstrom, *Insulin granule dynamics in pancreatic beta cells*. Diabetologia, 2003. **46**(8): p. 1029-45.
120. Rutter, G.A., *Nutrient-secretion coupling in the pancreatic islet beta-cell: recent advances*. Mol Aspects Med, 2001. **22**(6): p. 247-84.
121. Straub, S.G. and G.W. Sharp, *Glucose-stimulated signaling pathways in biphasic insulin secretion*. Diabetes Metab Res Rev, 2002. **18**(6): p. 451-63.
122. Olofsson, C.S., et al., *Fast insulin secretion reflects exocytosis of docked granules in mouse pancreatic B-cells*. Pflugers Arch, 2002. **444**(1-2): p. 43-51.

123. Shapiro, A.M. and J.R. Lakey, *Future trends in islet cell transplantation*. *Diabetes Technol Ther*, 2000. **2**(3): p. 449-52.
124. Brolen, G.K., et al., *Signals from the embryonic mouse pancreas induce differentiation of human embryonic stem cells into insulin-producing beta-cell-like cells*. *Diabetes*, 2005. **54**(10): p. 2867-74.
125. Kroon, E., et al., *Pancreatic endoderm derived from human embryonic stem cells generates glucose-responsive insulin-secreting cells in vivo*. *Nat Biotechnol*, 2008. **26**(4): p. 443-52.
126. D'Amour, K.A., et al., *Efficient differentiation of human embryonic stem cells to definitive endoderm*. *Nat Biotechnol*, 2005. **23**(12): p. 1534-41.
127. D'Amour, K.A., et al., *Production of pancreatic hormone-expressing endocrine cells from human embryonic stem cells*. *Nat Biotechnol*, 2006. **24**(11): p. 1392-401.
128. Stanford, W.L., J.B. Cohn, and S.P. Cordes, *Gene-trap mutagenesis: past, present and beyond*. *Nat Rev Genet*, 2001. **2**(10): p. 756-68.
129. Sauer, B. and N. Henderson, *The cyclization of linear DNA in Escherichia coli by site-specific recombination*. *Gene*, 1988. **70**(2): p. 331-41.
130. Hara-Kaonga, B., et al., *Variable recombination efficiency in responder transgenes activated by Cre recombinase in the vasculature*. *Transgenic Res*, 2006. **15**(1): p. 101-6.

