



LUND UNIVERSITY

Riverine sources of bioreactive macroelements and their impact on bacterioplankton metabolism in a recipient boreal estuary

Soares, Ana

2018

Document Version:

Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (APA):

Soares, A. (2018). *Riverine sources of bioreactive macroelements and their impact on bacterioplankton metabolism in a recipient boreal estuary*. [Doctoral Thesis (compilation), Centre for Environmental and Climate Science (CEC)]. Lund University, Faculty of Science, Center for Environmental and Climate Research / Department of Physical Geography and Ecosystem Science.

Total number of authors:

1

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Riverine sources of bioreactive macroelements and their impact on bacterioplankton metabolism in a recipient boreal estuary

ANA SOARES | CENTER FOR ENVIRONMENTAL AND CLIMATE RESEARCH
DEPARTMENT OF PHYSICAL GEOGRAPHY AND ECOSYSTEM SCIENCE | LUND UNIVERSITY



Riverine sources of bioreactive macroelements and their impact on bacterioplankton metabolism in a recipient boreal estuary

Ana Soares



LUND
UNIVERSITY

DOCTORAL DISSERTATION

by due permission of the Faculty of Science, Lund University, Sweden.
To be defended at Världen auditorium, Geocentrum I, Sölvegatan 10, Lund, on the
9th of March of 2018 at 10.00 for the degree of Doctor of Philosophy in
Environmental Science.

Faculty opponent

Professor Sybil Seitzinger
University of Victoria, Canada

Organization LUND UNIVERSITY	Document name DOCTORAL DISSERTATION	
Centre for Environmental and Climate Research Department of Geology Department of Physical Geography and Ecosystem Science	Date of issue: February 2018	
Author(s) Ana Soares	Sponsoring organization	
Title and subtitle: Riverine sources of bioreactive macroelements and their impact on bacterioplankton metabolism in a recipient boreal estuary		
<p>The loading of terrestrially derived macroelements, such as dissolved organic carbon (DOC), total nitrogen (N), and total phosphorus (P), to inland and estuarine waters is increasing in the northern hemisphere. These macroelements often limit heterotrophic bacterioplankton metabolism (production and respiration), which in turn influence food web structures, exchange of greenhouse gases between the atmosphere and aquatic systems and oxygen consumption in estuarine ecosystems. In order to predict the impact of increasing macroelement inflows on the function and structure of boreal aquatic ecosystems it is vital to understand the potential sources of macroelements and their regulation of bacterioplankton metabolism.</p> <p>In this thesis, I aimed to determine whether terrestrially derived macroelement loadings to inland and estuarine waters can be utilized by bacterioplankton. In addition, I assessed how riverine influxes of DOC, N and P, alone or combined, regulate estuarine bacterioplankton metabolism. For this purpose, I applied a bioassay approach to measure bacterioplankton respiration (BR) and production (BP) on samples from boreal freshwaters and estuarine systems. Terrestrially derived macroelement availability in brown-water lakes increased from DOC to N to P and averaged 2%, 31% and 49%, respectively. Although extremely low, relative DOC bioreactivity increased with downstream transit for rivers with long water residence time and high fractions of urban and agricultural land use. Riverine bioreactive DOC was not the primary source of energy supporting bacterioplankton respiration at the Öre estuary, confirming the low bioreactive DOC exports expected for the Öre river. Instead riverine nutrients showed a more important effect on BR through stimulation of primary production, which in turn derived the organic carbon utilized by the bacterioplankton in the estuary. However, BP at the Öre estuary was in general limited by P and BR by carbon. In light of expected increases in riverine loadings of DOC and P, and decreases of estuarine primary production, estuarine BP is expected to increase, as well as estuarine BR of riverine DOC inputs. These bacterioplankton responses may be exacerbated by combined increases of riverine inflows of DOC, N and P. In summary, terrestrially derived macroelements can have a substantial impact on freshwater and estuarine bacterioplankton metabolism. Given the large role of bacterioplankton on the structure and function of aquatic systems, continuous changes in the input of terrestrial macroelements may have large consequences for boreal aquatic ecosystems.</p>		
Key words: Dissolved organic carbon; aquatic ecology; bacterial respiration; bacterioplankton production; dissolved organic matter; nutrient bioavailability; DOC bioreactivity		
Classification system and/or index terms (if any)		
Supplementary bibliographical information		Language: English
ISSN and key title		ISBN:
Recipient's notes	Number of pages xxx	Price
	Security classification	

I, the undersigned, being the copyright owner of the abstract of the above-mentioned dissertation, hereby grant to all reference sources permission to publish and disseminate the abstract of the above-mentioned dissertation.

Signature _____



Date 2018-02-05

Riverine sources of bioreactive macroelements and their impact on bacterioplankton metabolism in a recipient boreal estuary

Ana Soares



LUND
UNIVERSITY

A doctoral thesis at the university in Sweden is produced either as a monograph or as a collection of papers. In the latter case the introductory part constitutes the formal thesis, which summarizes the accompanying papers already published or manuscripts at various stages (in press, submitted, or in preparation)

Copyright © Ana Soares

Coverphoto: The Öre estuary.

(Photo: A. Soares)

Faculty of Science, Center for Environmental and Climate Research/Department of Physical Geography and Ecosystem Science

ISBN 978-91-7753-588-1

Printed in Sweden by Media-Tryck, Lund University
Lund 2018



Abstract

The loading of terrestrially derived macroelements, such as dissolved organic carbon (DOC), total nitrogen (N), and total phosphorus (P), to inland and estuarine waters is increasing in the northern hemisphere. These macroelements often limit heterotrophic bacterioplankton metabolism (production and respiration), which in turn influence food web structures, exchange of greenhouse gases between the atmosphere and aquatic systems and oxygen consumption in estuarine ecosystems. In order to predict the impact of increasing macroelement inflows on the function and structure of boreal aquatic ecosystems it is vital to understand the potential sources of macroelements and their regulation of bacterioplankton metabolism.

In this thesis, I aimed to determine whether terrestrially derived macroelement loadings to inland and estuarine waters can be utilized by bacterioplankton. In addition, I assessed how riverine influxes of DOC, N and P, alone or combined, regulate estuarine bacterioplankton metabolism. For this purpose, I applied a bioassay approach to measure bacterioplankton respiration (BR) and production (BP) on samples from boreal freshwaters and estuarine systems. Terrestrially derived macroelement availability in brown-water lakes increased from DOC to N to P and averaged 2%, 31% and 49%, respectively. Although extremely low, relative DOC bioreactivity increased with downstream transit for rivers with long water residence time and high fractions of urban and agricultural land use. Riverine bioreactive DOC was not the primary source of energy supporting bacterioplankton respiration at the Öre estuary, confirming the low bioreactive DOC exports expected for the Öre river. Instead riverine nutrients showed a more important effect on BR through stimulation of primary production, which in turn derived the organic carbon utilized by the bacterioplankton in the estuary. However, BP at the Öre estuary was in general limited by P and BR by carbon. In light of expected increases in riverine loadings of DOC and P, and decreases of estuarine primary production, estuarine BP is expected to increase, as well as estuarine BR of riverine DOC inputs. These bacterioplankton responses may be exacerbated by combined increases of riverine inflows of DOC, N and P.

In summary, terrestrially derived macroelements can have a substantial impact on freshwater and estuarine bacterioplankton metabolism. Given the large role of bacterioplankton on the structure and function of aquatic systems, continuous changes in the input of terrestrial macroelements may have large consequences for boreal aquatic ecosystems.

Popular Science Summary

The loading of macroelements such as, dissolved organic carbon (DOC), total nitrogen (N) and total phosphorus (P) from terrestrial soils to aquatic systems is increasing in the northern hemisphere. This phenomenon has several ecological and biogeochemical consequences for inland and coastal water systems, which are linked to bacterial cycling of DOC, N and P. During growth, bacteria assimilate macroelements into biomass (a process known as bacterial production), removing DOC, N and P from the water column and channelling these elements to consumers. Bacteria can also utilize organic carbon as a source of energy for respiration. For this purpose they take up oxygen dissolved in the water and release CO₂ to the atmosphere. Increases in bacterial production have consequences for aquatic food web structures, whereas increases in bacterial respiration have implications for greenhouse gas emissions and for oxygen concentrations of estuarine waters. Currently, it is still unclear the degree to which land derived macroelements can be utilized by bacteria. Moreover, it is also not understood how bacterial metabolism (production and respiration) will respond to increased terrestrial macroelement fluxes.

In this thesis, I aimed to determine the fraction of the terrestrially derived macroelement loading that can be utilized by bacteria. Moreover, I investigated whether riverine inflows of DOC, N and P, alone or combined, limited the metabolism of bacteria in estuarine waters. I found that on average only 2% of the DOC exported from land was utilized by bacteria for production of biomass during a seven-day period. This share amounted to approximately 50% in the case P, which was thus the macroelement mostly available to bacteria among the three macroelements studied. Yet, the relatively low DOC quality increased downstream for rivers with long river water residence time and high catchment proportions of agricultural and urban land. These riverine catchment features are thus important to predict the export of oxygen consuming organic carbon (C) from rivers to estuaries.

The DOC transported in the Öre river, was less important for support of bacterial respiration than nutrients. These findings agree well with predictions of DOC quality for the Öre river, based on the its catchment features. Riverine nutrients stimulated primary production at the Öre estuary, which in turn supplied organic carbon to bacteria. In general, bacterial production in the Öre estuary was limited by P, while bacterial respiration was limited by organic C. In light of

predicted increases of riverine macroelement deliveries and expected reductions of estuarine primary production, bacterial production will likely increase, as well as the bacterial respiration of riverine delivered organic C. Both estuarine bacterial production and respiration are expected to increase the most in response to combined increases of riverine deliveries of DOC, N and P.

In summary, land derived macroelements can be substantially available to freshwater and estuarine bacteria and impact their metabolism. Given the large role of bacteria in food web structures and aquatic CO₂ emissions, continued changes in the input of terrestrial macroelements may have large implications for boreal aquatic ecosystems.

Contents

List of papers	10
Abbreviations	13
1. Introduction	15
1.1. Macroelement export from land to aquatic systems	15
1.2. Reactivity of land-derived macroelements to bacterioplankton	16
1.3. Impact of riverine inputs of macroelements on estuarine bacterioplankton metabolism	17
2. Aims of the thesis	19
3. Materials and methods	21
3.1. Study area and sampling	21
3.2. Experimental setup	23
4. Main results and discussion	25
4.1. Availability of dissolved organic carbon, nitrogen and phosphorus to bacterioplankton in brown-water lakes (paper I)	25
4.2. WRT and landscape influence on riverine bioreactive DOC (paper II)	26
4.3. Relative importance of direct and indirect effects of riverine dissolved organic matter on estuarine bacterioplankton respiration (paper III)	27
4.4. Single and interactive effects of increasing DOC, nitrogen and phosphorus loadings on bacterioplankton respiration and production in a boreal estuary (paper IV)	28
5. Concluding remarks	31
6. Outlook	33
References	35
Acknowledgements	41

List of papers

This thesis is based on the following manuscripts, which are referred to in the text by their Roman numerals:

- Paper I** Soares, A. R. A., Bergström, A.-K., Sponseller, R. A., Moberg, J. M., Giesler, R., Kritzberg, E. S., Jansson, M. & Berggren, M. 2017. New insights on resource stoichiometry: assessing availability of carbon, nitrogen, and phosphorus to bacterioplankton. *Biogeosciences* 14, 1527-1539.
- Paper II** Soares, A. R. A., Lapierre, J.-F., Selvam, B. P., Lindström, G. & Berggren, M. Controls on dissolved organic carbon bioreactivity in river systems. *Manuscript*.
- Paper III** Soares, A. R. A., & Berggren, M. Importance of riverine dissolved organic matter for bacterioplankton respiration in a boreal estuary. *Manuscript*.
- Paper IV** Soares, A. R. A., Kritzberg, E.S., Custelcean, I., & Berggren, M. 2017. Bacterioplankton responses to increased organic carbon and nutrient loading in a boreal estuary—separate and interactive effects on growth and respiration. *Microbial Ecology*.

The contribution of Ana Soares to the papers included in this thesis was as follows:

- I Had the main responsibility for running the experiment, data handling, data analyses, interpretations, and writing.
- II Planned the study together with co-authors. Performed the field work together with co-authors. Had the main responsibility for running the experiment, data handling, data analyses, interpretations, and writing.
- III Planned the study together with co-author. Had the main responsibility for running the experiment, data handling, data analyses, interpretations, and writing.
- IV Planned the study together with co-authors. Had the main responsibility for running the experiment, data handling, data analyses, interpretations, and writing.

Abbreviations

BP	Bacterioplankton production
BR	Bacterioplankton respiration
C	Carbon
CDOM	Colored dissolved organic matter
CO ₂	Carbon dioxide
DOC	Dissolved organic carbon
DOM	Dissolved organic matter
LTBR	Long-term bioreactivity
N	Nitrogen
P	Phosphorus
SE	Standard error
SEM	Structural equation modelling
STBR	Short-term bioreactivity
SUVA	Dissolved organic carbon specific UV absorbance at 254 nm
tDOC	Terrestrial dissolved organic carbon
tDOM	Terrestrial dissolved organic matter
TP	Total phosphorus
WRT	Water residence time

1. Introduction

1.1 Macroelement export from land to aquatic systems

Over the past decades a browning trend has been observed in inland and coastal surface waters of North America and Northern Europe (Monteith et al. 2007). This trend has been mainly ascribed to increased loadings of terrestrial dissolved organic matter (DOM), which consists of a mixture of plant, soil and organism remains. In Sweden, for example, terrestrial DOM concentrations in aquatic systems have increased as much as 60 % since the late 1980s (Kritzberg and Ekström 2012), and projections are that this increase will continue (de Wit et al. 2016). Several mechanisms have been suggested as drivers of the change in terrestrial DOM loadings to the aquatic environment, including increases in air temperature (Weyhenmeyer and Karlsson 2009), changes in land-cover patterns (Kritzberg 2017), increases in precipitation and runoff (Erlandsson et al. 2008), and a decline in atmospheric sulphate deposition (Monteith et al. 2007). The browning of aquatic systems has marked socio-environmental impacts on drinking water production costs (Eikebrokk et al. 2004), on global biogeochemical cycles (da Cunha et al. 2007), on the metabolism of aquatic organisms (Zwart et al. 2016), among others. Particularly, increased loadings of terrestrial DOM to aquatic systems imply an increase in the supply of elements of special importance to living organisms (i.e. macroelements), such as organic carbon (C), nitrogen (N) and phosphorus (P; Hessen et al. 2010). Since bacteria are the main biological agents of the assimilation and mineralization of macroelements (Cho and Azam 1988; Pomeroy et al. 1991), it is essential to understand the regulation of such processes in light of changes in macroelement loadings.

1.2 Reactivity of land-derived macroelements to bacterioplankton

Heterotrophic bacterioplankton (hereafter referred to as bacterioplankton), which consist of the bacteria in the water column that depend on external supply of chemical energy, represent an important entry point of C, N and P to aquatic heterotrophic food webs (Azam et al. 1983). They remove macroelements from the water column to support their growth (or production), and are in turn grazed upon, supplying macroelements to higher trophic levels of the food web (Jones 1992). Bacterioplankton production (BP) represents a major transfer of energy and nutrients in which the rest of the food web depends on, especially in northern aquatic systems where the amount of energy derived from primary production is low (Jansson et al. 2000). Despite the large amounts of terrestrially derived macroelements received by inland waters, macroelement availability is often a factor limiting BP (Jansson et al. 2012). In fact, only a variable share of land-derived macroelement flux can be accessed by bacteria for production (i.e. is bioavailable; Jansson 1998). For example, a fraction of the N and P transported in the terrestrial DOM may be bound to organic compounds and be thus unavailable to bacteria (Jansson et al. 2012; Stepanauskas et al. 2000). Despite the importance of macroelement bioavailability at the ecosystem level, knowledge on the topic is limited, mostly due to difficulties in determining the bioavailable pool size (Berggren et al. 2015). This knowledge makes it difficult to predict biological and ecological pathways of this increasing C, N and P flux (Eilola et al. 2011).

The role of bacterioplankton on the cycling of C has received special interest during the last decades. This is especially due to the increase of CO₂ concentrations in the atmosphere and associated impact on global climate (Ciais et al. 2014). Dissolved organic carbon (DOC) compromises the major pathway of C from terrestrial soils to freshwater systems, and ultimately to coasts (Cole et al. 2007). Dissolved organic carbon may be further added within aquatic systems by algae and macrophytes (Guillemette and del Giorgio 2011). The total amount of C annually exported from soils to aquatic systems is now well integrated into the global C cycle, and it is estimated to be approximately half of that sequestered by terrestrial primary producers every year (Battin et al. 2009). During transit through inland waters, a large amount of DOC can be respired by bacterioplankton (Cole et al. 2007). Bacterioplankton respiration (BR) is a process involving a series of redox reactions in order to produce energy, during which oxygen is consumed from water and CO₂ released to the atmosphere. The degree to which DOC can support BR is variable, and the mechanisms underpinning its bioreactivity at the end of inland waters remain unclear (Bauer et al. 2013; Regnier et al. 2013). This knowledge gap is mainly due to a limited understanding of how physical processes, such as water residence time (WRT), together with biogeochemical

macroelement cycling, as function of landscape and climate change, influence DOC bioreactivity properties at river mouths (Ward et al. 2017). Although traditionally DOC bioreactivity is known to decrease with increases in WRT (Middelburg 1989), recent literature suggests that other environmental factors may well play a large role for DOC bioreactivity (Evans et al. 2017). The inadequate understanding of the DOC pool that can potentially be mineralized has major implications for estimates of inland water CO₂ emissions and estimates of oxygen-consuming riverine DOC exported to estuarine areas, which are currently associated with a large degree of uncertainty (Ward et al. 2017).

1.3 Impact of riverine inputs of bioreactive macroelements on estuarine bacterioplankton metabolism

Increased loadings of riverine DOM are expected to have a profound effect on estuarine bacterioplankton production and respiration (Figueroa et al. 2016; Nydahl et al. 2013). This is expected to be the case for estuarine areas that are highly influenced by riverine DOM inflows, such as boreal estuaries (Andersson et al. 2015). In response to increased riverine DOM deliveries, BP will likely increase (Figueroa et al. 2016). This change in BP may alter estuarine food web structures, and overall productivity of estuaries, which are among the most productive ecosystems on Earth (Canuel et al. 2012). Moreover, increases in estuarine BR can lead to increased CO₂ emissions from estuarine waters and reduce the concentration of dissolved oxygen in the water column (Diaz and Rosenberg 2008; Panigrahi et al. 2013). While the importance of increasing riverine DOM influxes to estuarine systems is evident, little is known regarding the potential utilization of riverine transported macroelements inputs by bacterioplankton. Moreover, it is also unclear how riverine macroelements, alone or combined, regulate estuarine bacterioplankton metabolism (production and respiration; Andersson et al. 2015). This limited understanding hampers the prediction of how the structure and function of boreal estuaries will respond to changes in riverine macroelement deliveries.

2. Aims of the thesis

The overall purpose of this thesis is to determine the potential reactivity of the increasing terrestrially derived macroelement exports to bacterioplankton, and to determine its effect on estuarine bacterioplankton metabolism. The specific aims of the thesis were to:

- Determine the potential bioavailable size of the dissolved organic carbon, nitrogen and phosphorus pool in aquatic systems dominated by terrestrial dissolved organic matter inputs (paper **I**)
- Assess the effect of landscape patterns and water residence time on dissolved organic carbon bioreactivity in river systems (paper **II**)
- Determine the relative importance of direct and indirect effects of riverine dissolved organic matter for bacterioplankton respiration in a boreal estuary (paper **III**)
- Assess the single and interactive effects of increasing macroelement loadings on bacterioplankton respiration and production in a boreal estuary (paper **IV**)

3. Materials and methods

3.1 Study area and sampling

This study was conducted on brown-water lakes, rivers and on a boreal estuary within Sweden. Specifically, the study presented in Paper **I** was conducted in epilimnion samples of four lakes located in northern of Sweden (Övre Björntjärnen, Lillsjöleden, Struvtjärnen and Stortjärnen; Fig. 1). Samples were taken between September 2012 and September 2014. The lakes have a similar size and are located within the same region. The catchments of these lakes are mainly composed of different portions of coniferous forest (Scots Pine, *Pinus sylvestris*; Norway spruce, *Picea abies*) and wetlands (mires).

The study presented in Paper **II** was conducted on samples from 18 river outlets that drain to the Kattegat or into the Baltic Sea. Samples were taken once, during low flow conditions, between June and August 2013. River catchments fall between latitudes 68° N and 56° N and span a 1800 km climate gradient, which includes a subarctic climate in the north and a more temperate climate in the south. The size of the sampled catchments varied from 131 to 48136 km². Land use and land cover varied from mountainous, wetland and forested areas in the North, to more agricultural and urban areas in the South (Sponseller et al. 2014).

The research presented in papers **III** and **IV** was conducted at the Öre estuary, located in the northwestern part of the Bothnian Sea (Baltic Sea). The estuary consists of a semi-enclosed area with no tides, of roughly 50 km², an average depth of 16 m and a volume of 1.0×10^9 m³ (Wikner et al. 1999). For the study presented in paper **III**, samples were taken along a fresh-marine water transect, which included the Öre river (sampled at 1 m depth), the Öre estuary (sampled at 1 m and 15 m depth), and a marine site (sampled at 1 m). Samples for this study were taken between March and September of 2014. The study presented in paper **IV** includes samples taken between March 2014 and March 2015 at 1 m depth from two estuarine sites, which differ in terms of proximity to the Öre river mouth.




Figure 1. Geographic location of the lakes sampled for paper I (legended as I), the river outlets sampled for paper II (II), the freshwater-seawater transect sampled for paper III (III) and the Öre estuary sampled for paper IV (IV).

3.2 Experimental setup

Bioavailable concentrations of terrestrially derived C, N and P from lakes dominated by tDOM inputs were determined with enrichment bioassays, in which strong macroelement limitation was induced (paper **I**). The response of bacteria was measured by the leucine incorporation of ^3H -leucine according to the method developed by Kirchman et al. (1985) over seven days. Total amounts of ^3H -leucine incorporated were converted into concentrations of bioavailable C, N or P based on standard bacterial growth curves, which described the amount of leucine incorporated per unit of bioavailable limiting macroelement.

The DOC bioreactivity at the end of river outlets (paper **II**) was determined as the amount of DOC lost during short- (0-6 days) and long-term (23-365 days) dark bioassays. Losses of DOC were either calculated from differences in DOC or oxygen concentrations. Data on catchment WRT was derived following the method described in (Lindström et al. 2017), and proportions of land use and land cover were retrieved from the Swedish Meteorological and Hydrological Institute. Causal links between landscape and WRT and bioreactive DOC were tested using structural equation modeling (SEM). These relationships were mediated by other variables which were also included in the SEM: primary production (PP), specific ultraviolet absorption at 254 nm (SUVA_{254}), colored dissolved organic matter (CDOM) and total phosphorus (TP).

Single and interactive effects of increasing C, N and P on estuarine BR and BP were tested with a full-factorial experiment (paper **III**). Sample sub-volumes were added to glass vials and standard solutions of C (glucose), N (ammonium nitrate) and P (disodium phosphate) were amended. Responses of BR were determined from changes in oxygen concentration during three days and responses of BP were measured by leucine incorporation of ^3H -leucine (Kirchman et al. 1985).

For the purpose of assessing the direct and indirect impacts of riverine DOM on the source of organic C used for BR at the Öre estuary (paper **IV**), the stable isotopic composition ($\delta^{13}\text{C}$) of the bioreactive DOC was determined. Changes in concentration and $\delta^{13}\text{C}$ of DOC were recorded over time in long-term dark experiments, and the Keeling plot method (Keeling 1958) was used to estimate the $\delta^{13}\text{C}$ of the bioreactive organic C. The results were compared with estimates of $\delta^{13}\text{C}$ values of bioreactive organic C from terrestrial and marine sources.

4. Main results and discussion

4.1 Availability of dissolved organic carbon, nitrogen and phosphorus to bacterioplankton in brown-water lakes (paper I)

The concentration of bioavailable macroelement in tDOM-rich lakes increased from C to N and from N to P. Specifically, macroelement absolute concentrations varied between 104 and 692 $\mu\text{g C L}^{-1}$, 23 and 287 $\mu\text{g N L}^{-1}$, and 0 and 16 $\mu\text{g P L}^{-1}$. In relation to the total pool, the average share of bioavailable macroelement was 2 %, 31 % to 49 % for C, N and P, respectively (Fig. 2). Phosphorus was thus the most available macroelement, suggesting that in terrestrial DOM-rich systems P availability may be higher than previously thought (Jansson et al. 2012).

Organic N represented on average 80% (± 13 SE) of the total bioavailable N share, while approximately 61% (± 46 SE) of the bioavailable P pool was organic. Bacterioplankton were thus able to take up large fractions of organic macroelements and to utilize them for support of production. Total nutrient ratios were higher in comparison to bioavailable ratios (dependent t test, $p < 0.05$, $n = 26$). For example, average total carbon-to-nitrogen ratio (55 ± 9 SE) was ca. 13 times higher than the average bioavailable carbon-to-nitrogen ratio (4 ± 3 SE). Likewise, average total carbon-to-phosphorus ratio was (4774 ± 2135 SE) was ca. 12 times higher than the average bioavailable carbon-to-phosphorus ratio (369 ± 915 SE). These results highlight that total nutrient fractions are poor forecasters of resource bioavailability in boreal freshwater ecosystems. Due to a high degree of C recalcitrance, predictions of macroelement limitation based on total macroelement ratios may be inadequate when C is part of the ratio.




Figure 2. Share of organic non-bioavailable, organic bioavailable and inorganic nutrient pools of dissolved organic carbon (DOC), total dissolved nitrogen (TDN) and total phosphorus (TP)

4.2 WRT and landscape influence on riverine bioreactive DOC (paper II)

Bulk DOC concentrations measured in surface waters varied from 0.8 to 20.6 mg C L⁻¹. Concentrations of short-term bioreactive (STBR) DOC varied between 0.04 and 0.45 mg C L⁻¹ and concentrations of long-term bioreactive (LTBR) DOC varied between 0.52 and 7.75 mg C L⁻¹. While bulk DOC concentrations were negatively related to WRT, bioreactive pools were independent of WRT (Fig. 3).

The variation of the STBR was strongly related to land cover and land use classes such as forest, agriculture and urban, as function of surface water P concentrations. High P concentrations in turn stimulated the production of a short-term bioreactive C pool towards the end of riverine networks ($r^2=0.52$; $p<0.05$; Fig. 3). These findings agree well with those from paper I by showing that P derived from forested soils had a high potential to support aquatic productivity. In addition, they revealed that agricultural and urban land were also important sources of bioavailable P to adjacent waters, which in turn stimulated the phytoplankton supply of short-term bioreactive C, typically preferred by bacteria to support of metabolic processes (Amon and Benner 1996). Since these two land types are typically located close to river mouths (Weyhenmeyer and Conley 2017), they play an important role for the bioreactivity of the C and P exported that into estuarine zones.

Variation of the LTBR was explained by both land use and WRT, as function of SUVA₂₅₄ and CDOM ($r^2=0.78$; $p<0.05$; Fig 3). Long photochemical processing of the aromatic, recalcitrant C substrates (paper I) likely fueled the LTBR pool observed at river mouths. While these results agree with previous reports of

preferential loss of colored recalcitrant terrestrial DOM during water transit time (Kellerman et al. 2015; Weyhenmeyer et al. 2012), they also contradict the general notion that DOC bioreactivity decreases with increased WTR (Middelburg 1989). Although absolute DOC concentrations declined over time, the DOC measured at river mouths was relatively more bioreactive.




Figure 3. Structural equation model relating short- (STBR) and long-term bioreactive (LTBR) DOC pools with land use and water residence time (WTR). Red arrows show negative, while green positive relationships. The coefficients translate the rate at which the response variable changes in response to a change in its predictor. Significant links are considered at $p < 0.05$ level. Dashed grey arrows represent non-significant pathways. Forest, agricultural and urban land exhibited high negative loadings on principal component one (PC1).

4.3 Relative importance of direct and indirect effects of riverine dissolved organic matter on estuarine bacterioplankton respiration (paper III)

The average signature of the $\delta^{13}\text{C}$ of the bioreactive DOC consumed in the estuary ($-22 \text{‰} \pm 1\text{SE}$) resembled that of organic C derived from marine phytoplankton (-22 to -21‰ ; Peterson and Fry 1987). These results indicate that BR in the Öre estuary was not primarily supported by riverine DOC, but rather supported by organic C derived from marine phytoplankton production. Even during high flow

periods, such as during spring flood, when bioavailable riverine DOC exports to the estuary are highest (Wikner and Hagstrom 1999), marine-derived organic C was the main source of energy supporting BR. The isotopic signal of the bioreactive DOC was positively related to riverine DOC and nutrient concentrations, which in turn were positively related to riverine flow. These findings suggest that riverine discharge had an important indirect role as source of nutrients to marine primary producers, which was greater than its role as direct source of energy to estuarine bacterioplankton. High riverine nutrient loadings likely contributed to increased primary productivity which in turn derived the main pool of bioreactive organic C utilized by bacterioplankton in the estuary. These findings match well the predictions of low oxygen consuming organic C exports from the Öre river, which has short WRT, and catchment proportions of agricultural and urban land (paper II). Despite riverine nutrients being bound to organic matter, paper I showed that a large fraction of the total N and total P can be bioavailable. Moreover, the bioavailability of nutrients bound in the terrestrial DOM may increase when entering the saline coastal environment; for example, estuarine photolysis can increase organic nutrient bioavailability to phytoplankton (Bushaw et al. 1996).

Nutrient availability is currently the limiting factor to primary productivity at the Öre estuary during growing season (Andersson et al. 1996). However recent studies suggest that increased riverine DOM inflows are expected to decrease primary production due to a decrease in water column light attenuation (Andersson et al. 2013; Wikner and Andersson 2012). Moreover, increased riverine C loadings may decrease the reliance of bacterioplankton on organic carbon derived from primary production (Jansson et al. 2007), increasing bacteria competition with primary producers for N and P. Since bacteria are superior nutrient competitors (Vadstein 2000), bacteria will likely outcompete primary producers, which will in turn further contribute to a possible decrease of primary production in estuaries (Andersson et al. 2015; Carney et al. 2016; Meunier et al. 2017).

4.4 Single and interactive effects of increasing DOC, nitrogen, and phosphorus loadings on bacterioplankton respiration and production in a boreal estuary (paper IV)

Bacterioplankton respiration increased on average 2.2 times in response to single organic C amendments in both estuarine sites. In contrast, amendments of N and P alone did not have a significant effect on BR. These results are in line with the general idea that organic C is the main driver of BR, and that nutrients alone have a marginal impact on estuarine BR (Asmala et al. 2013; del Giorgio and Newell

2012). In light of predicted increases of riverine DOC supplies to estuarine areas, and considering the relatively high bioreactive DOC patterns found at Baltic Sea river mouths (paper **II**), an increases in estuarine BR of riverine DOC supplies is expected.

Bacterioplankton production varied between the two studied estuarine sites. At the site located close to the river mouth, BP increased on average 1.8 times in response to single P amendments, however it did not increase in response to amendments of organic C or N. These results highlight the role of P as limiting macroelement for BP in the estuary. Although resource limitation studies conducted in coastal and offshore areas of the Baltic Sea suggesting overall N-limitation (Kivi et al. 1993; Vaquer-Sunyer et al. 2015), N did not limit bacterioplankton metabolism in the estuary. This is most likely due to the riverine discharge received by the estuary being characterized by high total nitrogen to total phosphorus ratios, which thus lead to a P shortage (Graneli et al. 1990). Considering that P exported from terrestrial soils can be highly bioavailable (paper **I**), the predicted increases of riverine P deliveries are expected to increase BP. At the estuarine site located furthest away from the river mouth, BP was not limited by P but instead was weakly limited by organic C. These results suggest that the fractions of bioreactive organic C in the estuary (paper **III**) were likely depleted at the site close to the estuary, which then led to organic C-limitation of BP at site located furthest away from the from the river mouth.

The combined amendments of organic C, N and P led to the highest observed BP and BR responses in both estuarine sites. Organic C, N and P further showed an interactive effect on all variables, but on BP at the site near the river mouth, where P had a main effect. These results suggest that the three macroelements have a costimulating effect on bacterial metabolism. In light of the predicted increase of riverine macroelement inflows it is thus likely that estuarine BP and estuarine BR of riverine DOM will be in the future intensified.

5. Concluding remarks

This thesis showed that terrestrially derived macroelement loadings have large influence on bacterioplankton metabolism. The bioavailability of terrestrially derived macroelements increased from organic C to N to P in terrestrial DOM-rich lakes, suggesting that in such systems organic C bioavailability is extremely low. Yet, organic carbon bioavailability changed during water transit to river mouths, where a relatively high bioreactive organic C pool was observed. Changes in organic C bioreactivity occurred in river catchments with long water residence time and high portions of agricultural and urban land. These findings suggest that such catchment characteristics must be considered for improved forecasting of riverine inflows of oxygen-consuming organic carbon to estuarine ecosystems. The indirect effect of riverine bioavailable N and P inflows on estuarine bacterioplankton respiration was relatively more important than that of riverine bioreactive organic C. These results matched well the predictions of low bioreactive organic C deliveries from the Öre river based on its catchment characteristics. Instead, riverine nutrients stimulated estuarine primary production which in turn supplied organic carbon to estuarine bacterioplankton. Overall, estuarine bacterioplankton respiration was primarily limited by C and estuarine bacterioplankton production was primarily limited by P. In light of continued increases of riverine P supplies, bacterioplankton production will likely increase, thus shifting the estuary towards net-heterotrophy. Expected decreases of estuarine primary production combined with increased exports of riverine organic C, will likely increase the bacterioplankton respiration of riverine organic C, therefore increasing estuarine emissions of riverine organic carbon-based CO₂. Combined inputs of riverine organic C, N and P may further exacerbate the expected effects of bacterioplankton metabolism on the structure and function of boreal estuarine ecosystems.

6. Outlook

Northern aquatic systems are experiencing increased loading of terrestrially derived macroelements with ecological and biogeochemical significance linked to the bacterial cycling of DOC, N and P (Andersson et al. 2015). To forecast the impact of such increasing flux it is important to improve knowledge on the available C, N and P shares, and to consider the inclusion of this pool on water monitoring programmes.

Moreover, from this thesis it became clear that a high share of the terrestrially derived P is bioavailable, and that P was an important driver of aquatic productivity at river outlets and in the Öre estuary. However, one aspect not covered in this study but related to its findings is the impact of iron-phosphorus interactions on P bioavailability, since iron has the potential to immobilize P. Because terrestrial exports of iron are increasing (Kritzberg and Ekström 2012), this topic should be addressed in future studies.

While this thesis addressed the effect of C, N and P increases on bacterioplankton metabolism, other drivers associated with the increase of terrestrial DOM may also have an impact on bacterioplankton metabolism. For example, the increase of terrestrial DOM is expected to change surface water temperature due to its light absorbing properties and to decrease estuarine salinity (Nydahl et al. 2013). To account for the impact of multiple factors on bacterioplankton metabolic responses, more mesocosm experiments should be employed.

Paper IV proved the role of organic C derived from primary producers for support of estuarine BR. This shows that bacteria and phytoplankton can be coupled even in estuaries that are recipients of high loadings of terrestrial DOM. Due to the strong effect of terrestrial DOM on the bacterioplankton to phytoplankton ratio (Wikner and Andersson 2012), more knowledge is needed on the impact of the expected increases of terrestrial DOM loadings on overall basal production (bacteria and phytoplankton) and more emphasis should be given to secondary aquatic production by water monitoring programmes.

Finally, more importance should be given to the effects of land cover change and management on increasing export fluxes of terrestrial DOM. For example, afforestation and forest practices such as harvesting, clear-cutting, logging and ditching have a large influence on land export of macroelements to aquatic systems (Laudon et al. 2009; Oni et al. 2015; Rantakari et al. 2010), and should be better considered in the future management of boreal aquatic resources.

References

- Amon, R. M. W., and R. Benner. 1996. Bacterial utilization of different size classes of dissolved organic matter. *Limnol. Oceanogr.* **41**: 41-51.
- Andersson, A. and others 2013. Can humic water discharge counteract eutrophication in coastal waters? *Plos One* **8**.
- Andersson, A. and others 2015. Projected future climate change and Baltic Sea ecosystem management. *AMBIO* **44**: 345-356.
- Asmala, E., R. Autio, H. Kaartokallio, L. Pitkänen, C. A. Stedmon, and D. N. Thomas. 2013. Bioavailability of riverine dissolved organic matter in three Baltic Sea estuaries and the effect of catchment land use. *Biogeosciences* **10**: 6969-6986.
- Azam, F., T. Fenchel, J. G. Field, R. A. Meyer-Reil, and T. F. Thingstad. 1983. The ecological role of water-column microbes in the sea. *Marine Ecology-Progress Series* **10**: 257-263.
- Battin, T. J., S. Luyssaert, L. A. Kaplan, A. K. Aufdenkampe, A. Richter, and L. J. Tranvik. 2009. The boundless carbon cycle. *Nature Geoscience* **2**: 598-600.
- Bauer, J. E., W. J. Cai, P. A. Raymond, T. S. Bianchi, C. S. Hopkinson, and P. A. G. Regnier. 2013. The changing carbon cycle of the coastal ocean. *Nature* **504**: 61-70.
- Berggren, M., R. A. Sponseller, A. R. A. Soares, and A. K. Bergstrom. 2015. Toward an ecologically meaningful view of resource stoichiometry in DOM-dominated aquatic systems. *Journal of Plankton Research* **37**: 489-499.
- Bushaw, K. L. and others 1996. Photochemical release of biologically available nitrogen from aquatic dissolved organic matter. *Nature* **381**: 404-407.
- Canuel, E. A., S. S. Cammer, H. A. McIntosh, and C. R. Pondell. 2012. Climate Change Impacts on the Organic Carbon Cycle at the Land-Ocean Interface, p. 685-+. In R. Jeanloz [ed.], *Annual Review of Earth and Planetary Sciences*, Vol 40. Annual Review of Earth and Planetary Sciences.
- Carney, R. L., J. R. Seymour, D. Westhorpe, and S. M. Mitrovic. 2016. Lotic bacterioplankton and phytoplankton community changes under dissolved organic-carbon amendment: evidence for competition for nutrients. *Marine and Freshwater Research* **67**: 1362-1373.
- Cho, B. C., and F. Azam. 1988. Major role of bacteria in biogeochemical fluxes in the ocean's interior. *Nature* **332**: 441.

- Ciais, P. and others 2014. Current systematic carbon-cycle observations and the need for implementing a policy-relevant carbon observing system. *Biogeosciences* **11**: 3547-3602.
- Cole, J. J. and others 2007. Plumbing the global carbon cycle: Integrating inland waters into the terrestrial carbon budget. *Ecosystems* **10**: 171-184.
- da Cunha, L. C., E. T. Buitenhuis, C. Le Quere, X. Giraud, and W. Ludwig. 2007. Potential impact of changes in river nutrient supply on global ocean biogeochemistry. *Global Biogeochemical Cycles* **21**.
- de Wit, H. A. and others 2016. Current Browning of Surface Waters Will Be Further Promoted by Wetter Climate. *Environmental Science & Technology Letters* **3**: 430-435.
- del Giorgio, P. A., and R. E. I. Newell. 2012. Phosphorus and DOC availability influence the partitioning between bacterioplankton production and respiration in tidal marsh ecosystems. *Environmental Microbiology* **14**: 1296-1307.
- Diaz, R. J., and R. Rosenberg. 2008. Spreading dead zones and consequences for marine ecosystems. *Science* **321**: 926-929.
- Eikebrokk, B., R. D. Vogt, and H. Liltved. 2004. NOM increase in Northern European source waters: discussion of possible causes and impacts on coagulation/contact filtration processes, p. 47-54. *In* G. Newcombe and L. Ho [eds.], *Natural Organic Material Research: Innovations and Applications for Drinking Water*. Water Supply : The Review Journal of the International Water Supply Association.
- Eilola, K., B. G. Gustafsson, I. Kuznetsov, H. E. M. Meier, T. Neumann, and O. P. Savchuk. 2011. Evaluation of biogeochemical cycles in an ensemble of three state-of-the-art numerical models of the Baltic Sea. *Journal of Marine Systems* **88**: 267-284.
- Erlandsson, M. and others 2008. Thirty-five years of synchrony in the organic matter concentrations of Swedish rivers explained by variation in flow and sulphate. *Global Change Biology* **14**: 1191-1198.
- Evans, C. D., M. N. Futter, F. Moldan, S. Valinia, Z. Frogbrook, and D. N. Kothawala. 2017. Variability in organic carbon reactivity across lake residence time and trophic gradients. *Nature Geoscience* **10**: 832+.
- Figueroa, D., O. F. Rowe, J. Paczkowska, C. Legrand, and A. Andersson. 2016. Allochthonous Carbon—a Major Driver of Bacterioplankton Production in the Subarctic Northern Baltic Sea. *Microbial Ecology* **71**: 789-801.
- Graneli, E., K. Wallstrom, U. Larsson, W. Graneli, and R. Elmgren. 1990. NUTRIENT LIMITATION OF PRIMARY PRODUCTION IN THE BALTIC SEA AREA. *Ambio* **19**: 142-151.
- Guillemette, F., and P. A. del Giorgio. 2011. Reconstructing the various facets of dissolved organic carbon bioavailability in freshwater ecosystems. *Limnol. Oceanogr.* **56**: 734-748.

- Hessen, D. O. and others 2010. Input of organic carbon as determinant of nutrient fluxes, light climate and productivity in the Ob and Yenisey estuaries. *Estuarine, Coastal and Shelf Science* **88**: 53-62.
- Jansson, M. 1998. Nutrient Limitation and Bacteria — Phytoplankton Interactions in Humic Lakes, p. 177-195. *In* D. O. Hessen and L. J. Tranvik [eds.], *Aquatic Humic Substances: Ecology and Biogeochemistry*. Springer Berlin Heidelberg.
- Jansson, M., M. Berggren, H. Laudon, and A. Jonsson. 2012. Bioavailable phosphorus in humic headwater streams in boreal Sweden. *Limnol. Oceanogr.* **57**: 1161-1170.
- Jansson, M., A. K. Bergström, P. Blomqvist, and S. Drakare. 2000. Allochthonous organic carbon and phytoplankton/bacterioplankton production relationships in lakes. *Ecology* **81**: 3250-3255.
- Jansson, M., L. Persson, A. M. DeRoos, R. I. Jones, and L. J. Tranvik. 2007. Terrestrial carbon and intraspecific size-variation shape lake ecosystems. *Trends in Ecology & Evolution* **22**: 316-322.
- Jones, R. I. 1992. The influence of humic substances on lacustrine planktonic food-chains. *Hydrobiologia* **229**: 73-91.
- Keeling, C. D. 1958. The concentration and isotopic abundances of atmospheric carbon dioxide in rural areas. *Geochimica et Cosmochimica Acta* **13**: 322-334.
- Kellerman, A. M., D. N. Kothawala, T. Dittmar, and L. J. Tranvik. 2015. Persistence of dissolved organic matter in lakes related to its molecular characteristics. *Nature Geoscience* **8**: 454-U452.
- Kirchman, D., E. Knees, and R. Hodson. 1985. Leucine incorporation and its potential as a measure of protein-synthesis by bacteria in natural aquatic systems. *Applied and Environmental Microbiology* **49**: 599-607.
- Kivi, K. and others 1993. NUTRIENT LIMITATION AND GRAZING CONTROL OF THE BALTIC PLANKTON COMMUNITY DURING ANNUAL SUCCESSION. *Limnol. Oceanogr.* **38**: 893-905.
- Kritzberg, E. S. 2017. Centennial-long trends of lake browning show major effect of afforestation. *Limnology and Oceanography Letters* **2**: 105-112.
- Kritzberg, E. S., and S. M. Ekström. 2012. Increasing iron concentrations in surface waters – a factor behind brownification? *Biogeosciences* **9**: 1465-1478.
- Laudon, H., J. Hedtjarn, J. Schelker, K. Bishop, R. Sorensen, and A. Agren. 2009. Response of dissolved organic carbon following forest harvesting in a boreal forest. *Ambio* **38**: 381-386.
- Lindström, G., A. Bartosova, N. Hjerdt, and J. Strömqvist. 2017. Upphållstider i ytvatten i relation till vattenkvalitet - NET, ett generellt uppskalningsverktyg. SMHI Rapport Hydrologi Nr. 199.

- Meunier, C. L. and others 2017. Allochthonous carbon is a major driver of the microbial food web - A mesocosm study simulating elevated terrestrial matter runoff. *Mar. Environ. Res.* **129**: 236-244.
- Middelburg, J. J. 1989. A simple rate model for organic matter decomposition in marine sediments. *Geochim Cosmochim Acta* **53**: 1577-1581.
- Monteith, D. T. and others 2007. Dissolved organic carbon trends resulting from changes in atmospheric deposition chemistry. *Nature* **450**: 537-U539.
- Nydahl, A., S. Panigrahi, and J. Wikner. 2013. Increased microbial activity in a warmer and wetter climate enhances the risk of coastal hypoxia. *Fems Microbiology Ecology* **85**: 338-347.
- Oni, S. K. and others 2015. Local- and landscape-scale impacts of clear-cuts and climate change on surface water dissolved organic carbon in boreal forests. *J. Geophys. Res.-Biogeosci.* **120**: 2402-2426.
- Panigrahi, S., A. Nydahl, P. Anton, and J. Wikner. 2013. Strong seasonal effect of moderate experimental warming on plankton respiration in a temperate estuarine plankton community. *Estuarine Coastal and Shelf Science* **135**: 269-279.
- Peterson, B. J., and B. Fry. 1987. Stable isotopes in ecosystem studies. *Annual Review of Ecology and Systematics* **18**: 293-320.
- Pomeroy, L. R., W. J. Wiebe, D. Deibel, R. J. Thompson, G. T. Rowe, and J. D. Pakulski. 1991. Bacterial responses to temperature and substrate concentration during the Newfoundland spring bloom. *Marine Ecology-Progress Series* **75**: 143-159.
- Rantakari, M., T. Mattsson, P. Kortelainen, S. Piirainen, L. Finer, and M. Ahtiainen. 2010. Organic and inorganic carbon concentrations and fluxes from managed and unmanaged boreal first-order catchments. *Science of the Total Environment* **408**: 1649-1658.
- Regnier, P. and others 2013. Anthropogenic perturbation of the carbon fluxes from land to ocean. *Nature Geoscience* **6**: 597-607.
- Sponseller, R. A., J. Temnerud, K. Bishop, and H. Laudon. 2014. Patterns and drivers of riverine nitrogen (N) across alpine, subarctic, and boreal Sweden. *Biogeochemistry* **120**: 105-120.
- Stepanauskas, R., H. Laudon, and N. O. G. Jorgensen. 2000. High DON bioavailability in boreal streams during a spring flood. *Limnol. Oceanogr.* **45**: 1298-1307.
- Vadstein, O. 2000. Heterotrophic, planktonic bacteria and cycling of phosphorus - Phosphorus requirements, competitive ability, and food web interactions, p. 115-167. *In* B. Schink [ed.], *Advances in Microbial Ecology*, Vol 16. *Advances in Microbial Ecology*.

- Vaquer-Sunyer, R., D. J. Conley, S. Muthusamy, M. V. Lindh, J. Pinhassi, and E. S. Kritzberg. 2015. Dissolved Organic Nitrogen Inputs from Wastewater Treatment Plant Effluents Increase Responses of Planktonic Metabolic Rates to Warming. *Environmental science & technology* **49**: 11411-11420.
- Ward, N. D. and others 2017. Where Carbon Goes When Water Flows: Carbon Cycling across the Aquatic Continuum. *Frontiers in Marine Science* **4**.
- Weyhenmeyer, G. A., and D. J. Conley. 2017. Large differences between carbon and nutrient loss rates along the land to ocean aquatic continuum—implications for energy:nutrient ratios at downstream sites. *Limnol. Oceanogr.* **62**: S183-S193.
- Weyhenmeyer, G. A. and others 2012. Selective decay of terrestrial organic carbon during transport from land to sea. *Global Change Biology* **18**: 349-355.
- Weyhenmeyer, G. A., and J. Karlsson. 2009. Nonlinear response of dissolved organic carbon concentrations in boreal lakes to increasing temperatures. *Limnol. Oceanogr.* **54**: 2513-2519.
- Wikner, J., and A. Andersson. 2012. Increased freshwater discharge shifts the trophic balance in the coastal zone of the northern Baltic Sea. *Global Change Biology* **18**: 2509-2519.
- Wikner, J., R. Cuadros, and M. Jansson. 1999. Differences in consumption of allochthonous DOC under limnic and estuarine conditions in a watershed. *Aquatic Microbial Ecology* **17**: 289-299.
- Wikner, J., and A. Hagstrom. 1999. Bacterioplankton intra-annual variability: importance of hydrography and competition. *Aquatic Microbial Ecology* **20**: 245-260.
- Zwart, J. A. and others 2016. Metabolic and physiochemical responses to a whole-lake experimental increase in dissolved organic carbon in a north-temperate lake. *Limnology and Oceanography* **61**: 723-734.

Acknowledgments

Many have contributed to the work presented in this thesis. First and foremost, I would like to thank my supervisor, Martin Berggren, for all the help, advice, positiveness, encouragement, enthusiasm, and great supervision throughout these years. I would also like to thank my co-supervisor, Emma Kritzberg, for encouragement, optimism, and for many insightful thoughts which were very helpful to this project. Also, thanks to my co-supervisor, Andreas Persson, for being present at the meetings, and for showing interest on the development of this project.

The following have help with

—field and laboratory assistance: Siv Huseby, Anna Palmbo Bergman, Anne Deininger, Marcus Klaus, Marcin Jackowicz-Korczynski, Bala Selvam, Åsa Wallin, Julia Jakobsson, Karla Müzner, Lina Allesson, Joanna Moberg, Ioana Custelcean, Sohedul Islam

—comments and edits on the manuscripts: Ann-Kristin Bergström, Ryan Sponseller, Jean-François Lapierre

—freeze-drier sharing: Carla Nantke, Laurie Charrieau, Guillaume Fontorbe, Wim Clymans, Patrik Frings

—administrative/technical tasks: Rafael Przybyszewski, Ekaterina Volkova, Irma Habermann, Åsa-Katrin Erlandsson

—comments on the thesis summary: Katarina Hedlund, Pål Axel Olsson, Emma Kritzberg, Martin Berggren

—language editing on the thesis summary: Thomas Dowling, Carolina Funkey, Geert Hensgens

—preparing the map included in the thesis summary: Antonin Kusbach

—thesis formatting: Jan Blanke, Min Wang

Thanks to all those who are/were part of Martin's lab, for contributing to a helpful and encouraging work atmosphere: (those who have not yet been acknowledged) Pearl Mzobe, Enass Said, Carlos Arellano. Thanks to the INES PhD students, INES colleagues and CEC PhD students for the all meetings, lunches, activities and for good companionship. Thanks to the "LeChef Jam" for the many PhD tips and for all nice gatherings. Thanks to Jens, Irina and Maci for support during the final writing months. Lastly, thanks to my old friends and to my family for constant support throughout my studies, and for encouragement to pursue my interests.

This project was co-funded by the Centre for Environmental and Climate Research, the Multistressors (FORMAS), the Royal Physiographic Society of Lund, The Helge Ax:son Foundation, Åforsk and KSLA.

DOCTORAL THESES PUBLISHED IN ENVIRONMENTAL SCIENCE, LUND UNIVERSITY

1. Georg K.S. Andersson (2012) Effects of farming practice on pollination across space and time. Department of Biology/Centre for environmental and climate research
2. Anja M. Ödman (2012) Disturbance regimes in dry sandy grasslands – past, present and future. Department of Biology/ Centre for environmental and climate research
3. Johan Genberg (2013) Source apportionment of carbonaceous aerosol. Department of Physics/ Centre for environmental and climate research
4. Petra Bragée (2013) A palaeolimnological study of the anthropogenic impact on dissolved organic carbon in South Swedish lakes. Department of Geology/ Centre for environmental and climate research
5. Estelle Larsson (2013) Sorption and transformation of anti-inflammatory drugs during wastewater treatment. Department of Chemistry/ Centre for environmental and climate research
6. Magnus Ellström (2014) Effects of nitrogen deposition on the growth, metabolism and activity of ectomycorrhizal fungi. Department of Biology/ Centre for environmental and climate research
7. Therese Irminger Street (2015) Small biotopes in agricultural landscapes: importance for vascular plants and effects on management. Department of physical geography and ecosystem science/ Department of Biology/ Centre for environmental and climate research
8. Helena I. Hanson (2015) Natural enemies: Functional aspects of local management in agricultural landscapes. Department of Biology/ Centre for environmental and climate research
9. Lina Nikoleris (2016) The estrogen receptor in fish and effects of estrogenic substances in the environment: ecological and evolutionary perspectives and societal awareness Department of Biology/ Centre for environmental and climate research
10. Cecilia Hultin (2016) Estrogen receptor and multixenobiotic resistance genes in freshwater fish and snails: identification and expression analysis after pharmaceutical exposure. Centre for environmental and climate research

11. Annika M. E. Söderman (2016) Small biotopes: Landscape and management effects on pollinators. Department of Biology/ Centre for environmental and climate research
12. Wenxin Ning (2016) Tracking environmental changes of the Baltic Sea coastal zone since the mid-Holocene. Department of Geology/ Centre for environmental and climate research
13. Karin Mattsson (2016) Nanoparticles in the aquatic environment, Particle characterization and effects on organisms. Department of Chemistry/ Centre for environmental and climate research
14. Ola Svahn (2016) Tillämpad miljöanalytisk kemi för monitorering och åtgärder av antibiotika- och läkemedelsrester i Vattenriket. School of Education and Environment, Kristianstad University/ Centre for environmental and climate research
15. Pablo Urrutia Cordero (2016) Putting food web theory into action: Local adaptation of freshwaters to global environmental change. Department of Biology/ Centre for environmental and climate research
16. Lin Yu (2016) Dynamic modelling of the forest ecosystem: Incorporation of the phosphorous cycle. Centre for environmental and climate research
17. Behnaz Pirzamanbein (2016) Reconstruction of past European land cover based on fossil pollen data: Gaussian Markov random field models for compositional data. Centre for Mathematical Sciences/ Centre for environmental and climate research
18. Arvid Bolin (2017) Ecological interactions in human modified landscapes – Landscape dependent remedies for the maintenance of biodiversity and ecosystem services. Department of Biology/ Centre for environmental and climate research
19. Johan Martinsson (2017) Development and Evaluation of Methods in Source Apportionment of the Carbonaceous Aerosol. Department of Physics/ Centre for environmental and climate research
20. Emilie Öström (2017) Modelling of new particle formation and growth in the atmospheric boundary layer. Department of Physics/ Centre for environmental and climate research
21. Lina Herbertsson (2017) Pollinators and Insect Pollination in Changing Agricultural Landscapes. Centre for environmental and climate research
22. Sofia Hydbom (2017) Tillage practices and their impact on soil organic carbon and the microbial community. Department of Biology/ Centre for environmental and climate research

24. Erik Ahlberg (2017) Speeding up the Atmosphere: Experimental oxidation studies of ambient and laboratory aerosols using a flow reactor. Department of Physics/ Centre for environmental and climate research
23. Laurie M. Charrieau (2017) DISCO: Drivers and Impacts of Coastal Ocean Acidification. Department of Geology/ Centre for environmental and climate research
25. Kristin Rath (2018) Soil salinity as a driver of microbial community structure and functioning. Department of Biology/ Centre for environmental and climate research
26. Lelde Krumina (2018) Adsorption, desorption, and redox reactions at iron oxide nanoparticle surfaces. Department of Biology/ Centre for environmental and climate research



Center for Environmental and Climate Research
Department of Physical Geography
and Ecosystem Science
Faculty of Science

ISBN 978-91-7753-588-1