



LUND UNIVERSITY

Pathological Conditions Involving Extracellular Hemoglobin: Molecular Mechanisms, Clinical Significance, and Novel Therapeutic Opportunities for alpha(1)-Microglobulin

Gram, Magnus; Allhorn, Maria; Bülow, Leif; Hansson, Stefan; Ley, David; Olsson, Martin L; Schmidtchen, Artur; Åkerström, Bo

Published in:
Antioxidants & Redox Signaling

DOI:
[10.1089/ars.2011.4282](https://doi.org/10.1089/ars.2011.4282)

2012

[Link to publication](#)

Citation for published version (APA):

Gram, M., Allhorn, M., Bülow, L., Hansson, S., Ley, D., Olsson, M. L., Schmidtchen, A., & Åkerström, B. (2012). Pathological Conditions Involving Extracellular Hemoglobin: Molecular Mechanisms, Clinical Significance, and Novel Therapeutic Opportunities for alpha(1)-Microglobulin. *Antioxidants & Redox Signaling*, 17(5), 813-846. <https://doi.org/10.1089/ars.2011.4282>

Total number of authors:
8

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Comprehensive invited review

**PATHOLOGICAL CONDITIONS INVOLVING EXTRACELLULAR
HEMOGLOBIN: MOLECULAR MECHANISMS, CLINICAL SIGNIFICANCE, AND
NOVEL THERAPEUTIC OPPORTUNITIES FOR α_1 -MICROGLOBULIN**

Magnus G Olsson¹, Maria Allhorn¹, Leif Bülow², Stefan R Hansson³, David Ley⁴, Martin L
Olsson⁵, Artur Schmidtchen⁶, Bo Åkerström^{1*}

Divisions of ¹Infection Medicine, ²Pure and Applied Biochemistry, ³Obstetrics and
Gynecology, ⁴Pediatrics, ⁵Haematology and Transfusion Medicine, and ⁶Dermatology, at
Lund University, Lund, Sweden.

*Correspondence to:

Bo Åkerström, Division of Infection Medicine, Department of Clinical Sciences, Lund
University, 22184 Lund, Sweden.

Tel.: + 46 46 222 8578; Fax: + 46 46 157756; E-mail: bo.akerstrom@med.lu.se

Word count: 15 297

Reference numbers: 341

Greyscale illustrations: 13.

Color illustrations: 3 (online 3 and hardcopy 2).

Reviewing Editors: Abdu Alayash, Lawrence Druhan, Prasenjit Guchhait, John P Harrington,
Flávio Alves Lara and Masaki Otagiri

Abstract

Hemoglobin is the major oxygen-carrying system of the blood, but has many potentially dangerous side effects due to oxidation and reduction reactions of the heme-bound iron and oxygen. Extracellular hemoglobin, resulting from hemolysis or exogenous infusion, is shown to be an important pathogenic factor in a growing number of diseases. This review briefly outlines the oxidative/reductive toxic reactions of hemoglobin and its metabolites. It also describes physiological protection mechanisms that have evolved against extracellular hemoglobin, with a focus on the most recently discovered: the heme- and radical-binding protein α_1 -microglobulin (A1M). This protein is found in all vertebrates including man and operates by rapidly clearing cytosols and extravascular fluids of heme groups and free radicals released from hemoglobin. Five groups of pathological conditions with high concentrations of extracellular hemoglobin are described: hemolytic anemias and transfusion reactions, the pregnancy complication preeclampsia, cerebral intraventricular hemorrhage of premature infants, chronic inflammatory leg ulcers, and infusion of hemoglobin-based oxygen carriers as blood substitutes. Finally, possible treatments of these conditions are discussed, giving special attention to the described protective effects of A1M.

Table of Content

I. Introduction	5
II. Hemoglobin, heme and oxidative stress	7
A. Red blood cells and hemoglobin	7
1. Hemoglobin variants	8
B. Extracellular hemoglobin and its intrinsic toxicity	9
III. Protection mechanisms	12
A. Haptoglobin	12
B. Hemopexin	16
C. Heme oxygenase	17
D. Protective heme-binding mechanisms	18
E. Antioxidation mechanisms	20
F. α_1 -Microglobulin (A1M)	21
1. Structure	22
2. Synthesis	23
3. Gene	24
4. Expression and distribution	24
5. Immunoregulation	25
6. Cell receptor binding	26
7. Antioxidation mechanisms	26
a. Reduction	27
b. Heme binding and degradation	27
c. Radical scavenging	28
d. Up-regulation of the AMBP-gene	29
8. Protective effects	30
a. Hb- heme- and ROS induced cell- and tissue damage	30
b. Irradiated cell cultures	30
c. <i>Ex vivo</i> perfused placenta	31
d. Extracellular matrix and collagen fibrils	32
IV. Pathological conditions	32
A. Hemolytic anemias and transfusion reactions	33
1. Hemolytic anemia	33
2. Transfusion	35
3. Hemolysis and kidney function	37
4. Current treatment of hemolysis	37
B. Preeclampsia	38
1. Pathophysiology	38
2. Hemoglobin in PE placenta	40
3. Treatment of PE	42
C. Intraventricular hemorrhage in preterm infants	42
1. Mechanisms of white matter brain damage in the immature brain	43
2. Free hemoglobin and brain damage	44
3. Treatment of IVH	45
D. Chronic leg ulcers	45
1. Inflammation	45
2. Heme, iron and oxidative stress	46
E. Blood substitutes and oxygen therapeutics	48
V. α_1 -microglobulin – a novel therapeutic concept	52
A. Extravascular Hb-protective role of A1M	52

B. Studies in diseases	53
1. Chronic leg ulcers	53
2. Preeclampsia	54
3. Intraventricular hemorrhage in preterm infants	55
4. Hemolytic anemias and transfusion reactions	55
5. Blood substitutes	56
VI. Conclusion	56
Acknowledgments	58
Abbreviations	59
Figure Legends	60
References	67

I. Introduction

The transport of oxygen from the lungs to peripheral cells and tissues is performed by red blood cells (RBCs). To account for this task, the RBCs are packed with hemoglobin (Hb), a protein with a capacity to bind four oxygen molecules (O_2) each. OxyHb, i.e. the complex between O_2 and Hb, is a highly reactive molecule with many chemical and biological properties besides its oxygen-transporting function (21,47,50,235). Most of these reactions are dependent on the redox-activities of the iron atom, which is chelated by the heme-group in the center of each of the four globin-chains of a Hb-molecule. A somewhat simplified view of the role of Hb is that oxygen-binding/transport is the true physiological function, whereas all other reactivities are unwanted, potentially deleterious, side-effects. While encapsulated within the RBCs, the Hb-molecules are kept separated from the rest of the body and an arsenal of protective mechanisms inhibit the side-reactions and eliminate their products [cf. (64,248)]. However, Hb is a very abundant molecule, in fact, the concentration in blood is around 150 g/L (50) and an adult human body consequently contains almost one kg of the protein. Thus, significant amounts escape from the RBCs under healthy conditions and massive amounts can be released during pathological conditions involving hemolysis. Therefore, a number of defence mechanisms have evolved to counteract the chemical threat of extracellular Hb to exposed tissues.

The term "oxidative stress" is used to describe conditions with an abnormally high production of reductive and oxidative compounds and/or impaired antioxidative tissue defence systems (114). Major mediators of oxidative stress are reactive oxygen species (ROS) including free radicals, extremely reactive compounds due to the presence of unpaired electrons. ROS, oxidants, and free radicals react with proteins, DNA and other molecular components, which can lead to unwanted modifications of the target molecules and loss of their functions.

Formation of ROS and free radicals can be induced for example by metals and metal-containing compounds and proteins. Since Hb is a metal-containing protein, most of the deleterious reactions of Hb are reductive or oxidative, i.e. involves exchange of electrons, and Hb is therefore categorized as a major inducer of oxidative stress. In addition, the Hb-metabolites heme and free iron are also potent inducers of oxidative stress. Therefore, Hb-induced oxidative stress is associated with extensive cell- and tissue damage.

Many pathological situations involve abnormal levels of hemolysis and high levels of extracellular Hb. Below, five groups of such pathological conditions, diseases and iatrogenic conditions will be discussed. Although they have vastly different etiology, they share many symptoms and clinical sequelae as a consequence of the Hb-induced oxidative damage. 1) Various hemolytic anemias, including inherited RBC dysfunctions and deficiencies, autoimmune transfusion reactions and mechanical RBC disruption. 2) Preeclampsia is a pregnancy disease shown to be associated with elevated levels of fetal Hb (HbF), leaking from fetus to mother. 3) Intraventricular hemorrhage and subarachnoidal bleeding results in locally high concentrations of free Hb in cerebral cavities and these conditions are associated with severe cerebral damage and dysfunctions. 4) Chronic leg ulcers are inflammatory conditions that involve dermal and epidermal extravascularization of RBCs and hemolysis. 5) Massive amounts of cell-free Hb in the bloodstream are also found during clinical administration of "artificial blood components", i.e. Hb-based oxygen carriers, employed for treatment of blood-loss after trauma, etc.

It is the object of this article to give an overview of the toxic reactions of extracellular Hb focussing on oxidative stress-inducing reactions, and of defence mechanisms that protect human tissues against these reactions with a special emphasis on the ubiquitous plasma- and

tissue-protein α_1 -microglobulin (A1M)(82,220,339,341). We will also describe the five groups of diseases and iatrogenic conditions mentioned above, in which Hb-induced oxidative stress is a major pathological factor. Finally, possible therapies for these types of pathological conditions will be discussed, giving special attention to the heme-binding and radical-scavenging effects of A1M.

II. Hemoglobin, heme and oxidative stress.

A. Red blood cells and hemoglobin.

Human RBCs have a life-span of ~120 days in circulation, after which senescent RBCs are phagocytized by resident macrophages in the reticuloendothelial system, spleen and bone marrow (37,47). The main constituent of RBCs is Hb and the primary function of this important protein is to bind O_2 reversibly and hence to function as an O_2 -transport protein. The Hb molecule is an assembly of four globular protein subunits where each protein chain arranges into a set of α -helix structural segments connected together in a globin fold arrangement (117,276). This folding pattern creates a pocket that very efficiently binds a heme group. The heme group consists of an iron atom that is coordinated by four nitrogen atoms of a tetrapyrrole ring (constituting the heme moiety) and a nitrogen of a histidine imidazole ring, linking the heme to the globular protein structure. This leaves one coordination-site free to reversibly bind gaseous molecules, normally O_2 , but carbon monoxide (CO) and nitric oxide (NO) can also be bound to the iron (236). In adults, the most common Hb isoform is a tetramer denoted HbA, which consists of two α - and two β -subunits ($\alpha_2\beta_2$) (see Fig. 1). In the fetus, the Hb molecule is made up by two α -chains and two γ -chains ($\alpha_2\gamma_2$, denoted HbF). The γ -chains are gradually replaced by β -chains at birth, and as the infant grows (50). HbF has a higher affinity for oxygen and a lower dissociation constant

required for sufficient oxygenation of the fetus *in utero*. In healthy adults HbF is restricted to RBCs called F-cells, which constitute a limited proportion of the total red cell mass (approximately 1% of total adult Hb is HbF).

1. Hemoglobin variants

Besides HbA and HbF, there are also a number of Hb variants that are found in different disease conditions. Hb variants occur when genetic changes in the globin genes cause alterations in the amino acid sequences of the globin proteins. These changes may affect the structure of the Hb, its behavior, its production rate, and/or its stability. Thus, more than 1400 hemoglobinopathies, inherited disorders of Hb, and thalassemias, conditions with reduced Hb levels in blood, are caused by different mutations of α -chains and β -chains, respectively, and are listed at the HbVar database (93,101). Their severity is governed by the number of genes affected. β -chain Hb variants are inherited in an autosomal recessive fashion. Several hundred β -chain Hb variants have been documented, however, only a few are common and clinically significant. A well described β -chain related disease is sickle cell disease. This disease is caused by a substitution of a single amino acid (Glu6Val) in the β -chain, that results in the HbS variant, which is associated with malformation and destabilization of RBCs (301). HbE (Glu26Lys) is the most common β -chain variant which manifests in an wide range of clinical symptoms, from asymptomatic to severe thalassemias depending on the genetic background (246,313). HbC (Glu6Lys) is another common β -chain variant causing aggregation of intracellular Hb resulting in moderate reduction of RBC plasticity in sickle cell HbC disease (298).


B. Extracellular hemoglobin and its intrinsic toxicity.

Although Hb is vital for delivery of O₂, it needs to be strictly compartmentalized within the RBCs in order to be kept in a functional non-toxic state. The balance between normal physiology and toxicity is largely dependent on the physical and biochemical barrier, presented by RBCs, separating Hb from the extra-RBC environment. Furthermore, RBCs contain a unique setup of enzymatic mechanisms that preserves Hb in this functional non-toxic state (248). However, if these systems are disturbed, for instance as a consequence of intravascular hemolysis, a number of toxic, potentially dangerous side effects, originating from the rich chemistry associated with the free radical reactivity of the heme group, can occur (see Fig. 1). The key chemistry involved in mediating the intrinsic toxicity of Hb is outlined below. First the ferrous iron (Fe²⁺) in the oxyHb molecule spontaneously auto-oxidizes, producing the ferric protein (Fe³⁺, metHb) and the superoxide radical.



Ferrous heme \Rightarrow ferric heme + superoxide radical

The superoxide then dismutates to hydrogen peroxide (H₂O₂); and the ferric protein can react with this peroxide, generating a reactive ferryl iron (Fe⁴⁺) and free radical bound to the protein (R^{·+})(66).



Globin + ferric heme + peroxide \Rightarrow globin radical + ferryl heme + water

Both the protein-bound radical (R^{·+}) and the ferryl iron (Fe⁴⁺ = O²⁻) are strongly oxidizing molecules that have pro-inflammatory, pro-oxidant and toxic effects and are capable of

damaging and disrupting cell membranes (230,235,299). The metabolites of Hb, i.e. free heme and iron, have the ability to damage lipids, proteins and DNA through direct oxidation and/or generation of ROS (4,69,261). Due to the lipophilic nature of the heme-group, it intercalates membranes and has destabilizing effects on the cytoskeleton (134). Because of their high redox-potentials, both the protein-based radical and ferryl heme can induce a wide range of oxidation reactions. When these oxidation reactions occur *in vivo* they have the ability to induce a wide array of physiological and pathophysiological responses. It has become apparent that many of these responses involve the formation of signaling molecules, often derived from oxidation of lipids (229). Lipid oxidation products are of immense physiological importance because of their role in cell-signaling pathways. The central concept underlying this field of “redox cell signaling” is that changes in the reductive or oxidative capacity of the cell lead to posttranslational modification of proteins by ROS species (67,160,272). These posttranslational modifications, which can be induced by Hb redox-chemistry, are emerging as an important field of research in pathologies such as atherosclerosis, ischemia/reperfusion injuries, posthemolytic injuries, and blood substitutes. Hb also binds and reacts with nitric oxide (144). It has been described to react with NO in one of two possible reactions:



High-affinity binding of NO to deoxyferrous Hb



Conversion of NO to nitrate by oxyferrous Hb

Since NO is a highly vasoactive molecule, the binding of NO by Hb affects the vasomotor tone, and hemolysis and elevated extracellular Hb-concentrations are therefore associated with hypertension. This has been described in for example sickle cell disease (239) and preeclampsia [(218); see also below]. Normally, the compartmentalization of Hb in the erythrocyte limits NO scavenging *in vivo* and subsequent vasoconstriction (162,304). Interestingly, it has also been proposed that NO may function as an antioxidant, playing a role in the mechanism to detoxify high oxidation states of Hb under oxidative conditions (235).

An important group of oxidant molecules in humans and animals is ROS, which includes for instance H₂O₂, superoxide and the hydroxyl radical. Superoxide and the hydroxyl radical are also categorized as free radicals, a group of molecules that are highly reactive due to the presence of unpaired electrons in their outer electron shells. All these molecules have the potential to react with proteins, DNA and other molecular cell and tissue components, leading to unwanted modifications and ultimately loss of function (Fig. 2). The human body is constantly exposed to free radicals and oxidants, both exogenously via the environment (food, air, smoke, irradiation, etc.) and endogenously as by-products of normal metabolism. As these molecules are reactive and potentially dangerous they need to be balanced by the activity of antioxidants, protective factors that eliminate oxidants or prevent their oxidation reactions [(113), see also below]. Thus, it is extremely important that cells maintain a well-controlled redox-balance. An imbalance in the redox-system will induce oxidative stress, a condition characterized by a surplus of unrestrained oxidants, ROS and free radicals. Several physiological processes, such as inflammation, ischemia/reperfusion, diabetes, obesity and kidney insufficiency, are associated with oxidative stress. Furthermore, hemolytic diseases involving destruction of erythrocytes in an uncontrolled manner, leads to high levels of cell-free Hb, as described above. Additionally, neutrophil activation during infections results in

oxidative stress, mainly due to production of superoxide anions and hypochloric acid by the enzymes NADPH oxidase and myeloperoxidase (19,20,51,202,269).

III. Protection mechanisms.

In humans, there are several Hb- and heme-detoxification systems described (Fig. 3). Haptoglobin (Hp), perhaps the most well-investigated Hb-clearance system, binds extracellular Hb in blood (21,156) and the resulting Hp-Hb complex is cleared from blood by binding to the macrophage receptor CD163 (146). Excess oxyHb that is not cleared by Hp undergoes auto-oxidation reactions resulting in ROS and free heme-groups. A complex network of antioxidation mechanisms (such as superoxide dismutase (SOD), glutathione peroxidase (Gpx) and catalase) inhibit and eliminate the oxidative compounds and repair the oxidative damage caused by them. Free heme in blood is sequestered by albumin and hemopexin (Hpx) (21,72,149) and the Hpx-heme complex is cleared from the circulation by the hepatocyte receptor CD91 (127). Intracellularly, heme oxygenase (HO) is the most essential heme catabolic protein, converting heme to free iron, biliverdin and CO (290,300). The plasma and tissue protein A1M (333) was described to bind and degrade heme and has the ability to reduce metHb (8,9). Recent reports suggest that a truncated form of A1M is involved in extracellular heme catabolism (8). The heme-binding and putative heme cleavage properties of A1M provide an additional physiologic protection mechanism against extracellular heme (216,220).

A. Haptoglobin

Several proteins are able to protect against Hb toxicity, and one of the most powerful is Hp (235,312). Hp is synthesized predominantly in the liver by hepatocytes. It is an abundant

acute phase protein and its levels in the blood-circulation vary greatly within humans, ranging from 0.3 to 3.0 mg/ml. In addition, the synthesis of Hp can be rapidly increased in response to e.g. inflammatory signaling. If free Hb is released into the circulation, it binds immediately to Hp to form an Hp-Hb complex, which subsequently is removed by the CD163 receptor expressed mainly on monocytes, macrophages and Kupfer cells in the liver (107,146,161,206,207,311). The binding between Hp and Hb is among the strongest non-covalent interactions known in biological systems with an estimated K_d of 10^{-15} and with slow dissociation rates. If Hp becomes depleted as a consequence of hemolysis, Hb accumulates in the kidney and is secreted in the urine. Hp thereby offers an effective protection against kidney damage in the first phases of moderate Hb release.

The Hp monomeric protein consists of two chains; one α -chain and one β -chain, expressed as a single precursor protein (Fig. 4). Two human Hp-gene alleles, Hp 1 and Hp 2, encode two different α -chains, α -1 with $M_r \approx 9$ kDa, and α -2 with $M_r \approx 16$ kDa, whereas the β -chain has an $M_r \approx 33$ kDa in both alleles. In mammals, the α -chains and the β -chains are linked together by a disulfide bond and in most species a disulfide bond also crosslinks the α -chains to each other forming a dimer composed of two $\alpha\beta$ -subunits. Additionally, Hp can form larger aggregates where the stoichiometry of the polymer is genetically determined and controlled primarily by the α -chain (194,208,230). Thus, Hp 1-1 homozygous individuals, having the Hp 1-allele on both chromosomes, form dimers via a single disulfide bond between the α -chains, whereas Hp 1-2 and Hp 2-2 individuals form polymers of higher order because the α -2 chains can form two disulfide bonds each, linking together the polymers. Hp binds Hb $\alpha\beta$ -dimers (48). All monomeric forms of Hp, and dimeric Hp 1-1, bind the Hb-dimers with high affinity. The polymeric forms of Hp can also bind to Hb, though at much lower affinities than monomers and dimers.

Several other functional differences between Hp phenotypes have been reported that could have important biological and clinical consequences. Hp polymorphism, not only linked to different subunit interactions but also to different glycosylation profiles, is associated with the prevalence and clinical evolution of many inflammatory diseases, including infections, atherosclerosis, and autoimmune disorders [c.f. (70)]. These differences are explained by an Hp-dependent modulation of oxidative stress (see below) and prostaglandin synthesis. For example, the distribution of highly polymeric Hp proteins in extravascular fluids is restricted by their molecular mass. Consequently, the Hb-protective capacity in body fluids is less efficient in Hp 2-2 individuals. Other reported differences among phenotypes include antigenic activities as well as binding to macrophage receptors including CD163 (206). The Hb–Hp complex impairs filtration and clearance of Hb dimers by the kidney, and instead directs Hb to CD163 on macrophages for a process of endocytosis and final degradation (138). Within macrophages the HO enzyme (see below) breaks down the heme-group of Hb into bilirubin and CO which both have been shown to have antioxidant and vasodilatory benefits (206). Therefore, this clearance of Hb indirectly protects against heme-mediated oxidative damage as well as other oxidative enzymatic reactions of Hb, described above, and thus Hp can be viewed as part of a human antioxidation mechanism [c.f. (161)]. Direct binding of Hb to CD163 has also been reported as a secondary clearing mechanism in addition to the Hp–Hb complex formation and subsequent clearance by the macrophages (206). The binding of Hp-Hb to CD163 results in the production of several cytokines, antioxidative genes and non-inflammatory mediators. Particularly, the induction of HO-1 is notable (260).

The literature suggests that O₂-binding- and redox-reactions of Hb are not inhibited by Hp-binding. In addition, upon Hp-Hb complex formation, the Hb moiety retains its NO-binding

characteristics (23). Likewise, the Hp-Hb complex can still react with H₂O₂ (46). In spite of this, Hp has been shown to inhibit Hb-induced lipid peroxidation (112,194) and it may be speculated that Hp achieves such antioxidation by for instance interfering with the diffusion of radical species from the heme center to surrounding tissue components (46,161).

Interestingly, it has been proposed that Hp may also protect the Hb-molecule from oxidative attack, including auto-oxidation. In addition to a site-specific protection of amino acids that are prone to oxidation on the surface of the Hb molecule, Hp allows the heme active site to operate unhindered leading to the elimination of oxidants (46). Recent work on the interactions of Hp with Hb has produced a model which gives a structural basis of the Hp mode of action in protecting Hb against oxidative toxicity. Mutagenesis studies have thus confirmed that a tyrosine residue at the α -subunit of Hb (α Tyr42) acts as a conduit for electron transfer to and from the heme which facilitates the auto-reduction of the ferryl Hb (237). In addition, EPR studies have shown that this electron transfer is also diverted to another Tyr residue on the β -subunit, β Tyr145, when Hb is complexed with Hp. Upon addition of H₂O₂, Hp stabilizes the ferryl-induced free radicals on β Tyr145. This radical reactivity may ultimately be directed from Hb to Hp resulting in a safer and more stable redox inactive Hb molecule (7)(Cooper et al, submitted).

It has recently been shown in two animal models, dogs and guinea pigs, that Hp can limit the toxic effects of cell-free Hb after infusion (33). Data obtained from both models showed that the formation of Hp-Hb complexes reduced the hypertensive response during Hb exposure, and reduced Hb toxicity in the kidney. Infusion of Hp may thus be explored to control adverse blood pressure responses and tissue toxicity effects associated with hematological disorders and diseases. Indeed, it was suggested that Hp-supplementation may be employed to counter

the adverse effects of the high loads of free Hb associated with hemoglobin-based oxygen carrier (HBOC) therapy (33); see also below, section IV.E).

B. Hemopexin.

Heme scavenging by Hpx (together with HDL, LDL, plasma albumin and A1M) provides protection against oxidative damage, limits access of pathogens to heme, and contributes to iron homeostasis by recycling the heme iron. During the first seconds after heme appearance in plasma, more than 80% binds to HDL and LDL, and 20% binds to Hpx and plasma albumin. Heme then transits to Hpx or is slowly removed from HDL and LDL by Hpx (21,26,72,103,188,196). Human Hpx consists of a single polypeptide chain containing 439 amino acids residues with six intrachain disulfide bridges (12). The polypeptide chain contains two homologous domains of about 200 residues each, joined by a 20-residue linker (295). The crystal structure of isolated Hpx C-terminal domain (86) showed that these domains have a four-bladed β -propeller fold that is a variant of the larger β -propeller domains found in heterotrimeric G proteins, clathrin, and integrins (151,274,292,295). Such domains appear to be of major importance in cell biology, especially as mediators of protein–protein interactions (273). Each propeller blade comprises a four-stranded, antiparallel β -sheet, with the first and fourth blades tied together by disulfide bonds. The heme binds between the two propeller domains. Two histidines coordinate the heme iron, giving a stable bis-histidyl Fe(3+) complex.

A striking feature of the heme-binding site is the preponderance of aromatic and basic residues (224). The construction and location of the heme pocket clearly relates to the requirements of high affinity but reversible heme-binding by Hpx and fits perfectly with the design of other high-affinity binding proteins such as the bacterial periplasmic binding

proteins (232) and iron-binding transferrins (15). Heme-binding and release results from opening and closing of the heme binding pocket, through movement of the two β -propeller domains and/or the interdomain linker peptide (188,224). Hpx delivers the bound heme upon interaction with its specific receptor, identified as the LDL receptor-related protein CD91, where it is internalized by receptor-mediated endocytosis. The human Hpx-heme receptor is expressed in several cell types including macrophages, hepatocytes, neurons, and syncytiotrophoblasts. After delivering the heme, the heme-group is degraded and intact Hpx can be released back into the bloodstream (21,72,188,295). The Hpx-heme complex removal by a receptor-mediated pathway shows striking similarities to the CD163-mediated Hp-Hb clearance in macrophages (127).

C. Heme oxygenases.

HO are cytosolic enzymes that participate in heme-detoxification by binding and degrading the free heme-group [(290); recently reviewed in (1,105,316)]. Two isoforms of HO have been described: HO-1, the expression of which is induced by a variety of conditions of environmental stress (183,324), whereas HO-2 is constitutively expressed (184,296). A third genetic variant, HO-3, has been described, but is most likely a non-expressed pseudogene (120). HO-1 and HO-2 are separate gene products and have different amino acid sequences, structure and size, but have similar mechanisms of heme catalytic activity and specificity.

Both enzymes are anchored to the membrane of the endoplasmic reticulum via a C-terminal hydrophobic domain (192). The enzymes operate in concert with microsomal NADPH-cytochrome P450 reductase to convert heme to biliverdin, CO and Fe^{2+} , utilizing three molecules each of O_2 and NADPH for each molecule of heme (Fig. 5; (209,257,291)). The products of HO-activity have important beneficial physiological activities providing further

antioxidation effects besides the mere elimination of heme. Biliverdin is reduced to bilirubin by biliverdin reductase (150) and bilirubin is a powerful physiological antioxidant (279). CO has been reported to have both pro-oxidant and antioxidant effects, mostly as a result of binding to heme-proteins, replacing molecular O₂ [reviewed in (228)]. Indeed, CO had protective effects against lung injury (95,221). CO also induces vasodilation by binding to the heme-protein guanylyl cyclase (182). Finally, CO participates in a positive feed-forward loop by inducing an up-regulation of HO-1 (145). The release of free Fe²⁺ iron is closely linked to synthesis of ferritin, an iron-sequestering protein. Thus, HO-1 induction stimulates ferritin expression (81), and inhibition of HO-1 leads to ferritin down-regulation (314).

Under resting, non-stressed conditions, HO-2 is the predominating form in all organs except the spleen (36), and the highest activities are found in spleen, testes and brain (181). The expression of HO-1 is ubiquitously up-regulated by a multitude of stimuli related to oxidative stress, including high concentrations of free Hb and heme [(17,251) reviewed in (105,180,181,316)]. In fact, elevated levels of HO-1-messenger RNA is commonly used as a biomarker of oxidative stress. In contrast, transcription of the HO-2 gene is only responsive to stimuli from adrenal glucocorticoids (193,234).

D. Protective heme-binding mechanisms

Plasma albumin binds heme at a protein:heme ratio of 1:1 and 1:2, but with lower affinity than Hpx ($K_d \approx 10^{-8}$ M) (171,247). Since the plasma concentrations of albumin are high (30-50 mg/mL) the amounts of heme that can be bound by albumin are considerable, and the albumin-binding becomes physiologically important especially during conditions of heme-overload when the Hpx is depleted (265). This may explain the beneficial effects of albumin-infusion on cerebral malaria (185). Albumin partially inhibits the toxic effects of heme (109)

but there are no indications of a targeted transport function or specific enzymatic inhibitory actions. Thus, the general belief is that the protective effect of albumin is a result of the protein functioning as a depot of free heme-groups.

Free heme in plasma is also bound rapidly, and with high affinity, to the lipoproteins LDL and HDL (196). The heme-group catalyzes oxidation of the lipoprotein particle and subsequent endothelial damage, and probably contributes to the development of atherosclerosis (27,55).

The protein α_1 -microglobulin (A1M) binds heme in plasma, extravascular fluids and cells with a $K_d 10^{-6}$ M (8,11,153). The binding has been shown to yield protection from the toxic effects of heme in various *in vitro* molecular, cell-culture and *ex vivo* organ systems. The structure, physiology and antioxidative properties of A1M are described in some detail below.

As argued in previous publications, the heme-group is probably not freely diffusable across membranes, [cf (105,155)]. Several heme-binding and heme-transporting proteins that mediate the uptake of extracellular heme to the cytosol have been described (233). Two cytosolic proteins, as well as HO-1, have been shown to have protective effects against free heme toxicity, peroxiredoxin I (Prx1 or HBP23) (80,129,133,205) and A1M (220), both of which are upregulated in response to elevated concentrations of free heme (130,217).

Interestingly, intracellular A1M can be derived by local synthesis in most cells but also by uptake of extracellular A1M produced in the liver and transported to tissues via blood (see below).

E. Antioxidation mechanisms

As described above, unsequestered Hb, heme and iron generate free radicals, mostly ROS.

While the Hb- heme- and iron-binding and sequestering mechanisms described above generally operate to prevent the formation of radicals and ROS, a network of antioxidation molecules, acting with many different mechanisms, co-operate to protect cells and tissues against the deleterious reactions of free radicals and ROS once they have been formed.

Several high molecular weight enzymatic antioxidants have been described. SOD catalyzes the conversion of superoxide radicals ($O_2^{\cdot-}$) to H_2O_2 [reviewed in (96)]. Three variants of SOD exist in mammals and birds: cytoplasmic Cu/ZnSOD (=SOD1)(191), mitochondrial MnSOD (=SOD2)(306), and extracellular Cu/ZnSOD (=SOD3)(186). Catalase, belonging to the functional group of peroxidases, is a heme-carrying enzyme catalyzing the conversion of two molecules of H_2O_2 to water plus O_2 [c.f. (61,104)]. Catalase is localized to peroxisomes of nucleated cells (263,332) where its main function is to eliminate the H_2O_2 formed in this organelle and to keep cytosolic H_2O_2 concentrations low. Gpx, the second most abundant peroxidase, is a ubiquitously expressed cytosolic and mitochondrial enzyme (174) catalysing the reduction of H_2O_2 by employing glutathione as an electron donating substrate. Albumin is generally regarded as an important non-enzymatic antioxidation factor that is present in high concentrations and hence buffers deleterious oxidation reactions in plasma and interstitial fluids.

Several non-enzymatic, low molecular weight radical scavengers operate either by direct reactions with ROS and non-ROS radicals, or by acting as cofactors in reactions with the enzymatic antioxidants. The most important is perhaps the cytosolic tripeptide glutathione (GSH). GSH is synthesized in the cytosol and provides reducing equivalents via its free thiol group (187). Thus, the ratio between GSH and its oxidized form, GSSG, constitutes a

cytosolic redox buffer which regulates the balance between antioxidant capacity and beneficial ROS-activity (78,135). Thioredoxin (Trx) is a ubiquitous 12 kDa polypeptide with a Cys-X-X-Cys-motif that acts as an electron-reservoir via a disulfide-exchange between the two thiol groups (123,211). High ratios between reduced and oxidized GSH and Trx, and thus a high antioxidant capacity, are maintained by the regenerating enzymes GSH reductase and Trx reductase, respectively, drawing reducing equivalents from cytosolic NADH and NADPH (18,147,199). The essential nutrients vitamin C (ascorbic acid) and vitamin E (tocopherol) are also generally believed to serve important functions as antioxidants (122,222). The water-soluble vitamin C is both a reducing agent which is transformed to dehydroascorbic acid when reacting with ROS/radicals, and a co-factor of several enzyme reactions, whereas the lipid-soluble vitamin E is an important lipid peroxidation radical scavenger. In addition, a large number of nutritional low molecular weight radical scavengers with antioxidant properties have been described, such as turmeric curcumin (330) and red wine resveratrol (137).

F. α_1 -Microglobulin (AIM)

AIM was discovered in human urine in the early 1970's and was named in the tradition of plasma proteins, reflecting its small size (26 kDa) and its electrophoretic migration slightly behind albumin (82). The protein was characterized by several research groups and is also known under the alternative names protein HC, "human complex-forming protein, heterogeneous in charge" (289) and α_1 -microglycoprotein (267). Two striking biochemical properties were described early: a yellow-brown colour and a pronounced charge heterogeneity (82,338).

A physiological role of A1M in heme- and radical-binding was only recently proposed. First, immunoregulatory (mainly suppressive) properties of A1M were identified during the 1980's and 1990's, but did not seem distinct or strong enough to constitute the major function of the molecule *in vivo*. Then several reports during 2000's showed enzymatic reductase activities, as well as heme- and radical-scavenging properties. Recent investigations have described strong protective effects of A1M in molecular, cell- and organ systems and it has become clear that the protein has a role as an antioxidant employing several chemical mechanisms. The precise reactions of this protein/enzyme are still largely unknown, however, as is the relationship between its protective and immunoregulatory biological effects.

1. Structure

Human A1M has a molecular weight of 26 kDa, and consists of a peptide chain with 183 amino acids (82,140,283). The peptide is glycosylated with oligosaccharides in three positions, two sialylated complex type, diantennary and triantennary carbohydrates linked to Asn17 and Asn96 and one O-linked oligosaccharide linked to Thr5 (13,83). The protein is well conserved during evolution and an A1M homologue has been isolated and sequenced from humans and other mammals, birds, amphibians and fish (58,115,128,141,168,169,200,286,320,334,335,340). No crystal structure has yet been obtained but a model of the three-dimensional structure was developed based on its membership in the lipocalin protein family (Fig. 6). The lipocalins are a group of 40-50 proteins found in all branches of life (bacteria, plants, fungi, animals) (92,337) and twelve human genes are known. The lipocalins share a common protein fold: a single polypeptide, 150-190 amino acid residues, forms an antiparallel eight-stranded β -barrel with one closed and one open end. The interior of the barrel often serves as a binding site for small hydrophobic compounds (99). The Cys34 side-chain of A1M (Fig. 6), found in all species,

was suggested to be located on a flexible loop near the lipocalin pocket and is solvent exposed (318), and involved in complex formation with other proteins, redox-reactions and radical binding (8,9,31,110,341) (see also below). Covalent modifications of the Lys 69, Lys 92, Lys118, Lys 130 and Cys 34 side-chains (Fig. 6) have been found in human A1M isolated from urine, and these "chromophores" have been proposed to cause both the yellow-brown colour and the charge-heterogeneity of the protein (29,83). The modifications probably represent the oxidized degradation products of organic radicals covalently trapped by A1M in vivo (see further below).

2. Synthesis

The gene for A1M is called *AMB*P (Alpha-1-Microglobulin-Bikunin Precursor gene) because it also encodes bikunin, another plasma protein (140) (Fig. 7). Bikunin is a Kunitz-type protease inhibitor (271) and a structural component of extracellular matrix (63). Bikunin is the common subunit of a group of protein/carbohydrate complexes that constitute plasma proteins of the inter- α -inhibitor family (94,254). Transcription of the *AMB*P gene produces an mRNA that is translated into a precursor protein consisting of the A1M and bikunin proteins connected by a linker tripeptide (166). The last amino acid of A1M, Arg, plus the linker tripeptide constitute a basic cleavage site, R-V-R-R, that conforms to the consensus recognition sequence of furin and other proprotein convertases (PCs)(266). Before secretion, the A1M- and bikunin-components are therefore separated by proteolytic cleavage by furin or another PC in the late phase of post-translational processing in the Golgi-system (39). No physical or functional connection has been found between A1M and bikunin after secretion and the reason for their co-synthesis is not understood. It has also been shown in different expression systems that both A1M and bikunin can be expressed alone (38,294). In spite of

this, the A1M/bikunin genetic construction is conserved in all species where A1M has been found.

3. Gene

The *AMBP* gene has been cloned from man (77) and mouse (167). In both species, the gene contains ten exons, of which the six first code for A1M and the four last code for the linker peptide and bikunin (Fig. 7). The *AMBP* gene has been mapped to the 9q32-33 region in man (76) and to chromosome 4 in mouse (256), in both species in regions where other lipocalin genes have been found to be clustered. The A1M-exons are separated from the bikunin-exons by intron F, which contains retroposons and other repeated structures, suggesting that it is a recombinatorial hot-spot (167). This could have provided the basis for a fusion between an ancestral lipocalin gene (A1M) with an ancestral Kunitz inhibitor gene (bikunin). The pronounced expression of A1M in liver is dependent on a strong liver-specific enhancer located about 2500 nt upstream of the start-codon, and contains eight binding sites for hepatocyte-enriched nuclear factors, HNFs (249,250).

4. Expression and distribution.

Liver, blood plasma, and kidney (Fig. 8) are the major sites of A1M localization (333). The liver is the major site of synthesis in adult tissues (288,334). The protein is then secreted to the blood, where it exists in free form as well as in disulfide-linked high molecular weight complexes with other plasma proteins. The total concentration of A1M in human plasma is approximately 2 μM (74), and approximately 50% of the total plasma A1M is linked to IgA via a reduction-resistant disulfide bond, 7% to albumin and 1% to prothrombin (31,54,110,111). High molecular weight forms of A1M have been found in all species examined. In rat serum, A1M can be found covalently linked by a disulfide link to fibronectin

(88) and by a reduction resistant bond to the proteinase inhibitor α_1 -inhibitor-3, a homologue of human α_2 -macroglobulin (89). A1M complexes were also detected in plaice serum (169). Thus, the complex forming ability of A1M is conserved from fish to man, although the identities of the complex partners are not conserved. Both free A1M and the complexed forms are rapidly equilibrated between the intra- and extra-vascular compartments and their half-lives in blood are only 2-3 min (154,307). Free monomeric A1M passes almost unhindered through the glomerular membranes out into the primary urine, where most of it is reabsorbed by the proximal tubular cells and catabolized (210). Significant amounts, however, are excreted in the urine and the A1M-concentration in urine is a sensitive indicator of tubular renal damage (82).

Besides the liver, A1M mRNA has been detected in adult kidney (139,157), pancreas (30,132), stomach (286), placenta (218) and blood cells (157,217). The A1M protein has been identified in the perivascular connective tissue of most organs (30,212,213) and is especially abundant in epidermis of skin (11,34,216) and epithelium of the gut (35,154). It is often co-localized with elastin and collagen (190,212,213,216) and binds to collagen *in vitro* (216,258). A distribution of A1M at various interfaces between the cells of the body and the external environment (blood/tissue, air/tissue, intestinal lumen/villi), as well as at the interface between maternal blood and fetal tissues in placenta (30,190), is consistent with a protective role of A1M *in vivo* (see below).

5. Immunoregulation.

A1M inhibits central events of the immune response *in vitro*. Thus, the antigen-induced cell division and IL-2 production of lymphocytes were inhibited by A1M (178,179,308).

Inflammatory responses of blood cells were inhibited by A1M; these included migration

(177,178) and chemotaxis (195) of neutrophil granulocytes and the production of free radicals and IL-1 β by peripheral lymphocytes/monocytes (258). It is possible that the immunoregulatory effects of A1M are related to its antioxidant and reductase activities, as ROS have been shown to be involved in cell signalling during lymphocyte activation [reviewed in (78,317)].

6. Cell receptor binding.

A1M has been shown to bind to the surface of various cells, including blood lymphocytes, neutrophils, blood cell lines and keratinocytes (24,90,195,216,220,307,308). The binding is specific for A1M, saturable and trypsin-sensitive, suggesting the presence of an A1M-receptor on these cells. The affinity of A1M for its receptor on T cells and mouse peripheral lymphocytes was estimated to 10^4 - 10^5 M $^{-1}$ (25,308) and somewhat higher on the histiocyte cell line U937, 10^7 M $^{-1}$ (90). The receptor has not yet been identified, however. Internalization of A1M from medium to the cytosol has been shown in erythroid K562 cells (220), hepatoma HepG2 cells; (217), and primary skin keratinocytes (216).

7. Antioxidation mechanisms of A1M.

As mentioned above, several reports during the last 10 years have shown enzymatic reductase activities, as well as heme- and radical-binding properties, of A1M. Recent investigations have also described strong protective effects of A1M in molecular, cellular and organ systems and it has become clear that the protein has a role as an antioxidant employing several chemical mechanisms. Figure 9 summarizes the reported mechanistic properties of A1M which may contribute to its possible role as a physiological antioxidant. These include *a*) reductase and dehydrogenase activity, *b*) heme-binding and degradation, *c*) covalent radical-scavenging, and *d*) physiological up-regulation of the AMBP-gene.

a. Reduction. Enzymatic reductase/dehydrogenase properties were recently described for A1M with a wide spectrum of organic and inorganic substrates (9,341). The protein was capable of reducing heme-proteins, free iron and the synthetic compounds nitroblue tetrazolium (NBT), using the electron donors ascorbate and NADH/NADPH as co-factors (9). Using mutated A1M-variants, it was shown that the reductase and dehydrogenase activities were dependent on the thiol group of Cys34 and the three lysyl residues of K92, K118 and K130 (9). According to the hypothetical structural model of A1M, these four side-chains are located in the vicinity of each other at the open end of the lipocalin barrel, Cys34 most likely on a flexible Ω -loop and Lys92, 118 and 130 close to the rim of the pocket (Fig.6). It was speculated that the role of the three Lys residues in the reaction may be to create a positive electrostatic environment around the Cys 34, thus lowering the pK_a of the thiol group and favouring its oxidation (9). This model of the three-dimensional structure of A1M suggest that it may be possible for Cys34 to interact both with the Lys-groups and the electron-accepting substrates. A1M also rapidly reduced the synthetic radical 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid) (ABTS), again in a reaction that was dependent on the free thiol group of C34 (341). Finally, the cell cytosol and thiol groups of cytosolic proteins of the erythroid cell line K562 were reduced by A1M (220).

b. Heme-binding and degradation. A1M binds heme with $K_d=10^{-6}$ M (153,216). The heme-binding was shown to occur to A1M in plasma from man, mouse, rat, guinea pig, cow, chicken and plaice (153), suggesting that this property is evolutionarily conserved. Both free, monomeric A1M and the IgA-A1M complex could bind to the heme-group (8,153). Apart from this non-covalent, relatively weak interaction, a stronger binding was seen which remained after boiling in SDS, suggestive of a covalent interaction (8,153). Interestingly, a processed form of A1M, called t-A1M (t=truncated), lacking the C-terminal tetrapeptide

(leucine-isoleucine-proline-arginine; LIPR) of full-length A1M, was generated by a reaction with lysed RBCs or with purified Hb (8). The processed t-A1M also binds heme, but was capable of degrading the heme-group (8). The t-A1M-heme complex showed a time-dependent spectral rearrangement suggestive of degradation of heme concomitantly with formation of a heterogeneous chromophore associated with the protein (8). The t-A1M form is found in urine, skin and placenta and is thus formed *in vivo* (8,11,173,218). Together, these reports suggest that A1M is involved in extracellular heme catabolism by binding to free heme-groups, and that a proteolytically activated, heme-degrading form of A1M is induced by free, extracellular Hb. The degradation mechanisms are as yet unknown, but may involve C34-dependent reductase reactions and covalent binding of degradation products to side-chains of A1M.

c. Radical scavenging. A1M reacts with organic radicals in a reaction that involves C34-catalyzed reduction and results in covalent trapping of radical metabolites to several amino acid side-chains on A1M (Fig. 10). To study the radical scavenging mechanism, the reaction between A1M and the synthetic free radical ABTS was characterized in detail (341). It was shown that A1M reduced 5-6 molecules of the ABTS-radical in a rapid semi-catalytic reaction, and covalently trapped 3 additional ABTS-radicals by attaching them onto tyrosyl side-chains in a modified, oxidized form giving the protein a purple colour. Two of the modified Tyr-residues were identified and localized on the A1M-molecule by mass-spectrometry (Tyr22, Tyr132). A possible reaction scheme involving an intra-molecular transfer of electrons between the C34 thiol and the tyrosyl side-chains is shown in Fig 10. Besides these two Tyr-residues, previous reports suggest that several Lys-side-chains may be modified by radical trapping. Lysyl side-chains in urinary A1M from hemodialysis patients were shown to be modified by the kynurenine derivate 3-hydroxy-kynurenin (253). 3-

hydroxy-kynurenin is a tryptophan metabolite that has a propensity to form free radicals (131,215,305) and is found in elevated concentrations in plasma of for example hemodialysis patients (226). Furthermore, although the structures of the modifications could not be identified, the side-chains of Lys69 (336), Lys92, Lys118 and Lys130 were shown to be covalently modified in human A1M isolated from urine (29,83). The latter three Lys-residues may therefore serve a dual role both in activation of the Cys34 thiol by lowering its pKa (9) and as electron-donors in the radical-trapping mechanism. In addition, the heme-degradation products strongly attached to A1M, and especially to its t-A1M form described above, could be formed by reactions that are related this reductase/covalent trapping mechanism. The term "radical sink" was used to describe the radical binding property of A1M (341), referring to the fact that both the radicals and the A1M protein itself are electroneutral after the reactions and therefore do not present any oxidative stress to tissue components.

d. Up-regulation of the AMBP-gene. The expression of the A1M-gene (*AMBP*) was increased by exposure of hepatocyte-, histiocyte- and erythrocyte cell lines to oxy- and metHb, heme and ROS (217). The up-regulation, both on the mRNA- and protein levels, was reversed by reacting Hb with cyanide which prevents redox reactions (217). An up-regulation of A1M was also seen in primary keratinocytes in response to heme and ROS (216). The latter may be important for the antioxidation defense in the skin where keratinocytes constitute the major cell type. Recent results (218) show a correlation between the concentrations in plasma and placenta tissue, of A1M and Hb and between A1M and biomarkers of oxidative stress (protein carbonyl groups), respectively, in the human diseases preeclampsia (PE) (see also below), supporting the notion of an up-regulation *in vivo* of the AMBP-gene by elevated concentrations of free Hb and conditions of oxidative stress.

8. Protective effects of A1M.

Employing the antioxidation mechanisms described above, A1M can protect cells and tissues from Hb-, heme- and ROS-induced cell-damage [Fig. 9; (190,216,218,220)]. A unifying model of the physiological role of A1M is that it functions as a “radical sink”, continuously cleaning tissues from free radicals and oxidants, including free heme and ROS generated by extracellular Hb, heme and iron, and delivering the products to the kidneys for degradation and/or excretion. Below, findings of protective effects of A1M are summarized.

a. Protection against Hb- heme- and ROS induced cell- and tissue damage. Several reports describe *in vitro* protective actions of A1M in cell cultures stressed by exposure to Hb-, heme- and ROS. A1M prevented intracellular oxidation and up-regulation of HO-1 induced by heme, H₂O₂ and Fenton reaction-generated hydroxyl radicals in primary keratinocytes (216) and in the erythroid cell line K562 (220). Silencing of the endogenous A1M expression by addition of siRNA led to an increased cytosol oxidation (220). A1M also inhibited cell lysis of K562 cells caused by heme and cleared the cells from bound heme (220). It was shown that 200 μM heme killed 50% of the cells but adding 2 μM A1M lowered the number of dead cells to 15%, i.e. by approximately 70% (220). Similar effects were described in heme- and ROS-stressed skin explants (216). It was shown that A1M is localized ubiquitously in the dermal and epidermal layers of skin, and that the A1M-gene, which is expressed in keratinocytes, was up-regulated after exposure to heme and ROS. Exogeneously added A1M inhibited the heme- and ROS-induced ultrastructural damage, up-regulation of antioxidation- and cell cycle regulatory genes, and formation of protein carbonyl groups, a marker of oxidative stress, in skin (216).

b. Protection of irradiated cell cultures. A1M was also shown to inhibit the propagation of

cell-death induced by charged particle irradiation of HepG2 cell monolayers (219). The cells were irradiated with a low dose of alpha-particles at a small, restricted area. The directly hit cells were killed and the cells surrounding the irradiation area, not directly hit by the particles (= "bystander" cells), showed a delayed and slowly accumulating necrosis up to 3 days after the irradiation. Furthermore, a significant increase in apoptosis, protein carbonyl groups, and expression of the stress response genes HO-1, p21 and p53 was observed in the whole cell population. Addition of A1M reduced the amount of dead cells by approximately 50 % in irradiated cells and 100 % in bystander cells, and completely inhibited the irradiation-induced apoptosis, formation of carbonyl groups and upregulation of HO-1, p21 and p53. Irradiation induced an upregulation of endogeneous synthesis and secretion of A1M and an increased uptake of A1M from the medium. This study suggests that A1M protects against bystander cell killing by preventing the propagation of oxidants and ROS produced in the irradiated cells.

c. Protection of *ex vivo* perfused placenta. To study Hb-induced tissue damage in a more complex system than cell cultures, the dual placenta perfusion model (262) was employed. In this model, the fetal and maternal placental circulations are perfused separately and it is possible to study cell- and matrix structure and functions including the fetomaternal barrier. It was shown that *in vitro* perfusion of placentas with Hb in the fetal circulation led to a significant increase of the perfusion pressure, fetomaternal leakage of free Hb, morphological damage, and up-regulation of genes related to immune response, apoptosis, and oxidative stress (190). Simultaneous addition of A1M to the maternal circulation inhibited the Hb leakage, morphological damage and gene up-regulation. This study suggests that A1M may have therapeutic potential in pregnancy diseases involving placenta malfunction (see below).

d. Protection of extracellular matrix and collagen fibrils. The skin and placenta studies (190,216) also demonstrated a protective effect of the protein *vis-à-vis* extracellular matrix and collagen fibrils. In fact, disruption of collagen fibrils by Hb-, heme- and ROS-incubation was inhibited by A1M (Fig. 11). The fibrils could also be repaired by addition of A1M after the destruction already had taken place. It was speculated that oxidative lesions of the collagen monomers, induced by reactions with Hb, heme or ROS, may shift the equilibrium between monomers and fibrils towards the monomers by preventing their interaction, whereas A1M, by removing and/or reducing the oxidative modifications, promotes a shift of the equilibrium back towards fibril formation (216). Interestingly, perfusion of placentas with A1M induced a significant up-regulation of extracellular matrix genes, and a remarkable increase was observed in the number of collagen fibrils *vis-à-vis* Hb-perfusion, as monitored by electron microscopy (190).

IV. Pathological conditions

Hemolysis and elevated concentrations of free, extracellular Hb in blood and extravascular fluids are seen in many different pathological conditions. Five diseases/groups of such conditions will be described in this section. All are associated with a unique set of clinical scenarios and have a variety of underlying causes but a common denominator is the increased oxidative stress that the extracellular Hb induces, and with a few exceptions hypertension and renal insufficiency are a common theme. There is still a great need for development of treatment options targeting the oxidative environment that affects patients with hemolytic disease independent of its underlying causes.

A. Hemolytic anemias and transfusion reactions

The clinical definition of hemolysis is a reduction of the life span of RBCs to less than the normal range of 100-120 days (227). This premature and uncontrolled destruction of RBCs can lead to symptoms and laboratory signs with different degrees of severity, from none or very mild to fatal reactions. As outlined above, the extracellular Hb released from rupturing RBCs is associated with generation of various ROS which can induce oxidative damage to matrix molecules, cell membranes and other tissue components. Thus, in general, one of the key features of the hemolytic state, independent of its origin, is that it puts the individual under heavy oxidative stress (91). In this section, we will briefly outline various clinical situations where hemolysis is associated not only with increased extracellular Hb but also with anemia, i.e. loss of total blood Hb, including transfusion-induced hemolysis.

1. Hemolytic anemia

The diagnosis of hemolytic anemia is justified when shortened RBC survival constitutes a major underlying mechanism in explaining a low blood Hb-value. The first case of hemolytic anemia was described 140 years ago and the report includes the hallmark observation of splenomegaly without hepatomegaly and acute attacks of jaundice followed by the passing of reddish brown urine (303). There are many ways to categorize hemolytic disease but a major distinction is made between hereditary and acquired hemolytic anemias. The former includes various genetic abnormalities of intracellular RBC proteins including enzymes (287) and structural proteins (73), additionally alterations in proteins exposed on the extracellular surface of the cell membrane can lead to shortened circulation of RBCs (14,44). The latter group of acquired hemolytic anemias encompasses many different kinds of cell destruction including mechanical lysis. This can be observed in population groups as diverse as marathon runners where RBCs are destroyed due to repeated trauma when passing through the blood

vessels in the athletes' feet, and patients whose RBCs are destroyed by non-biological heart valves or during extracorporeal circulation. Yet another classification targets the location of hemolysis, whether the RBCs are lysed in the blood vessel (intravascular) or outside the circulation (extravascular), see Fig. 12. Obviously, the mechanical lysis described above occurs intravascularly while spherocytosis, a state in which RBCs do not exist in their normal biconcave discoid shape but are spherical due to errors in the assembly of intracellular structural proteins (73), will mostly result in extravascular destruction and lead to an enlarged spleen. Finally, a very important categorization of immune hemolytic anemias (AIHA) is made on the basis of auto- vs. alloimmunity and the nature of the immunoglobulins involved in the lytic process. AIHA occurs when antibodies are directed against the individual's own RBCs. From a clinical perspective, this is one of the most common patient groups related to hemolysis in the Western world. AIHA can be categorized as being of the *warm* or *cold type*, so named because the autoantibodies in each type react optimally at high or low temperature, respectively. In the warm type of AIHA, the targets of the autoimmune response are protein epitopes (the vast majority of which are Rh-related) on the RBC surface and the antibodies are usually of IgG type. These antibodies may sometimes activate complement but usually do not. Instead, the main mechanism behind the accelerated RBC destruction in this form is extravascular lysis following Fc receptor recognition of the IgG-coated RBC by immune cells residing in the spleen (Fig. 12). Conversely, the cold type of AIHA is due to IgM antibodies directed at RBC auto-epitopes (mostly carbohydrate structures) and the result is intravascular lysis due to the ability of IgM to activate the complement cascade (Fig. 12). It should be noted that there are cases with mixed type AIHA where both IgM and IgG mechanisms are at play. There are also other kinds of intravascular hemolysis that are not antibody-mediated but still involve the complement system. For instance, paroxysmal nocturnal hemoglobinuria (PNH) is due to acquired somatic mutations in the *PIG-A* gene which normally codes for an enzyme

that is necessary for formation of glycosylphosphatidylinositol anchors between proteins and the lipid bilayer of the cell membrane (40). When an abnormal bone marrow progenitor clone is formed where this capacity is lost, the progeny RBCs will lack a whole range of surface proteins including several blood-group-carrying molecules and most importantly in this perspective a selection of complement regulatory proteins. In their absence, the RBC becomes sensitive to complement-induced intravascular lysis and a key symptom is the passing of dark urine when the patient wakes up. In addition, there is a poorly understood relationship between PNH in both leukaemia development and the formation of life-threatening clots in the large vessels.

In addition to the above-mentioned causes of hemolysis, microbial agents are known to be able to induce hemolysis. This will not be discussed further here but some of the clinically very significant diseases (and agents) involved include malaria (*Plasmodium* species), hemolytic uremic syndrome (*E.coli*), dysentery (*Shigella*), pneumonia (*Mycoplasma*), fifth disease (*Parvovirus* B19) and hemorrhagic fevers (various viruses).

2. Transfusion

When RBCs from a blood donor are transfused to a patient, the donor blood unit could have been stored at 4 degrees for a maximum of 35-42 days depending on local regulations. Storage of blood units results in storage-time dependent suboptimal function and stability of the cellular elements in the stored blood. Moreover, even if there are regulations to ensure an optimal quality of the blood components transfused, a certain amount of Hb leakage from the cells is unavoidable and expected. A progressive oxidation of cytoskeletal proteins and accumulation of denatured Hb also occurs in the stored RBCs. In addition, a fair percentage of damaged RBCs will be lysed or cleared immediately from the circulation in the recipient upon

infusion. Thus, even if the best possible matching of blood groups is ensured by crossmatching, etc, at the blood center, the recipient will suffer an increased load of extracellular Hb following transfusion and consequently also the negative effects of oxidative stress. Despite this, millions of RBC units are given annually around the globe and transfusions are considered a prerequisite for many medical procedures, including surgery, safe obstetrics, chemotherapy treatment for cancer, as well as stem cell or organ transplantation. If chronic transfusions are given, the additional risk of iron overload is yet another threat which is also associated with increased oxidative stress and needs to be treated (40).

The most severe form of iatrogenic hemolysis can be observed if ABO-incompatible blood is transfused to a patient by mistake. This acute reaction is based on alloimmune RBC destruction and the presence of naturally-occurring, pre-existing IgM antibodies against the ABO antigens lacking in the recipient. Even though blood transfusion is a safe treatment in general today, ABO incompatibility unfortunately still results in death or injury of patients following transfusion of allogeneic blood (259,275). However, the consequences of this mismatched transfusion can vary from asymptomatic laboratory signs to death within minutes (165). The reasons why some patients react violently, with disseminated intravascular coagulation and shock, while other patients have no noticeable reaction remain obscure. However, it is known that if the incompatibility involves blood groups other than ABO, for instance Rh, Duffy or Kell, then the adverse reaction is typically, but not always, milder and slower. Thus, a delayed hemolytic transfusion reaction is much more likely to be due to an anamnestic IgG response to protein epitopes, which will result in opsonisation of the allogeneic RBCs with IgG against the foreign, non-self blood group antigen and extravascular removal of the targeted cells. A transfusion-unrelated variant of this phenomenon is the

hemolytic disease of the fetus and newborn (HFDN), where maternal IgG against paternal blood group antigens in the offspring passes the placenta and covers the fetal RBCs. This is a potentially lethal disease, since the macrophages of the reticuloendothelial system in the fetus will start phagocytosis of sensitised RBCs, with severe anemia and hepatosplenomegaly. If the Hb levels reach dangerously low values, *hydrops fetalis* may develop if treatment in the form of intrauterine, antigen-negative RBC units is not started. Hydrops is a critical state in which the fetus accumulates water because of the relative lack of RBCs able to circulate Hb.

3. Hemolysis and kidney function

Depending on what type of hemolysis (as discussed above) is encountered, elevated levels of Hb can be observed in the blood and/or urine. Plasma Hb is filtered through the glomeruli of the kidneys and re-absorbed by tubular cells where it may lead to precipitate formation termed hemosiderin during conditions with Hb-overload, causing oxidative damage and glomerular and tubular necrosis (85,164,285). These effects have mainly been reported in intravascular (136,158,214) but also in extravascular hemolysis (210). A particularly dramatic scenario is the renal consequences of an acute hemolytic transfusion reaction, after which a possible outcome is that the patient's life is saved but the massive release of extracellular Hb leads to kidney failure and the need for dialysis and potentially a subsequent kidney transplant.

4. Current treatment of hemolysis

There are various means of treating patients undergoing hemolysis but none of these are specifically aimed at reducing the oxidative stress or inactivating the toxic free Hb molecules. Following massive intravascular lysis associated with ABO-incompatible transfusion, plasmapheresis may be the most efficacious way to remove excessive free Hb from the circulation before it damages the glomeruli permanently. However, because this procedure

takes some time to arrange, massive doses of loop diuretics in combination with large volumes of clear fluids are often used to dilute the toxic effects of Hb. In the cold type of AIHA, it is customary to increase the temperature in the patient's room and use warm blankets to prevent hemolysis when the blood circulates to fingers, toes and nose. Steroids are mainly used in warm type AIHA but the effect can be slow. Monoclonal antibodies against the antibody-producing B lymphocytes are used increasingly (268). A relatively new tool used in PNH treatment is a monoclonal antibody against one of the final components in the complement cascade (176) and it has been suggested that this approach could be applicable in other hemolytic situations as well.

B. Preeclampsia

PE has been named the disease of theories (241) and was described as early as 3000 years ago by the ancient Egyptians (277). PE is still one of the dominating obstetric complications that cause perinatal and maternal morbidity and mortality. The worldwide prevalence ranges 3 to 7% of pregnancies.

1. Pathophysiology.

There is a documented risk of developing PE when there is a maternal and paternal family history of PE (163). Discordance between monozygotic twins indicates that hereditary factors that follow classical Mendelian hereditary traits do not exist (293,297). Black women are known to have a higher risk of developing PE than Caucasians (116). Several genes are likely to be involved in development of PE (42,119,241,284). Mainly primigravidae are affected indicating that the maternal immune system may play a role. It was recently shown that early and late onset PE, as well as PE with intra uterine growth restriction (IUGR) all have different inflammatory plasma protein profiles (75). Furthermore, systemic diseases such as diabetes,

obesitas, essential hypertension, and renal diseases are all predisposing factors for PE. These data illustrate the complex, multi-factorial nature of PE.

Removal of the placenta is crucial for the resolution of the symptoms (241), a fact that has led to the theory of a placenta-derived factor as a culprit. Although the etiology and pathophysiology are still not known, PE is suggested to evolve in two stages (Fig. 13)(243). Stage one occurs during formation of the placenta, and a defective and shallow invasion of the placental cells, trophoblasts, into the muscle layers of the spiral arteries, has been shown (41). Stage two, the clinical manifestations, i.e. hypertension and proteinuria, appear from 20 weeks of gestation onwards. Early onset PE (< week 34) is often more severe than late onset PE (> week 34). PE is defined as blood pressure of > 140/90 mmHg and proteinuria of > 0.03 g/l or a rise of diastolic blood pressure > 20 mmHg during pregnancy.

As the disease progresses, angiospasmus in the brain and brain oedema may cause severe epileptic seizures – eclampsia (170). The renal disturbances seen in PE lead to reduced glomerular filtration rate and proteinurea, as discussed above in relation to the hemolytic anemias. Glomerular endotheliosis is characteristic for PE, in fact it is the most common glomerular disease in the world (278). General endothelial dysfunction and diminished ability to vasodilatation are vascular hallmarks of PE (240). Severe PE also involves pathological activation of the coagulation and fibrinolytic systems that further compromise the blood circulation in general and the utero-placental blood flow specifically (41,223). The reduced utero-placental blood flow results in IUGR, seen in one out of every four PE cases. The inadequate blood flow also gives rise to a reduced O₂ delivery, which results in an ischemia/reperfusion-like situation and oxidative stress. A growing body of evidence suggests

that this oxidative stress further aggravates vascular function in the placenta (116,126,242,270).

Decreased placental perfusion, in combination with oxidative stress, causes general endothelial cell damage. An inflammation of the endothelial cells in the vascular system, in kidneys in particular, is a typical histological finding in PE (71,106,241,245,280).

Vasoconstriction and elevated resistance to blood flow follows as a consequence. The increased vascular resistance can be evaluated clinically by using pulsed Doppler techniques on the uterine arteries as a high pulsatility index and/or occurrence of early diastolic notch in the maximum velocity waveform. Patients with persistent bilateral notching in the uterine arteries have a significantly elevated risk of developing PE and IUGR (22,159).

A severe form of PE, the HELLP syndrome, is characterized by low platelets, elevated liver enzymes and hemolysis with high levels of free maternal HbA. Patients with HELLP often present with acute manifestations, possibly due to release of large amounts of free HbA.

Oxidative stress is considered a key factor of the HELLP syndrome, and significant increased levels of antioxidants in placentas have been shown (100).

2. Hemoglobin in PE placenta

Recently, gene and protein profiling studies have revealed a new potential etiology for PE (Fig. 13). The first stage, caused by a defect formation of the placenta, gives rise to uneven blood perfusion, oxidative stress, production and accumulation of free HbF (56). The mechanisms of the generation of free, extracellular HbF in placenta blood vessels are not known but it can be speculated that hypoxic conditions induce an up-regulation of the HbF-genes, and that a microenvironment characterized by insufficient blood-flow and oxidative

stress promote hemolysis (218). Interestingly, it was shown that PE is associated with decreased glucose-6-phosphate dehydrogenase activity and impaired redox regulation in erythrocytes and fetal endothelial cells (5), which may contribute to increased hemolysis in PE. Free HbF further aggravates the oxidative stress which damages the blood-placenta barrier causing a leakage between the fetal and maternal circulations (189). Recent results show that the free HbF leaks over to the maternal blood stream from the damaged placenta as early as the first trimester (16). Free HbF may also contribute to the general systemic endothelial damage and inflammation that is characteristic of stage two. In addition, maternal plasma levels of HbA are elevated in women with PE at term of the pregnancy as compared to control, and both HbF and HbA are correlated with disease parameters (218). Possible mechanistic explanations of the role of HbF in the early PE-development are, as described above, that the free HbF may generate ROS and free heme-groups, which damage the placenta and its barrier function. This may cause a leakage of fetal and placental factors, including HbF itself, from the fetus to the mother, thus linking Stage 1 and Stage 2 of the disease (Fig. 13). In fact, free heme, bilirubin, and biliverdin have been identified among 14 metabolites in a metabolomic signature of PE using first trimester plasma (142).

The hypothesis that Hb may be an important pathological factor in PE by causing placenta damage is further supported by a recent report that *ex vivo* perfusion of placentas with Hb in the fetal circulation led to a significant increase of the perfusion pressure, feto-maternal leakage of free Hb, morphological damage, and up-regulation of genes related to immune response, apoptosis, and oxidative stress (57,190).

Likewise, the elevated concentrations of free maternal plasma HbF and HbA may have a central role in causing the clinical symptoms of the fully developed PE by inducing 1)

oxidative stress, 2) NO-scavenging and vasoconstriction, 3) further hemolysis. Indeed, in term pregnancies, elevated biomarkers of oxidative stress and up-regulation of the heme-scavenger and antioxidant A1M were found (218), supporting an etiological role of free Hb in PE. Free HbF and A1M were also shown to be useful as prognostic biomarkers for PE as early as in the first trimester, several weeks before clinical manifestations were present (16).

3. Treatment of PE

Only symptomatic treatment of PE is available and includes treatment of the high blood-pressure with anti-hypertensives and prevention of seizures with magnesium sulfate-infusion (79,264). Several studies have focused on preventing PE by using antioxidative treatment with vitamin C, E, melatonin and even dark chocolate, however these treatments have not been able to prevent development of PE (244). Thus, delivery of the fetus, by labour induction or Caesarian section, is the only cure today.

C. Intraventricular hemorrhage in the preterm infant

Intraventricular hemorrhage (IVH) is a well characterized hemorrhagic lesion of the brain mainly occurring in premature infants (Fig. 14). The hemorrhage typically initiates in the germinal matrix, a structure located at the head of the caudate nucleus and underneath the ventricular ependyma, which is rich in vasculature and consists of glial and neuronal precursor cells. The hemorrhage can either remain stationary within the germinal matrix or rupture through the ependymal wall resulting in a collection of blood within the cerebral ventricles. Rupture of the germinal matrix vasculature leading to IVH occurs almost without exception during the first 72 hours after birth and is easily detected by bed-side ultrasound. Severe cerebral IVH, with more than 50 % of the ventricular area filled by blood, occurs in about 15 to 20% of very preterm infants (310,321,322) and the incidence of IVH has not

changed over the last decade (319). Fig 13 B-E illustrates severe IVH in a preterm infant as determined by cranial ultrasound.

Children with severe IVH have a severe short- and long-term prognosis. The mortality in the neonatal period has been reported to be 20-50% and of the surviving infants over 50% develop post-hemorrhagic ventricular dilatation (PHVD) leading to hydrocephalus and life-long requirement of a ventriculo-peritoneal shunt. At 2 years of age, 40-80% of children show signs of severe neurological impairment, mainly cerebral palsy and mental retardation (3,43,97,98,198).

1. Mechanisms of white matter brain damage in the immature brain

The mechanisms leading to injury of the periventricular white matter following IVH are incompletely understood. The pathology in white matter following IVH is characterised by oligodendroglial cell death and immune cell activation, *i.e.* astrogliosis and microglial activation. The principal cellular target within the white matter of the preterm infant is the developing oligodendrocyte. These early lineages of oligodendrocytes, the pre-oligodendrocytes, possess an extremely heightened vulnerability to insults associated with increased production of ROS. This is at least partly dependent on a maturational delay in the production of endogenous antioxidants such as SOD, Gpx and catalase (143). Experimental studies have to date primarily focused on hypoxia/ischemia and inflammation as prime upstream initiators of ROS production. Microglial activation has been shown to play a key role and activation of microglial toll-like receptor (TLR)-2 and TLR-4 by both pathogen-related and endogenous molecules results in generation of ROS and cytokines causing either death or maturational arrest of the pre-oligodendrocytes. The resulting hypo-myelination of cortico-spinal tracts forms the basis for cerebral palsy detectable during early childhood, most

frequently observed as spastic diplegia with a pre-dominant impairment of lower extremity motor function.

2. Free hemoglobin and brain damage

The following will discuss a plausible and potentially treatable mechanism whereby IVH causes white matter brain damage (Fig. 14A). Following rupture of germinal matrix vasculature and of the ventricular ependyma, extravasated blood is mixed with intraventricular cerebrospinal fluid (CSF). Ensuing hemolysis of erythrocytes causes release of free Hb. In adults with subarachnoidal hemorrhage (SAH), Hb reaches its peak by day 2 following hemorrhage. Hb, primarily present as oxyHb, will with time be auto-oxidized to metHb. Subsequently metHb may be decompose into heme and globin. The rates of hemolysis and resulting levels of free oxyHb and metHb in intraventricular CSF have yet to be characterised in preterm infants following IVH. Free Hb is known to be deleterious to the central nervous system and is cytotoxic in neuronal cultures (152,238). One obvious mechanism is through ROS formation related to free heme release from Hb break-down and subsequently of free iron from the catabolism of heme by HO-1 or the constitutive HO-2. Clinical and experimental studies of adult SAH show convincing signs of immune activation with intracerebral glial activation, macrophage activation, neutrophil recruitment and increased production of pro-inflammatory cytokines and adhesion molecules. A recent study showed that induced SAH *in vivo* was associated with up-regulation of endothelial TLR-4 and a subsequent *in vitro* study revealed that exposure to oxyHb activated TLR-4 in vascular smooth muscle cells (323,331). In summary, available data show that released free Hb and its downstream metabolites exert neurotoxic and pro-inflammatory effects highly relevant for the previously described pathways of white matter brain damage in the preterm infant.

3. Treatment of IVH

There is currently no established therapy to prevent infants from developing neurological disability following detection of severe IVH. Infants who develop post-hemorrhagic hydrocephalus receive a life-long ventriculo-peritoneal shunt which is an efficient means of preventing progressive ventricular distension but does not reduce neurological impairment. Recent data show that an intervention aimed at removing blood clots from the intraventricular space, *i.e.* perfusing the intraventricular compartment via inserted catheters with artificial CSF, can reduce the requirement of life-long ventriculoperitoneal shunt and improve neuro-developmental outcome at early childhood (309). These data support the concept that blood components within the intraventricular space constitute risk factors for development of hydrocephalus and periventricular tissue damage.

D. Chronic leg ulcers.

Chronic venous disease and chronic leg ulcers are common in Europe and the USA, and the point prevalence for non-healing venous ulcers has been estimated to about 0.3 % (204). The social and economic impact of these non-healing wounds constitute a great burden on both patients and the health care system (203,204). Although much evidence for a dysregulated inflammatory process in non-healing venous ulcers has been presented, the exact mechanisms underlying the disease, as well as the relative importance of the so far identified pathogenetic steps, are still not clarified (6). Current evidence however, indicate that the ulcers are characterized by bacterial colonization, ongoing and unregulated inflammation and excessive proteolysis, chemotactic activity and endothelial leakage (6).

1. Inflammation

Based on both *in vitro* experiments and clinical studies, several causative mechanisms

underlying the chronic inflammation have been proposed (Fig. 15). The sustained venous pressure of these patients leads to a reduced pressure differential between the arterial and venous systems, and this has been proposed to underlie the fact that these ulcers contain significant amounts of extravasated or “trapped” leukocytes in the connective tissue. Extravasation of leukocytes also occurs in patients with venous disease only, and this in turn, may lead to ulcer formation (315). Extravasation of fibrin leads to the formation of fibrin depositions, possibly trapping and also inactivating growth factors involved in wound healing in the perivascular areas (87). The high numbers of activated leukocytes lead to release of matrix metalloproteinases (MMPs), which further degrade extracellular matrix components as well as growth factors (6). Furthermore, all chronic ulcers are constantly colonized or infected by various bacteria such as *Pseudomonas aeruginosa*, *Enterococcus faecalis*, *Staphylococcus aureus*, and *Proteus mirabilis* and evidence supports the view that high bacterial loads contribute to non-healing (68). These findings, together with the observation that keratinocytes display a non-migratory phenotype in non-healing venous ulcers suggest a failure of re-epithelialization and thus poor healing.

2. Heme, iron and oxidative stress

Recent studies demonstrate a pivotal role for accumulation of iron in the tissue surrounding non-venous ulcers, leading to induction of the high inflammation noted in this disease. Iron deposits (Fig. 16) in non-healing ulcers cause the readily visible brown-colored areas which often precede, and typically are found around non healing, venous ulcers. The origin of these increased iron stores is extravasation of erythrocytes due to the venous stasis combined with excessive capillary leakage. Iron, in free form or complex-bound to protoporphyrin IX in heme, has been implicated as an important pathogenic factor in the initiation of the high inflammation in leg ulcers (2,10,300). As described above, the oxidized ferric forms of heme

and free iron are strong oxidative agents and the ferrous iron can generate O₂-derived free radicals by the Fenton reaction (172,282). These compounds present considerable oxidative stress that leads to tissue damage and cell destruction, and have been proposed to participate in the pathogenesis of chronic ulcers by initiating local lipid peroxidation of polyunsaturated fatty acids in cell membranes as well as enhancing the synthesis and activation of MMPs (10). Interestingly, heme was identified in significant amounts in fluids from non-healing ulcers (11). Heme was also found particularly in tissue sections from chronic ulcers as opposed to acute wound sections or normal skin sections [(11); Fig. 16]. It thus seems likely that, dependent on venous hypertension in chronic ulcers, heme diffuses from plasma to the extravascular compartments across the endothelium and basement membranes in non-healing ulcers. Heme may therefore be trapped and accumulated in pericapillary fibrin cuffs of venous ulcers. Thus, excess accumulation of heme causes an increased vasopermeability, increased expression of endothelial adhesion molecules and infiltration of leukocytes, changes typically characterizing venous ulcers (10). Interestingly, high amounts of A1M were also found in these locals, supporting the view of A1M as a heme scavenger in non-healing ulcers [(11); Fig. 16]. It may be speculated that A1M has a similar effect as HO, promoting iron storage; however, additional studies are required in order to elucidate a role of A1M in iron metabolism in non-healing ulcers. Unlike HO, A1M functions extracellularly, possibly providing the main source of protection against unsequestered heme in the extracellular space. Interestingly, and supporting a role for heme and iron in chronic ulcer pathogenesis, mutations in the human hemochromatosis protein (*HFE*)-gene seem to be linked to a higher risk of developing ulcers (326-328).

The causes of non-healing ulcers are summarized in Fig. 15. Taken together, the data presented so far support the concept of heme and iron as important mediators of the high inflammation in non-healing ulcers, with A1M as part of a previously undisclosed

endogenous defense mechanism against heme-induced oxidative stress.

E. Blood substitutes and oxygen therapeutics

The attempts to develop “artificial blood” or blood substitutes have been ongoing with various intensities for more than 70 years. These research efforts have mostly been focused on the ability of Hbs to carry and deliver O₂. Most of the products that have been passed into advanced-phase clinical trials are also derivatives of Hb. The phrase ‘hemoglobin-based oxygen carrier’ (with the acronym ‘HBOC’) has been applied to these materials, as blood contains many molecules and proteins that provide other functions than O₂ transport (49,60,62,325).

There are several important driving forces behind the development of RBC substitutes. For instance, effective trauma care, an increased number of elective surgeries and the prevalent but small risk of transmission of blood-borne pathogens stimulate this development. In addition, these products can function as organ preservatives to prevent or decrease reperfusion injury to donor organs. Various religious and ethnic groups with concerns regarding the use of human blood products may accept these substances.

The biologic properties of an optimal blood substitute should include a reasonable level of O₂ delivery when compared to normal human RBCs. An ideal blood substitute should fulfill several additional requirements, including lack of any antigenicity, readily availability, a long half-life, stability allowing storage at ambient temperature and elimination of risks to transmit infections. However, to date, no oxygen-carrying blood substitutes have been approved for use by the US Food and Drug Administration. This fact clearly demonstrates the difficulties that exist in formulating an effective blood substitute. Therefore, most current efforts with

these agents do not primarily aim at substituting blood but rather focus on mediating O₂ therapeutics in severe clinical settings. HBOCs can thus be viewed as an attractive alternative O₂ delivery system which can partly augment whole blood during a severe blood loss situation or, alternatively, for use when whole blood simply is not available for treatment.

One of the major obstacles in the clinical development of HBOCs has been the uncontrollable and sudden elevation in systematic and pulmonary blood pressure in trauma victims or patients undergoing elective surgery when infused with these products. Furthermore, native free Hb, when infused rapidly, readily splits into dimers followed by its clearance by glomerular filtration. The initial attempts of transfusing Hb-based products thus resulted in renal dysfunction and hypertension. The instability and reactivity of the Hb tetramers have been overcome by altering human (or bovine) Hb by three general approaches: the use of chemical cross-linking reagents that stabilize proteins along with modifications to adjust O₂ affinity; exploring reagents specifically designed to increase the size of the Hb product and modifying its reduction to notably NO; protein engineering and expression of modified versions of Hb with desirable properties.

Over the years several different Hb based products have been produced. For instance, a so called first generation product is Hemopure, which is a polymerized form of bovine Hb with an O₂ affinity that is close to human Hb. It has an intravascular half-life of 8-23 hours and a shelf life of 36 months at room temperature. Hemopure is approved in South Africa for the treatment of adult surgical patients who are acutely anemic with the intention of eliminating or reducing the need for RBC transfusions (121). Hb binds NO readily at the same site at which it binds O₂ and at the thiol of cysteine (see also above, section II.B). It was apparent at an early phase that the elevating effects on blood pressure was most likely related to strong

binding of NO to HBOCs. This inherent property of free Hb would be one of the major causes of vasoconstriction, leading to hypertension. In 2004, the first modified Hb that did not elevate blood pressure was reported. In 2004, the first modified Hb product that did not elevate blood pressure was reported. The Hb molecule was conjugated to polyethylene glycol (PEG) and a safe product could be considered to be a real possibility. PEG-modified Hb, such as MP4, is based on human Hb produced from out-dated blood. In animal models MP4 has been shown to be effective in cases of hemorrhagic shock. Adverse effects associated with the vasoactive properties of first-generation blood substitutes are not seen with MP4. At relatively low concentrations, MP4 is capable of transporting large amounts of O₂ (65).

Besides PEGylation, several lines of developments have been described. For instance, HBOCs have been cross-linked with enzymes, in efforts to synthesize compounds that not only perform the function of carrying O₂, but also harbor some of the enzyme activity that normal RBCs possess. Polymerized Hb has thus been chemically cross-linked with catalase, SOD and carbonic anhydrase to form a compound that, in animal models can not only carry O₂ but also remove O₂ radicals that are responsible for ischemia reperfusion injuries (32). Such conjugates can be prepared by chemical cross-linking or by gene fusion of the corresponding structural genes in order to generate more homogenous preparations (108). Attempts have also been made to encapsulate Hb within a lipid-membrane to create a compound capable of carrying O₂ while not being associated with significant vasoconstriction. These liposomes appear to be retained in plasma for a significant period, but they are difficult to produce and can activate the reticuloendothelial system, the complement pathway, and platelets. The ultimate RBC substitute would need to contain an artificial membrane mimicking the native counterpart (252). However, production of such a product would be extremely challenging. Efforts have also been made to use the biopolymer

polylactide, which is biodegradable and converted to lactic acid in the body. These artificial RBCs should contain Hb along with SOD, metHb reductase, and catalase (59).

Adverse effects associated with HBOCs include beside hypertension, abdominal pain, skin rash, diarrhea, jaundice, hemoglobinuria, oliguria, fever, stroke, and laboratory anomalies such as an elevation in lipase levels. Although most of these side effects are transient and clinically asymptomatic, many clinical trials involving these agents have been discontinued or withheld due to the associated adverse effects. In 2008 a meta-analysis of the results of clinical trials of blood substitutes led to the striking conclusions that none are effective and some are not safe. After that report appeared, the US Food and Drug Administration announced that trials of products under consideration would not proceed further (84,201). With existing documentation of clinical problems, it is clear that new designs and application areas for HBOCs are needed. Ongoing research has improved our understanding of the RBC and its interactions with the surrounding cellular environment. This has facilitated the development of newer products that do not have significant vasoactive properties, as did the first-generation compounds. Several of these recent approaches are promising and involve linking Hb with itself or nonreactive species in order to increase size, emulating the size-expanding effects of PEG (53,124,125,175). For instance, Bucci et al. produce larger materials by combining Hb tetramers under dehydrating conditions to give a 'zero-link' product (45), while Zal's group has used protein engineering to express a multi-Hb ensemble that occurs in marine organisms (118). Marden has developed linked Hb tetramers, called octamers, and has also attached Hb to nanoparticles as a means to increase size (28). Many of the current polymerized preparations still have a small fraction of free tetrameric Hb in solution. This fact has often been proposed to be a major reason for the observed side effects associated with transfusion. Therefore, novel chemical strategies have been developed to

increase the degree of polymerization and minimize free Hb in solution (329). In addition, auto-oxidation and shelf-life are two critical parameters in the design of HBOCs. By storing the product in the the carboxy state (COHb) potentially valuable improvements have been observed (302). The use of COHb to release a limited amount of CO may also have potential benefits in ischemic tissues. Small molecules that release CO have been shown to exert vasodilatory and anti-inflammatory properties in a variety of models in the peripheral circulation (197). Transfusion of RBCs containing high COHb was thus shown to be beneficial in limiting microvascular markers of apoptosis after hemorrhagic shock (52).

V. α_1 -microglobulin – a novel therapeutic concept

The biochemical and physiological properties of A1M are suggestive of an *in vivo* protective role of the protein against radicals and oxidative stress. The findings that A1M can bind heme and that Hb induces a proteolytic activation of heme-degradation activities of A1M, suggest that it may have a specific role during hemolytic conditions in protecting tissues exposed to Hb- and heme-induced oxidative attack. Does A1M have properties that allow, and even favour, therapeutic use in clinical situations with a high load of free, extracellular Hb? Below, we speculate on a possible extravascular role of A1M to complement an intravascular role of Hp and Hpx. We will also propose tentative A1M-based therapies in the diseases discussed in this article.

A. Extravascular Hb-protective role of A1M

In a therapeutic situation, what are the possible advantages of A1M compared to the other Hb-protective systems? Available data today points to a specific protective role of Hp and Hpx intravascularly, whereas A1M seems to operate mostly extravascularly. The two former

proteins are found in high concentrations in blood and promote clearance of Hb and heme, respectively, from the circulation. However, it should be remembered that the Hp levels fall towards zero very rapidly following the release of free Hb into the plasma, due to scavenger receptors for the Hp-Hb complex and it is possible that other protective mechanisms are desirable and needed in this situation. The literature suggests that A1M has a role as an antioxidant and heme-scavenger mainly in extravascular fluids and cytosol rather than in blood. It is secreted to blood from the liver but its concentration in blood is comparatively low, 1-2 μM , due to a very rapid equilibrium ($T_{1/2}=2-3$ min) between intra- and extravascular compartments (see section III.F.4). Likewise, the heme-degrading HO-proteins mainly have a cytosolic location whereas A1M is found both in cytosolic and extracellular locations. Finally, in relation to the vast network of more or less monospecific antioxidants that protect tissues from Hb- and heme-induced oxidative damage/stress, A1M seems to be multispecific, binding and clearing cytosols and extravascular fluids from ROS as well as small organic radicals.

B. Studies in diseases

1. Chronic leg ulcers

The possible use of A1M in treatment of chronic leg ulcers was suggested by Goldsmith (102). The localization and heme-binding activities of the protein in inflamed tissue and fluid from wounds (11), the heme-binding activities (11), and the *in vitro* experiments with keratinocytes, skin and collagen fibrils ((216)see also section III.F.8) support this notion. The *AMBIP* gene was also up-regulated in keratinocytes and skin by heme- and ROS-exposure suggesting physiological advantages from increased concentrations of the protein in skin tissue during oxidative stress (Olsson et al., PLoS One 2011). Furthermore, as described above (section III.F.8.d) several lines of evidence suggest that A1M may have repair-

/rebuilding effects on extracellular matrix, promoting increased fibrillation of collagen I and an up-regulation of elastin- and collagen genes. However, to prove a clinically beneficial effect of A1M in chronic leg wounds or in other oxidative stress-related skin disorders such as during UV-irradiation, atopic dermatitis and psoriasis, *in vivo* experiments in humans or animals have yet to be performed.

2. Preeclampsia

Another target of A1M-based therapy may be PE, which, as described above, is associated with higher-than normal plasma and placenta concentrations of HbA and HbF as well as markers of oxidative stress. It was hypothesized that PE manifests as a result of an Hb-dependent oxidative insult and that A1M is involved in the (futile) defense (56,218). Indeed, elevated serum concentrations of A1M were found in the first trimester of women who later develop PE (16) and in the placenta, urine and blood from women with diagnosed PE at term of the pregnancy, and evidence of t-A1M formation in placenta (148,218,225). Furthermore, as described above, *ex vivo* perfusion of placenta showed that the PE-like damage induced by perfusion of Hb was effectively inhibited by A1M. Antioxidative treatment of PE has been suggested and tried before. Several studies have evaluated the use of vitamin C and E in high-risk pregnancies in order to prevent the PE-associated oxidative stress. These studies failed to show a reduction in the rate of adverse maternal or perinatal outcomes related to pregnancy-associated hypertension (231,244). This does not disprove the oxidative nature of the disease, however, since the scavenging capacity of vitamin C is limited and the oxidized form, dihydroascorbate, which is formed by its oxidation, may also present an oxidative challenge in the tissues during the disease.

3. Intraventricular hemorrhage in preterm infants

Treatment of IVH and/or SAH with A1M may be a possible option if Hb- and heme-induced toxicity and oxidative stress actually are important pathological factors, as described above.

The heme- and radical-binding of A1M, the rapid turnover rate, and the fact that it is an endogeneous, non-immunogenic molecule makes this protein an attractive therapeutic option for these diseases.

4. Hemolytic anemias and transfusion reactions

Hemolytic anemias and adverse reactions following blood-transfusion may also be future targets of A1M-treatment, but both scenarios present a much greater challenge than for instance chronic leg ulcers or IVH since the whole body is affected. Thus, systemic, rather than local administration has to be considered. Intravascular hemolysis, such as that occurring in autoimmune hemolytic anemia of the cold (IgM-mediated) type, paroxysmal nocturnal hemoglobinuria, and paroxysmal cold hemoglobinuria may be diagnoses in which a benefit from systemic A1M administration could be possible since these states are often associated with depletion of Hp as discussed above. Similarly, we know that patients with extravascular hemolysis, *e.g.* diagnoses including autoimmune hemolytic anemia of the warm (IgG-mediated) type and drug-induced hemolytic anemia, acquire impairment of their kidney functions due to detrimental effects of Hb. It is likely that they also suffer from increased oxidative stress based on the large amounts of extracellular Hb and heme they have to process. It is thus probable that also these patients could benefit from antioxidative/radical scavenging therapy by A1M. Today's storage solutions for RBC concentrates for transfusion do not take into account the problem with oxidative damage during storage. To increase the quality of RBCs for transfusion one may therefore propose to add A1M before and/or after storage. Thus, A1M could be added separately 1) before the storage period is started

(proactive); 2) as a rejuvenation agent after a certain time period had passed (interactive); or 3) just prior to issuing the blood unit to the patient/ward (cleaning/therapeutic approach). In analogy with data from other systems, A1M would have the ability to prevent oxidation and, in the latter two cases, also repair oxidative modifications of the blood components. This approach may be especially appealing in the situation of gamma irradiation of blood to prevent graft-versus-host disease, since those units tend to be more damaged with membrane disturbances causing potassium leakage, compared to untreated blood for transfusion.

5. Blood substitutes

As discussed above, many of adverse side-effects of treatment with blood substitution products are related to generation of ROS, free radicals, and free heme, and may therefore also be future targets of A1M-intervention. For example, A1M may be administered as an addition to the HBOC itself, given to patients as a means of coping with the negative consequences and oxidative stress induced by free Hb and heme. An interesting alternative could perhaps be to cross-link A1M with Hb or to create a genetically modified fusion protein between Hb and A1M. Another approach would be to encapsulate Hb and A1M together in liposomes. Both ways, A1M would do the job of keeping the Hb-molecule in a reduced state and eliminate Hb-generated ROS as soon as they are produced. However, no investigations on *in vivo* trials with HBOC in combination with A1M have been reported yet.

VI. Conclusion

Toxic effects of extracellular Hb are major components in the pathogenesis of many diseases and iatrogenic situations, including hemolytic anemias and transfusion-induced intravascular hemolysis, preeclampsia, intraventricular hemorrhage, chronic inflammatory leg ulcers, and

infusion of recombinant hemoglobin. The pathogenesis involves one-electron reactions between oxy-Hb, its downstream metabolites heme, iron, ROS and free radicals and exposed tissue components. Many mechanisms have evolved to protect organisms against damage from extracellular Hb. One of these, the ubiquitous protein A1M, is found in all vertebrates including man and operates by rapidly clearing cytosols and extravascular fluids from heme-groups and free radicals released from the oxy-Hb molecule. A1M has been shown to protect cells and matrix from Hb-, heme- and ROS-induced damage and is a promising therapeutic alternative in diseases with Hb-induced pathology.

Acknowledgements

Part of the work shown in this paper was supported by the Swedish Research Council, governmental ALF research grants to Lund University and Lund University Hospital, the Royal Physiographic Society in Lund, the Foundations of Greta and Johan Kock and Alfred Österlund, and the Blood and Defence Network, Lund University. The authors also wish to acknowledge Mattias Mörgelin and Maria Baumgarten for outstanding electron microscopy work.

Abbreviation

α_1 -microglobulin	- A1M
Red blood cell	- RBC
Hemoglobin	- Hb
Oxygen	- O ₂
Reactive oxygen species	- ROS
Fetal hemoglobin	- HbF
Carbon monoxide	- CO
Nitric oxide	- NO
Adult hemoglobin	- HbA
Ferrous iron	- Fe ²⁺
Ferric iron	- Fe ³⁺
Hydrogen peroxide	- H ₂ O ₂
Ferryl iron	- Fe ⁴⁺
Haptoglobin	- Hp
Superoxide dismutase	- SOD
Glutathione peroxidase	- Gpx
Hemopexin	- Hpx
Heme oxygenase	- HO
Peroxiredoxin I	- Prx or HBP23
Superoxide radical	- O ₂ ⁻
Cytoplasmic SOD	- Cu/Zn SOD (=SOD1)
Mitochondrial SOD	- MuSOD (=SOD2)
Extracellular SOD	- Cu/ZnSOD (SOD3)
Glutathione	- GSH
Oxidized glutathione	- GSSG
Thioredoxin	- Trx
Alpha-1-Microglobulin Bikunin Precursor gene	- AMBP gene
Proprotein convertase	- PC
Hepatocyte-enriched nuclear factor	- HNF
Nitroblue tetrazolium	- NBT
2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid)	- ABTS
Truncated A1M (lacking C terminal tetrapeptide LIPR)	- t-A1M
Preeclampsia	- PE
Autoimmune hemolytic anemias	- AIHA
Paroxysmal nocturnal hemoglobinuria	- PNH
Hemolytic disease of the fetus and newborn	- HFDN
Intra uterine growth restriction	- IUGR
Intraventricular hemorrhage	- IVH
Post hemorrhagic ventricular dilatation	- PHVD
Toll-like receptor	- TLR
Cerebrospinal fluid	- CSF
Subarachnoidal hemorrhage	- SAH
Matrix metalloproteinases	- MMP
Polyethylene glycol	- PEG
Carboxy hemoglobin	- COHb
Hemoglobin-based oxygen carrier	- HBOC

Figure legends

Figure 1. Hb-structures, auto-oxidation products and toxicity. When the RBCs rupture (hemolysis) Hb is released. Adult Hb, HbA, consists of 2 α -chains and 2 β -chains, and fetal Hb, HbF consists of 2 α -chains and 2 γ -chains. All chains contain an Fe-containing heme-group which binds molecular oxygen, O₂. OxyHb spontaneously decomposes via rearrangement of electrons in a process called auto-oxidation, leading to formation of ROS and degradation products like heme and metHb. All of these molecules are highly reactive and have the ability to damage lipids, protein and DNA and are therefore pro-inflammatory, pro-oxidant and toxic (oxidative stress). Furthermore, Hb also binds and reacts with NO, thereby affecting the vasomotor tone. Thus hemolysis and elevated extracellular Hb concentrations is associated with hypertension. Finally, the metabolites formed by the Hb-degradation are hemolytic and thus, oxidative stress accelerates the hemolysis, in a positive feed-back loop.

Figure 2. Redox-balance in health and disease. Human health is dependent upon the balance between oxidants and antioxidants. The human body is exposed to oxidants constantly from exogenous sources like air, food, radiation exposure etc. and from endogenous metabolism. The production of oxidants, under normal healthy conditions, is counteracted by production of antioxidants from the body and uptake from food. However, a disturbance in the balance between oxidants and antioxidants, caused by for instance increased uptake of oxidants, increased production of oxidants from metabolism or decreased production of endogenous antioxidants, will result in damage to cells, membranes, DNA and proteins.

Figure 3. Overview of protection mechanisms against extracellular Hb-toxicity. Cell-free Hb is released under pathological conditions involving hemolysis or infused during blood

substitution treatment. The Hb-molecule is bound by haptoglobin and cleared by receptor-mediated uptake in monocytes and macrophage-cells. Excess oxy-Hb undergoes auto-oxidation reactions resulting in reactive oxidative compounds, represented by reactive oxygen species (ROS) and free heme-groups. A complex network of antioxidation mechanisms inhibit and eliminate the ROS-species, and repair the oxidative damage caused by them. Free heme-groups are bound by hemopexin and albumin in plasma, metabolized by heme oxygenases (HO) in cells, and bound/degraded by α_1 -microglobulin (A1M) in plasma, cells, and extracellular fluids. A1M also participate in non-heme targeted antioxidation mechanisms.

Figure 4. Schematic representation of Hp-Hb complex formation, endocytosis mediated by CD163 receptor and degradation of heme in the lysosome.

Figure 5. Heme degradation by heme oxygenase-1. Cytosolic HO-1 catalyzes the degradation of heme-groups to biliverdin, carbon monoxide (CO), and iron atoms ($\text{Fe}^{2+/3+}$). The reaction takes place on the cytosolic face of the ER-membrane and involves oxidative breakage of the protoporphyrin IX-ring in concert with microsomal NADPH-cytochrome P450. Reduction of biliverdin to bilirubin is catalyzed by the enzyme biliverdin reductase. Liberated $\text{Fe}^{2+/3+}$ atoms are stored as Fe^{2+} by cytosolic ferritin H. CO diffuses freely and participates in several physiological reactions, mostly by binding to heme-proteins.

Figure 6. Structural model of α_1 -microglobulin (A1M). The model was generated using Swiss-Pdb Viewer and Swiss Model by alignment of the human A1M amino acid sequence with rat epididymal retinoic acid-binding protein (E-RABP) (281). β -strands are shown in

yellow, α -helix in orange, Cys-34 in red and Lys-92, Lys-118 and Lys-130 in blue. Viewed from the side of the β -barrel (left) and looking down into the pocket (right).

Figure 7. *AMB*P-gene and synthesis of A1M and bikunin. The α_1 -microglobulin/bikunin precursor gene (*AMB*P), located on chromosome 9q34 (77,255), codes for the precursor-protein A1M-bikunin (*AMB*P). The signal peptide and the A1M-part are encoded by exons 1-6 and a linker peptide and bikunin-part by exons 7-10. The A1M- and bikunin-parts are separated in the Golgi apparatus via proteolytic cleavage and then secreted separately (38). Post-translational modification of the precursor includes glycosylation of both the A1M and bikunin parts (blue). Bikunin is also cross-linked to various heavy chains via a glykosaminoglycan chain, forming the inter- α -inhibitor family (254).

Figure 8. Metabolism and distribution of A1M. 1) The main location of A1M-synthesis is the liver. After secretion to the blood as described in Fig. 7, the protein is found in plasma in free form and bound to IgA, albumin and prothrombin at a total concentration of approximately 2 μ M. 2) Free and complexed A1M are rapidly transported across the capillary walls to the extravascular compartment. 3) A1M is also synthesized and secreted from epithelial cells in all examined organs and tissues. 4) A1M is found in interstitial fluids, in association with collagen fibers and other extracellular matrix fibers, and in association with perivascular basement membranes. 5) A1M is cleared from the blood-circulation by filtration in the renal glomeruli. Most of the filtrated A1M is degraded in the proximal tubular cells but a small part is excreted in urine (appr 5 mg/day). Thus, the major metabolic pathway of A1M is Liver→Blood→Interstitial fluid→Blood→Kidneys, minor pathways are Epithelial cells→Blood→Kidneys, and the turnover rate in blood is $T_{1/2}$ =2-3 min. A1M is depicted as small yellow Lipocalin barrels.

Figure 9. Protective properties of A1M. Oxidative damage is induced by hemolysis and subsequent release of free hemoglobin, followed by generation of free heme-groups, iron, and ROS, or by non-hemolytic events such as infection, inflammation, trauma, and ischemia/reperfusion. Four reported biochemical and physiological properties of A1M contributes to protection against the oxidative damage. 1) Reductase and dehydrogenase activity, 2) heme-binding by A1M and heme-degradation by a proteolytically activated form (t-A1M), 3) radical-scavenging, and 4) up-regulation of the AMBP-gene in hepatic and non-hepatic cells by elevated concentrations of hemoglobin, heme and ROS.

Figure 10. Tentative radical trapping mechanism of A1M. The figure is modified from (341) and based on the reactions with the synthetic radical 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid) (ABTS). A1M is represented by a Lipocalin barrel with Y132 and the thiolate group of C34 high-lighted. The latter is located on a large, flexible (Ω) loop. The pK_a of the C34 thiolate is lowered by the proximity of the three positively charged side-chains of K92, K118 and K130, located near the Ω -loop. Reaction (1): The C34 thiolate group reacts with a radical (R^\bullet) and a thioly radical and a reduced radical (R^-) are formed. Reaction (2): The thiolate is then regenerated by intramolecular reaction with Y132, producing a tyrosyl radical. Reaction (3): The tyrosyl radical reacts with another R^\bullet , forming a stable Tyr-R adduct. The same reaction is also proposed for Y22, identified in the paper, and an additional tyrosine with an unknown location.

Figure 11. A1M protects against destruction of collagen fibrils. Collagen S (5 $\mu\text{g}/\text{mL}$) was incubated with buffer (50 mM Tris-HCl, 0.15 M NaCl, pH 7.6) for 24 hours at room temperature to allow fibrillation. Fibrils were then incubated for a second 24 hour period at room temperature with buffer only (**A**), OxyHb (**B**, 30 μM) or OxyHb and A1M (**C**, 30 μM)

Hb, 10 μM A1M). The samples were then adsorbed for 1 min onto carbon-coated grids, briefly washed with water, and stained with 0.75% (w/w) uranyl formate. The grids had been rendered hydrophilic by glow discharge at low pressure in air beforehand. The samples were analyzed in a Jeol JEM 1230 electron microscope, operated at 80 kV accelerating voltage. Images were recorded with a Gatan Multiscan 791 charge-coupled device camera. The scale bar indicates 0.2 μm . Similar results were obtained with heme (20 μM) and the Fenton-reaction mixture (100 μM Fe^{3+} , 1.0 mM ascorbate, 200 μM H_2O_2) as described (216).

Figure 12. Schematic representation of the principles differentiating intravascular and extravascular hemolysis. When immunoglobulin M (IgM) binds to the red blood cell (RBC) surface, complement (C) is easily activated and membrane attack complexes (MAC) are formed. This leads to membrane rupture intravascularly and release of free Hb into the blood plasma. This phenomenon is often observed in so-called cold-type autoimmune hemolytic anemia and following ABO-incompatible blood transfusions due to the IgM antibodies made against carbohydrate targets on the RBC surface. If immunoglobulin G (IgG) binds instead, immediate and complement-dependent lysis is rarely observed. Instead, IgG-coated RBCs are captured in the reticulo-endothelial system, e.g. when passing the sinusoids of the spleen as indicated in the figure. Based on Fc receptor recognition of the Fc portion of IgG, phagocytosis/lysis of the RBCs will take place extravascularly but intracellularly. This phenomenon is often observed in the warm type of autoimmune hemolytic disorder and is in most cases due to IgG against epitopes on the Rh blood group proteins.

Figure 13. Two-stage model of preeclampsia. Stage 1, during which the placenta is affected, begins with a defect placentation, most likely involving reduced perfusion and hypoxia (243). Oxidative stress is an essential component damaging the placental structure. These placental


lesions induce increased synthesis and accumulation of free Hb in the PE placenta (56,218). Free Hb causes further damage to the blood-placenta-barrier leading to leakage of free Hb into the maternal circulation (16,190,218). Increasing levels of free Hb induces oxidative stress in the maternal circulation, aggravating the endothelial damage seen in PE. Stage two is characterized by the maternal clinical manifestations that appears from gestational week 20. Typical symptoms are hypertension, oedema and proteinuria, kidney and general organ damage that is based on endothelial inflammation.


Figure 14. A) Schematic representation of the event following intraventricular hemorrhage (IVH). Right panels: Coronal ultrasound sections of the preterm rabbit brain showing evolution of (IVH). B) Normal brain without hemorrhage. Normal size of the left lateral ventricle is delineated; C) Acute IVH with both lateral ventricles filled and distended by high-echogenic blood. Size of left lateral ventricle is delineated; D) Resorption of intraventricular blood with clot formation and remaining distention of lateral ventricles at 48h of age; E) Established post-hemorrhagic ventricular dilatation at 72h of age. Ventricular walls exhibit characteristic increase in echogenicity.


Figure 15. Proposed causes of non-healing leg wounds. The pathophysiology of chronic leg ulcers is complex and can be described as a balance between factors/processes leading to wound healing and opposed factors/processes preventing wound healing and hence promoting non-healing wounds. The most common factors/processes of each kind are shown in the figure and are discussed in the text.


Figure 16. A. Venous leg ulcer, positioned adjacent to the medial malleolus (©Ragnar Mårtensson, Department of Dermatology, Lund). B. (x40) and C. (x40) show porphyrins and


iron depositions respectively in skin from a patient with chronic ulcers. Porphyrins were stained with Fouchet technique and iron with Perls' Prussian blue stain. High density depositions are marked with arrows. FC=fibrin cuffs; V=vessel. D. Immunohistochemical localization of A1M (x10) in the tissue from chronic leg ulcers. The section was immunostained using polyclonal antiserum against A1M followed by peroxidase/DAB incubation. A1M appears in high density in basement membranes and fibrin cuffs surrounding the blood vessels (arrows). The Figure is modified from Allhorn et al. (11).












Heme





Biliverdin + CO + Fe^{2+/3+}


Biliverdin reductase
↓


Bilirubin



↓
Storage by ferritin



AMBP-gene







Hemoglobin

Heme

ROS,
free radicals

Oxidative
damage

1. Reduction

2. Heme binding


3. Radical scavenging




4. Up-regulation
of synthesis


Liver

A1M

Peripheral
organs







Placenta

Early defect
placentation
(hypoxia?)

Oxidative
stress

HbF

Placenta
injury

A1M

Stage 1
week < 20


Mother



HbF
HbA

A1M

Preeclampsia:
Endothelial inflammation
Hypertension
Proteinuria
Edema


Stage 2
week > 20






Matrix degradation
Growth factor inactivation
Migration inhibition
Bacterial influence
Hemolysis
High oxidative stress

Matrix production
Reepithelialisation
Angiogenesis
Cell migration
No hemolysis
Controlled oxidative stress




No healing


Healing



Heme



Iron



A1M

References

1. Abraham NG and Kappas A. Pharmacological and clinical aspects of heme oxygenase. *Pharmacol Rev* 60: 79-127, 2008.
2. Ackerman Z, Seidenbaum M, Loewenthal E and Rubinow A. Overload of iron in the skin of patients with varicose ulcers. Possible contributing role of iron accumulation in progression of the disease. *Arch Dermatol* 124: 1376-8, 1988.
3. Adams-Chapman I, Hansen NI, Stoll BJ and Higgins R. Neurodevelopmental outcome of extremely low birth weight infants with posthemorrhagic hydrocephalus requiring shunt insertion. *Pediatrics* 121: e1167-77, 2008.
4. Aft RL and Mueller GC. Hemin-mediated DNA strand scission. *J Biol Chem* 258: 12069-72, 1983.
5. Afzal-Ahmed I, Mann GE, Shennan AH, Poston L and Naftalin RJ. Preeclampsia inactivates glucose-6-phosphate dehydrogenase and impairs the redox status of erythrocytes and fetal endothelial cells. *Free Radic Biol Med* 42: 1781-90, 2007.
6. Agren MS, Eaglstein WH, Ferguson MW, Harding KG, Moore K, Saarialho-Kere UK and Schultz GS. Causes and effects of the chronic inflammation in venous leg ulcers. *Acta Derm Venereol Suppl (Stockh)* 210: 3-17, 2000.
7. Alayash AI. Haptoglobin: Old protein with new functions. *Clin Chim Acta* 412: 493-8, 2011.
8. Allhorn M, Berggård T, Nordberg J, Olsson ML and Åkerström B. Processing of the lipocalin α_1 -microglobulin by hemoglobin induces heme-binding and heme-degradation properties. *Blood* 99: 1894-901, 2002.
9. Allhorn M, Klapayta A and Åkerström B. Redox properties of the lipocalin α_1 -microglobulin: reduction of cytochrome c, hemoglobin, and free iron. *Free Radic Biol Med* 38: 557-67, 2005.
10. Allhorn M, Lundqvist K, Schmidtchen A and Akerstrom B. Heme-scavenging role of alpha1-microglobulin in chronic ulcers. *J Invest Dermatol* 121: 640-6, 2003.
11. Allhorn M, Lundqvist K, Schmidtchen A and Åkerström B. Heme-scavenging role of α_1 -microglobulin in chronic ulcers. *J Invest Dermatol* 121: 640-6, 2003.
12. Altruda F, Poli V, Restagno G, Argos P, Cortese R and Silengo L. The primary structure of human hemopexin deduced from cDNA sequence: evidence for internal, repeating homology. *Nucleic Acids Res* 13: 3841-59, 1985.
13. Amoresano A, Minchiotti L, Cosulich ME, Campagnoli M, Pucci P, Andolfo A, Gianazza E and Galliano M. Structural characterization of the oligosaccharide chains of human alpha1-microglobulin from urine and amniotic fluid. *Eur J Biochem* 267: 2105-12, 2000.
14. An X and Mohandas N. Disorders of red cell membrane. *Br J Haematol* 141: 367-75, 2008.
15. Anderson BF, Baker HM, Norris GE, Rumball SV and Baker EN. Apolactoferrin structure demonstrates ligand-induced conformational change in transferrins. *Nature* 344: 784-7, 1990.
16. Anderson UD, Olsson MG, Rutardóttir S, Centlow M, Kristensen KH, Isberg PE, Thilaganathan B, Åkerström B and Hansson SR. Fetal hemoglobin and α_1 -microglobulin as first- and early second-trimester predictive biomarkers for preeclampsia. *Am J Obstet Gynecol* 204: 520 e1-5, 2011.
17. Applegate LA, Luscher P and Tyrrell RM. Induction of heme oxygenase: a general response to oxidant stress in cultured mammalian cells. *Cancer Res* 51: 974-8, 1991.
18. Arner ES and Holmgren A. Physiological functions of thioredoxin and thioredoxin reductase. *Eur J Biochem* 267: 6102-9, 2000.
19. Arnhold J. Properties, functions, and secretion of human myeloperoxidase. *Biochemistry (Mosc)* 69: 4-9, 2004.
20. Arnhold J and Flemmig J. Human myeloperoxidase in innate and acquired immunity. *Arch Biochem Biophys* 500: 92-106, 2010.

21. Ascenzi P, Bocedi A, Visca P, Altruda F, Tolosano E, Beringhelli T and Fasano M. Hemoglobin and heme scavenging. *IUBMB Life* 57: 749-59, 2005.
22. Axt-Fliedner R. Second trimester uterine artery Doppler ultrasound as a screening test for adverse pregnancy outcome. *Clin Exp Obstet Gynecol.* 31: 9-11, 2004.
23. Azarov I, He X, Jeffers A, Basu S, Ucer B, Hantgan RR, Levy A and Kim-Shapiro DB. Rate of nitric oxide scavenging by hemoglobin bound to haptoglobin. *Nitric Oxide* 18: 296-302, 2008.
24. Babiker-Mohamed H, Olsson ML, Boketoft A, Lögdberg L and Åkerström B. α_1 -microglobulin is mitogenic to human peripheral blood lymphocytes. Regulation by both enhancing and suppressive serum factors. *Immunobiology* 180: 221-34, 1990.
25. Babiker-Mohamed H, Åkerström B and Lögdberg L. Mitogenic effect of alpha-1-microglobulin on mouse lymphocytes. Evidence of T- and B-cell cooperation, B-cell proliferation, and a low-affinity receptor on mononuclear cells. *Scand J Immunol* 32: 37-44, 1990.
26. Baker HM, Anderson BF and Baker EN. Dealing with iron: common structural principles in proteins that transport iron and heme. *Proc Natl Acad Sci U S A* 100: 3579-83, 2003.
27. Balla G, Jacob HS, Eaton JW, Belcher JD and Vercellotti GM. Hemin: a possible physiological mediator of low density lipoprotein oxidation and endothelial injury. *Arterioscler Thromb* 11: 1700-11, 1991.
28. Baudin-Creuzat V, Chauvierre C, Domingues E, Kiger L, Leclerc L, Vasseur C, Celier C and Marden MC. Octamers and nanoparticles as hemoglobin based blood substitutes. *Biochim Biophys Acta* 1784: 1448-53, 2008.
29. Berggård T, Enghild JJ, Badve S, Salafia CM, Lögdberg L and Åkerström B. Histologic distribution and biochemical properties of α_1 -microglobulin in human placenta. *Am J Reprod Immunol* 41: 52-60, 1999.
30. Berggård T, Oury TD, Thogersen IB, Åkerström B and Enghild JJ. α_1 -microglobulin is found both in blood and in most tissues. *J Histochem Cytochem* 46: 887-94, 1998.
31. Berggård T, Thelin N, Falkenberg C, Enghild JJ and Åkerström B. Prothrombin, albumin and immunoglobulin A form covalent complexes with α_1 -microglobulin in human plasma. *Eur J Biochem* 245: 676-83, 1997.
32. Bian Y, Rong Z and Chang TM. Polyhemoglobin-superoxide dismutase-catalase-carbonic anhydrase: a novel biotechnology-based blood substitute that transports both oxygen and carbon dioxide and also acts as an antioxidant. *Artif Cells Blood Substit Immobil Biotechnol* 39: 127-36, 2011.
33. Boretti FS, Buehler PW, D'Agnillo F, Kluge K, Glaus T, Butt OI, Jia Y, Goede J, Pereira CP, Maggiorini M, Schoedon G, Alayash AI and Schaer DJ. Sequestration of extracellular hemoglobin within a haptoglobin complex decreases its hypertensive and oxidative effects in dogs and guinea pigs. *J Clin Invest* 119: 2271-80, 2009.
34. Bouic P, Kanitakis J, Schmitt D, Vincent C, Revillard JP and Thivolet J. Alpha 1-microglobulin: a new antigenic component of the epidermo-dermal junction in normal human skin. *Br J Dermatol* 112: 35-41, 1985.
35. Bouic P, Vincent C and Revillard JP. Localization of alpha 1-microglobulin (HC protein) in normal human tissues: an immunohistochemical study using monoclonal antibodies. *Histochem J* 16: 1311-24, 1984.
36. Braggins PE, Trakshel GM, Kutty RK and Maines MD. Characterization of two heme oxygenase isoforms in rat spleen: comparison with the hematin-induced and constitutive isoforms of the liver. *Biochem Biophys Res Commun* 141: 528-33, 1986.
37. Bratosin D, Mazurier J, Tissier JP, Estaquier J, Huart JJ, Ameisen JC, Aminoff D and Montreuil J. Cellular and molecular mechanisms of senescent erythrocyte phagocytosis by macrophages. A review. *Biochimie* 80: 173-95, 1998.
38. Bratt T, Cedervall T and Åkerström B. Processing and secretion of rat α_1 -microglobulin-bikunin expressed in eukaryotic cell lines. *FEBS Lett* 354: 57-61, 1994.

39. Bratt T, Olsson H, Sjöberg EM, Jergil B and Åkerström B. Cleavage of the α_1 -microglobulin-bikunin precursor is localized to the Golgi apparatus of rat liver cells. *Biochim Biophys Acta* 1157: 147-54, 1993.
40. Brittenham GM. Iron-chelating therapy for transfusional iron overload. *N Engl J Med* 364: 146-56, 2011.
41. Brosens JJ, Pijnenborg R and Brosens IA. The myometrial junctional zone spiral arteries in normal and abnormal pregnancies. *Am J Obstet Gynecol.* 187: 1416-1423, 2002.
42. Broughton Pipkin F. What is the place of genetics in the pathogenesis of pre-eclampsia? *Biol Neonate.* 76: 325-330, 1999.
43. Brouwer A, Groenendaal F, van Haastert IL, Rademaker K, Hanlo P and de Vries L. Neurodevelopmental outcome of preterm infants with severe intraventricular hemorrhage and therapy for post-hemorrhagic ventricular dilatation. *J Pediatr* 152: 648-54, 2008.
44. Bruce LJ. Red cell membrane transport abnormalities. *Curr Opin Hematol* 15: 184-90, 2008.
45. Bucci E, Kwansa H, Koehler RC and Matheson B. Development of zero-link polymers of hemoglobin, which do not extravasate and do not induce pressure increases upon infusion. *Artif Cells Blood Substit Immobil Biotechnol* 35: 11-8, 2007.
46. Buehler PW, Abraham B, Vallelian F, Linnemayr C, Pereira CP, Cipollo JF, Jia Y, Mikolajczyk M, Boretti FS, Schoedon G, Alayash AI and Schaer DJ. Haptoglobin preserves the CD163 hemoglobin scavenger pathway by shielding hemoglobin from peroxidative modification. *Blood* 113: 2578-86, 2009.
47. Buehler PW and D'Agnillo F. Toxicological consequences of extracellular hemoglobin: biochemical and physiological perspectives. *Antioxid Redox Signal* 12: 275-91, 2010.
48. Buehler PW, Vallelian F, Mikolajczyk MG, Schoedon G, Schweizer T, Alayash AI and Schaer DJ. Structural stabilization in tetrameric or polymeric hemoglobin determines its interaction with endogenous antioxidant scavenger pathways. *Antioxid Redox Signal* 10: 1449-62, 2008.
49. Buehler PW, Zhou Y, Cabrales P, Jia Y, Sun G, Harris DR, Tsai AG, Intaglietta M and Palmer AF. Synthesis, biophysical properties and pharmacokinetics of ultrahigh molecular weight tense and relaxed state polymerized bovine hemoglobins. *Biomaterials* 31: 3723-35, 2010.
50. Bunn H (Ed) *Hemoglobin* Weinheim; 1992.
51. Bylund J, Brown KL, Movitz C, Dahlgren C and Karlsson A. Intracellular generation of superoxide by the phagocyte NADPH oxidase: how, where, and what for? *Free Radic Biol Med* 49: 1834-45, 2010.
52. Cabrales P, Tsai AG and Intaglietta M. Hemorrhagic shock resuscitation with carbon monoxide saturated blood. *Resuscitation* 72: 306-18, 2007.
53. Caccia D, Ronda L, Frassi R, Perrella M, Del Favero E, Bruno S, Pioselli B, Abbruzzetti S, Viappiani C and Mozzarelli A. PEGylation promotes hemoglobin tetramer dissociation. *Bioconjug Chem* 20: 1356-66, 2009.
54. Calero M, Escribano J, Grubb A and Mendez E. Location of a novel type of interpolypeptide chain linkage in the human protein HC-IgA complex (HC-IgA) and identification of a heterogeneous chromophore associated with the complex. *J Biol Chem* 269: 384-9, 1994.
55. Camejo G, Halberg C, Manschik-Lundin A, Hurt-Camejo E, Rosengren B, Olsson H, Hansson GI, Forsberg GB and Ylhen B. Hemin binding and oxidation of lipoproteins in serum: mechanisms and effect on the interaction of LDL with human macrophages. *J Lipid Res* 39: 755-66, 1998.
56. Centlow M, Carninci P, Nemeth K, Mezey E, Brownstein M and Hansson SR. Placental expression profiling in preeclampsia: local overproduction of hemoglobin may drive pathological changes. *Fertil Steril* 90: 1834-43, 2008.
57. Centlow M, Wingren C, Borrebaeck C, Brownstein MJ and Hansson SR. Differential gene expression analysis of placentas with increased vascular resistance and pre-eclampsia using whole-genome microarrays. *J Pregnancy* 2011: 472354, 2011.
58. Chan P and Salier JP. Mouse alpha-1-microglobulin/bikunin precursor: cDNA analysis, gene evolution and physical assignment of the gene next to the orosomuroid locus. *Biochim Biophys Acta* 1174: 195-200, 1993.

59. Chang TM. Future generations of red blood cell substitutes. *J Intern Med* 253: 527-35, 2003.
60. Chang TM. Blood substitutes based on nanobiotechnology. *Trends Biotechnol* 24: 372-7, 2006.
61. Chelikani P, Fita I and Loewen PC. Diversity of structures and properties among catalases. *Cell Mol Life Sci* 61: 192-208, 2004.
62. Chen JY, Scerbo M and Kramer G. A review of blood substitutes: examining the history, clinical trial results, and ethics of hemoglobin-based oxygen carriers. *Clinics (Sao Paulo)* 64: 803-13, 2009.
63. Chen L, Mao SJ and Larsen WJ. Identification of a factor in fetal bovine serum that stabilizes the cumulus extracellular matrix. A role for a member of the inter-alpha-trypsin inhibitor family. *J Biol Chem* 267: 12380-6, 1992.
64. Cimen MY. Free radical metabolism in human erythrocytes. *Clin Chim Acta* 390: 1-11, 2008.
65. Cole RH and Vandegriff KD. MP4, a vasodilatory PEGylated hemoglobin. *Adv Exp Med Biol* 701: 85-90, 2011.
66. Cooper CE (Ed) *Radical reactions of haem proteins*. London: BMJ Books/Intensive Care Society.; 2002. 66-79 p.
67. Cooper CE, Patel RP, Brookes PS and Darley-Usmar VM. Nanotransducers in cellular redox signaling: modification of thiols by reactive oxygen and nitrogen species. *Trends Biochem Sci* 27: 489-92, 2002.
68. Davies CE, Wilson MJ, Hill KE, Stephens P, Hill CM, Harding KG and Thomas DW. Use of molecular techniques to study microbial diversity in the skin: chronic wounds reevaluated. *Wound Repair Regen* 9: 332-40, 2001.
69. Davies KJ. Protein damage and degradation by oxygen radicals. I. general aspects. *J Biol Chem* 262: 9895-901, 1987.
70. de Albuquerque Wobeto VP, Zaccariotto TR and Sonati MF. Polymorphism of human haptoglobin and its clinical importance. *Genetics and Molecular Biology* 31: 602-620, 2008.
71. de Groot CJM and Taylor RN. New Insights into the etiology of pre-eclampsia. *Annals of Medicine* 25: 243-249, 1993.
72. Delanghe JR and Langlois MR. Hemopexin: a review of biological aspects and the role in laboratory medicine. *Clin Chim Acta* 312: 13-23, 2001.
73. Delaunay J. The molecular basis of hereditary red cell membrane disorders. *Blood Rev* 21: 1-20, 2007.
74. DeMars DD, Katzmann JA, Kimlinger TK, Calore JD and Tracy RP. Simultaneous measurement of total and IgA-conjugated alpha-1-microglobulin by a combined immunoenzyme/immunoradiometric assay technique. *Clin Chem* 35: 766-72, 1989.
75. Dexlin-Melby L, Sandström A, Centlow M, Nygren S, Hansson SR, Borrebaeck CAK and Wingren C. Tissue proteome profiling of preeclamptic placenta using recombinant antibody microarrays. *Proteomics Clin. Appl.* in press, 2010.
76. Diarra-Mehrpour M, Bourguignon J, Sesboue R, Mattei MG, Passage E, Salier JP and Martin JP. Human plasma inter-alpha-trypsin inhibitor is encoded by four genes on three chromosomes. *Eur J Biochem* 179: 147-54, 1989.
77. Diarra-Mehrpour M, Bourguignon J, Sesboue R, Salier JP, Leveillard T and Martin JP. Structural analysis of the human inter-alpha-trypsin inhibitor light-chain gene. *Eur J Biochem* 191: 131-9, 1990.
78. Droge W. Free radicals in the physiological control of cell function. *Physiol Rev* 82: 47-95, 2002.
79. Duley L. Pre-eclampsia, eclampsia, and hypertension. *Clin Evid (Online)* 2011, 2011.
80. Egler RA, Fernandes E, Rothermund K, Sereika S, de Souza-Pinto N, Jaruga P, Dizdaroglu M and Prochownik EV. Regulation of reactive oxygen species, DNA damage, and c-Myc function by peroxiredoxin 1. *Oncogene* 24: 8038-50, 2005.

81. Eisenstein RS, Garcia-Mayol D, Pettingell W and Munro HN. Regulation of ferritin and heme oxygenase synthesis in rat fibroblasts by different forms of iron. *Proc Natl Acad Sci U S A* 88: 688-92, 1991.
82. Ekström B and Berggård I. Human α_1 -microglobulin. Purification procedure, chemical and physicochemical properties. *J Biol Chem* 252: 8048-57, 1977.
83. Escribano J, Grubb A, Calero M and Mendez E. The protein HC chromophore is linked to the cysteine residue at position 34 of the polypeptide chain by a reduction-resistant bond and causes the charge heterogeneity of protein HC. *J Biol Chem* 266: 15758-63, 1991.
84. Estep T, Bucci E, Farmer M, Greenburg G, Harrington J, Kim HW, Klein H, Mitchell P, Nemo G, Olsen K, Palmer A, Valeri CR and Winslow R. Basic science focus on blood substitutes: a summary of the NHLBI Division of Blood Diseases and Resources Working Group Workshop, March 1, 2006. *Transfusion* 48: 776-82, 2008.
85. Everse J and Hsia N. The toxicities of native and modified hemoglobins. *Free Radic Biol Med* 22: 1075-99, 1997.
86. Faber HR, Groom CR, Baker HM, Morgan WT, Smith A and Baker EN. 1.8 Å crystal structure of the C-terminal domain of rabbit serum haemopexin. *Structure* 3: 551-9, 1995.
87. Falanga V, Kirsner R, Katz MH, Gould E, Eaglstein WH and McFalls S. Pericapillary fibrin cuffs in venous ulceration. Persistence with treatment and during ulcer healing. *J Dermatol Surg Oncol* 18: 409-14, 1992.
88. Falkenberg C, Enghild JJ, Thogersen IB, Salvesen G and Åkerström B. Isolation and characterization of fibronectin- α_1 -microglobulin complex in rat plasma. *Biochem J* 301 (Pt 3): 745-51, 1994.
89. Falkenberg C, Grubb A and Åkerström B. Isolation of rat serum α_1 -microglobulin. Identification of a complex with alpha 1-inhibitor-3, a rat alpha 2-macroglobulin homologue. *J Biol Chem* 265: 16150-7, 1990.
90. Fernandez-Luna JL, Leyva-Cobian F and Mollinedo F. Identification of the protein HC receptor. *FEBS Lett* 236: 471-4, 1988.
91. Fibach E and Rachmilewitz E. The role of oxidative stress in hemolytic anemia. *Curr Mol Med* 8: 609-19, 2008.
92. Flower DR. The lipocalin protein family: structure and function. *Biochem J* 318 (Pt 1): 1-14, 1996.
93. Forget BH and Higgs DR (Eds). *Disorders of Hemoglobin: Genetics, Pathophysiology, and Clinical Management* New York: Cambridge University Press; 2009.
94. Fries E and Blom AM. Bikunin--not just a plasma proteinase inhibitor. *Int J Biochem Cell Biol* 32: 125-37, 2000.
95. Fujita T, Toda K, Karimova A, Yan SF, Naka Y, Yet SF and Pinsky DJ. Paradoxical rescue from ischemic lung injury by inhaled carbon monoxide driven by derepression of fibrinolysis. *Nat Med* 7: 598-604, 2001.
96. Fukai T and Ushio-Fukai M. Superoxide dismutases: role in redox signaling, vascular function, and diseases. *Antioxid Redox Signal* 15: 1583-606, 2011.
97. Futagi Y, Suzuki Y, Toribe Y, Nakano H and Morimoto K. Neurodevelopmental outcome in children with posthemorrhagic hydrocephalus. *Pediatr Neurol* 33: 26-32, 2005.
98. Futagi Y, Toribe Y, Ogawa K and Suzuki Y. Neurodevelopmental outcome in children with intraventricular hemorrhage. *Pediatr Neurol* 34: 219-24, 2006.
99. Ganfornina L, Sanchez D, Greene LH and Flower DR. The lipocalin protein family. Protein sequence, structure and relationship to calycin superfamily. In: *Lipocalins*. edited by Åkerström B, Borregaard N, Flower DR and Salier JP. Georgetown, TX.: Landes Bioscience; 2006. pp. 17-27.
100. Giannubilo SR, Tranquilli AL, Santolini C, Principi F, Mancinelli R and Littarru GP. Placental CoQ10 levels in HELLP syndrome. *Biofactors* 25: 159-63, 2005.

101. Giardine B, van Baal S, Kaimakis P, Riemer C, Miller W, Samara M, Kollia P, Anagnou NP, Chui DH, Wajcman H, Hardison RC and Patrinos GP. HbVar database of human hemoglobin variants and thalassemia mutations: 2007 update. *Hum Mutat* 28: 206, 2007.
102. Goldsmith LA. Clinical Snippets. *Journal of Investigative Dermatology* 121: V, 2003.
103. Goswami T, Rolfs A and Hediger MA. Iron transport: emerging roles in health and disease. *Biochem Cell Biol* 80: 679-89, 2002.
104. Goyal MM and Basak A. Human catalase: looking for complete identity. *Protein Cell* 1: 888-97, 2010.
105. Gozzelino R, Jeney V and Soares MP. Mechanisms of cell protection by heme oxygenase-1. *Annu Rev Pharmacol Toxicol* 50: 323-54, 2010.
106. Granger JP, Alexander BT, Bennett WA and Khalil RA. Pathophysiology of pregnancy-induced hypertension. *Am J Hypertens.* 14: 178S-185S, 2001.
107. Graversen JH, Madsen M and Moestrup SK. CD163: a signal receptor scavenging haptoglobin-hemoglobin complexes from plasma. *Int J Biochem Cell Biol* 34: 309-14, 2002.
108. Grey M, Yainoy S, Prachayasittikul V and Bulow L. A superoxide dismutase-human hemoglobin fusion protein showing enhanced antioxidative properties. *FEBS J* 276: 6195-203, 2009.
109. Grinberg LN, O'Brien PJ and Hrkal Z. The effects of heme-binding proteins on the peroxidative and catalytic activities of hemin. *Free Radic Biol Med* 27: 214-9, 1999.
110. Grubb A, Mendez E, Fernandez-Luna JL, Lopez C, Mihaesco E and Vaerman JP. The molecular organization of the protein HC-IgA complex (HC-IgA). *J Biol Chem* 261: 14313-20, 1986.
111. Grubb AO, Lopez C, Tejler L and Mendez E. Isolation of human complex-forming glycoprotein, heterogeneous in charge (protein HC), and its IgA complex from plasma. Physicochemical and immunochemical properties, normal plasma concentration. *J Biol Chem* 258: 14698-707, 1983.
112. Gutteridge JM. The antioxidant activity of haptoglobin towards haemoglobin-stimulated lipid peroxidation. *Biochim Biophys Acta* 917: 219-23, 1987.
113. Halliwell B and Gutteridge JM. The definition and measurement of antioxidants in biological systems. *Free Radic Biol Med* 18: 125-6, 1995.
114. Halliwell B and Gutteridge JMC. *Free Radicals in Biology and Medicine.* (4th ed): Oxford University Press; 2007.
115. Hanley S and Powell R. Sequence of a cDNA clone encoding the Atlantic salmon alpha 1-microglobulin/bikunin protein. *Gene* 147: 297-8, 1994.
116. Hansson SR, Chen Y, Brodzki J, Chen M, Hernandez-Andrade E, Inman JM, Kozhich OA, Larsson I, Marsal K, Medstrand P, Xiang CC and Brownstein MJ. Gene expression profiling of human placentas from preeclamptic and normotensive pregnancies. *Mol Hum Reprod* 12: 169-79, 2006.
117. Hardison RC. A brief history of hemoglobins: plant, animal, protist, and bacteria. *Proceedings of the National Academy of Sciences of the United States of America* 93: 5675-9, 1996.
118. Harnois T, Rousselot M, Rogniaux H and Zal F. High-level production of recombinant *Arenicola marina* globin chains in *Escherichia coli*: a new generation of blood substitute. *Artif Cells Blood Substit Immobil Biotechnol* 37: 106-16, 2009.
119. Harrison GA, Humphrey KE, Jones N, Badenhop R, Guo G, Elakis G, Kaye JA, Turner RJ, Grehan M, Wilton AN, Brennecke SP and Cooper DW. A genomewide linkage study of preeclampsia/eclampsia reveals evidence for a candidate region on 4q. *Am J Hum Genet.* 60: 1158-1167, 1997.
120. Hayashi S, Omata Y, Sakamoto H, Higashimoto Y, Hara T, Sagara Y and Noguchi M. Characterization of rat heme oxygenase-3 gene. Implication of processed pseudogenes derived from heme oxygenase-2 gene. *Gene* 336: 241-50, 2004.
121. Henkel-Honke T and Oleck M. Artificial oxygen carriers: a current review. *AANA J* 75: 205-11, 2007.

122. Herrera E and Barbas C. Vitamin E: action, metabolism and perspectives. *J Physiol Biochem* 57: 43-56, 2001.
123. Holmgren A. Thioredoxin and glutaredoxin systems. *J Biol Chem* 264: 13963-6, 1989.
124. Hu D and Kluger R. Functional cross-linked hemoglobin bis-tetramers: geometry and cooperativity. *Biochemistry* 47: 12551-61, 2008.
125. Hu T, Li D, Manjula BN, Brenowitz M, Prabhakaran M and Acharya SA. PEGylation of Val-1(α) destabilizes the tetrameric structure of hemoglobin. *Biochemistry* 48: 608-16, 2009.
126. Hung TH, Skepper JN, Charnock-Jones DS and Burton GJ. Hypoxia-reoxygenation: a potent inducer of apoptotic changes in the human placenta and possible etiological factor in preeclampsia. *Circ Res* 90: 1274-1281, 2002.
127. Hvidberg V, Maniecki MB, Jacobsen C, Hojrup P, Moller HJ and Moestrup SK. Identification of the receptor scavenging hemopexin-heme complexes. *Blood* 106: 2572-9, 2005.
128. Ide H, Itoh H and Nawa Y. Sequencing of cDNAs encoding α 1-microglobulin/bikunin of Mongolian gerbil and Syrian golden hamster in comparison with man and other species. *Biochim Biophys Acta* 1209: 286-92, 1994.
129. Immenschuh S and Baumgart-Vogt E. Peroxiredoxins, oxidative stress, and cell proliferation. *Antioxid Redox Signal* 7: 768-77, 2005.
130. Immenschuh S, Nell C, Iwahara S, Katz N and Muller-Eberhard U. Gene regulation of HBP 23 by metalloporphyrins and protoporphyrin IX in liver and hepatocyte cultures. *Biochem Biophys Res Commun* 231: 667-70, 1997.
131. Ishii T, Iwahashi H, Sugata R and Kido R. Oxidation of 3-hydroxykynurenine catalyzed by methemoglobin with hydrogen peroxide. *Free Radic Biol Med* 13: 17-20, 1992.
132. Itoh H, Tomita M, Kobayashi T, Uchino H, Maruyama H and Nawa Y. Expression of inter-alpha-trypsin inhibitor light chain (bikunin) in human pancreas. *J Biochem* 120: 271-5, 1996.
133. Iwahara S, Satoh H, Song DX, Webb J, Burlingame AL, Nagae Y and Muller-Eberhard U. Purification, characterization, and cloning of a heme-binding protein (23 kDa) in rat liver cytosol. *Biochemistry* 34: 13398-406, 1995.
134. Jarolim P, Lahav M, Liu SC and Palek J. Effect of hemoglobin oxidation products on the stability of red cell membrane skeletons and the associations of skeletal proteins: correlation with a release of hemin. *Blood* 76: 2125-31, 1990.
135. Jones DP, Carlson JL, Mody VC, Cai J, Lynn MJ and Sternberg P. Redox state of glutathione in human plasma. *Free Radic Biol Med* 28: 625-35, 2000.
136. Jose MD and Lynn KL. Acute renal failure in a patient with paroxysmal nocturnal hemoglobinuria. *Clin Nephrol* 56: 172-4, 2001.
137. Kalantari H and Das DK. Physiological effects of resveratrol. *Biofactors* 36: 401-6, 2010.
138. Kapralov A, Vlasova, II, Feng W, Maeda A, Walson K, Tyurin VA, Huang Z, Aneja RK, Carcillo J, Bayir H and Kagan VE. Peroxidase activity of hemoglobin-haptoglobin complexes: covalent aggregation and oxidative stress in plasma and macrophages. *J Biol Chem* 284: 30395-407, 2009.
139. Kastern W, Björck L and Åkerström B. Developmental and tissue-specific expression of α ₁-microglobulin mRNA in the rat. *J Biol Chem* 261: 15070-4, 1986.
140. Kaumeyer JF, Polazzi JO and Kotick MP. The mRNA for a proteinase inhibitor related to the HI-30 domain of inter-alpha-trypsin inhibitor also encodes alpha-1-microglobulin (protein HC). *Nucleic Acids Res* 14: 7839-50, 1986.
141. Kawahara A, Hikosaka A, Sasado T and Hirota K. Thyroid hormone-dependent repression of α ₁-microglobulin/bikunin precursor (AMBIP) gene expression during amphibian metamorphosis. *Dev Genes Evol* 206: 355-362, 1997.
142. Kenny LC, Broadhurst DI, Dunn W, Brown M, North RA, McCowan L, Roberts C, Cooper GJ, Kell DB and Baker PN. Robust early pregnancy prediction of later preeclampsia using metabolomic biomarkers. *Hypertension* 56: 741-9, 2010.
143. Khwaja O and Volpe JJ. Pathogenesis of cerebral white matter injury of prematurity. *Arch Dis Child Fetal Neonatal Ed* 93: F153-61, 2008.

144. Kim-Shapiro DB, Schechter AN and Gladwin MT. Unraveling the reactions of nitric oxide, nitrite, and hemoglobin in physiology and therapeutics. *Arteriosclerosis, thrombosis, and vascular biology* 26: 697-705, 2006.
145. Kim KM, Pae HO, Zheng M, Park R, Kim YM and Chung HT. Carbon monoxide induces heme oxygenase-1 via activation of protein kinase R-like endoplasmic reticulum kinase and inhibits endothelial cell apoptosis triggered by endoplasmic reticulum stress. *Circ Res* 101: 919-27, 2007.
146. Kristiansen M, Graversen JH, Jacobsen C, Sonne O, Hoffman HJ, Law SK and Moestrup SK. Identification of the haemoglobin scavenger receptor. *Nature* 409: 198-201, 2001.
147. Krohne-Ehrich G, Schirmer RH and Untucht-Grau R. Glutathione reductase from human erythrocytes. Isolation of the enzyme and sequence analysis of the redox-active peptide. *Eur J Biochem* 80: 65-71, 1977.
148. Kronborg C, Vittinghus E, Allen J and Knudsen UB. Excretion patterns of large and small proteins in pre-eclamptic pregnancies. *Acta Obstet Gynecol Scand* 90: 897-902, 2011.
149. Kumar S and Bandyopadhyay U. Free heme toxicity and its detoxification systems in human. *Toxicol Lett* 157: 175-88, 2005.
150. Kutty RK and Maines MD. Purification and characterization of biliverdin reductase from rat liver. *J Biol Chem* 256: 3956-62, 1981.
151. Lambright DG, Sondek J, Bohm A, Skiba NP, Hamm HE and Sigler PB. The 2.0 Å crystal structure of a heterotrimeric G protein. *Nature* 379: 311-9, 1996.
152. Lara FA, Kahn SA, da Fonseca AC, Bahia CP, Pinho JP, Graca-Souza AV, Houzel JC, de Oliveira PL, Moura-Neto V and Oliveira MF. On the fate of extracellular hemoglobin and heme in brain. *J Cereb Blood Flow Metab* 29: 1109-20, 2009.
153. Larsson J, Allhorn M and Åkerström B. The lipocalin α_1 -microglobulin binds heme in different species. *Arch Biochem Biophys* 432: 196-204, 2004.
154. Larsson J, Wingårdh K, Berggård T, Davies JR, Lögdberg L, Strand SE and Åkerström B. Distribution of iodine ¹²⁵-labeled α_1 -microglobulin in rats after intravenous injection. *J Lab Clin Med* 137: 165-75, 2001.
155. Latunde-Dada GO, Simpson RJ and McKie AT. Recent advances in mammalian haem transport. *Trends Biochem Sci* 31: 182-8, 2006.
156. Laurell CB and Nyman M. Studies on the serum haptoglobin level in hemoglobinemia and its influence on renal excretion of hemoglobin. *Blood* 12: 493-506, 1957.
157. Leaver MJ, Wright J and George SG. Conservation of the tandem arrangement of alpha 1-microglobulin/bikunin mRNA: cloning of a cDNA from plaice (*Pleuronectes platessa*). *Comp Biochem Physiol Biochem Mol Biol* 108: 275-81, 1994.
158. Lee JW, Kim SH and Yoon CJ. Hemosiderin deposition on the renal cortex by mechanical hemolysis due to malfunctioning prosthetic cardiac valve: report of MR findings in two cases. *J Comput Assist Tomogr* 23: 445-7, 1999.
159. Lees CC, Brown AS, Harrington KF, Beacon HJ, Martin JF and Campbell S. A cross-sectional study of platelet volume in healthy normotensive women with bilateral uterine artery notches. *Ultrasound Obstet Gynecol*: 277-281, 1997.
160. Levonen AL, Patel RP, Brookes P, Go YM, Jo H, Parthasarathy S, Anderson PG and Darley-Usmar VM. Mechanisms of cell signaling by nitric oxide and peroxynitrite: from mitochondria to MAP kinases. *Antioxid Redox Signal* 3: 215-29, 2001.
161. Levy AP, Asleh R, Blum S, Levy NS, Miller-Lotan R, Kalet-Litman S, Anbinder Y, Lache O, Nakhoul FM, Asaf R, Farbstein D, Pollak M, Soloveichik YZ, Strauss M, Alshiek J, Livshits A, Schwartz A, Awad H, Jad K and Goldenstein H. Haptoglobin: basic and clinical aspects. *Antioxid Redox Signal* 12: 293-304, 2010.
162. Liao JC, Hein TW, Vaughn MW, Huang KT and Kuo L. Intravascular flow decreases erythrocyte consumption of nitric oxide. *Proc Natl Acad Sci U S A* 96: 8757-61, 1999.

163. Lie RT, Rasmussen S, Brunborg H, Gjessing HK, Lie-Nielsen E and Irgens LM. Fetal and maternal contributions to risk of pre-eclampsia: population based study. *BMJ* 316: 1343-1347, 1998.
164. Lim SK, Kim H, bin Ali A, Lim YK, Wang Y, Chong SM, Costantini F and Baumman H. Increased susceptibility in Hp knockout mice during acute hemolysis. *Blood* 92: 1870-7, 1998.
165. Linden JV, Wagner K, Voytovich AE and Sheehan J. Transfusion errors in New York State: an analysis of 10 years' experience. *Transfusion* 40: 1207-13, 2000.
166. Lindqvist A, Bratt T, Altieri M, Kastern W and Åkerström B. Rat α_1 -microglobulin: co-expression in liver with the light chain of inter-alpha-trypsin inhibitor. *Biochim Biophys Acta* 1130: 63-7, 1992.
167. Lindqvist A, Rouet P, Salier JP and Åkerström B. The α_1 -microglobulin/bikunin gene: characterization in mouse and evolution. *Gene* 234: 329-36, 1999.
168. Lindqvist A and Åkerström B. Bovine α_1 -microglobulin/bikunin. Isolation and characterization of liver cDNA and urinary α_1 -microglobulin. *Biochim Biophys Acta* 1306: 98-106, 1996.
169. Lindqvist A and Åkerström B. Isolation of plaice (*Pleuronectes platessa*) α_1 -microglobulin: conservation of structure and chromophore. *Biochim Biophys Acta* 1430: 222-33, 1999.
170. Lipstein H, Lee CC and Crupi RS. A current concept of eclampsia. *Am J Emerg Med*. 21: 223-226, 2003.
171. Little HN and Neilands JB. Binding of haematin by human serum albumin. *Nature* 188: 913-5, 1960.
172. Lloyd RV, Hanna PM and Mason RP. The origin of the hydroxyl radical oxygen in the Fenton reaction. *Free Radic Biol Med* 22: 885-8, 1997.
173. Lopez C, Grubb A and Mendez E. Human protein HC displays variability in its carboxyl-terminal amino acid sequence. *FEBS Letter* 144: 349-353, 1982.
174. Lubos E, Loscalzo J and Handy DE. Glutathione peroxidase-1 in health and disease: from molecular mechanisms to therapeutic opportunities. *Antioxid Redox Signal* 15: 1957-97, 2011.
175. Lui FE and Kluger R. Enhancing nitrite reductase activity of modified hemoglobin: bis-tetramers and their PEGylated derivatives. *Biochemistry* 48: 11912-9, 2009.
176. Luzzatto L, Gianfaldoni G and Notaro R. Management of paroxysmal nocturnal haemoglobinuria: a personal view. *Br J Haematol* 153: 709-20, 2011.
177. Lögdberg L and Wester L. Immunocalins: a lipocalin subfamily that modulates immune and inflammatory responses. *Biochim Biophys Acta* 1482: 284-97, 2000.
178. Lögdberg L and Åkerström B. Immunosuppressive properties of α_1 -microglobulin. *Scand J Immunol* 13: 383-90, 1981.
179. Lögdberg L, Åkerström B and Shevach EM. α_1 -microglobulin is mitogenic for guinea pig lymphocytes. *Scand J Immunol* 24: 575-81, 1986.
180. Maines MD. New developments in the regulation of heme metabolism and their implications. *Crit Rev Toxicol* 12: 241-314, 1984.
181. Maines MD. Heme oxygenase: function, multiplicity, regulatory mechanisms, and clinical applications. *FASEB J* 2: 2557-68, 1988.
182. Maines MD. The heme oxygenase system: a regulator of second messenger gases. *Annu Rev Pharmacol Toxicol* 37: 517-54, 1997.
183. Maines MD and Kappas A. Cobalt induction of hepatic heme oxygenase; with evidence that cytochrome P-450 is not essential for this enzyme activity. *Proc Natl Acad Sci U S A* 71: 4293-7, 1974.
184. Maines MD, Trakshel GM and Kutty RK. Characterization of two constitutive forms of rat liver microsomal heme oxygenase. Only one molecular species of the enzyme is inducible. *J Biol Chem* 261: 411-9, 1986.

185. Maitland K, Pamba A, English M, Peshu N, Marsh K, Newton C and Levin M. Randomized trial of volume expansion with albumin or saline in children with severe malaria: preliminary evidence of albumin benefit. *Clin Infect Dis* 40: 538-45, 2005.
186. Marklund SL. Extracellular superoxide dismutase and other superoxide dismutase isoenzymes in tissues from nine mammalian species. *Biochem J* 222: 649-55, 1984.
187. Masella R, Di Benedetto R, Vari R, Filesi C and Giovannini C. Novel mechanisms of natural antioxidant compounds in biological systems: involvement of glutathione and glutathione-related enzymes. *J Nutr Biochem* 16: 577-86, 2005.
188. Mattu RK, Trevelyan J, Needham EW, Khan M, Adiseshiah MA, Richter D, Murray RG and Betteridge DJ. Lipoprotein lipase gene variants relate to presence and degree of microalbuminuria in Type II diabetes. *Diabetologia* 45: 905-13, 2002.
189. May K, Rosenlof L, Olsson MG, Centlow M, Morgelin M, Larsson I, Cederlund M, Rutardottir S, Siegmund W, Schneider H, Akerstrom B and Hansson SR. Perfusion of human placenta with hemoglobin introduces preeclampsia-like injuries that are prevented by alpha(1)-microglobulin. *Placenta*, 2011.
190. May K, Rosenlöf L, Olsson MG, Centlow M, Mörgelin M, Larsson I, Cederlund M, Rutardóttir S, Siegmund W, Schneider H, Åkerström B and Hansson SR. Perfusion of human placenta with hemoglobin introduces preeclampsia-like injuries that are prevented by α_1 -microglobulin. *Placenta* 32: 323-32, 2011.
191. McCord JM and Fridovich I. Superoxide dismutase. An enzymic function for erythrocyte (hemocuprein). *J Biol Chem* 244: 6049-55, 1969.
192. McCoubrey WK, Jr. and Maines MD. Domains of rat heme oxygenase-2: the amino terminus and histidine 151 are required for heme oxidation. *Arch Biochem Biophys* 302: 402-8, 1993.
193. McCoubrey WK, Jr. and Maines MD. The structure, organization and differential expression of the gene encoding rat heme oxygenase-2. *Gene* 139: 155-61, 1994.
194. Melamed-Frank M, Lache O, Enav BI, Szafrank T, Levy NS, Ricklis RM and Levy AP. Structure-function analysis of the antioxidant properties of haptoglobin. *Blood* 98: 3693-8, 2001.
195. Mendez E, Fernandez-Luna JL, Grubb A and Leyva-Cobian F. Human protein HC and its IgA complex are inhibitors of neutrophil chemotaxis. *Proc Natl Acad Sci U S A* 83: 1472-5, 1986.
196. Miller YI and Shaklai N. Kinetics of hemin distribution in plasma reveals its role in lipoprotein oxidation. *Biochim Biophys Acta* 1454: 153-64, 1999.
197. Motterlini R. Carbon monoxide-releasing molecules (CO-RMs): vasodilatory, anti-ischaemic and anti-inflammatory activities. *Biochem Soc Trans* 35: 1142-6, 2007.
198. Murphy BP, Inder TE, Rooks V, Taylor GA, Anderson NJ, Mogridge N, Horwood LJ and Volpe JJ. Posthaemorrhagic ventricular dilatation in the premature infant: natural history and predictors of outcome. *Arch Dis Child Fetal Neonatal Ed* 87: F37-41, 2002.
199. Mustacich D and Powis G. Thioredoxin reductase. *Biochem J* 346 Pt 1: 1-8, 2000.
200. Nakata J, Nakahari A, Kato Y, Yamachi N, Sato N, Yonezawa T, Takahashi C, Chikazawa S, Hori Y, Kanai K, Ito N, Higuchi S and Hoshi F. Molecular cloning and expression analysis of feline alpha1-microglobulin. *Vet Immunol Immunopathol* 139: 79-82, 2011.
201. Natanson C, Kern SJ, Lurie P, Banks SM and Wolfe SM. Cell-free hemoglobin-based blood substitutes and risk of myocardial infarction and death: a meta-analysis. *JAMA* 299: 2304-12, 2008.
202. Nauseef WM, Metcalf JA and Root RK. Role of myeloperoxidase in the respiratory burst of human neutrophils. *Blood* 61: 483-92, 1983.
203. Nelzen O, Bergqvist D and Lindhagen A. Leg ulcer etiology--a cross sectional population study. *J Vasc Surg* 14: 557-64, 1991.
204. Nelzen O, Bergqvist D, Lindhagen A and Hallbook T. Chronic leg ulcers: an underestimated problem in primary health care among elderly patients. *J Epidemiol Community Health* 45: 184-7, 1991.
205. Neumann CA, Cao J and Manevich Y. Peroxiredoxin 1 and its role in cell signaling. *Cell Cycle* 8: 4072-8, 2009.

206. Nielsen MJ and Moestrup SK. Receptor targeting of hemoglobin mediated by the haptoglobins: roles beyond heme scavenging. *Blood* 114: 764-71, 2009.
207. Nielsen MJ, Moller HJ and Moestrup SK. Hemoglobin and heme scavenger receptors. *Antioxid Redox Signal* 12: 261-73, 2010.
208. Nielsen MJ, Petersen SV, Jacobsen C, Thirup S, Enghild JJ, Graversen JH and Moestrup SK. A unique loop extension in the serine protease domain of haptoglobin is essential for CD163 recognition of the haptoglobin-hemoglobin complex. *J Biol Chem* 282: 1072-9, 2007.
209. Noguchi M, Yoshida T and Kikuchi G. Identification of the product of heme degradation catalyzed by the heme oxygenase system as biliverdin IX alpha by reversed-phase high-performance liquid chromatography. *J Biochem* 91: 1479-83, 1982.
210. Nordberg J, Allhorn M, Winqvist I, Åkerström B and Olsson ML. Quantitative and qualitative evaluation of plasma and urine α_1 -microglobulin in healthy donors and patients with different haemolytic disorders and haemochromatosis. *Clin Chim Acta* 386: 31-7, 2007.
211. Nordberg J and Arner ES. Reactive oxygen species, antioxidants, and the mammalian thioredoxin system. *Free Radic Biol Med* 31: 1287-312, 2001.
212. Odum L and Nielsen HW. Human protein HC (alpha 1-microglobulin) and inter-alpha-trypsin inhibitor in connective tissue. *Histochem J* 26: 799-803, 1994.
213. Odum L and Nielsen HW. Bikunin and alpha 1-microglobulin in human zona pellucida and connective tissue. *Histochem J* 29: 199-203, 1997.
214. Ohsawa I, Uehara Y, Hashimoto S, Endo M, Fujita T and Ohi H. Autoimmune hemolytic anemia occurred prior to evident nephropathy in a patient with chronic hepatitis C virus infection: case report. *BMC Nephrol* 29: 31-37, 2003.
215. Okuda S, Nishiyama N, Saito H and Katsuki H. Hydrogen peroxide-mediated neuronal cell death induced by an endogenous neurotoxin, 3-hydroxykynurenine. *Proc Natl Acad Sci U S A* 93: 12553-8, 1996.
216. Olsson MG, Allhorn M, Larsson J, Cederlund M, Lundqvist K, Schmidtchen A, Sorensen OE, Mörgelin M and Åkerström B. Up-Regulation of A1M/ α_1 -Microglobulin in Skin by Heme and Reactive Oxygen Species Gives Protection from Oxidative Damage. *PLoS One* 6: e27505, 2011.
217. Olsson MG, Allhorn M, Olofsson T and Åkerström B. Up-regulation of α_1 -microglobulin by hemoglobin and reactive oxygen species in hepatoma and blood cell lines. *Free Radic Biol Med* 42: 842-51, 2007.
218. Olsson MG, Centlow M, Rutardóttir S, Stenfors I, Larsson J, Hosseini-Maaf B, Olsson ML, Hansson SR and Åkerström B. Increased levels of cell-free hemoglobin, oxidation markers, and the antioxidative heme scavenger α_1 -microglobulin in preeclampsia. *Free Radic Biol Med* 48: 284-91, 2010.
219. Olsson MG, Nilsson EJ, Rutardottir S, Paczesny J, Pallon J and Åkerström B. Bystander Cell Death and Stress Response is Inhibited by the Radical Scavenger α_1 -Microglobulin in Irradiated Cell Cultures. *Radiat Res*, 2010.
220. Olsson MG, Olofsson T, Tapper H and Åkerström B. The lipocalin α_1 -microglobulin protects erythroid K562 cells against oxidative damage induced by heme and reactive oxygen species. *Free Radic Res* 42: 725-36, 2008.
221. Otterbein LE, Mantell LL and Choi AM. Carbon monoxide provides protection against hyperoxic lung injury. *Am J Physiol* 276: L688-94, 1999.
222. Padayatty SJ, Katz A, Wang Y, Eck P, Kwon O, Lee JH, Chen S, Corpe C, Dutta A, Dutta SK and Levine M. Vitamin C as an antioxidant: evaluation of its role in disease prevention. *J Am Coll Nutr* 22: 18-35, 2003.
223. Page EW. The relation between hydatid moles, relative ischemia of the gravid uterus, and the placental origin of eclampsia. *Am J Obstet Gynecol.* 37: 291-293, 1939.

224. Paoli M, Anderson BF, Baker HM, Morgan WT, Smith A and Baker EN. Crystal structure of hemopexin reveals a novel high-affinity heme site formed between two beta-propeller domains. *Nat Struct Biol* 6: 926-31, 1999.
225. Park J, Cha DH, Lee SJ, Kim YN, Kim YH and Kim KP. Discovery of the serum biomarker proteins in severe preeclampsia by proteomic analysis. *Exp Mol Med* 43: 427-35, 2011.
226. Pawlak D, Pawlak K, Malyszko J, Mysliwiec M and Buczko W. Accumulation of toxic products degradation of kynurenine in hemodialyzed patients. *Int Urol Nephrol* 33: 399-404, 2001.
227. Petz LD and Garatty G. *Immune Hemolytic Anemias.* (2nd ed) Philadelphia, PA, USA: Churchill Livingstone; 2004.
228. Piantadosi CA. Carbon monoxide, reactive oxygen signaling, and oxidative stress. *Free Radic Biol Med* 45: 562-9, 2008.
229. Poli G, Leonarduzzi G, Biasi F and Chiarpotto E. Oxidative stress and cell signalling. *Curr Med Chem* 11: 1163-82, 2004.
230. Polticelli F, Bocedi A, Minervini G and Ascenzi P. Human haptoglobin structure and function-- a molecular modelling study. *FEBS J* 275: 5648-56, 2008.
231. Polyzos NP, Mauri D, Tsappi M, Tzioras S, Kamposioras K, Cortinovis I and Casazza G. Combined vitamin C and E supplementation during pregnancy for preeclampsia prevention: a systematic review. *Obstet Gynecol Surv* 62: 202-6, 2007.
232. Quijcho FA. Atomic structures of periplasmic binding proteins and the high-affinity active transport systems in bacteria. *Philos Trans R Soc Lond B Biol Sci* 326: 341-51; discussion 351-2, 1990.
233. Rajagopal A, Rao AU, Amigo J, Tian M, Upadhyay SK, Hall C, Uhm S, Mathew MK, Fleming MD, Paw BH, Krause M and Hamza I. Haem homeostasis is regulated by the conserved and concerted functions of HRG-1 proteins. *Nature* 453: 1127-31, 2008.
234. Raju VS, McCoubrey WK, Jr. and Maines MD. Regulation of heme oxygenase-2 by glucocorticoids in neonatal rat brain: characterization of a functional glucocorticoid response element. *Biochim Biophys Acta* 1351: 89-104, 1997.
235. Reeder BJ. The redox activity of hemoglobins: from physiologic functions to pathologic mechanisms. *Antioxid Redox Signal* 13: 1087-123, 2010.
236. Reeder BJ. The redox activity of hemoglobins: from physiologic functions to pathologic mechanisms. *Antioxidants & Redox Signaling* 13: 1087-123, 2010.
237. Reeder BJ, Grey M, Silaghi-Dumitrescu RL, Svistunenko DA, Bulow L, Cooper CE and Wilson MT. Tyrosine residues as redox cofactors in human hemoglobin: implications for engineering nontoxic blood substitutes. *J Biol Chem* 283: 30780-7, 2008.
238. Regan RF and Panter SS. Neurotoxicity of hemoglobin in cortical cell culture. *Neurosci Lett* 153: 219-22, 1993.
239. Reiter CD, Wang X, Tanus-Santos JE, Hogg N, Cannon RO, 3rd, Schechter AN and Gladwin MT. Cell-free hemoglobin limits nitric oxide bioavailability in sickle-cell disease. *Nat Med* 8: 1383-9, 2002.
240. Roberts JM. Endothelial dysfunction in preeclampsia. *Semin Reprod Endocrinol* 16: 5-15, 1998.
241. Roberts JM and Cooper DW. Pathogenesis and genetics of pre-eclampsia. *Lancet* 357: 53-56, 2001.
242. Roberts JM and Hubel CA. Is oxidative stress the link in the two-stage model of pre-eclampsia? *Lancet* 354: 788-789, 1999.
243. Roberts JM and Hubel CA. The two stage model of preeclampsia: variations on the theme. *Placenta* 30 Suppl A: S32-7, 2009.
244. Roberts JM, Myatt L, Spong CY, Thom EA, Hauth JC, Leveno KJ, Pearson GD, Wapner RJ, Varner MW, Thorp JM, Jr., Mercer BM, Peaceman AM, Ramin SM, Carpenter MW, Samuels P, Sciscione A, Harper M, Smith WJ, Saade G, Sorokin Y and Anderson GB. Vitamins C and E to prevent complications of pregnancy-associated hypertension. *N Engl J Med* 362: 1282-91, 2010.

245. Roberts JM, Taylor RN, Musci TJ, Rodgers GM, Hubel CA and McLaughlin MK. Preeclampsia: An endothelial cell disorder. *Am J Obstet Gynecol* 161: 1200-1204, 1989.
246. Roche CJ, Malashkevich V, Balazs TC, Dantsker D, Chen Q, Moreira J, Almo SC, Friedman JM and Hirsch RE. Structural and functional studies indicating altered redox properties of hemoglobin E: implications for production of bioactive nitric oxide. *J Biol Chem* 286: 23452-66, 2011.
247. Rosenfeld M and Surgenor DM. Methemalbumin, interaction between human serum albumin and ferriprotoporphyrin IX. *Journal Biological Chemistry* 183: 663-677, 1950.
248. Rother RP, Bell L, Hillmen P and Gladwin MT. The clinical sequelae of intravascular hemolysis and extracellular plasma hemoglobin: a novel mechanism of human disease. *JAMA : the journal of the American Medical Association* 293: 1653-62, 2005.
249. Rouet P, Raguenez G, Ruminy P and Salier JP. An array of binding sites for hepatocyte nuclear factor 4 of high and low affinities modulates the liver-specific enhancer for the human alpha1-microglobulin/bikunin precursor. *Biochem J* 334 (Pt 3): 577-84, 1998.
250. Rouet P, Raguenez G, Tronche F, Yaniv M, N'Guyen C and Salier JP. A potent enhancer made of clustered liver-specific elements in the transcription control sequences of human alpha 1-microglobulin/bikunin gene. *J Biol Chem* 267: 20765-73, 1992.
251. Ryter SW and Choi AM. Heme oxygenase-1: redox regulation of a stress protein in lung and cell culture models. *Antioxid Redox Signal* 7: 80-91, 2005.
252. Sakai H, Takeoka S and Kobayashi K. Gas bioengineering using hemoglobin-vesicles for versatile clinical applications. *Curr Pharm Des* 17: 2352-9, 2011.
253. Sala A, Campagnoli M, Perani E, Romano A, Labo S, Monzani E, Minchiotti L and Galliano M. Human alpha-1-microglobulin is covalently bound to kynurenine-derived chromophores. *J Biol Chem* 279: 51033-41, 2004.
254. Salier JP, Rouet P, Raguenez G and Daveau M. The inter-alpha-inhibitor family: from structure to regulation. *Biochem J* 315 (Pt 1): 1-9, 1996.
255. Salier JP, Simon D, Rouet P, Raguenez G, Muscatelli F, Gebhard W, Guenet JL and Mattei MG. Homologous chromosomal locations of the four genes for inter-alpha-inhibitor and pre-alpha-inhibitor family in human and mouse: assignment of the ancestral gene for the lipocalin superfamily. *Genomics* 14: 83-8, 1992.
256. Salier JP, Verga V, Doly J, Diarra-Mehrpour M and Erickson RP. The genes for the inter-alpha-inhibitor family share a homologous organization in human and mouse. *Mamm Genome* 2: 233-9, 1992.
257. Sano S, Sano T, Morishima I, Shiro Y and Maeda Y. On the mechanism of the chemical and enzymic oxygenations of alpha-oxyprotohemin IX to Fe.biliverdin IX alpha. *Proc Natl Acad Sci U S A* 83: 531-5, 1986.
258. Santin M and Cannas M. Collagen-bound alpha1-microglobulin in normal and healed tissues and its effect on immunocompetent cells. *Scand J Immunol* 50: 289-95, 1999.
259. Sazama K. Transfusion errors: scope of the problem, consequences, and solutions. *Curr Hematol Rep* 2: 518-21, 2003.
260. Schaer CA, Schoedon G, Imhof A, Kurrer MO and Schaer DJ. Constitutive endocytosis of CD163 mediates hemoglobin-heme uptake and determines the noninflammatory and protective transcriptional response of macrophages to hemoglobin. *Circ Res* 99: 943-50, 2006.
261. Schmitt TH, Frezzatti WA, Jr. and Schreier S. Hemin-induced lipid membrane disorder and increased permeability: a molecular model for the mechanism of cell lysis. *Arch Biochem Biophys* 307: 96-103, 1993.
262. Schneider H and Huch A. Dual in vitro perfusion of an isolated lobe of human placenta: method and instrumentation. *Contrib Gynecol Obstet* 13: 40-7, 1985.
263. Schrader M and Fahimi HD. Peroxisomes and oxidative stress. *Biochim Biophys Acta* 1763: 1755-66, 2006.

264. Schroeder BM. ACOG practice bulletin on diagnosing and managing preeclampsia and eclampsia. *Am Fam Physician*. 66: 330-331, 2002.
265. Sears DA. Disposal of plasma heme in normal man and patients with intravascular hemolysis. *J Clin Invest* 49: 5-14, 1970.
266. Seidah NG, Mayer G, Zaid A, Rousselet E, Nassoury N, Poirier S, Essalmani R and Prat A. The activation and physiological functions of the proprotein convertases. *Int J Biochem Cell Biol* 40: 1111-25, 2008.
267. Seon BK and Pressman D. Unique human glycoprotein, alpha1-microglycoprotein: isolation from the urine of a cancer patient and its characterization. *Biochemistry* 17: 2815-21, 1978.
268. Seve P, Philippe P, Dufour JF, Broussolle C and Michel M. Autoimmune hemolytic anemia: classification and therapeutic approaches. *Expert Rev Hematol* 1: 189-204, 2008.
269. Shatwell KP and Segal AW. NADPH oxidase. *Int J Biochem Cell Biol* 28: 1191-5, 1996.
270. Shennan AH, Poston L, Chappell LC and Seed PT. Prevention of pre-eclampsia. *Lancet* 357: 1534, 2001.
271. Shigetomi H, Onogi A, Kajiwara H, Yoshida S, Furukawa N, Haruta S, Tanase Y, Kanayama S, Noguchi T, Yamada Y, Oi H and Kobayashi H. Anti-inflammatory actions of serine protease inhibitors containing the Kunitz domain. *Inflamm Res* 59: 679-87, 2010.
272. Shiva S, Moellering D, Ramachandran A, Levenon AL, Landar A, Venkatraman A, Ceaser E, Ulasova E, Crawford JH, Brookes PS, Patel RP and Darley-Usmar VM. Redox signalling: from nitric oxide to oxidized lipids. *Biochem Soc Symp*: 107-20, 2004.
273. Smith TF, Gaitatzes C, Saxena K and Neer EJ. The WD repeat: a common architecture for diverse functions. *Trends Biochem Sci* 24: 181-5, 1999.
274. Springer TA. Folding of the N-terminal, ligand-binding region of integrin alpha-subunits into a beta-propeller domain. *Proc Natl Acad Sci U S A* 94: 65-72, 1997.
275. Stainsby D, Jones H, Asher D, Atterbury C, Boncinelli A, Brant L, Chapman CE, Davison K, Gerrard R, Gray A, Knowles S, Love EM, Milkins C, McClelland DB, Norfolk DR, Soldan K, Taylor C, Revill J, Williamson LM and Cohen H. Serious hazards of transfusion: a decade of hemovigilance in the UK. *Transfus Med Rev* 20: 273-82, 2006.
276. Steinberg MH (Ed) *Disorders of Hemoglobin: Genetics, Pathophysiology, and Clinical Management*: Cambridge University Press; 2001.
277. Stevens JM. Gynaecology from ancient Egypt: The papyrus Kahun: A translation of the oldest treatise on gynaecology that has survived from the ancient world. *Med J Aust*. 2: 949-952, 1975.
278. Stillman IE and Karumanchi SA. The glomerular injury of preeclampsia. *J Am Soc Nephrol* 18: 2281-4, 2007.
279. Stocker R, Yamamoto Y, McDonagh AF, Glazer AN and Ames BN. Bilirubin is an antioxidant of possible physiological importance. *Science* 235: 1043-6, 1987.
280. Strevens H, Wide-Svensson D, Hansen A, Horn T, Ingemarsson I, Larsen S, Willner J and Olsen S. Glomerular endotheliosis in normal pregnancy and pre-eclampsia. *Br J Obstet Gynaecol*. 110: 831-836, 2003.
281. Sundaram M, Sivaprasadarao A, Aalten DM and Findlay JB. Expression, characterization and engineered specificity of rat epididymal retinoic acid-binding protein. *Biochem J* 334 (Pt 1): 155-60, 1998.
282. Sutton HC and Winterbourn CC. On the participation of higher oxidation states of iron and copper in Fenton reactions. *Free Radic Biol Med* 6: 53-60, 1989.
283. Takagi T, Takagi K and Kawai T. Complete amino acid sequence of human alpha 1-microglobulin. *Biochem Biophys Res Commun* 98: 997-1001, 1981.
284. Talosi G, Endreffy, E., Turi, S., Nemeth, I. Molecular and Genetic Aspects of Preeclampsia: State of the Art. *Mol Genet Metab*. 71: 565-572, 2000.
285. Tam SC and Wong JT. Impairment of renal function by stroma-free hemoglobin in rats. *J Lab Clin Med* 111: 189-93, 1988.

286. Tavakkol A. Molecular cloning of porcine alpha 1-microglobulin/HI-30 reveals developmental and tissue-specific expression of two variant messenger ribonucleic acids. *Biochim Biophys Acta* 1088: 47-56, 1991.
287. Tavazzi D, Taher A and Cappellini MD. Red blood cell enzyme disorders: an overview. *Pediatr Ann* 37: 303-10, 2008.
288. Tejler L, Eriksson S, Grubb A and Astedt B. Production of protein HC by human fetal liver explants. *Biochim Biophys Acta* 542: 506-14, 1978.
289. Tejler L and Grubb AO. A complex-forming glycoprotein heterogeneous in charge and present in human plasma, urine, and cerebrospinal fluid. *Biochim Biophys Acta* 439: 82-94, 1976.
290. Tenhunen R, Marver HS and Schmid R. The enzymatic conversion of heme to bilirubin by microsomal heme oxygenase. *Proc Natl Acad Sci U S A* 61: 748-55, 1968.
291. Tenhunen R, Marver HS and Schmid R. Microsomal heme oxygenase. Characterization of the enzyme. *J Biol Chem* 244: 6388-94, 1969.
292. ter Haar E, Musacchio A, Harrison SC and Kirchhausen T. Atomic structure of clathrin: a beta propeller terminal domain joins an alpha zigzag linker. *Cell* 95: 563-73, 1998.
293. Thornton JG and Macdonald AM. Twin mothers, pregnancy hypertension and pre-eclampsia. *Br J Obstet Gynaecol.* 106: 570-575, 1999.
294. Thuveson M and Fries E. Intracellular proteolytic processing of the heavy chain of rat pre-alpha-inhibitor. The COOH-terminal propeptide is required for coupling to bikunin. *J Biol Chem* 274: 6741-6, 1999.
295. Tolosano E and Altruda F. Hemopexin: structure, function, and regulation. *DNA Cell Biol* 21: 297-306, 2002.
296. Trakshel GM, Kutty RK and Maines MD. Purification and characterization of the major constitutive form of testicular heme oxygenase. The noninducible isoform. *J Biol Chem* 261: 11131-7, 1986.
297. Treloar SA, Cooper, D.W., Brennecke, S.P., Grehan, M.M., Martin, N.G. An Australian twin study of the genetic basis of preeclampsia and eclampsia. *Am J Obstet Gynecol.* 184: 374-381, 2001.
298. Tripette J, Alexy T, Hardy-Dessources MD, Mouguel D, Beltan E, Chalabi T, Chout R, Etienne-Julan M, Hue O, Meiselman HJ and Connes P. Red blood cell aggregation, aggregate strength and oxygen transport potential of blood are abnormal in both homozygous sickle cell anemia and sickle-hemoglobin C disease. *Haematologica* 94: 1060-5, 2009.
299. Tsemakhovich VA, Bamm VV, Shaklai M and Shaklai N. Vascular damage by unstable hemoglobins: the role of heme-depleted globin. *Archives of Biochemistry and Biophysics* 436: 307-15, 2005.
300. Wagener FA, Eggert A, Boerman OC, Oyen WJ, Verhofstad A, Abraham NG, Adema G, van Kooyk Y, de Witte T and Figdor CG. Heme is a potent inducer of inflammation in mice and is counteracted by heme oxygenase. *Blood* 98: 1802-11, 2001.
301. Walter PB, Fung EB, Killilea DW, Jiang Q, Hudes M, Madden J, Porter J, Evans P, Vichinsky E and Harmatz P. Oxidative stress and inflammation in iron-overloaded patients with beta-thalassaemia or sickle cell disease. *Br J Haematol* 135: 254-63, 2006.
302. Vandegriff KD, Young MA, Lohman J, Bellelli A, Samaja M, Malavalli A and Winslow RM. CO-MP4, a polyethylene glycol-conjugated haemoglobin derivative and carbon monoxide carrier that reduces myocardial infarct size in rats. *Br J Pharmacol* 154: 1649-61, 2008.
303. Vanlair CF and Masius JR. De la microcythemie. *Bull Acad Med Belg 3e Ser* 5: 515-613, 1871.
304. Vaughn MW, Huang KT, Kuo L and Liao JC. Erythrocytes possess an intrinsic barrier to nitric oxide consumption. *J Biol Chem* 275: 2342-8, 2000.
305. Vazquez S, Garner B, Sheil MM and Truscott RJ. Characterisation of the major autoxidation products of 3-hydroxykynurenine under physiological conditions. *Free Radic Res* 32: 11-23, 2000.
306. Weisiger RA and Fridovich I. Superoxide dismutase. Organelle specificity. *J Biol Chem* 248: 3582-92, 1973.

307. Wester L, Fast J, Labuda T, Cedervall T, Wingårdh K, Olofsson T and Åkerström B. Carbohydrate groups of α_1 -microglobulin are important for secretion and tissue localization but not for immunological properties. *Glycobiology* 10: 891-900, 2000.
308. Wester L, Michaelsson E, Holmdahl R, Olofsson T and Åkerström B. Receptor for α_1 -microglobulin on T lymphocytes: inhibition of antigen-induced interleukin-2 production. *Scand J Immunol* 48: 1-7, 1998.
309. Whitelaw A, Jary S, Kmita G, Wroblewska J, Musialik-Swietlinska E, Mandera M, Hunt L, Carter M and Pople I. Randomized trial of drainage, irrigation and fibrinolytic therapy for premature infants with posthemorrhagic ventricular dilatation: developmental outcome at 2 years. *Pediatrics* 125: e852-8, 2010.
310. Whitelaw A, Thoresen M and Pople I. Posthaemorrhagic ventricular dilatation. *Arch Dis Child Fetal Neonatal Ed* 86: F72-4, 2002.
311. Wicher KB and Fries E. Haptoglobin, a hemoglobin-binding plasma protein, is present in bony fish and mammals but not in frog and chicken. *Proc Natl Acad Sci U S A* 103: 4168-73, 2006.
312. Wicher KB and Fries E. Evolutionary aspects of hemoglobin scavengers. *Antioxid Redox Signal* 12: 249-59, 2010.
313. Vichinsky E. Hemoglobin E syndromes. *Hematology Am Soc Hematol Educ Program*: 79-83, 2007.
314. Vile GF, Basu-Modak S, Waltner C and Tyrrell RM. Heme oxygenase 1 mediates an adaptive response to oxidative stress in human skin fibroblasts. *Proc Natl Acad Sci U S A* 91: 2607-10, 1994.
315. Wilkinson LS, Bunker C, Edwards JC, Scurr JH and Smith PD. Leukocytes: their role in the etiopathogenesis of skin damage in venous disease. *J Vasc Surg* 17: 669-75, 1993.
316. Wilks A. Heme oxygenase: evolution, structure, and mechanism. *Antioxid Redox Signal* 4: 603-14, 2002.
317. Williams MS and Kwon J. T cell receptor stimulation, reactive oxygen species, and cell signaling. *Free Radic Biol Med* 37: 1144-51, 2004.
318. Villoutreix BO, Åkerström B and Lindqvist A. Structural model of human α_1 -microglobulin: proposed scheme for the interaction with the Gla domain of anticoagulant protein C. *Blood Coagul Fibrinolysis* 11: 261-75, 2000.
319. Wilson-Costello D, Friedman H, Minich N, Fanaroff AA and Hack M. Improved survival rates with increased neurodevelopmental disability for extremely low birth weight infants in the 1990s. *Pediatrics* 115: 997-1003, 2005.
320. Vincent C, Bouic P and Revillard JP. Characterization of rat alpha 1 microglobulin. *Biochem Biophys Res Commun* 116: 180-8, 1983.
321. Vohr BR, Wright LL, Dusick AM, Mele L, Verter J, Steichen JJ, Simon NP, Wilson DC, Broyles S, Bauer CR, Delaney-Black V, Yolton KA, Fleisher BE, Papile LA and Kaplan MD. Neurodevelopmental and functional outcomes of extremely low birth weight infants in the National Institute of Child Health and Human Development Neonatal Research Network, 1993-1994. *Pediatrics* 105: 1216-26, 2000.
322. Volpe JJ. Perinatal brain injury: from pathogenesis to neuroprotection. *Ment Retard Dev Disabil Res Rev* 7: 56-64, 2001.
323. Wu Y, Zhao XD, Zhuang Z, Xue YJ, Cheng HL, Yin HX and Shi JX. Peroxisome proliferator-activated receptor gamma agonist rosiglitazone attenuates oxyhemoglobin-induced Toll-like receptor 4 expression in vascular smooth muscle cells. *Brain Res* 1322: 102-8, 2010.
324. Yoshida T, Takahashi S and Kikuchi G. Partial purification and reconstitution of the heme oxygenase system from pig spleen microsomes. *J Biochem* 75: 1187-91, 1974.
325. Yu B, Bloch KD and Zapol WM. Hemoglobin-based red blood cell substitutes and nitric oxide. *Trends Cardiovasc Med* 19: 103-7, 2009.
326. Zamboni P, Izzo M, Tognazzo S, Carandina S, De Palma M, Catozzi L, Caggiati A, Scapoli G and Gemmati D. The overlapping of local iron overload and HFE mutation in venous leg ulcer pathogenesis. *Free Radic Biol Med* 40: 1869-73, 2006.

327. Zamboni P, Scapoli G, Lanzara V, Izzo M, Fortini P, Legnaro R, Palazzo A, Tognazzo S and Gemmati D. Serum iron and matrix metalloproteinase-9 variations in limbs affected by chronic venous disease and venous leg ulcers. *Dermatol Surg* 31: 644-9; discussion 649, 2005.
328. Zamboni P, Tognazzo S, Izzo M, Pancaldi F, Scapoli GL, Liboni A and Gemmati D. Hemochromatosis C282Y gene mutation increases the risk of venous leg ulceration. *J Vasc Surg* 42: 309-14, 2005.
329. Zhang N and Palmer AF. Polymerization of human hemoglobin using the crosslinker 1,11-bis(maleimido)triethylene glycol for use as an oxygen carrier. *Biotechnol Prog* 26: 1481-5, 2010.
330. Zhou H, Beevers CS and Huang S. The targets of curcumin. *Curr Drug Targets* 12: 332-47, 2011.
331. Zhou ML, Shi JX, Hang CH, Zhang FF, Gao J and Yin HX. Expression of Toll-like receptor 4 in the brain in a rabbit experimental subarachnoid haemorrhage model. *Inflamm Res* 56: 93-7, 2007.
332. Zhou Z and Kang YJ. Cellular and subcellular localization of catalase in the heart of transgenic mice. *J Histochem Cytochem* 48: 585-94, 2000.
333. Åkerström B. Tissue distribution of guinea pig α_1 -microglobulin. *Cell Mol Biol* 29: 489-95, 1983.
334. Åkerström B. Immunological analysis of α_1 -microglobulin in different mammalian and chicken serum. α_1 -Microglobulin is 5-8 kilodaltons larger in primates. *J Biol Chem* 260: 4839-44, 1985.
335. Åkerström B and Berggård I. Guinea-pig alpha 1 microglobulin. Isolation and properties in comparison with human α_1 - microglobulin. *Eur J Biochem* 101: 215-23, 1979.
336. Åkerström B, Bratt T and Enghild JJ. Formation of the α_1 -microglobulin chromophore in mammalian and insect cells: a novel post-translational mechanism? *FEBS Lett* 362: 50-4, 1995.
337. Åkerström B, Flower DR and Salier JP. Lipocalins: unity in diversity. *Biochim Biophys Acta* 1482: 1-8, 2000.
338. Åkerström B and Lögdberg L. An intriguing member of the lipocalin protein family: α_1 -microglobulin. *Trends Biochem Sci* 15: 240-3, 1990.
339. Åkerström B and Lögdberg L. α_1 -microglobulin. In: *In: Lipocalins*. edited by Åkerström B, Borregaard N, Flower DR and Salier J-P. Georgetown, TX, USA: Landes Bioscience; 2006. pp. 110-20.
340. Åkerström B, Lögdberg L, Babiker-Mohamed H, Lohmander S and Rask L. Structural relationship between α_1 -microglobulin from man, guinea-pig, rat and rabbit. *Eur J Biochem* 170: 143-8, 1987.
341. Åkerström B, Maghzal GJ, Winterbourn CC and Kettle AJ. The lipocalin α_1 -microglobulin has radical scavenging activity. *J Biol Chem* 282: 31493-503, 2007.