



LUND UNIVERSITY

Can post-sternotomy mediastinitis be prevented by a closed incision management system?

Dohmen, Pascal M; Markou, Thanasie; Ingemansson, Richard; Rotering, Heinrich; Hartman, Jean M; van Valen, René; Brunott, Maaike; Kramer, Axel; Segers, Patrique

Published in:
GMS hygiene and infection control

DOI:
[10.3205/dgkh000239](https://doi.org/10.3205/dgkh000239)

2014

[Link to publication](#)

Citation for published version (APA):

Dohmen, P. M., Markou, T., Ingemansson, R., Rotering, H., Hartman, J. M., van Valen, R., Brunott, M., Kramer, A., & Segers, P. (2014). Can post-sternotomy mediastinitis be prevented by a closed incision management system? *GMS hygiene and infection control*, 9(3), Doc19. <https://doi.org/10.3205/dgkh000239>

Total number of authors:
9

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Can post-sternotomy mediastinitis be prevented by a closed incision management system?

Kann Mediastinitis bei Zustand nach Sternotomie durch einen Vakuumverband verhindert werden?

Abstract

Post-sternotomy mediastinitis is a serious complication after cardiothoracic surgery and contribute significantly to post-operative morbidity, mortality, and healthcare costs. Negative pressure wound therapy is today's golden standard for post-sternotomy mediastinitis treatment. A systematic literature search was conducted at PubMed until October 2012 to analyse whether vacuum-assisted closure technique prevents mediastinitis after clean surgical incisions closure. Today's studies showed reduction of post-sternotomy mediastinitis including a beneficial socio-economic impact. Current studies, however included only high-risk patients, hence furthermore, larger randomised controlled trials are warranted to clarify the benefit for using surgical incision vacuum management systems in the general patient population undergoing sternotomy and clarify risk factor interaction.

Keywords: incisional negative pressure therapy, cardiothoracic surgery, surgical wounds, surgical site infection, closed incision management, negative pressure wound therapy (NPWT)

Zusammenfassung

Mediastinitis bei Zustand nach Sternotomie ist eine ernste Komplikation nach herz- und thoraxchirurgischen Eingriffen, die durch eine Vielzahl von Faktoren beeinflusst wird. Sie ist mit erhöhter Morbidität, Mortalität und behandlungsassoziierten Kosten verbunden.

Die Vakuumtherapie ist der heutige Standard zur Mediastinitisbehandlung bei Zustand nach Sternotomie. Im vorliegenden Minireview wird analysiert, ob der Vakuumverband auch in die Lage ist, einer Mediastinitis vorzubeugen.

Im Ergebnis einer systematischen Literaturrecherche in PubMed wurde festgestellt, dass die Anwendung des Vakuum-assoziierten Verschlusses eine Reduktion der Mediastinitis zur Folge hatte. Ebenso konnten die behandlungsassoziierten Kosten gesenkt werden. Allerdings waren in allen Studien nur Hoch-Risiko-Patienten eingeschlossen, sodass dringend weitere Studien zur Absicherung der Ergebnisse einschließlich der Abklärung von Risikofaktoren benötigt werden.

Pascal M. Dohmen¹

Thanasie Markou²

Richard Ingemansson³

Heinrich Rotering⁴

Jean M. Hartman⁵

René van Valen⁶

Maaike Brunott⁶

Axel Kramer⁷

Patrique Segers⁸

1 Department of Cardiac Surgery, Heart Center Leipzig, University of Leipzig, Germany

2 Department of Cardiothoracic Surgery, Isala Klinieken Zwolle, The Netherlands

3 Department of Cardiothoracic Surgery, University Hospital of Lund, Sweden

4 Department of Cardiothoracic Surgery, University Clinic Münster, Germany

5 Department of Cardiothoracic Surgery, University Medical Center Groningen, The Netherlands

6 Department of Cardiothoracic Surgery, Erasmus University Rotterdam, The Netherlands

7 Institute of Hygiene and Environmental Medicine, University Medicine Greifswald, Germany

8 Department of Cardiothoracic Surgery, Academic Medical Center Amsterdam, The Netherlands

Introduction

Post-sternotomy mediastinitis is a serious complication after cardiothoracic surgery and contribute significantly to post-operative morbidity, mortality, and healthcare costs [1]. Negative pressure wound therapy is today's golden standard for post-sternotomy mediastinitis treatment due to oedema reduction, removing exudation, increase wound perfusion, stimulate granulation formation, and decrease microbial colonization [1]. This positive effect on wound healing triggered the interest in using negative pressure wound therapy after closure of clean surgical incisions in contrast with traditional wound care such as gauze dressings, hydrocolloids, growth factors or cultured skin to prevent surgical site infections, especially post-sternotomy mediastinitis. The surgical incision management system (Prevena™ Incision Management System, Kinetic Concepts, Inc. USA, San Antonio, TX, USA) is a single-use, battery-powered therapy unit that delivers negative pressure of -125 mmHg [2]. Surgical incision management holds the incision edges together, reduces lateral tension and oedema, stimulates perfusion, and protects the surgical site from external infectious sources [3]. The skin interface layer containing 0.019% ionic silver, allows no direct contact of the foam, wicks fluid from the skin surface, and reduces bacterial colonisation within the fabric.

High-quality, multi-centre or single-centre, randomised controlled trials (level I based on the Evidence Rating Scale for Therapeutic Studies developed by the American Society of Plastic Surgeons) [4] in other clean surgery fields [5] showed positive outcomes by using incisional negative pressure wound therapy. These results encourage using surgical incision management, which is functionally equivalent to incisional negative pressure wound therapy, in high-risk patients undergoing median sternotomy to prevent post-sternotomy mediastinitis. Therefore a systematic of the literature was performed.

Method

The systematic literature search at PubMed was conducted through October 30th 2012. The following key words were included: "negative pressure wound therapy" and "sternotomy wound infection prevention" and "cardiac surgery". Excluded were case reports and articles not preventing surgical site infections but treated.

Results

The largest prospective comparative study (Level II) was performed by Grauhan et al. [6] including 150 consecutive obese (body mass index >30 kg/m²) patients with median sternotomy. Standard wound dressings were applied in the control group (n=75), while the treatment group received surgical incision management (n=75). This study showed significant reduction of surgical site infections

(SSI) after median sternotomy, respectively 16% versus 4% (OR 4.57, CI 95% 1.23-16.94; p=0.0266).

Atkins et al. [7] examined 57 adult cardiac surgery patients at higher risk for sternal wound infection, who were treated with incisional negative pressure wound therapy (Level III). Patient population included morbid obesity (77.2%), diabetics (54.4%), and obese plus diabetic (50.9%). Overall, 50.9% of patients underwent coronary artery bypass graft with one internal mammary artery and 14% with bilateral mammary artery use. Approximately 20% underwent coronary artery bypass graft with concomitant procedures. Since this study included no control group the estimated risk for post-sternotomy mediastinitis was based on risk scores, which predict SSI. Based on this system, the estimated average risk for developing post-operative post-sternotomy mediastinitis in this group of high-risk cardiac surgery patients was 6.1 ± 4.0%; therefore, at least three cases of post-sternotomy mediastinitis were expected in this study population. Ten patients (17.5%) required readmission within the first 30 days after discharge; however, no admissions were due to sternal wound complications. Therefore the authors recommend that incision negative pressure wound therapy should be strongly considered for patients with increased risk of surgical site infection.

Finally Colli et al. [8] used surgical incision management over the surgical incisions of a small case series of ten patients at high risk for post-sternotomy mediastinitis following coronary artery bypass graft surgery (Level III). Surgical incision management was used for 5 continuous days immediately following sternal wound closure. This study also included no control group, and therefore the authors again utilised a risk score system finding a prediction of 6.4 ± 4.4% for post-sternotomy mediastinitis. This high-risk cardiac surgery population included, diabetes in all, peripheral vessel disease in 90%, morbid obesity in 50%, chronic obstructive pulmonary disease in 30%, and renal failure in 20%. The left internal mammary artery was used in 100% and bilateral mammary artery grafting was performed in 50%. The system was well tolerated and all patients experienced complete wound healing with no evidence of early or late wound infections. The authors again recommend the use of surgical incision management in high-risk cardiac patients. These preliminary findings demonstrate the favourable efficacy and safety of surgical incision management systems in preventing wound complications after cardiac surgery in high-risk patients.

Discussion

The remarkable infection-preventive effect raises the question of the possible causes. The bacterial density in wounds predicts the risk of wound infection with subsequent healing [9], [10], [11]. Therefore, it was obvious that negative pressure affects the bacterial colonization. But some studies on acute and chronic wounds refutes that bacterial bioburden is consistently lessened during

VAC therapy [12], [13]. In an in vitro wound model bacterial load of sponges with or without negative pressure did not differ [14]. The reduction in bacteria demonstrated in previous studies appears to be caused by other effects than physical suction alone. Since primary closed sternotomy wounds are not critically colonized, other mechanisms must be responsible. From a theoretical considerations point of view, following factors could affect a favourable outcome: (i) improved circulation and increased of vascularity [15], (ii) stimulation of cell proliferation [15], (iii) immediate tight aseptic wound closure, (iv) microbio-static activity of polyurethane-coated polyester fabric with silver and (v) optimal wound edges adaptation. Using a computer model, the hypothesis was generated, that micromechanical forces may be stimulate wound healing through promotion of cell division, angiogenesis, and local elaboration of growth factors [16]. These questions should be analyzed in further studies i.e. in sensitive animal models.

Conclusion

All studies showed reduction of expected post-sternotomy mediastinitis in high-risk cardiac surgery patients for SSI such as morbid obesity, including obesitas per magna, insulin-dependent diabetes, chronic renal failure, and bilateral mammary artery grafting. Other risk-factors including low body mass index ($<18 \text{ kg/m}^2$), long-term immunosuppressive therapy, high age and female gender, need to be investigated. Furthermore, larger randomised controlled trials are warranted to clarify exact benefit for the use of a surgical incision management system. These studies should not only include reduction of surgical site infection, including patient's morbidity and mortality, but also investigate the social economic impact. Due to the complexity of SSI and multiple factors influencing post-sternotomy mediastinitis, additional studies are needed to improve wound healing in cardiac surgery patients bringing prevention measures in proportion to risk factors.

Notes

Acknowledgements

The authors would like to thank Alice Goodwin for editorial assistance.

Conflict of interest

All authors declare that they have no conflicts of interest.

Role of the funding source

Supporting grant provided by KCI Europe Holding B.V. All authors had final responsibility for the decision to submit the article for publication.

References

1. Ennker IC, Pietrowski D, Vöhringer L, Kojcici B, Albert A, Vogt PM, Ennker J. Surgical debridement, vacuum therapy and pectoralis plasty in poststernotomy mediastinitis. *J Plast Reconstr Aesthet Surg.* 2009 Nov;62(11):1479-83. DOI: 10.1016/j.bjps.2008.05.017
2. Dohmen PM, Misfeld M, Borger MA, Mohr FW. Closed incision management with negative pressure wound therapy. *Expert Rev Med Devices.* 2014 Jul;11(4):395-402. DOI: 10.1586/17434440.2014.911081
3. Morykwas MJ, Argenta LC, Shelton-Brown EI, McGuirt W. Vacuum-assisted closure: a new method for wound control and treatment: animal studies and basic foundation. *Ann Plast Surg.* 1997 Jun;38(6):553-62. DOI: 10.1097/00000637-199706000-00001
4. Sullivan D, Chung KC, Eaves FF 3rd, Rohrich RJ. The level of evidence pyramid: indicating levels of evidence in Plastic and Reconstructive Surgery articles. *Plast Reconstr Surg.* 2011 Jul;128(1):311-4. DOI: 10.1097/PRS.0b013e3182195826
5. Stannard JP, Volgas DA, McGwin G 3rd, Stewart RL, Obremskey W, Moore T, Anglen JO. Incisional negative pressure wound therapy after high-risk lower extremity fractures. *J Orthop Trauma.* 2012 Jan;26(1):37-42. DOI: 10.1097/BOT.0b013e318216b1e5
6. Grauhan O, Navasardyan A, Hofmann M, Müller P, Stein J, Hetzer R. Prevention of poststernotomy wound infections in obese patients by negative pressure wound therapy. *J Thorac Cardiovasc Surg.* 2013 May;145(5):1387-92. DOI: 10.1016/j.jtcvs.2012.09.040
7. Atkins BZ, Wooten MK, Kistler J, Hurley K, Hughes GC, Wolfe WG. Does negative pressure wound therapy have a role in preventing poststernotomy wound complications? *Surg Innov.* 2009 Jun;16(2):140-6. DOI: 10.1177/1553350609334821
8. Colli A, Camara ML. First experience with a new negative pressure incision management system on surgical incisions after cardiac surgery in high risk patients. *J Cardiothorac Surg.* 2011;6:160. DOI: 10.1186/1749-8090-6-160
9. Elek SD. Experimental staphylococcal infections in the skin of man. *Ann N Y Acad Sci.* 1956 Aug;65(3):85-90. DOI: 10.1111/j.1749-6632.1956.tb36626.x
10. Liedberg NC, Reiss E, Artz CP. The effect of bacteria on the take of split-thickness skin grafts in rabbits. *Ann Surg.* 1955 Jul;142(1):92-6. DOI: 10.1097/00000658-195507000-00011
11. Krizek TJ, Robson MC. Evolution of quantitative bacteriology in wound management. *Am J Surg.* 1975 Nov;130(5):579-84. DOI: 10.1016/0002-9610(75)90516-4
12. Weed T, Ratliff C, Drake DB. Quantifying bacterial bioburden during negative pressure wound therapy: does the wound VAC enhance bacterial clearance?. *Ann Plast Surg.* 2004 Mar;52(3):276-9; discussion 279-80. DOI: 10.1097/01.sap.0000111861.75927.4d
13. Mouës CM, Vos MC, van den Bemd GJ, Stijnen T, Hovius SE. Bacterial load in relation to vacuum-assisted closure wound therapy: a prospective randomized trial. *Wound Repair Regen.* 2004 Jan-Feb;12(1):11-7. DOI: 10.1111/j.1067-1927.2004.12105.x
14. Assadian O, Assadian A, Stadler M, Diab-Elschahawi M, Kramer A. Bacterial growth kinetic without the influence of the immune system using vacuum-assisted closure dressing with and without negative pressure in an in vitro wound model. *Int Wound J.* 2010 Aug;7(4):283-9. DOI: 10.1111/j.1742-481X.2010.00686.x
15. Scherer SS, Pietramaggiori G, Mathews JC, Prsa MJ, Huang S, Orgill DP. The mechanism of action of the vacuum-assisted closure device. *Plast Reconstr Surg.* 2008 Sep;122(3):786-97. DOI: 10.1097/PRS.0b013e31818237ac

16. Saxena V, Hwang CW, Huang S, Eichbaum Q, Ingber D, Orgill DP. Vacuum-assisted closure: microdeformations of wounds and cell proliferation. *Plast Reconstr Surg.* 2004 Oct;114(5):1086-96; discussion 1097-8. DOI: 10.1097/01.PRS.0000135330.51408.97

Please cite as

Dohmen PM, Markou T, Ingemansson R, Rotering H, Hartman JM, van Valen R, Brunott M, Kramer A, Segers P. Can post-sternotomy mediastinitis be prevented by a closed incision management system? *GMS Hyg Infect Control.* 2014;9(3):Doc19. DOI: 10.3205/dgkh000239, URN: urn:nbn:de:0183-dgkh0002396

Corresponding author:

Prof. Dr. Pascal M. Dohmen

Department of Cardiovascular Surgery, Charité Hospital,
Medical University Berlin, Chariteplatz 1, 10117 Berlin,
Germany, Phone: +49 30 450 522092, Fax: +49 30 450
522921
pascal.dohmen@yahoo.de

This article is freely available from

<http://www.egms.de/en/journals/dgkh/2014-9/dgkh000239.shtml>

Published: 2014-09-30

Copyright

©2014 Dohmen et al. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by-nc-nd/3.0/deed.en>). You are free: to Share – to copy, distribute and transmit the work, provided the original author and source are credited.