

LUND UNIVERSITY

Pedon-scale silicate weathering : comparison of the PROFILE model and the depletion method at 16 forest sites in Sweden

Stendahl, Johan; Akselsson, Cecilia; Melkerud, Per-Arne; Belyazid, Salim

Published in: Geoderma

DOI: 10.1016/j.geoderma.2013.07.005

2013

Document Version: Peer reviewed version (aka post-print)

Link to publication

Citation for published version (APA):

Stendahl, J., Akselsson, C., Melkerud, P.-A., & Belyazid, S. (2013). Pedon-scale silicate weathering : comparison of the PROFILE model and the depletion method at 16 forest sites in Sweden. Geoderma, 211-212, 65-74. https://doi.org/10.1016/j.geoderma.2013.07.005

Total number of authors: 4

General rights

Unless other specific re-use rights are stated the following general rights apply:

- Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the
- legal requirements associated with these rights

· Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

You may not further distribute the material or use it for any profit-making activity or commercial gain
 You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: https://creativecommons.org/licenses/

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117 221 00 Lund +46 46-222 00 00

Pedon-scale silicate weathering: comparison of the PROFILE model and the depletion method at 16 forest sites in Sweden

- 3 Stendahl, Johan^{a,*}, Akselsson, Cecilia^b, Melkerud Per-Arne^a, Belyazid, Salim^c
- 4 5

1

2

- ^a Department of Soil and Environment, Swedish University of Agricultural Sciences, Sweden
- ^b Department of Earth and Ecosystem Sciences, Lund University, Sweden
- 7 ^c Belyazid Consulting and Communication
- 8
- 9 * Corresponding author. Address: Department of Soil and Environment, SLU, Box 7014, S-750 07 Uppsala,
 10 Sweden. Phone: +46 18 673801
- 11 E-mail addresses: johan.stendahl@slu.se (J. Stendahl), cecilia.akselsson@nateko.lu.se (C. Akselsson), per-
- 12 arne.melkerud@slu.se (P.-A. Melkerud), salim@belyazid.com (S. Belyazid)
- 13

14 Abstract

15 Weathering of soil minerals is important for the recovery from acidification and for the

- 16 sustainability of forestry. However, there is still substantial uncertainty about its absolute rate.
- 17 This study presents a harmonized comparison of field weathering rates estimated with the
- 18 mechanistic model PROFILE and the depletion method for 16 intensively sampled soil
- 19 profiles across Sweden representing different site conditions. In general, a correspondence in
- 20 total weathering rates was found between the two methods except in rare cases where either
- 21 method yielded deviating results. The weathering rate was higher according to the depletion
- 22 method than according to PROFILE for Mg, while PROFILE produced higher weathering
- 23 rates for the other base cations. The Spearman rank correlation (ρ) between the two methods
- indicated significant correlation for Ca (ρ =0.44, p=0.04) and non-significant correlation for Mg (ρ =0.51, p=0.09), Na (ρ =0.25, p=0.34), K (ρ =0.07, p=0.80), and the sum of the base
- 26 cations (p=0.11, p=0.67). The variation in weathering rates with depth showed opposite
- 27 gradients in the upper 50 cm, which reflects the conceptual differences between the methods.
- 28 This study shows the potential of using multiple methods to identify a probable weathering
- 29 rate, if harmonized input data is used. Furthermore, it highlights the importance of making
- 30 comparisons for individual elements in order to interpret differences between methods.
- 31 Regardless of the method used, weathering rates were below or at the same level as the losses
- 32 caused by whole-tree harvesting, particularly in southern Sweden, indicating a risk of
- 33 negative effects on soils and waters.

34

35 **1. Introduction**

- 36 Chemical weathering is important for the nutrient sustainability of forests, the neutralization
- 37 of acidifying compounds and the quality of water transported from upland soils to surface
- 38 water systems downstream. The historical deposition of acid compounds, which culminated
- around 1980 (Schöpp et al., 2003), caused substantial leaching of base cations from soils to
- 40 waters, leading to base cation depletion and acidification of soils. A reduction of acid
- 41 deposition to levels below the total of base cation weathering and base cation deposition is a
- 42 prerequisite for the recovery of soils from acidification. However, recovery will also depend
- 43 on the export of nutrients from the forest ecosystem through harvesting, which has increased
- 44 recently owing to the increased focus on using harvest residues for bioenergy, e.g. whole-tree
- 45 harvesting. While active measures, such as ash recycling, are recommended in e.g. Sweden
- 46 (Swedish Forest Agency, 2008) a basis for such activities is an assessment of the mass
- 47 balance of input and output of mineral nutrients in the ecosystem. Thus, there is a demand for
- 48 robust estimates of the release rate of mineral nutrients by weathering for predictions of future
- 49 recovery from acidification and for optimizing forest management policies.

50 Proposed methods for determining weathering rates include the mass balance approach using

51 experimental data (Bain et al., 1994), the mass balance approach using catchment modelling

52 (Cosby et al., 2001), the depletion method using an immobile element as an internal standard

- 53 (Brimhall & Dietrich, 1987; Olsson & Melkerud, 1989), and the process oriented model
- 54 PROFILE where weathering rates are estimated for soil pedons (Sverdrup & Warfvinge,
- 55 1993). The various methods are conceptually different and are intended for different spatial
- scales; from the soil pedon at site level to the entire soil deposit at the catchment level.
- 57 Furthermore, the estimates consider different time perspectives; from long-term historical
- 58 averages to present day weathering and steady state weathering.
- 59 Several attempts have been made to quantify weathering rate uncertainties through an
- 60 ensemble approach (Kolka et al., 1996; Starr et al. 1998; Sverdrup et al., 1998; Whitfield et al.
- 61 2006; Koseva et al., 2010; Klaminder et al., 2011; Futter et al., 2012). Sverdrup et al. (1998)
- 62 compared rates from six different approaches in two catchments in Gårdsjön, Sweden. The
- variation in base cation weathering rates (Ca, Mg, Na and K) was 30-39 and 47-81 meq m^{-2} y⁻
- ⁶⁴ ¹ in the two catchments respectively, indicating relatively low uncertainties. On the other
- hand, Klaminder et al. (2011) demonstrated a wide span for Ca and K weathering rates, 7-150
- $meq m^{-2} y^{-1}$ and 2-32 meq m⁻² y⁻¹ respectively, at a site in northern Sweden where ten studies
- 67 were compared, and concluded that estimated weathering rates were too uncertain to be used
- 68 in assessments of sustainable harvesting. They revised their conclusions slightly in Futter et
- al. (2012), where they pointed out that much of the variation was related to input data and
- 70 known conceptual differences between the methods, for example that different estimations
- 71 were valid for different soil depths. They concluded that at least three independent estimates
- should be used when making management decisions. This is in line with the conclusions by
- 73 Whitfield et al. (2006) who compared weathering estimates from five approaches in five

- 74 catchments in Canada. The three soil profile-based approaches gave similar results, with low
- 75 weathering rates ranging from 3 to 13 meq $m^{-2} y^{-1}$. However, the two catchment-based
- 76 methods gave one order of magnitude higher rates. Koseva et al. (2010) compared PROFILE
- 77 weathering estimated with catchment mass balance calculations for 19 sites. The PROFILE
- 78 weathering rates were in most cases within the range of the catchment mass balance
- weathering rates, but PROFILE weathering rates were generally lower, as expected whencomparing soil pedon weathering rates with catchment weathering rates. Although they did
- comparing soil pedon weathering rates with catchment weathering rates. Although they did
 not present any direct uncertainty measures based on this, they used it to demonstrate the
- 81 not present any direct uncertainty measures based on this, they used it to demonstrate the 82 reliability of PROFILE. A critical aspect when comparing weathering estimates is the quality
- 83 of the input data, boundary conditions and assumptions made, e.g. the maximum soil depth
- 84 considered and corrections for coarse fragments and organic matter. The stone and boulder
- 85 content may amount to >50% vol in many forest soils (Stendahl et al., 2009) and the way it is
- 86 treated will have a strong influence on estimated pedon-scale weathering rates. It is crucial
- 87 that comparisons of weathering estimation methods are harmonized with regard to
- assumptions and input data, and they should be carefully evaluated when published results
- 89 from different studies are synthesized.
- 90 In this study we estimated the base cation weathering at the pedon-scale by the depletion
- 91 method and PROFILE for 16 intensively sampled soil profiles representing a wide range of
- 92 soil conditions in Sweden. Both methods have been frequently used to estimate weathering
- 93 rates on local, regional and national scale for assessments of sustainable harvesting (Olsson et
- 94 al., 1993; Sverdrup & Rosén, 1998; Akselsson et al., 2007). The depletion method quantifies
- 95 the loss of mobile elements since the parent material was deposited, whereas PROFILE
- 96 estimates the release of elements due to the dissolution of soil minerals at steady state. Hence,
- 97 the two methods consider very different time perspectives. Despite the different time
- 98 perspectives of the methods the comparison was justified by the relatively young age of the
- 99 profiles $(10\ 000 16\ 000\ years)$. The methods were applied in a harmonized way using input
- 100 data from the same forest sites and the same pits, as well as using the same assumptions.
- Furthermore, the estimated weathering rates were compared with estimated base cation losses at whole-tree harvesting on a selection of the sites, in the same way as in e.g. Olsson et al.
- 103 (1993) and Klaminder et al. (2011). This simplified mass balance calculation puts the
- 104 difference in weathering rates between the two methods in a sustainability perspective. The
- 105 objectives of this study were to: (i) compare how the two methods ranked the sites with
- 106 regards to Ca, Mg, K, and Na weathering, (ii) compare how weathering intensity vary with
- 107 depth in the uppermost soil profile for the two methods, (iii) investigate the causes for
- 108 deviating results between the two methods, and (iv) put the results in a sustainability
- 109 perspective by comparing the estimated weathering rates with base cation losses associated
- 110 with whole-tree harvesting.
- 111 The comparison can shed light on how the total weathering rate and weathering intensity
- 112 within the soil profile evolve over time and can be seen as a robustness test of the two
- 113 methods.

114

115 **2. Material and Methods**

116 2.1. Sites and soil profile data

117 The 16 sites, which are part of the NORDSOIL database (Raulund-Rasmussen & Callesen, 1999), are located at latitudes 56-68 °N across Sweden on podzolised glacial till without 118 119 stratigraphic layering (Fig. 1, Table 1). All sites are located above the highest Quaternary 120 shoreline and it was assumed that no redistribution of soil material has occurred since 121 deglaciation. The parent material is of granitic composition with varying mineralogical 122 composition. Among the sites 12 were used previously in weathering studies by Olsson & 123 Melkerud (1989, 1991, 2000) and Olsson et al. (1993). For each site, a profile was used that had complete soil chemical and physical data necessary to apply the two methods. Sampling 124 125 was made at 10 cm depth intervals from the top of the mineral soil to the maximum mineral 126 soil depth (42-263 cm, Table 1). Volumetric samples were taken by a core sampler for each 127 10 cm layer in the A, B and uppermost C horizon and bulk density was determined after 128 drying and weighing. Grain size distribution data was determined for 10 cm layers, although it 129 was missing for the A horizons and ca 25% of the other layers. In those cases the grain size 130 distribution was estimated based on the layers above and below. For each site the stone 131 content was estimated based on the soil texture class to 10-40% vol. An analysis of the total 132 geochemical content of the bulk soil material <2 mm was made for the samples from each 10 133 cm layer throughout the profile by extracting a representative subsample (20 g) that was 134 homogenized by dry grinding in an agate mortar after which 0.125 g was fused with 0.375 g lithium metaborate (LiBO₂) at 1000 °C and dissolved in nitric acid. The content of major 135 136 elements and trace elements was quantified by plasma emission spectrometry analysis (ICP-

AES). The analytical data were normalized with respect to dry matter weight.

138

139 2.2. The depletion method

140 The basic principle of the method is the long-term depletion of mobile (weatherable) elements 141 and the concurrent enrichment of immobile (inert) elements in weathered soil layers during soil development. The concentration of the mobile and the immobile element in a weathered 142 143 and an unweathered soil layer are utilized to estimate the loss of the mobile element in the 144 weathered layer. The idea to use immobile elements to estimate element losses was first proposed by Marshall & Haseman (1942) and the theoretical framework was formalized by 145 146 Brimhall & Dietrich (1987) and Brimhall et al. (1991). Their framework focuses on the 147 relative mass loss (or gain) of elements from layers in the original soil pedon and they define 148 the fractional mass loss as:

149
$$\tau_{j,w} = \frac{C_{j,w}C_{i,p}}{C_{j,p}C_{i,w}} - 1$$
(1)

- 150 where C denote concentrations (%), *i* denotes the immobile element, *j* denotes the mobile
- 151 element, *w* denotes a weathered horizon, and *p* denotes the unweathered parent material.
- 152 Values of $\tau < 0$ indicates net mass loss, $\tau > 0$ net mass gain, and $\tau = 0$ no change.

153 The weathered amount of the mobile element from a soil layer $(g m^{-2})$ can be expressed based 154 on the same principles (Olsson & Melkerud 1989):

155
$$W_{j,w} = \frac{d_w \rho_w}{100} \left(C_{j,p} \left(\frac{C_{i,w}}{C_{i,p}} \right) - C_{j,w} \right)$$
 (2)

156 where d is layer depth (m), and ρ is bulk density (g m⁻³).

157 Zirconium, which is predominantly found in the weathering resistant mineral zircon (ZrSiO₄),

158 was used in this study as the immobile element, since its redistribution within the profile is

159 negligible (Hodson, 2002). The depletion method assumes that there is no weathering of the

160 immobile element, that there is no weathering below a certain soil depth, and that the soil

161 profile was developed from homogenous regolith. One critical aspect of the method is to

162 define a reference soil layer to represent the unweathered parent material.

163 In this study we used equation (2) to estimate the loss of mobile base cations. The reference

soil layer used (Table 2) was the uppermost layer that met the following criteria: (1) it was

165 located in the soil C horizon, (2) the ratio between the mobile element and Zr was stable

below this layer, and (3) the Zr gradient was stable below this layer. The long-term

167 weathering rate was calculated by dividing the total weathered amount by the soil age, i.e. the

age since deglaciation given by the National Atlas of Sweden (Fredén, 2009; Table 1).

169

170 2.3. The PROFILE model and input data preparation

PROFILE is a steady state soil chemistry model, originally developed to calculate the effect 171 172 of acid rain on soil chemistry (Sverdrup & Warfvinge, 1993). PROFILE includes process 173 oriented descriptions for chemical weathering of minerals, leaching and accumulation of 174 dissolved chemical components and solution equilibrium reactions. The weathering module is 175 a central part of the model. Weathering rates are calculated using transition state theory, based 176 on the geochemical properties of the soil system, climate, deposition and tree uptake. Soil 177 physical properties, such as soil mineralogy, soil moisture content, bulk density and exposed 178 mineral surface area, have been identified as the input parameters with the highest influence on 179 the output in several studies (Jönsson et al., 1995; Hodson et al., 1996; Zak et al., 1997).

- 180 In PROFILE, the soil is divided into soil layers with different properties, preferably based on
- 181 the naturally occurring soil horizons. Soil chemistry and weathering rates are calculated for
- 182 each layer, and the soil chemistry of each layer is affected by the layer above, through vertical
- 183 water flow.
- 184 The estimated soil chemistry and weathering rates describe the situation at steady state, i.e.
- 185 the soil chemistry and weathering rates that will eventually evolve, with the deposition, tree
- 186 uptake, climate conditions and soil properties that are given as input to the model (Sverdrup &
- 187 Warfvinge, 1993). Thus, the model ignores development over time, e.g. changes in soil
- 188 properties due to weathering.
- 189 Detailed soil data was available for the chosen profiles from the NORDSOIL database and
- 190 was used in the input data preparation for the PROFILE model runs. Total elemental content
- 191 for the different soil layers was used, along with information about predominant minerals and
- 192 their compositions compiled in previous studies (Warfvinge & Sverdrup, 1995; Akselsson et
- al., 2004), for calculating soil mineralogy with the A2M model (Posch & Kurz, 2007). The
- 194 modeled mineralogical composition for the sites is given in Table 3. The specific surface area
- 195 of minerals, i.e. the weatherable area, was calculated using an empirical algorithm (Warfvinge
- 196 & Sverdrup, 1995; Holmqvist et al., 2002) using data on the grain size distribution and bulk
- 197 densities for each layer (data from 50 cm given in Table 1). Data on gravel content were only
- available for a few of the sites, for the rest of the sites the gravel content was estimated based
- 199 on grain size distribution (Table 1) and sieving curves (SGU, 1994). The estimated specific
- surface area was corrected for the stone volume in the soil by subtracting the stone volumefrom the soil volume, assuming that the weathering from stones is negligible due to the low
- 202 specific surface area as compared with fine soil. Soil moisture class was given for each site
- 203 according to a classification system with seven classes (Table 1). Soil moisture was allocated
- to the soil moisture classes, according to methods in Warfvinge & Sverdrup (1995).
- 205 Concentration of DOC, CO₂ pressure and Al solubility constants for different layers were
- 206 derived from Martinson et al. (2003).
- 207 Information on tree species and mean annual production from the NORDSOIL database
- 208 (Table 1) was used for estimating net uptake according to methods described in Akselsson et
- al. (2007), using base cation concentrations in stem biomass compiled in Akselsson (2005).
- 210 Mean annual temperature and precipitation was derived from the NORDSOIL database (Table
- 211 1), whereas runoff was derived from national maps from the Swedish Meteorological and
- 212 Hydrological Institute (SMHI) (Raab & Vedin, 1995). Deposition values were derived from
- 213 modeled deposition values for 1998 by the MATCH model (Langner et al., 1996).
- 214
- 215 2.4. Comparison with base cation losses at whole-tree harvesting
- 216 The base cation losses were estimated for stem (log) only and whole-tree harvesting for the
- 217 eight spruce sites. The calculations were based on the mean annual stem biomass production

- 218 for each site (Table 1) and the amount of stems, branches and needles was estimated using
- 219 biomass expansion functions (Marklund, 1988), which are based on empirical relationships
- between the diameter at breast height and empirical biomass data for a large number of trees
- in Sweden. The amount of stems, branches and needles was multiplied with concentrations ofCa, Mg, K and Na in the different tree compartments compiled in Akselsson (2005). Stem
- harvesting was defined as removal of the stem only, whereas whole-tree harvesting was
- defined as the harvest of the stem together with 60% of the branches in thinnings and in final
- felling (Swedish Forest Agency, 2008). Furthermore, it was assumed that 75% of the needles
- on 60 % of the branches, i.e. in total 45% of the needles, were removed. The methodology is
- described in more detail in Akselsson et al. (2007).
- 228

3. Results

- 230 The comparison of the total base cation weathering rates in the upper 50 cm of the mineral
- soil showed significantly (p=0.006, t-test) higher rates for PROFILE (with an interval of 12-
- 135 meq m⁻² yr⁻¹ and a median of 39 meq m⁻² yr⁻¹) than for the depletion method (with a
- corresponding interval of 8-41 meq m⁻² yr⁻¹ and a median of 19 meq m⁻² yr⁻¹). There were substantial deviations in estimated weathering rates for three sites. The rate estimated by the
- 235 PROFILE model was extremely high in Nunasvaara for Ca (35 meq m⁻² yr⁻¹), Mg (40 meq m⁻²
- 236 yr^{-1}) and Na (51 meq m⁻² yr⁻¹); 5 times higher than the rates derived from the depletion
- 237 method. In Målaskog the PROFILE-modeled Ca weathering rate was also high (31 meq m^{-2})
- $238 ext{ yr}^{-1}$) compared to the depletion method (6.6 meq m⁻² yr⁻¹). In Vålberget it was the other way
- around, with high rates of Ca (21 meq m⁻² yr⁻¹) and Mg (13 meq m⁻² yr⁻¹) weathering from the
- depletion method compared to PROFILE (4.8 and 2.3 meq $m^{-2} yr^{-1}$ for Ca and Mg
- respectively).
- For Ca and Mg there was correlation between the results from the two methods (Fig. 2). The
- 243 weathering of Mg was generally higher according to the depletion method, whereas it was the
- other way around for Ca. For K and Na there was weak or no correlation between the two
- 245 methods. PROFILE gave generally higher weathering rates for Na, whereas there was no
- such difference for K. The Spearman rank correlation between the depletion method and PPOFILE (T, t) = 0.041 + 0.041
- PROFILE (Table 4) indicated significant correlation for Ca (ρ =0.44, p=0.04), while it was non-significant for Mg (ρ =0.51, p=0.09), Na (ρ =0.25, p=0.34) and K (ρ =0.07, p=0.80). For
- the sum of the base cations the correlation was non-significant (ρ =0.11, p=0.67). The
- correlation coefficients for Ca and Mg were strongly influenced by the three extreme sites
- where either method gave highly deviating results. Disregarding these sites resulted in highly
- significant correlations for Ca (ρ =0.71, p=0.007) and Mg (ρ =0.82, p=0.001). For Ca and Na
- 253 weathering estimated with the same method there was a highly significant correlation; $\rho=0.80$
- (p<0.001) for PROFILE and $\rho=0.73$ (p=0.001) for the depletion method.

- 255 The average mass loss estimated by the depletion method (Fig. 3) increased towards the
- surface in the investigated profiles, but the gradients varied for the different base cations. The
- largest average mass loss in the uppermost mineral soil was found for Mg (66%), followed by
- Ca (43%), Na (38%) and K (29%) as compared to the reference soil layers. The gradient in weathering rate with depth was the opposite for the two methods (Fig. 4). The depletion
- 260 method, reflecting the average rates since the last glaciation, showed high rates in the upper
- 261 layer, and decreasing rates with depth, reaching ca 12 meq m^{-3} yr⁻¹ at the 50 cm level. The
- 262 PROFILE results showed, on the other hand, an increasing weathering rate with depth,
- 263 reaching ca 110 meq m^{-3} yr⁻¹ at the 50 cm level.
- 264 The calculation of harvest losses of base cations showed that harvesting of branches and
- 265 needles almost doubled the losses compared with stem harvest only (Fig. 5). The comparison
- between harvest losses and weathering rates showed that the losses at whole-tree harvesting
- 267 were of the same order of magnitude as PROFILE modeled weathering rates for the three
- 268 northern sites, whereas the weathering rates from the depletion method were about half the
- size. For the southern sites, the harvest losses at whole-tree harvesting were substantially
- 270 higher than the weathering rates from either approach.
- 271

4. Discussion

273 4.1. Comparison between results and concepts

274 The weathering rates estimated by the two methods agreed in some respects, but not in others 275 and the observed differences may be due to a number of factors. The depletion method gives 276 the average weathering rate since the deglaciation, but the historical development of the 277 weathering in the soil is not fully known. Steady state weathering, on the other hand, is a 278 theoretical concept used in PROFILE, which is based on steady state calculations where the 279 input data are kept constant at today's levels, and the steady state soil chemistry and 280 weathering are estimated under the assumption that the present minerals are not depleted. The 281 difference in time perspective for the two methods makes it crucial to consider the historical 282 development of the weathering rate in the interpretation of the results. The influence of soil 283 age on the weathering rate has been investigated in chronosequence studies (Bain et al., 1993; Taylor & Blum, 1995; White et al., 1996; Hodson & Langan, 1999) and by modelling 284 285 (Warfvinge et al., 1995). The results show high initial weathering rates followed by a decline, 286 which is most pronounced in the first centuries following soil exposure, but the results varies 287 greatly between sites. The decline is caused by the depletion of easily weathered minerals and decrease in reactive mineral surface area (Taylor & Blum, 1995). Attempts have been made to 288 289 correct the long-term weathering rate to today's levels (Taylor & Blum, 1995) using a power-290 law function fitted to depletion weathering rates from chronosequence data. These results together with simulations of historical weathering rates (Warfvinge et al., 1995) indicate that 291

292 the ratio between today's weathering rate and the long-term historical weathering rate is 293 approximately 0.3-0.5.

294 Based on this, the depletion method was expected to give higher values than the PROFILE 295 model. For Mg our results agreed with the theory; the depletion method gave higher 296 weathering rates than PROFILE. However, for Ca, Na and K the PROFILE model gave 297 higher values than the depletion method, although for K there was large non-systematic variation between the two approaches for many sites. Thus, the results for Ca, Na and K 298 299 contradict the theory about decreasing weathering rates over time. A contributing factor to our 300 observed differences between PROFILE and the depletion method is that the parent material 301 for our profiles is relatively poor in easily weathered minerals that may give rise to such 302 extreme initial weathering rates reported elsewhere (Bain et al., 1993; Taylor & Blum, 1995). 303 Furthermore, the soil profiles were developed in glacial till that partly consists of soil material 304 that was newly formed during the Weichselian glaciation and partly includes redistributed old 305 till deposits from previous glaciations. Consequently, the parent material in the investigated profiles was not completely unweathered at the time of deglaciation. The relatively elevated 306 307 acid deposition at present could also contribute to increased weathering (April et al. 1986), as 308 well as the increased forest growth and harvest levels during the last century, leading to 309 increased acidification and removal of weathering products. It is, however, difficult to design 310 experiments aiming at quantifying these effects. A sensitivity analysis of PROFILE (Hodson 311 et al., 1996) showed that increased sulphur deposition led to increased or unchanged 312 weathering rates for most minerals, but the effect was generally relatively small.

313 The correlation analysis for Ca and Mg showed that the coefficients were negatively 314 influenced by three sites where one of the two methods yielded highly deviating results. As 315 discussed below, there are good reasons to believe that these deviating results were caused by 316 conditions that caused either model to fail. Hence, the result indicates significant correlations 317 for Ca and Mg weathering when conditions are suitable for both methods. For K and Na the 318 correlation was weak and it could be hypothesized that K and Na weathering in PROFILE 319 was more sensitive to variation in other input data than mineralogy than for Ca and Mg. This 320 means that uncertainties in other input data would have a greater impact on the results. For K, 321 this is to some extent supported by Hodson et al. (1996), who identified K-feldspar, one of the 322 main sources of K, as a mineral particularly sensitive to input variations. To be able to fully 323 explain the difference, even more in-depth studies are required, including an accurate 324 determination of the actual minerals present. The Ca and Na weathering rates estimated with 325 the same method were significantly correlated for both methods. This indicates a proportional 326 release of Ca and Na that could be explained by the occurrence of both elements in the same 327 minerals e.g. the plagioclases, which are rather weatherable and very common in the 328 investigated area. The correlation study showed very different results for the different base 329 cations, which highlights the importance to compare individual base cations rather than

330 merely the sum of them.

- 331 Uncertainties related to the specific surface area are generally one of the main sources of
- 332 uncertainties to the mineral weathering rates estimated with PROFILE. This can be due to
- lack of detailed data on grain size distribution, and due to the generalized equations
- transforming grain size distribution to specific surface area. Hodson et al. (1998) questioned
- the equation for estimating specific surface area from grain size distributions, based on a
- 336 study in Scotland. However, since the equation was developed based on data from Sweden,
- the applicability can be expected to be higher in Sweden. Hodson and Langan (1999)
- highlighted the fact that changes over time of specific surface area and mineral reactivity are
- not taken into account in the PROFILE model, leading to uncertainties in the estimated
- 340 weathering rates. The uncertainties introduced by the data on specific surface area can be 341 reduced by using detailed input data on grain size distribution and densities for several layers
- reduced by using detailed input data on grain size distribution and densities for several layers,
 as was done in this study. However, although detailed data is used, the uncertainties will be
- 343 substantial, due to the problems discussed above.
- 344 The depletion method showed very large mass loss of base cations in the upper part of the soil
- profile considering the relatively young age of the investigated soils (ca $10\ 000 16\ 000$
- 346 years). Particularly the loss of Mg was considerable. It has been recognized that Mg is one of
- the most mobile elements in similar soils (Bain et al., 1993; Lång, 1995; Land et al., 1999;
- 348Olsson & Melkerud, 2000). In Land et al. (1999) they found a depletion level of 70% for Mg
- in the E horizon of Podzol soils in northern Sweden. The Mg loss is probably associated with
- depletion of easily weathered minerals such as chlorite and hornblende, which are important
- 351 Mg-bearing minerals in the soils of the Svecofennian granitic bedrock and important for forest
- soil productivity (Stendahl et al., 2002). The high depletion levels for all base cations indicate
- a general loss of easily weathered minerals.
- Important drivers for the weathering include processes at the soil surface such as biological production and deposition of acid compounds, which explain the pattern with higher historical weathering intensity towards the surface. However, there is an on-going depletion of easily weathered minerals over time in the uppermost part of the soil, which limits the weathering
- rate towards the surface. The weathering intensity is gradually shifted towards lower depths,
 where there are more easily weathered minerals present. Since less weathering agents (i.e.
- 357 where there are more easily weathered innerals present. Since less weathering agents 360 organic acids, HCO₃) are being consumed in the uppermost part of the soil, they are
- 361 transported down in the profile and contribute to the weathering at lower depths. In our
- 362 results, the steady-state weathering rate estimated with PROFILE exceeds the historical rates
- 363 estimated by the depletion method in deeper soil layers (Fig. 4).
- 364

365 *4.2. Applicability of the two approaches for different conditions*

The three sites that were outliers in the comparison can be useful to analyze critical conditions for the two approaches. In Nunasvaara the PROFILE weathering rates of Ca, Mg and Na were much higher than for any of the other sites, and also several times higher than the weathering

rates estimated with the depletion method. The depletion method indicated moderate losses 369 370 despite the seemingly good mineralogy indicated by the base cation content. The high 371 modeled weathering rates by PROFILE can be explained by a combination of high contents of 372 minerals rich in Ca, Mg and Na in the soil, e.g. plagioclase, hornblende, biotite and augite 373 (Table 3), high soil density and a high fraction of fine-grained material (Table 1). A possible 374 explanation to the deviation between the methods could be that the simplified handling of 375 hydrology in PROFILE does not take into account that the high soil density may lead to 376 saturation of the soil and thus substantially increased product inhibition, slowing down 377 weathering rates. In Vålberget the situation was the other way around; the depletion method 378 gave many times higher weathering rates for Ca and Mg than PROFILE. The Ca, Na and K 379 weathering rates in Vålberget were the lowest of the 16 sites according to PROFILE and the 380 Mg weathering rate was among the lowest. The depletion method gave by far the highest 381 weathering rates of the 16 sites for Ca and Mg, whereas the K weathering was one of the 382 lowest and the Na weathering was intermediate compared to the other sites. The fractional 383 mass loss of this site was also very large; on average 55% of the Ca and 44% of the Mg has 384 been lost down to 50 cm depth, while in the uppermost layer the result showed that as much 385 as 81% of the Ca and 87% of the Mg has been lost by weathering. An indication that the Ca 386 weathering rate in Vålberget could be problematic was that the site deviated completely from 387 the otherwise strong relationship between Ca and Na weathering estimated by the depletion 388 method. This indicate that the systematic Ca gradient in the profile was caused by 389 heterogeneities rather than soil development and that the weathering rate thus was overestimated. In Målaskog the PROFILE model resulted in much higher weathering rates 390 391 than the depletion method for Ca and Na. Although the soil in Målaskog was relatively 392 coarse, its high Ca content led to the second highest Ca weathering rate according to 393 PROFILE. The soil profile in Målaskog showed opposite gradients both for mobile and 394 immobile ions indicating vertical heterogeneity in parent material composition, which means 395 that the method assumptions for the depletion method were not fulfilled.

396 *4.3.* The results in a sustainability perspective

For detailed sustainability assessments the complete mass balance has to be taken intoaccount, i.e. base cation deposition, weathering, harvest losses and leaching (Sverdrup et al.,

399 2006; Akselsson et al., 2007). Base cation deposition in Sweden is in the same order of

400 magnitude as weathering rates (Akselsson et al., 2007), but the input of base cations through

401 weathering and deposition needs to be substantially higher than the harvest losses, to avoid

- 402 soil acidification. However, simplified mass balance including only weathering and harvest
- 403 losses is often used as an indication of the sustainability (Olsson et al., 1993; Klaminder et al.,
- 404 2011). If weathering rates are substantially higher than base cation removal at whole-tree
- 405 harvesting, the harvesting effects on the nutrient budget can be expected to be small.
- 406 However, if the weathering rates are in the same order of magnitude or lower than the base
- 407 cation removal, there is an obvious risk of negative effects on the soil nutrient status. Our
- 408 results show that the removal at whole-tree harvesting were in the same order of magnitude or

- 409 higher level than the weathering rate, no matter which of the approaches that was used. For
- 410 the sites in southern Sweden, the losses at harvesting were substantially larger than the
- 411 weathering rates. This area also contains the most acidic soils, due to high historical levels of
- 412 acid deposition (Pihl Karlsson et al., 2011). Thus, the risk of negative effects associated with
- 413 base cation losses at harvesting is largest in these regions. The northern sites have better soil
- 414 status to start with from an acidification point of view, and the losses at whole tree harvesting
- 415 are smaller. These areas are associated with lower risks, but since the losses at harvesting are
- 416 as big as the weathering rates, and since the base cation deposition is relatively low in
- 417 northern Sweden, there is not much left for the runoff water, which may lead to negative
- 418 effects for surface waters on a long term.

419 **5. Conclusions**

- 420 Where conditions are suitable for both methods there is a correspondence in weathering rates
- 421 between the depletion method and PROFILE. For Mg, the depletion method yield higher rates
- 422 than PROFILE, probably due to strong depletion of main Mg-bearing minerals, such as
- 423 chlorite and hornblende. For Ca, K and Na the weathering rates are generally higher
- 424 according to PROFILE than according to the depletion method. This indicates that initial
- 425 weathering rates of these elements at the time of soil formation were less extreme in the
- 426 investigated sites compared to other regions. The weathering rates in the upper 50 cm
- 427 decrease with depth according to the depletion method, but increase with depth according to
- 428 PROFILE. The pattern can be explained by the methods difference in time perspective;
- 429 historically most of the weathering has taken place in the upper soil, which is now depleted of
- 430 easily weathered minerals leading to a lower current rate as represented by PROFILE. The
- 431 application of multiple methods on a large selection of sites can help identify deviating results
- 432 for the methods included. The losses from whole-tree harvesting exceed weathering for both
- the depletion method and PROFILE, especially for sites in southern Sweden.

434 Acknowledgements

- 435 This study was financed by the Formas strong research environment QWARTS, the Swedish
- 436 Environmental Protection Agency and the environmental monitoring and assessment
- 437 programme at the Swedish University of Agricultural Sciences.

438 **References**

- 439 Akselsson, C., 2005. Regional nutrient budgets in forest soils in a policy perspective. Doctoral
- 440 Thesis, Department of Chemical Engineering, Lund University. ISBN 91-7422-076-4.
- 441 Akselsson, C., Holmqvist, J., Alveteg, M., Kurz, D. and Sverdrup, H., 2004: Scaling and
- 442 mapping regional calculations of soil chemical weathering rates in Sweden. Water, Air, &
- 443 Soil Pollution: Focus 4: 671-681.
- 444 Akselsson, C., Westling, O., Sverdrup, H., Gundersen, P., 2007. Nutrient and carbon budgets
- in forest soils as decision support in sustainable forest management. Forest Ecology and
- 446 Management 238(1-3), 167-174.
- 447 Akselsson, C., Westling, O., Sverdrup, H., Holmqvist, J., Thelin, G., Uggla, E. and Malm, G.,
- 448 2007. Impact of harvest intensity on long-term base cation budgets in Swedish forest soils.
- 449 Water, Air, and Soil Pollution: Focus 7: 201-210.
- April, R., Newton, R., Coles, L.T., 1986. Chemical-Weathering in 2 Adirondack Watersheds Past and Present-Day Rates. Geological Society of America Bulletin 97, 1232-1238.
- 452 Bain, D.C., Mellor, A., Robertson-Rintoul, M.S.E., Buckland, S.T., 1993. Variations in
- 453 weathering processes and rates with time in a chronosequence of soils from Glen Feshie,
- 454 Scotland. Geoderma 57, 275-293.
- 455 Bain, D.C., Mellor, A., Wilson, M.J., Duthie, D.M.L., 1994. Chemical and Mineralogical
- Weathering Rates and Processes in an Upland Granitic Till Catchment in Scotland. Water,Air, and Soil Pollution 73, 11-27.
- 458 Brimhall, G.H., Dietrich, W.E., 1987. Constitutive Mass Balance Relations between
- 459 Chemical-Composition, Volume, Density, Porosity, and Strain in Metasomatic
- 460 Hydrochemical Systems Results on Weathering and Pedogenesis. Geochimica et
- 461 Cosmochimica Acta 51, 567-587.
- 462 Brimhall, G.H., Lewis, C.J., Ford, C., Bratt, J., Taylor, G., Warin, O., 1991. Quantitative
- 463 Geochemical Approach to Pedogenesis Importance of Parent Material Reduction,
- 464 Volumetric Expansion, and Eolian Influx in Lateritization. Geoderma 51, 51-91.
- 465 Cosby, B.J., Ferrier, R.C., Jenkins, A., Wright, R.F., 2001. Modelling the effects of acid
- 466 deposition: refinements, adjustments and inclusion of nitrogen dynamics in the MAGIC
- 467 model. Hydrology and Earth System Sciences 5(3), 499-517.
- 468 Fredén, C., 2009. Geology. The National Atlas of Sweden. 3rd edition. SNA Publishing469 House, Stockholm.

- 470 Futter, M.N., Klaminder, J., Lucas, R.W., H., Laudon, H., Köhler, S.J., 2012. Uncertainty in
- 471 silicate mineral weathering rate estimates: source partitioning and policy implications.
- 472 Environmental Research Letters 7, 1-8.
- Hodson, M.E., 2002. Experimental evidence for mobility of Zr and other trace elements in
 soils. Geochimica et Cosmochimica Acta 66(5), 819-828.
- 475 Hodson, M.E., Langan, S.J., Wilson, M.J., 1996. A sensitivity analysis of the PROFILE
- 476 model in relation to the calculation of soil weathering rates. Applied Geochemistry 11, 835-477 844.
- Hodson, M.E., Langan, S.J., 1999. The influence of soil age on calculated mineral weathering
 rates. Applied Geochemistry. 14, 387-394.
- Hodson, M.E., Langan, S.J., Meriau, 1998. Determination of mineral surface area in relation
 to the calculation of weathering rates. Geoderma 83, 35-54.
- 482 Holmqvist, J., Thelin, G., Rosengren, U., Stjernquist, I., Wallman, P., Sevrdrup, H., 2002.
- 483 Assessment of sustainability in the Asa Forest Park. In H. Sverdrup, I. Stjernquist (Eds):
- 484 Developing Principles and Models for Sustainable Forestry in Sweden. Kluwer Academic
- 485 Publishers. ISBN 1-4020-0999-2.
- 486 Jönsson, C., Warfvinge, P. and Sverdrup, H.: 1995. Uncertainty in predicting weathering rates
- 487 and environmental stress factors with the profile model. Water, Air, and Soil Pollut. 81,
- 488 1.Klaminder, J., Lucas, R.W., Futter, M.N., Bishop, K.H., Köhler, S.J., Egnell, G., Laudon,
- 489 H., 2011. Silicate mineral weathering rate estimates: Are they precise enough to be useful
- 490 when predicting the recovery of nutrient pools after harvesting? Forest Ecology and
- 491 Management 261, 1-9.
- Kolka, R.K., Grigal, D.F., Nater, E.A., 1996. Forest soil mineral weathering rates: Use of
 multiple approaches. Geoderma 73, 1-21.
- 494 Koseva, I., Watmough, S., Aherne, J., 2010. Estimating base cation weathering rates in
- 495 Canadian forest soils using a simple texture-based model. Biogeochemistry 101, 183-196.
- Land, M., Ingri, J., Öhlander, B., 1999. Past and present weathering rates in northern Sweden.
 Applied Geochemistry. 14(6), 761-774.
- 498 Lång, L.-O., 1995. Mineral weathering rates and primary mineral depletion in forest soils, SW
 499 Sweden. Ecological Bulletins 44, 100-113.
- 500 Langner, J., Persson, C., Robertson, L., Ullerstig, A., 1996. Air Pollution Assessment Study
- 501 Using the MATCH Modelling System. Application to Sulphur and Nitrogen Compounds over
- 502 Sweden 1994. Swedish Meteorological and Hydrological Institute Report, 69. Swedish
- 503 Meteorological and Hydrological Institute, Norrköping, Sweden.

- 504 Marklund, L.-G., 1988. Biomass functions for pine, spruce, and birch in Sweden. Department
- 505 of Forest Taxation, Swedish University of Agricultural Sciences, SLU, Report 45:1988. (in
- 506 Swedish with English summary)
- 507 Marshall, C.E., Haseman, J.F., 1942. The quantitative evaluation of soil formation and
- development by heavy mineral studies: a Grundy silt loam profile. Soil Science Society of
 America Proceedings. 7, 448-453.
- 510 Martinson, L., Alveteg, M., Mörth, C-M., Warfvinge, P., 2003. The effect of changes in
- 511 natural and antropogenic deposition on modelling recovery from acidification. Hydrology and
- 512 Earth System Sciences 7(5), 766-776.
- 513 Olsson, M., Melkerud, P.-A., 1989. Chemical and Mineralogical Changes during Genesis of a514 Podzol from Till in Southern Sweden. Geoderma 45, 267-287.
- 515 Olsson, M. Melkerud, P.-A., 1991. Determination of weathering rates based on geochemical
- 516 properties of the soil. Environmental Geochemistry of Northern Europe. Geological Survey of
- 517 Finland, Special Paper 9, 69-78.
- 518 Olsson, M., Melkerud, P.-A., 2000. Weathering in three podzolized pedons on glacial deposits 519 in northern Sweden and central Finland. Geoderma 94, 149-161.
- 520 Olsson, M., Rosén, K., Melkerud, P.-A., 1993. Regional modeling of base cation losses from
- 520 Swedish forest soils due to whole-tree harvesting. Applied Geochemistry, Supplementary
- 522 Issue No 2, January 1993: Environmental Geochemistry, 189-194.
- 523 Pihl Karlsson, G., Akselsson, C., Hellsten, S., Karlsson, P.E., 2011. Reduced European
- 524 emissions of S and N Effects on air concentrations, deposition and soil water chemistry in
- 525 Swedish forests. Environmental Pollution 159, 3571-3582.
- Posch, M., Kurz, D., 2007. A2M A program to compute all possible mineral modes from
 geochemical analyses. Computers and Geosciences 33, 563-572.
- Raab, B., Vedin, H. (Eds.), 1995. Climate, lakes and rivers. The National Atlas of Sweden.
 SNA Publishing House, Stockholm.
- 530 Raulund-Rasmussen, K., Callesen, I., 1999. Nordsoil database. Datafile. Danish Forest and
- 531 Landscape Research Institute, Hørsholm, Denmark.
- 532 Schöpp, W., Posch, M., Mylona, S., Johansson, M., 2003. Long-term development of acid
- deposition (1880-2030) in sensitive freshwater regions in Europe. Hydrology and Earth
 System Sciences 7(4), 436-446.

- 535 SGU, 1994. Jordartskartor. Metodik och jordartsindelning (Quaternary map. Methodology
- and classification of Quaternary deposits). Särtryck ur SGU serie Ae. Uppsala ISBN 91-7158539-7. (in Swedish).
- 538 Starr, M., Lindroos, A.-J., Tarvainen, T., Tanskanen, H., 1998. Weathering rates in the
- 539 Hietajärvi Integrated Monitoring catchment. Boreal Environment Research 3, 275–285.
- 540 Stendahl, J., Olsson, M., Snäll, S., Holmgren, P., 2002. Influence of soil mineralogy and
- chemistry on site quality within geological regions in Sweden. Forest Ecology andManagement 170 (1-3), 75-88.
- 543 Stendahl, J., Lundin, L., Nilsson, T., 2009. The stone and boulder content of Swedish forest
 544 soils. Catena 77(3), 285-291.
- 545 Sverdrup, H., Warfvinge, P., 1993. Calculating field weathering rates using a mechanistic
 546 geochemical model PROFILE. Applied Geochemistry 8, 273-283.
- 547 Sverdrup, H., Rosén, K., 1998. Long-term base cation mass balances for Swedish forests and
 548 the concept of sustainability. Forest Ecology and Management 110, 221-236.
- 549 Sverdrup, H., Warfvinge, P., Wickman, T., 1998. Estimating the weathering rate at Gårdsjön
- 550 using different methods, In: Experimental Reversal of Acid Rain Effects: The Gårdsjön Roof
- 551 Project. John Wiley & Sons Ltd.
- 552 Sverdrup, H., Thelin, G., Robles, M., Stjernquist, I., Sörensen, J., 2006. Assessing Nutrient
- 553 Sustainability of Forest Production for Different Tree Species Considering Ca, Mg, K, N and
- P at Björnstorp Estate, Sweden. Biogeochemistry 81(2), 219-238.
- 555 Swedish Forest Agency, 2008. Rekommendationer vid uttag av avverkningsrester och
- askåterföring (Recommendations for the extraction of harvest residues and ash recycling).
- 557 Meddelanden 2, 2008. Swedish Forest Agency, Jönköping. (in Swedish).
- Taylor, A., Blum, J.D., 1995. Relation between soil age and silicate weathering rates
 determined from the chemical evolution of a glacial chronosequence Geology 23, 979-982.
- 560 Warfvinge, P., Sverdrup, H., 1995. Critical Loads of Acidity to Swedish Forest Soils. Reports
- 561 in ecology and environmental engineering 5: 1995. Department of Chemical Engineering II,
- 562 Lund University. ISSN 1104-2877.
- 563 Warfvinge, P., Sverdrup, H., Alveteg, M., Rietz, F., 1995. Modelling geochemistry and lake
- pH since glaciation at Lake Gårdsjon. Water, Air, and Soil Pollution 85, 713-718.
- 565 White, A.F., Blum, A.E., Schulz, M.S., Bullen, T.D., Harden, J.W., Peterson, M.L., 1996.
- 566 Chemical weathering rates of a soil chronosequence on granitic alluvium .1. Quantification of

- 567 mineralogical and surface area changes and calculation of primary silicate reaction rates.
- 568 Geochimica et Cosmochimica Acta 60, 2533-2550.
- 569 Whitfield, C.J., Watmough, S.A., Aherne, J., Dillon, P.J., 2006. A comparison of weathering
- 570 rates for acid-sensitive catchments in Nova Scotia, Canada and their impact on critical load
- 571 calculations. Geoderma 136(3–4), 899-911.
- 572 Zak, S. K., Beven, K. and Reynolds, B.: 1997. Uncertainty in the estimation of critical loads:
- 573 A practical methodology.Water, Air, and Soil Pollut. 98, 297.
- 574

575 Figure captions

576 Fig. 1. Location of the 16 sampling sites.

577 Fig. 2. Weathering rate (meq $m^{-2} yr^{-1}$) for Ca, Mg, K and Na estimated by the PROFILE and 578 the depletion method for 16 soil profiles.

579 Fig. 3. The fractional mass loss of Ca, Mg, K and Na at different soil depths (cm) estimated 580 by the depletion method. The error bars represent the 95% confidence limits.

- 581 Fig. 4. Weathering intensity (meq $m^{-3} yr^{-1}$) of the sum of base cations vs. soil depth (cm)
- estimated by the PROFILE and the depletion method. The error bars represent the 95%confidence limits.
- 584 Fig. 5. Weathering rates of base cations estimated by the two approaches compared with
- 585 harvest losses of base cations at eight spruce sites. Harvest losses are calculated for a whole-
- tree harvesting scenarios where all tree stems, 60% of the branches and 75% of the needles on
- the branches are harvested. The dashed lines in the harvesting bars indicate the level when
- 588 only tree stems are harvested.

Tables

Site	Lat.	Long.	MAT	MAP	Clay	Silt	Sand	Density ^b	Soil depth (cm)	Soil age	Soil	Tree species	Stem volume
			(°C)	(mm)	$(\%_{\rm wt})$	(% _{wt})	$(\%_{\rm wt})$	$(g dm^{-3})$		(calendar	moisture class ^c		production
										years)			$(m^3 ha^4 yr^4)$
Bodafors	57.6	14.7	5.4	729	2	38	60	1400	192	14 000	3	Norway spruce	10.7
Gårdsjön	58.0	12.0	6.6	952	12	33 ^a	55 ^a	1055	72	14 340	3	Norway spruce	9.5
Hjärtasjö	59.0	15.3	5.2	650	5	31 ^a	64 ^a	992	113	11 280	3	Norway spruce	9.5
Hässlen	60.3	16.3	4.3	629	9	41 ^a	50^{a}	1442	130	10 830	3	Other broadleaves	8.4
Kloten	59.9	15.3	4.2	807	2	48	50	1590	125	10 910	4	Scots pine	6.1
Kullarna	62.0	16.6	3.8	552	4	12^{a}	84^{a}	1390	102	10 400	3	Norway spruce	5.2
Lammhult	57.2	14.8	5.5	688	2	35	63	1376	178	14 260	3	Norway spruce	9.7
Målaskog	56.8	14.3	6.4	707	0	23 ^a	77^{a}	1072	175	14 360	3	Other broadleaves	11.3
Nunasvaara	67.8	21.4	-2.0	456	4	43 ^a	53 ^a	1852	227	9 880	3	Scots pine	1.7
Risfallet	60.3	16.2	4.3	629	7	35 ^a	58^{a}	1560	136	10 810	3	Scots pine	7.0
Skånes Värsjö	56.3	13.5	6.4	765	3	25 ^a	72^{a}	1291	112	14 700	3	Norway spruce	11.6
Stöde	62.3	16.4	2.5	707	2	33 ^a	65 ^a	1271	263	10 330	3	Norway spruce	5.5
Svartberget	64.3	19.7	1.0	573	2	26^{a}	72^{a}	1534	217	10 120	4	Norway spruce	3.6
Söderåsen	56.1	13.0	7.2	795	6	24^{a}	70^{a}	1350	126	15 900	3	Beech	12.6
Vindeln	64.2	19.6	1.2	523	4	37	59	1407	207	10 130	3	Scots pine	3.6
Vålberget	60.8	13.8	2.6	701	3	17^{a}	80^{a}	1243	176	10 580	3	Clearcut (spruce)	6.2

Table 1. Site characteristics and soil properties at 50 cm depth in the mineral soil for the 16 investigated sites.

^a The fraction 0.02-0.06 mm, which according to common definitions is a part of the silt fraction, was included in sand fraction in the original data for many sites. In this study, the sand and silt fractions were recalculated according to common definitions to be compatible with the PROFILE model. The calculations were based on relationships from the other sites, where complete grain size distributions were available.

^b Volume weight of < 2 mm fraction including coarser dead organic material (volume of live roots, gravel and stones deducted from sampled volume).

^c Drainage class according to the NORDSOIL database: 1. Excessively, 2. Somewhat excessively, 3. Well, 4. Moderately, 5. Imperfectly, 6. Poorly 7. Very poorly.

Site	Ref. depth	Ca	Mg	K	Na	Zr
	(cm)	(%)	(%)	(%)	(%)	(ppm)
Bodafors	62	1.47	0.65	2.31	2.03	298
Gårdsjön	57	1.60	0.68	2.54	2.29	299
Hjärtasjö	57	1.13	0.88	1.99	1.62	189
Hässlen	82.5	1.19	0.79	3.10	1.72	250
Kloten	48.5	0.94	0.28	2.47	1.76	247
Kullarna	62	1.14	0.58	2.99	1.84	343
Lammhult	98	1.54	0.59	2.34	2.31	232
Målaskog	60.0	2.39	1.07	2.47	2.66	261
Nunasvaara	54.5	2.95	2.04	1.88	3.29	192
Risfallet	76	1.01	0.61	2.55	2.05	181
Skånes Värsjö	62	1.00	0.24	3.10	2.33	466
Stöde	55.5	1.47	0.86	2.32	1.87	229
Svartberget	154	1.37	0.41	2.41	2.35	241
Söderåsen	101	0.75	0.36	2.84	1.8	341
Vindeln	129	1.00	0.92	2.47	1.67	279
Vålberget	117	1.57	0.77	1.36	0.79	186

Table 2. Element concentrations for the < 2 mm soil fraction at the reference layer depth used in the depletion method estimates.

Table 3. Normative mineralogy at 50 cm depth in the mineral soil based on the A2M model (Qu=Quartz, Or=Orthoclase, Pl=Plagioclase, Ho=Hornblende, Ep=Epidote, Bi=Biotite, Au=Augite, Ap=Apatite, Mu=Muscovite, Ch=Chlorite, Il=Illite, Ve=Vermiculite, An/Ab=ratio between Anorthite and Albite in Plagioclase).

]	Minera	l conte	nts (%)				
	Qu	Or	P1	Ho	Ep	Bi	Au	Ар	Mu	Ch	I1	Ve	An/Ab
Bodafors	41.0	13.2	25.8	1.2	3.8	0.0	0.0	0.5	3.6	2.5	1.6	5.5	0.12
Gårdsjön	36.5	4.7	26.4	1.0	3.6	0.0	0.0	0.2	8.4	1.4	12.2	3.5	0.14
Hjärtasjö	49.3	3.0	20.4	0.8	1.9	0.0	0.0	0.2	9.4	1.9	9.1	4.1	0.12
Hässlen	40.2	15.1	21.8	1.0	2.5	0.0	0.0	0.2	6.3	2.1	6.6	3.6	0.13
Kloten	51.5	13.6	21.8	0.5	3.0	0.0	0.0	0.2	3.4	1.0	3.0	1.3	0.10
Kullarna	39.1	15.3	25.6	1.0	2.9	0.0	0.0	0.2	5.5	1.8	4.8	2.5	0.15
Lammhult	37.9	14.6	29.0	1.5	4.2	0.0	0.0	0.4	2.4	2.0	1.2	4.9	0.12
Målaskog	28.5	15.7	31.8	2.6	7.5	0.0	0.0	0.6	2.6	2.7	1.3	5.2	0.11
Nunasvaara	16.7	10.2	36.0	11.1	2.1	2.7	8.1	0.5	0.0	1.3	4.5	4.7	0.05
Risfallet	45.4	13.5	26.2	0.9	2.6	0.0	0.0	0.2	3.8	2.1	3.4	3.2	0.10
Skånes Värsjö	38.8	16.5	29.7	0.7	2.4	0.0	0.0	0.3	4.3	1.0	2.2	1.4	0.10
Stöde	40.5	4.7	23.4	1.9	3.7	0.0	0.0	0.4	9.9	3.6	9.3	2.5	0.13
Svartberget	42.4	15.1	29.1	1.2	2.9	1.6	0.5	0.2	0.0	0.8	4.5	1.9	0.15
Söderåsen	48.9	10.2	21.3	0.5	1.4	0.0	0.0	0.3	9.9	0.7	4.9	0.9	0.08
Vindeln	39.1	9.8	24.3	1.0	2.3	1.2	0.5	0.3	0.0	0.7	17.1	5.4	0.09
Vålberget	66.3	6.4	8.9	2.6	2.5	0.0	0.0	0.3	3.6	1.6	0.8	5.2	0.22

			Depletion method									
		Ca	Mg	K	Na	Σ						
	Ca	0.44	0.20	0.31	0.59	0.36						
		0.09	0.46	0.24	0.02	0.16						
	Mg	0.12	0.51	0.23	0.22	0.24						
[T]		0.65	0.04	0.40	0.42	0.36						
ILI	Κ	-0.13	-0.01	0.07	0.18	-0.07						
SOF		0.63	0.97	0.80	0.50	0.80						
ΡF	Na	0.05	0.07	0.15	0.25	0.04						
		0.85	0.80	0.57	0.34	0.88						
	Σ	0.19	0.11	0.14	0.34	0.11						
1		0.49	0.69	0.62	0.19	0.67						

Table 4. Spearman rank correlation coefficients between weathering estimates by PROFILE and the depletion method where \sum indicates the sum of base cations (n=16, p-values in italic).

Figures



Fig.1







Fig. 2b







Fig. 2d















Na









Fig. 4b



Fig. 5