



# LUND UNIVERSITY

## Drosophila Sensory Neuroethology

Mansourian, Suzan

2018

*Document Version:*

Publisher's PDF, also known as Version of record

[Link to publication](#)

*Citation for published version (APA):*

Mansourian, S. (2018). *Drosophila Sensory Neuroethology*. Lund University, Faculty of Science, Department of Biology.

*Total number of authors:*

1

### General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY


PO Box 117  
221 00 Lund  
+46 46-222 00 00



# *Drosophila* Sensory Neuroethology

SUZAN MANSOURIAN

DEPARTMENT OF BIOLOGY | FACULTY OF SCIENCE | LUND UNIVERSITY



- I. Mansourian S and Stensmyr MC. (2015). The chemical ecology of the fly. *Current Opinion in Neurobiology* 34: 95-102. doi.org/10.1016/j.conb.2015.02.006
- II. Mansourian S, Corcoran J, Enjin A, Löfstedt Ch, Dacke M, and Stensmyr MC. (2016). Fecal derived phenol induces egg-laying aversion in *Drosophila*. *Current Biology* 26: 2762-2769. doi.org/10.1016/j.cub.2016.07.065
- III. Enjin A, Zaharieva EE, Frank D, Mansourian S, Suh GSB, Gallio M, and Stensmyr MC. (2016). Humidity sensing in *Drosophila*. *Current Biology* 26: 1352-1358. doi.org/10.1016/j.cub.2016.03.049
- IV. Mansourian S, Enjin A, Jirle E, Ramesh V, Pool E, and Stensmyr MC. Wild African *Drosophila melanogaster* are seasonal specialists on marula fruit. *Manuscript*.



# *Drosophila* Sensory Neuroethology

Suzan Mansourian



**LUND**  
UNIVERSITY

DOCTORAL DISSERTATION


by due permission of the Faculty Science, Lund University, Sweden.  
To be defended at the Blue Hall, Ecology Building, Sölvegatan 37, Lund.  
On 1<sup>st</sup> of June 2018 at 10.00.

*Faculty opponent*

Professor Matthew Cobb, University of Manchester, UK

Organization LUND UNIVERSITY	Document name	
	Date of issue May 8, 2018	
Author(s) Suzan Mansourian	Sponsoring organization	
Title and subtitle <i>Drosophila</i> Sensory Neuroethology		
<p>Abstract</p> <p>Animals, like humans, need to perceive their surroundings via their senses in order to make sensible behavioral decisions, reproduce successfully, and survive. Animals are equipped with audition, vision, thermosensation, hygosensation, mechanosensation, magnetoception, gustation, and olfaction, which detects physical and chemical changes in their habitats. Among these senses, olfaction is likely the most ancient sensory modality. Insects, the most abundant and successful group of the animal kingdom, predominantly use olfaction to find food, mates, breeding sites, and to avoid dangers. Moreover, hygosensation is vital for insects to find a suitable habitat and to avoid risks of dehydration. Our understanding of the molecular, neuronal, and morphological organization of the insect olfactory system is today substantial, in large parts thanks to <i>Drosophila melanogaster</i> (vinegar fly) and the wealth of sophisticated genetic tools available in this classic model system. Our knowledge regarding the functional and molecular basis of insect hygosensation, is, however, limited.</p> <p>In this thesis, I show that the vinegar fly olfactory system do not detect odor molecules randomly, but capture and process specific odors associated with needs and dangers. I demonstrate how the olfactory system cope with toxic and harmful matters in the natural habitat and I identify an olfactory circuit that mediates repellency towards phenol, which is produced by pathogenic bacteria, predominantly present in carnivore feces. Furthermore, I show that flies have an innate and species-specific ability to find suitable humidity levels, related to their native habitat. Vinegar flies can sense humidity changes in their environment through a trio of ionotropic receptors expressed in the sacculus of the antennae. Although <i>D. melanogaster</i> is known as a generalist, I show that wild populations of <i>D. melanogaster</i> from a mopane forest within the potential ancestral habitat have a strong breeding preference towards marula fruit. This fruit is seasonally abundant, native to Southern Africa, and is presumably the ancestral host of the vinegar fly. I also argue that marula drove the <i>D. melanogaster</i> to become a human commensal.</p> <p>In summary, the research presented in my thesis enhances our understanding of how the olfactory system operates, the behavior of wild flies, and introduces the genetic and neural basis underlying humidity sensation in insects. These findings might lead us to better strategies for controlling insect pests, as well as human disease vectors.</p>		
Key words: <i>Drosophila melanogaster</i> , vinegar fly, olfactory system, hygosensation, receptor, ancestral habitat.		
Classification system and/or index terms (if any)		
Supplementary bibliographical information		Language: English
ISSN and key title		ISBN: 978-91-7753-691-8
Recipient's notes	Number of pages 122	Price
	Security classification	

I, the undersigned, being the copyright owner of the abstract of the above-mentioned dissertation, hereby grant to all reference sources permission to publish and disseminate the abstract of the above-mentioned dissertation.

Signature 

Date: 2018-04-19

# *Drosophila* Sensory Neuroethology

Suzan Mansourian



**LUND**  
UNIVERSITY

Cover illustration at the front by *Sahar & Amirhossein Ashari*

Cover painting at the back by *Elnaz Saeedi*

Background Photo: Suzan Mansourian

© Suzan Mansourian

Faculty of Science  
Department of Biology

ISBN 978-91-7753-691-8 (print)

ISBN 978-91-7753-692-5 (pdf)

Printed in Sweden by Media-Tryck, Lund University  
Lund 2018



MADE IN SWEDEN 



Intertek

Media-Tryck is an environmental-  
ly certified and ISO 14001 certified  
provider of printed material.  
Read more about our environmental  
work at [www.mediatryck.lu.se](http://www.mediatryck.lu.se)

*To my kindest mami and baba.  
This journey began with your endless support.*



“I encourage all of us, whatever our beliefs, to question the basic narratives of our world, to connect past developments with present concerns, and not to be afraid of controversial issues.”

*Yuval Noah Harari*

# Table of Contents

Table of Contents .....	7
List of papers in the thesis.....	9
Author contributions .....	10
Popular science summary.....	11
Populärvetenskaplig sammanfattning .....	13
Persian popular summary .....	15
The Scope of this Thesis.....	17
Background of the study.....	19
<i>Drosophila melanogaster</i> – Lord of the flies .....	19
The lord’s close relatives.....	21
Olfaction – Lord of the senses.....	23
The peripheral olfactory system of the fly.....	23
The larval chemosensory system.....	24
Odorant receptors .....	25
Ionotropic receptors.....	26
Gustatory receptors.....	26
The central olfactory system of the fly.....	27
Antenna - a multifunctional sensory appendage.....	28
Hygrosensation - The sixth sense.....	29
Functional anatomy of hygrosensation in insects.....	29
Outcome and discussion of the studies.....	31
I. The chemical ecology of the fly.....	31
Host seeking .....	31
Volatile chemicals effecting oviposition .....	33
II. Foul feces fool flies .....	33
Phenol signals pathogenic bacteria to the fly .....	34
III. Humidity sensation in the fly .....	36
Early processing of hygrosensory information in the fly brain .....	37
IV. Marula – The ancestral host of the lord of flies .....	38
Or22a detects the marula volatile ethyl isovalerate.....	40

Concluding remarks.....	43
References .....	45
Acknowledgement.....	57

## List of papers in the thesis

- I. **Mansourian S** and Stensmyr MC. (2015). The chemical ecology of the fly. *Current Opinion in Neurobiology* 34: 95-102. doi.org/10.1016/j.conb.2015.02.006
  
- II. **Mansourian S**, Corcoran J, Enjin A, Löfstedt Ch, Dacke M, and Stensmyr MC. (2016). Fecal derived phenol induces egg-laying aversion in *Drosophila*. *Current Biology* 26: 2762-2769. doi.org/10.1016/j.cub.2016.07.065
  
- III. Enjin A, Zaharieva EE, Frank D, **Mansourian S**, Suh GSB, Gallio M, and Stensmyr MC. (2016). Humidity sensing in *Drosophila*. *Current Biology* 26: 1352-1358. doi.org/10.1016/j.cub.2016.03.049
  
- IV. **Mansourian S**, Enjin A, Jirle E, Ramesh V, Pool E, and Stensmyr MC. Wild African *Drosophila melanogaster* are seasonal specialists on marula fruit. *Manuscript*.

Papers I, II and III are reprinted with permission of the publishers.

## Author contributions

- I. **S.M** wrote the manuscript together with M.C.S.
- II. Conceptualization, **S.M.** and M.C.S.; Methodology, **S.M.**, M.D., and M.C.S.; Investigation, **S.M.**, A.E., J.C., and M.C.S.; Resources; C.L. and M.C.S.; Writing – Original Draft, M.C.S.; Writing – Review and Editing, all authors; Funding Acquisition, M.C.S., M.D., and C.L.; Supervision, C.L., M.D., and M.C.S.
- III. A.E., G.S.B.S., M.G., and M.C.S. conceived and designed the study. A.E. performed all humidity experiments and, with assistance from **S.M.**, anatomical studies by confocal microscopy. A.E. performed two-photon Ca<sup>2+</sup> imaging of humidity responses. E.E.Z. carried out all temperature-preference and two-photon anatomy experiments. D.D.F. performed two-photon Ca<sup>2+</sup> imaging for temperature responses. A.E., M.G., and M.C.S. analyzed the data and wrote the manuscript with input from all authors.
- IV. **S.M.** and M.C.S. conceived and designed the study. **S.M.**, E.J., and M.C.S. performed the field work. **S.M.** performed all behavioral experiments and analyzed the data. **S.M.** and A.E. performed the Ca<sup>2+</sup> imaging. V.R. and J.P. conducted the genetic analysis. M.C.S. and **S.M.** wrote the manuscript, with input from all authors.

# Popular science summary

You have most likely seen the tiny pesky flies sitting on your fruit basket or flying around in your kitchen, especially during late summer. Have you ever wondered from where these flies come from and how they find their way to your fruits? These insects, known as vinegar flies or fruit flies, use all kinds of overripe and fermenting fruit – rich in yeast and sugar – for feeding and breeding. Vinegar flies, similar to other insects, rely on their strong sense of smell (olfaction) to find their way to fruits, to find suitable partners for mating and good breeding spots. Vinegar flies sense odor molecules in the environment by their organs of smell, the antenna and maxillary palps, which have a similar function to the human nose. The fly “nose”, like the human nose, is equipped with dedicated receptors that capture odor molecules. These receptors do not detect odor molecule randomly, but capture odors associated with needs and dangers. For example, the fly has several smell receptors dedicated to odors coming from alcoholic fermentation and activation of these receptors guides the fly to suitable food sources rich in sugar and yeast (**Paper I**). Flies are also able to avoid toxic molecules and harmful matters by their sensitive “nose”. Vinegar flies avoid carnivore feces as a breeding site because of the smell of phenol, which is produced by pathogenic bacteria presence in carnivore feces. Phenol accordingly alarms vinegar flies about the looming danger of pathogenic bacteria (**Paper II**).

The fly “nose” has, however, more tricks up its sleeve. Flies, like other insects, are in high risk of dehydration due to their small body size and small capacity to store water, therefore, flies have an innate ability to find suitable humidity levels. Flies can sense humidity changes in their environment through humidity receptors, located in a invagination of the antenna. Humidity information is then processed together with thermal information in a dedicated region in the fly brain (**Paper III**).

The vinegar fly is today a cosmopolitan species, which means you can find them on all continents. However, the fly originated within southern central Africa (including Zambia and Zimbabwe). Although flies are known as generalists, wild populations of flies from Zimbabwe have a strong breeding preference towards marula fruit. This fruit is seasonally abundant, native to Southern Africa, and is presumably the ancestral host of the fruit fly (**Paper IV**).

This pesky fly has been a superhero in biological science for decades and has contributed to an astonishing amount of knowledge in genetics, neuroscience, embryology and modern biomedical research. The findings of my thesis further enhances the basic science behind olfaction and its mechanisms, the wild behavior of flies, and introduces the genetic and neural basis underlying humidity sensation in insects for the first time. These basic understandings may lead us to better strategies for controlling insect pests, as well as human disease vectors such as mosquitoes.

# Populärvetenskaplig sammanfattning

Du har säkert sett de små irriterande flugorna sitta i din fruktskål och flyga runt i köket, särskilt på sensommaren. Har du någonsin undrat varifrån dessa flugor kommer och hur de hittar till din frukt? Dessa insekter, vanligen kallade bananflugor eller vinägerflugor, använder all slags övermogen eller ruttnande frukt – rik på jäst och socker – till föda och för att föda upp larver.

Bananflugor är, liksom andra insekter, beroende av sitt känsliga luktsinne för att hitta föda, lämpliga parningspartners och bra ställen att lägga ägg på så larverna får tillgång till mat. Bananflugor känner doftmolekyler i omgivningen med hjälp av sina luktorgan, nämligen antennerna och maxillarpalperna, som har samma funktion som våra näsor.

Flugans luktorgan är liksom människans näsa utrustad med dedikerade receptorer som fångar upp doftmolekylerna. Dessa receptorer detekterar inte doftmolekylerna slumpmässigt, utan fångar upp dofter associerade med behov och med faror. Exempelvis har flugan flera luktreceptorer dedikerade till dofter från alkoholjäsning och aktivering av dessa receptorer leder flugan till lämpliga födokällor rika på socker och jäst (**Artikel I**).

Bananflugorna kan också undvika giftiga molekyler och skadliga ämnen genom sina känsliga luktreceptorer. De undviker köttätande rovdjurs exkrement som ägglägningsställen eftersom det luktar fenol, som produceras av patogena bakterier i exkrementerna. Tack vare fenolen varnas flugorna för den hotande faran med patogena bakterier (**Artikel II**).

Bananflugan har fler ess i rockärmen. Liksom andra insekter löper den stor risk för uttorkning beroende på liten kroppsstorlek och låg kapacitet att lagra vatten. Därför har flugorna en medfödd förmåga att hitta lämplig nivå på den relativa luftfuktigheten. Detta är ytterligare en orsak till att kök är en favoritplats för flugorna. De föredrar en relativ luftfuktighet på omkring 70%, vilket är precis vad kök ofta håller. Bananflugorna kan registrera förändringar i den relativa luftfuktigheten med hjälp av fuktreceptorer, vilka är belägna i en räfflad del av antennen. Den registrerade informationen bearbetas sedan tillsammans med temperaturinformation i en dedikerad del av hjärnan (**Artikel III**).

Bananflugan är idag en kosmopolitisk art, vilket betyder att den återfinns på alla kontinenter. Ursprungligen kommer den emellertid från de södra delarna av



centrala Afrika (exempelvis Zambia och Zimbabwe). Även om bananflugan är känd som en generalist globalt har vilda populationer av arten i Zimbabwe en stark förkärlek för marulafrukter när det gäller att föda upp sina larver. Marula är ett vanligt inhemskt träd i hela södra Afrika och dess frukter finns i överflöd när det är säsong. Förmodligen är marula den ursprungliga värdväxten för bananflugan (**Artikel IV**).

Den lilla irriterande bananflugan har visat sig vara en superhjärte inom biologi under flera decennier och har bidragit till en förbluffande mängd ny kunskap inom genetik, neurovetenskaplig embryologi och modern biomedicinsk forskning.

Resultaten i min avhandling ökar också de grundläggande kunskaperna bakom luktsinnet och dess mekanismer, beteendet hos vilda populationer av bananflugan och påvisar för första gången den genetiska och neurala bakomliggande basen för luktsinnet hos insekter. Dessa grundläggande kunskaper kan komma att leda till bättre bekämpningsstrategier mot skadeinsekter och mot smittbärande insekter som t.ex. myggor.

# Persian popular summary

## چکیده:

حشرات کوچک مزاحمی که بر روی سید میوه‌های شما می‌نشینند یا درون آشپزخانه تان پرسه می‌زنند، بله! بی‌شک با آن‌ها به ویژه هنگام پایان تابستان مواجه شده‌اید. آیا تا کنون اندیشیده‌اید که این حشرات از کجا آمده‌اند و چگونه راه خود را برای دسترسی به میوه‌های شما پیدا کرده‌اند؟

این حشرات به مگس سرکه مشهورند و از انواع میوه‌های رسیده یا در حال فساد که سرشار از قند و مخمر هستند، تغذیه می‌کنند. مگس‌های سرکه همانند سایر حشرات جهت رسیدن به میوه‌ها و یافتن جفت برای تولید مثل و همچنین پیدا کردن محل مناسب تخم‌ریزی، به سیستم بویایی قدرتمند خود متکی هستند.

مگس‌های سرکه ملکول‌های بوی موجود در محیط را از طریق اندام‌های بویایی خود که همان شاخک‌ها و پالپ‌ها هستند، تشخیص می‌دهند. به بیانی دیگر می‌توان این اندام‌ها را با بینی انسان مقایسه کرد.

“بینی” حشره مانند بینی انسان به گیرنده‌های اختصاصی مجهز شده که به یاری آن‌ها ملکول‌های بوی موجود در محیط را دریافت می‌کند. اما نکته‌ی بارز درخصوص این گیرنده‌ها این است که آن‌ها این بوها را به صورت تصادفی جذب نمی‌کنند بلکه بوهای مرتبط با نیاز و بقاء، توسط حشره دریافت می‌شوند. برای نمونه بینی حشره به چندین گیرنده‌ی بویایی ویژه مجهز است که برای تشخیص رایحه‌ی پراکنده شده از فرایند تخمیر الکلی تخصص یافته‌اند و با فعال شدن این گیرنده‌ها حشره راه خود را به سوی منابع غذایی مناسب و سرشار از قند و مخمر پیدا می‌کند. (نگاشته‌ی ۱)

حشره‌ها همچنین قادر به دوری از ملکول‌های سمی و مواد مضر به وسیله گیرنده‌های بویایی خود هستند.

مگس‌های سرکه از مدفوع جانوران گوشتخوار جهت تخم‌ریزی دوری می‌کنند، دلیل این امر وجود فنول بوده که حاصل فعالیت باکتری‌های پاتوژن موجود در مدفوع این جانوران است. فنول برای مگس

سرکه به معنی هشدار می‌کند که نمایانگر وجود باکتری پاتوژن، درون مدفوع است. (نگاشته‌ی ۲)

”بینی“ حشره همچنان شگفتی‌های دیگری دارد. مگس‌ها مانند سایر حشرات به دلیل اندازه‌ی کوچک و توان کم در ذخیره‌ی آب، همواره در معرض خطر از دست دادن آب بدن و مرگ قرار دارند. از این رو حشرات توانایی ذاتی در یافتن سطوح رطوبتی مناسب را دارند. مگس‌ها قادرند توسط گیرنده‌های رطوبتی که درون حفره‌ای در شاخک‌ها قرار گرفته، تغییرات رطوبت محیط را تشخیص دهند. سپس اطلاعات مربوط به رطوبت همراه با اطلاعات مرتبط با دمای محیط در بخشی از مغز مگس سرکه که برای این امر تخصص یافته، تحلیل می‌شوند. (نگاشته‌ی ۳)

امروزه مگس سرکه گونه‌ای همه جایی است. بدین معنی که در تمام قاره‌ها یافت می‌شود، هر چند که موطن اصلی آن‌ها جنوب آفریقای مرکزی (نواحی زامبیا و زیمبابوه) است.

گرچه این مگس‌ها گیاهخوار عمومی هستند ولی جمعیت‌های وحشی آن‌ها در زیمبابوه گرایش شدیدی برای تخم‌ریزی روی میوه مارولا دارند. این میوه در فصل بهار به فراوانی یافت می‌شود و بومی جنوب آفریقا است. تفکر و تحلیل در این باب می‌تواند مبتنی بر این ادعای مهم باشد که سرآغاز مسیر تکاملی مگس سرکه جنوب آفریقا، و مارولا اولین میزبان این حشره بوده است. (نگاشته‌ی ۴)

این حشره‌ی مزاحم در طی دهه‌ها در بستر علوم زیستی همواره یک ابر قهرمان بوده به گونه‌ای که همواره جزیی از دستاوردهای شگرف در علم ژنتیک، عصب‌شناسی، رویان‌شناسی و تحقیقات نوین پزشکی بوده است.

یافته‌های پایان نامه‌ی من موجب ارتقاء دانش پیشین در مورد سیستم بویایی و نحوه عملکرد آن و همچنین برای نخستین بار منتج به معرفی ساز و کار ژنتیکی و عصبی گیرنده‌های حس رطوبت در حشرات شده است.

امید است این یافته‌های بنیادین ما را به سمت تدابیر بهتر جهت کنترل حشرات آفت و همچنین مدیریت ناقلین بیماری‌های انسانی مانند پشه‌ها رهنمون سازد.

# The Scope of this Thesis

Animals, like humans, need to perceive their surroundings via their senses in order to make sensible behavioral decisions, reproduce successfully, and survive. Most animals do not only have more sensitive senses in comparison with humans but have also evolved more sensory systems enabling them to cope with special conditions in their natural habitat. Animals are equipped with audition, vision, thermosensation, humidity sensation (hygrosensation), mechanosensation, magnetoception, gustation, and olfaction which detects physical and chemical changes in their habitats. Among these senses, olfaction is likely the most ancient sensory modality. Insects, the most abundant and successful group of the animal kingdom, predominantly use olfaction to find food, mates, breeding sites, and to avoid dangers. Moreover, hygrosensation is vital for insects to find a suitable habitat and to avoid risks of dehydration.

Our understanding of the molecular, neuronal, and morphological organization of the insect olfactory system is today substantial, in large parts thanks to *Drosophila melanogaster* and the wealth of sophisticated genetic tools available in this classic model system. Our knowledge regarding the functional and molecular basis of insect hygrosensation, is, however, limited.

The aim of this thesis is to investigate how the olfactory system of *Drosophila* translates sensory inputs into behavioral responses (Paper I and II). Moreover, I have also studied the organization of the hygrosensory system and how *Drosophila* senses changes in humidity levels (Paper III).

Every biologist wonders about the origin of the creatures of earth and origin of life. This central question stimulated another aim in this thesis, the search of wild *Drosophila melanogaster* and its ancestral habitat (Paper IV).

This thesis begins with an overview of the fly and its astonishing contribution to the field of genetics and biology. Then the organization and function of the olfactory and hygrosensory systems are described. The thesis continues with the results and discussions from the four papers dealing with aforementioned aims. Finally, a conclusion and contributions to the field along with some ideas for future studies are described.



# Background of the study

## *Drosophila melanogaster* – Lord of the flies

*Drosophila melanogaster* or the vinegar fly (**Figure 1A**) has been one of the primary laboratory model organisms for over a century. The use of the fly as a genetic model organism goes back to the beginning of the 1900s with the work of Thomas Hunt Morgan (**Figure 1B**) at Columbia University, where he and his students were searching for a suitable animal model to study the laws of heredity. Morgan settled on the fly, and soon isolated a white-eyed mutant (wild-type flies have red eyes), and with the help of this sex-linked mutant he could show that the chromosomes carries the hereditary material (Morgan, 1910). Morgan was awarded the Nobel Prize in physiology or medicine for this discovery in 1933. Apart from this important discovery, Morgan together with Calvin Bridges, Alfred H Sturtevant, and Herman J Muller established most of the major principle of classic genetics such as the nature of genetic linkage and genetic maps, the genetic behavior of chromosome aberrations, the induction of gene and chromosome mutations by radiation, and the discovery of mitotic recombination (Bridges, 1921; Sturtevant, 1965). Calvin Bridges experimental evidence of genes and chromosomes linkage and the term “nondisjunction” was published as the first paper of the first issue of the journal *Genetics* (Bridges, 1916). Herman J Muller won the Nobel Prize in 1946 for using X-ray radiation to induce mutations in the fly (Muller, 1927). Since then, research using *D. melanogaster* has yielded astounding insights into processes of cell and molecular biology, neurobiology, embryonic development, learning and memory, behavior, and physiology. Fly research has received six Nobel Prizes to date, with the most recent one in 2017 for the discoveries concerning the molecular mechanism underlying circadian rhythms (Bargiello and Young, 1984; Bargiello et al., 1984; Zehring et al., 1984; Siwicki et al., 1988; Hardin et al., 1990).



**Figure 1. *Drosophila melanogaster*, Lord of the flies.**

A) A female *Drosophila melanogaster* perched on one of its favorite fruits, orange. Photo: Marcus Stensmyr B) Thomas Hunt Morgan (Columbia University, New York, USA), who introduced the fly as a genetic model organism. Photo: Wikimedia.

Today, the whole genome of *D. melanogaster* together with numerous tools to manipulate the genome is available, which allow for e.g. functional dissections of specific genes, as well as neuronal circuits (Adams et al., 2000). Therefore, mutant flies can be engineered through selective removal or replacement of gene sequences (Rong et al., 2002), the expression level of any gene can be reduced using RNA interference (Dietzl et al., 2007; Ni et al., 2009), any gene can be expressed in almost any tissue or cell using the yeast Gal4-UAS system (Brand and Perrimon, 1993; Pfeiffer et al., 2008). It is also possible to visualize the expression of genes of interest and monitor neural activity by introducing reporter genes, such as calcium indicators and green fluorescent proteins via the UAS system (Nakai et al., 2001; Marella et al., 2006). Moreover, with the UAS system one can also introduce genes that can modify neuronal activity, such as the tetanus-toxin light chain protein (TNT), which blocks synaptic transmission (Sweeney et al., 1995), or shibire that leads to depletion of vesicles in the synaptic termini (Kitamoto, 2001), or even light sensitive proteins, such as CsChrimson, which is a red-shifted channelrhodopsin enabling light activation of neurons (Klapoetke et al., 2014). With the aid of these tools, scientists have even succeeded in studying genes that are involved in neurodegenerative human diseases and their underlying mechanism such as Alzheimer and Parkinson (Lu and Vogel, 2009) in the fly.

*D. melanogaster* is an opportunistic human commensal species with a cosmopolitan distribution. This cosmopolitan species has originated within sub-Saharan Africa as evident from biogeography and genetic variation studies (Lachaise et al., 1988; Pool and Aquadro, 2006). A recent genomic study revealed that southern central African populations (e.g. Zambia and Zimbabwe) have the highest genetic diversity, suggesting that the ancestral range of *D. melanogaster* may be located within this area (**Figure 2A**) (Pool et al., 2006 and 2012). Most of Zambia and Zimbabwe are covered by seasonally dry miombo and mopane

woodlands, which have subtropical climate with a hot dry season from August through October and a hot wet season from November to March. Interestingly, African *D. melanogaster* in comparison with its close relatives has higher tolerance to dehydration and severe high temperatures (Stanley et al., 1980) that would fit its ancestral subtropical climate conditions.


*D. melanogaster* is considered a generalist, and can utilize a wide range of decaying vegetables, fruits and other plant matter as feeding and breeding substrate. Knowledge about the breeding substrates in the wild natural habitat of the fly is scant, at best, since essentially all reported populations even within Africa have been caught from sites within or near human settlements (Lachaise and Silvain, 2004). *D. melanogaster*, however, could maintain permanent populations in wild habitats after human activity has diminished and produce wild non-commensal generations. Lachaise (1988) reported 25 host plant species used as larval breeding resource by *D. melanogaster* in the Afrotropical region, e.g. *Lobelia* inflorescences in montane forests or *Pistia stratiotes* (water salad) from Kivu in Congo (Lachaise et al., 1988). 16 out of 25 identified host plants were native, which is high ratio for a domestic species like *D. melanogaster*.

Although vinegar flies are known as a generalist, they show strong egg-laying preference towards *Citrus* spp. – especially orange – over other type of fruits (Dweck et al, 2013). *Citrus* spp. originated in South-east Asia, hence, this fruit cannot be the hosts with which the fly has evolved. The ancestral host fruit of *D. melanogaster*, along with the history of this species evolution to a human commensal species is accordingly unknown.

## The lord's close relatives

*D. melanogaster* belongs to the melanogaster species group of the *Sophophora* subgenus (or genus pending whom you ask). The *melanogaster* group is geographically widespread and contains over 170 species (Schawaroch, 2002). The *melanogaster* species subgroup comprises nine closely related species, occupying different ecological habitats and has probably evolved 13-15 million years ago (**Figure 2B**). Species within this subgroup are ranging from single host specialists to generalists (Lachaise et al., 2000). *Drosophila sechellia* is the most highly specialized member of this subgroup, and preferentially lay eggs on noni fruit (*Morinda citrifolia*), a fruit native to the Seychelles islands (**Figure 2C**). *Morinda* fruit, which ripens throughout the year, is toxic, or at least not palatable, to other members of the *melanogaster* subgroup because of its high content of octanoic and hexanoic acid (Lachaise et al., 1988; Dekker et al., 2006).





**Figure 2. *Drosophila melanogaster* and its close relatives.**

A) Putative ancestral range of *D. melanogaster* showed in the dashed circle. B) Phylogeny of the nine related species in the *melanogaster* species subgroup. C) Noni fruit (*Morinda citrifolia*), host of *D. sechellia* Photo: E. Guinther. D) *Pandanus* cone, seasonal host of *D. erecta*. Photo: B. Navez.

*Drosophila erecta* and *Drosophila orena* are the two known seasonal specialist of this subgroup. *D. erecta*, which is endemic to gallery forests of West-central Africa, is a seasonal specialist on ripe fruits of *Pandanus* spp. (**Figure 2D**). *Pandanus* cones ripen once a year for two months, during which time *D. erecta* is exclusively found on this fruit. While *D. orena* for a long time was only known from a single location on a sole mountain in Cameroon (Tsacas and David, 1978), a recent discovery of another population on the West African island of Bioko revealed that the species has a close, but seasonal association with waterberrys *Syzygium staudtii* (Comeault et al., 2017).


On the other hand, *Drosophila simulans* and *D. melanogaster* are the only two cosmopolitan species of this species subgroup. As opportunistic human commensals, these two species have effectively spread out in the world by human transportation (David and Capy, 1988). *D. simulans*, however, is less associated with human activities and does not readily enter buildings, therefore, it is not found in all places in the world (Rouault and David, 1982). *D. melanogaster* is more tolerant to broader temperature variation and is accordingly more abundant in temperate regions than *D. simulans*. These two species are so similar morphologically that *D. simulans* was initially described as *D. melanogaster* in the beginning of the 20<sup>th</sup> century (Capy and Gibert, 2004).

# Olfaction – Lord of the senses

Animals in their natural habitat are surrounded by volatile chemicals, some of these chemicals will convey important information to the perceiving animal regarding e.g. presence of food, sexual partners, and potential dangers. Most if not all animals are also equipped with peripheral detection systems for volatile chemicals, and dedicated brain centers, which translate the detected volatiles, or odors, into appropriate behaviors. The whole procedure of these transformations is called olfaction. Olfaction is probably the most ancient sensory modality in the animal kingdom (Strausfeld and Hildebrand, 1999). Over 2000 years ago, Aristotle noticed that animals rely more on their sense of smell for their survival than humans do, however, the principle and function of the olfactory system in general, is quite similar in all organisms ranging from a little fly to a large mammal.

## The peripheral olfactory system of the fly

Flies are able to detect an array of volatile chemicals via two pairs of olfactory appendages on the head, the antennae and maxillary palps (**Figure 3A**). These organs are covered by sensory hairs, porous cuticular structures called sensilla (**Figure 3B**). Each sensillum houses dendrites of olfactory sensory neurons (OSNs), which express a distinct olfactory chemoreceptor (**Figure 3C**). These chemoreceptors come in three flavors, odorant receptors (OR), ionotropic receptors (IR) and gustatory receptors (GR), with each capturing specific environmental volatiles.



**Figure 3. The peripheral olfactory system of the adult fly.**

A) Head of the fly. B) Different types of sensilla (sensory hairs) are covering the two olfactory organs. C) Each sensilla house dendrites of 1-4 OSNs and axon of these OSNs projects to the antenna lobe.

The antennal olfactory sensilla of the fly are divided into four different morphological and functional types: basiconic, trichoid, intermediate, and coeloconic sensilla (**Figure 3B**). Each of these sensilla houses one to four OSNs. The maxillary palp has one type of basiconic sensilla (**Figure 3B**), each housing two OSNs (Vosshall and Stocker, 2007). Antennal sensilla of different morphological types have special functions to some degree; OSNs housed in basiconic sensilla are tuned to fermentation, microbial, and plant volatiles, whereas coeloconic OSNs primarily detect amines and acids (Benton et al., 2009; Yao et al., 2005). Trichoid OSNs are mainly responding to pheromones, such as (Z)-11-cis-vaccenylacetate (cVA), which mediates courtship behaviors via activation of specific trichoid neurons (Clyne et al., 1997; Ha and Smith, 2006; Kurtovic et al., 2007). The function of intermediate sensilla remains somewhat unclear, but at least one type of OSN housed in this type of sensilla is involved in oviposition site selection (Dweck et al., 2013). Most of the maxillary palps' OSNs detect only one single chemical and mediate short and long range attraction to specific chemicals such as 4-ethylguaiaicol, furaneol and methylether (Dweck et al., 2016). In total there are 1100-1250 OSNs on the antennae (Stocker, 2001) and 120 on the maxillary palps, which in total are expressing 62 ORs, 16 IRs and 12-14 GRs (Fishilevitch et al., 2005; Couto et al., 2005; Hallem et al., 2004; Benton et al., 2009; Menuz et al, 2014).

### **The larval chemosensory system**

The structure of the adult and larval olfactory pathways is quite similar even though the number of OSNs is less in larvae in comparison with adults (21 OSNs in larvae (Cobb, 1999). The main larval chemosensory organs are located in the cephalic lobe, and include the dorsal organ, ventral organ and terminal organ (Singh and Singh, 1984). There are three other organs located on the thoracic and abdominal segments, which are probably involved in contact chemoreception (Scott et al., 2001). The dorsal organ expresses ORs and GRs and acts as both an olfactory and gustatory organ, while the ventral and terminal organs only have a taste function (Singh and Singh, 1984; Singh, 1997). OSNs on the dorsal organ respond to a wide variety of chemicals, such as alcohols, acetates, aldehydes, esters, and ketones (Cobb et al., 1992; Cobb, 1999; Heimbeck et al., 1999; Cobb and Domain, 2000; Boyle and Cobb, 2005). The olfactory response of larvae is age dependent; with stronger responses to chemical stimuli observed by increased age from first instar larvae to third instar, however, late third instar larvae shows very weak olfactory responses (Kaiser and Cobb, 2008). The olfactory information detected at the peripheral level, converge into the larval antennal lobe (LAL) (Tissot et al., 1997).

## Odorant receptors

Odorant receptors (ORs) are the most prominent component of the olfactory pathway. Genes encoding olfactory receptors were first discovered in the rat by Buck and Axel in a 1991 milestone study (Buck and Axel, 1991), which led to the 2004 Nobel Prize in medicine or physiology. The olfactory receptor genes in vertebrates (and nematodes) belong to the family of seven transmembrane G-protein-coupled receptors (GPCRs). In 1999, with the identification of OR genes in *D. melanogaster* (Clyne et al., 1999; Gao and Chess, 1999; Vosshall et al., 1999), brought up an important distinction between the mammalian and insect olfactory systems, namely that the *Drosophila* OR genes showed no sequence similarity to the GPCRs of vertebrates and nematodes, but form a unique family. The insect ORs has moreover, a reversed transmembrane topology to that of GPCRs (Benton, 2006). The number of ORs varies between species, e.g. from 47 in the cotton leafworm (*Spodoptera littoralis*) (Poivet et al., 2013) to around 400 in the Indian jumping ants (*Harpegnathos saltator*) (Zhou et al., 2012), whereas mammalian genomes contain around 250 ORs in platypus to over 2000 in the African elephant (Niimura and Nei, 2007; Niimura et al., 2014).

The odorant receptor co-receptor, Orco, is highly conserved among insects, showing over 90% sequence homology between closely related species (Vosshall et al., 1999). Orco is co-expressed with the conventional receptor in both adults and larvae (Fig. 3b). Orco is involved in dendritic localization and critical for the function of the tuning ORs, but does itself not directly respond to odorants (Larsson et al., 2004). Orthologs of Orco from other animals than insects has not been reported yet.

Individual ORs has been examined using loss-of-function studies. Flies lacking specific ORs show a reduction in both behavioral and electrophysiological responses to distinct odors (Störtkuhl and Kettler, 2001; Dobritsa et al., 2003; Jones et al., 2007; Kreher et al., 2008; Kurtovic et al., 2007; Semmelhack and Wang, 2009). For example, mutant flies lacking a functional *Or67d* receptor revealed that the neurons in which this receptor is expressed regulate mating behavior. Both female and male *Or67d* mutant flies show defective mating behavior, including male-male courting (Kurtovic et al., 2007).

In *Drosophila*, like mammals, each OSN typically expresses a single OR gene apart from Orco, although, there are six classes of OSNs that express more than one OR. In four of these classes, the co-expressed ORs are closely linked and are probably the result of a recent gene duplication. An example of this case is found in the so-called ab3A OSNs, which co-express *Or22a* and *Or22b* (Hallem et al., 2004; Couto et al., 2005), but where only *Or22a* appears to confer odor sensitivity (Dobritsa and et al., 2003).

## **Ionotropic receptors**

Another family of insect olfactory receptors are the IRs, which belongs to the highly divergent family of Ionotropic Glutamate Receptors (iGluR) (Benton et al., 2009). IRs do not only have a role in olfaction but also act as sensory receptors for humidity, temperature, and taste. *Drosophila* adults express at least 57 IR genes. At least 16 of these are expressed in the antenna (Benton et al., 2009; Croset et al., 2010). IRs expressed in antennal neurons are found in the arista (a feather-like protrusion from the third antennal segment), the sacculus (a three-chambered pit on the third segment), and in neurons housed in coeloconic sensilla. Up to three IRs are usually co-expressed in one sensory neuron together with Ir8a, Ir25a, or Ir76b, which are broadly co-expressed and likely act as co-receptors, required for proper IR function (Benton et al., 2009; Abuin et al., 2011; Hussain et al., 2016). The ligands of antennal IRs differ from ORs. For instance, Ir8a positive neurons in coeloconic sensilla respond mainly to carboxylic acids and aldehydes, while Ir25a or Ir76b expressing neurons predominantly detects amines (Silbering et al., 2011). For example, Ir41a is co-expressed with Ir76b. Activation of these two IRs enables the fly to detect polyamine volatiles from a long range, and induces egg-laying (Hussain et al., 2016).

In the first two chambers of the sacculus, combinations of Ir25a, Ir40a, Ir68a, and Ir93a are expressed, which are all required for humidity sensing in the fly (Enjin et al., 2016; Frank et al., 2017; Knecht et al., 2016 and 2017). In the third chamber of the sacculus, Ir64a is expressed with Ir8a as a co-receptor and mediate acid sensing (Ai et al., 2010). In the arista, Ir21a is expressed with Ir25a where they mediate cool sensing (Ni et al., 2016). Ir21a and Ir25a have a similar function in larvae (Ni et al., 2016). Ir25a and Ir93a are also expressed in the thermosensory cells of the arista (Benton et al., 2009).

IRs are also expressed on the proboscis, pharynx and legs of adults and on the taste organ of the larvae. Ir76b is not only expressed in OSNs but also in gustatory receptor neurons (GRNs). Activation of Ir76b together with Gr66a in the taste neurons, helps the fly to evaluate polyamine presence in decaying fruits (Hussain et al., 2016). IRs expressed in the male foreleg (Ir52c and Ir52d) become activated in contact with the female and mediate courtship in male flies (Koh et al., 2014). In summary, IRs perform a number of different functions, most of which likely unrelated to olfaction.

## **Gustatory receptors**

GRs are expressed in sensory neurons on the mouthparts, maxillary palps, legs, wings, as well as the ovipositor (Stocker 1994; Vosshall and Stocker 2007; Ling et al., 2014; Depetris-Chauvin et al., 2015). A few are also expressed in OSNs on the

antennae (Jones et al., 2007; Kwon et al., 2007). The GR genes in insects have no sequence similarity to mammalian taste receptors, but are closely related to the insect ORs. Around 12-14 GRs are expressed in OSNs on the antenna (Menuz et al., 2014), including Gr21a and Gr63a, which together function as CO<sub>2</sub> detectors (Jones et al., 2007; Kwon et al., 2007). Another GR with high expression on the antenna is Gr10a, which is expressed in neurons in a large basiconic sensillum, but has unknown function (Hallem et al., 2004). The remaining GRs expressed on the antenna are presumably sugar detectors (Dahanukar et al., 2007).


GRs expressed on the labellum are mainly sweet and bitter detectors e.g. Gr5a, which together with a subset of related receptors detect sugar and trigger feeding, whereas, Gr66a and related genes, confer responses to bitter compounds and causes aversive behavior (Dahanukar et al., 2001; Chyb et al., 2003; Weiss et al., 2011). For example, flies detect and avoid caffeine via a combination of the bitter receptors Gr33a, Gr66a, and Gr93a (Moon et al., 2006; Moon et al., 2009; Lee et al., 2009). Male flies avoid courting other males by tasting (Z)-7-tricosene, a male sex pheromone and bitter stimuli via activation of Gr66a on the labellum (Lacaille et al., 2007).

Around 39 GRs are expressed in the larval taste organs, and most of them are supposed to be bitter receptors (Colomb et al., 2007; Kwon et al., 2011). Gr21 and Gr63 are also expressed in larvae and activation of these receptors cause aversive behavior in larvae as well (Jones et al., 2007; Kwon et al., 2007).

Gustatory sensory neurons (GSNs) from proboscis, mouth parts, and legs transmit gustatory information to the subesophageal ganglion (SOG), a dedicated taste center in the brain (Wang et al., 2004).

## **The central olfactory system of the fly**

The OSNs, after detecting and discriminating odor molecules, send their axons to the brain. In the antennal lobes (AL), these axons coalesce and form glomeruli, which are spheroidal structures (**Figure 4**). The AL is the insect counterpart to the mammalian olfactory bulb. Like olfactory bulbs, ALs pre-process and integrate olfactory information (Strausfeld and Hildebrand, 1999; Ache and Young, 2005). Each AL of the fly consists of 49 glomeruli. Each single class of OSNs expressing the same OR genes projects to a distinct glomerulus in the ALs, where axonal branches synapse with dendrites of the corresponding class of projection neurons (PNs) (Vosshall et al., 2000; Goa et al., 2000; Bhalerao et al., 2003). PNs can be uniglomerular (branch in one glomerulus) or multiglomerular (branch in many glomeruli) (Galizia, 2014). The glomeruli of the AL are also innervated by local interneurons (LNs), which regulate the activities of PNs and assist communication



**Figure 4. The *Drosophila* brain.**

Axons of OSNs project to the antennal lobe (AL), which is the primary olfactory center in the fly brain (dashed yellow circles). Projection neurons transfer the integrated olfactory information from the ALs to the mushroom body (MB) and terminate in the lateral horn (LH).

within and between glomeruli (Stocker et al., 1990; Wilson and Laurent, 2005; Seki et al., 2010; Wilson, 2013).

The axons from uniglomerular excitatory PNs relay olfactory information to the mushroom body (MB) (**Figure 4**), a center for olfactory learning and memory (Davis, 2005; Heisenberg, 2003), and to the lateral horn (LH), a less-understood higher-order center presumed to direct olfaction-mediated innate behaviors (Heimbeck et al., 2001). MBs are also involved in locomotor activity (Martin et al. 1998), male courtship behavior (Sakai and Kitamoto, 2006), and sleep regulation (Joiner et al., 2006; Pitman et al. 2006).

### **Antenna - a multifunctional sensory appendage**

The insect antenna is a multifunctional sensory appendage that does not only detect volatile chemicals, but it is also involved in a number of other tasks, such as gustation, tactile sensation, thermosensation and hygro-sensation (Schneider, 1964 ; Schafer and Sanchez, 1973; Norris and Chu 1974 ; Toh, 1977 ; Schaller, 1978; Altner and Prillinger, 1980 ; Chapman, 1982; Keil and Steinbrecht, 1984 ; Steinbrecht 1984; Lee and Strausfeld 1990). For instance, the nocturnal cockroach determines objects in the dark by the tactile sense on the antenna (Okada and Toh,

2000 and 2006; Okada et al., 2002). Cockroaches can also detect changes in the temperature by cold and hot receptor neurons on the antenna inhibited by heat and excited by cold (Yokohari, 1999; Mizunami et al., 2016). Moreover, some of the antennal sensilla of cockroaches carry moist and dry detecting neurons that respond to the changes in humidity as well as changes in air pressure (Tichy and Kallina, 2010). However, the molecular basis and the neural mechanism underlying hygrosensation is not well understood.

## Hygrosensation - The sixth sense

Humidity is a major environmental factor affecting animal health, their reproductive success, and their geographic distribution (Shelford, 1918). Small-bodied insects have a small storage capacity for water and a large surface area for losing it (Kuhnel et al., 2016), they are therefore more sensitive to humidity variation in comparison with mammals and birds (Ludwig, 1945). To avoid rapid dehydration, the insects body is covered by a layer of wax and long-chain cuticular hydrocarbons (CHs). A recent study on *Drosophila* showed that the proportion of certain cuticular hydrocarbons (CHs) enables *Drosophila* to resist dehydration (Ferveur et al., 2018) and a gene named *dsat1*, which regulate CHs production in the fly, is involved in dehydration resistance (Ferveur et al., 2018).

More notably, unlike mammals and worms, insects have dedicated hygrosensory receptor neurons (HRNs). Insects use humidity cues to locate oviposition sites, identify hosts and nectar-bearing flowers, and to find a suitable habitat based on humidity range. For example the female mosquito *Anopheles gambiae* – the vector of malaria – use humidity cues to find stagnant water to lay her eggs (Okal et al., 2013). Moreover, the humid and warm breath of animals along with olfactory and vision cues guides *A. gambiae* and other hematophagous species to favorable hosts for biting and feeding (van Breugel et al., 2015). The hummingbird moth *Hyles lineata* likely identifies flowers abundant in nectar based on humidity cues (von Arx et al., 2012).

### Functional anatomy of hygrosensation in insects

Studies of honeybees, locusts, cockroaches and stick insects revealed that their antennae are equipped with a small number of hygrosensory sensilla (Altner and Loftus, 1985). Unlike olfactory sensilla, the hygrosensory sensilla are poreless, but some carry a plug on the apical surface, called a molten pore, the function of which is unknown. Hygrosensory sensilla are in most insects located in an invagination of the antenna where they are protected from wind, unlike the



olfactory sensilla, which are in a position with highest exposure to the surrounding air (Tichy and Loftus, 1996; Enjin, 2017). The hygrosensory sensilla house the HRNs, reported for the first time from the honeybee *Apis mellifera* (Lacher, 1964). The layout of hygrosensory sensilla is similar between most insect species; one moist neuron together with one dry neuron and one hygrocool neuron are located in a single hygro-sensillum forming a triad (Altner and Loftus, 1985).


# Outcome and discussion of the studies

## I. The chemical ecology of the fly

Chemical compounds are involved in most interactions between animals and their environment. Chemical ecology is the study of these interactions from a chemical, genetic, neural, behavioral, and ecological perspective. This fascinating field was commenced when the sex pheromone of the silk moth *Bombyx mori* was discovered (Karlson and Butenandt, 1959). Since then, we have gained a fairly detailed understanding about the ecological relevance of insect's chemical interactions with their natural habitat. In **Paper I** of this thesis, I review the chemical ecology of *D. melanogaster* and discuss the ecologically relevant functions of many of the fly's ORs, IRs, and GRs (Mansourian and Stensmyr, 2015). I argue that the vinegar fly's receptors do not sample volatile chemicals randomly but are tuned to subsets of volatiles associated with its ecological needs by bringing up several examples in host seeking, oviposition, and avoiding toxic matters.

### Host seeking

Fermented fruit - rich in yeast and sugar – is the favorite food of *D. melanogaster* (Carson, 1971). In the process of fruit fermentation, yeast convert sugars into ethanol and CO<sub>2</sub>. This process also generates a large number of other volatiles, which are irresistible to *D. melanogaster*. These volatiles activates several receptors e.g. Ir31a that detects 2-oxopropionic acid and pyruvic acid generated from sucrose (Silbering et al., 2011), Gr21a/Gr63a detect CO<sub>2</sub> released from the decarboxylation of pyruvate into acetaldehyde (Jones et al., 2007), Or92a detects diacetyl formed from acetaldehyde, and Or42b detects acetal generated from the reaction of acetaldehyde with ethanol (Mathew et al., 2013). Activation of these receptors guide flies towards suitable food sources. Yeast is the major source of nutrition for adults and larvae of most *Drosophila* (Begon, 1982), and affects larval growth and survival (Starmer and Aberdeen, 1990). Yeast itself primarily releases two groups of volatiles, acetate esters and phenolics, which activate several olfactory receptors in the fly (Hallem and Carlson, 2004 and 2006; Stökl et al., 2010) (**Figure 5A, B**).



**Figure 5. Alcoholic fermentation and orang smells**

A) Volatiles emitted through the activity of yeast and fruit fermentation (e.g fermented apple) and B) the fly's receptors that detect fermented volatiles (from A) and volatiles from orange ( limonene and Valencene).

Fruit volatiles are also important for flies for host seeking and oviposition site selection. Even though *D. melanogaster* shows a preference towards *Citrus* spp. (Dweck et al., 2013), *D. melanogaster* is an opportunist and is able to utilize a wide range of fruit. Accordingly, their peripheral olfactory detectors are also equipped with receptors that are broadly tuned to common fruit volatiles such as Or43b, Or47a and Or85a, which detect fruity acetate esters, e.g. ethyl acetate, isoamyl acetate, and amyl acetate (Hallem and Carlson 2004, 2006). Or22a is another important broadly tuned receptor detecting fruit esters, primarily ethyl hexanoate (de Bruyne et al., 2001; Stensmyr et al., 2006).

## Volatile chemicals effecting oviposition


The fly larvae's mobility is limited, hence the selection of a suitable oviposition sites is a critical decision the female fly has to make. Olfaction guides the female in making the right decision in this matter. Examples of chemical compounds that trigger egg laying in the female are flavo terpenes, such as valencene and limonene released by *Citrus* spp and other fruit with a thick epicarp, which activates Or19a (Dweck et al., 2013). The thick epicarp protects the fly's larvae from endoparasitoid wasps (Dweck et al., 2013) (**Figure 5B**).

The microbial composition of the substrates is of critical importance in the oviposition site selection process. The smell of acetic acid, produced by acetobacteria during the fermentation process, activates Ir75a and induces egg-laying behavior (Joseph et al., 2009). *Brettanomyces* yeast on the surface of ripe fruit produces volatiles that stimulate oviposition, as well as feeding via activation of Or71a (Dweck et al., 2015), while the presence of harmful microbes on fermented fruit abolishes oviposition (Stensmyr et al., 2012). These harmful microbes release, among other compounds, a volatile called geosmin. Geosmin is detected by a single receptor, Or56a, and activation of this pathway alerts flies to the presence of e.g. *Penicillium* molds and triggers aversive behavior (Stensmyr et al., 2012).

## II. Foul feces fool flies

Feces is an abundant source of energy in nature, used by many organisms, not least by members of the order Diptera. Adult Mediterranean fruit flies (*Ceratitis capitata*) need to feed on animal fecal matter in order to develop their eggs (Hendrichs et al., 1993; Lauzon, 2003). Houseflies and blowflies feed and breed on animal feces (Amendt et al., 2004). In **Paper II** of this thesis, I study how *D. melanogaster* reacts to fecal matter.

I show that *D. melanogaster* displays aversive oviposition towards carnivore feces, while indifference or even attraction towards herbivore dung (**Figure 6A**). This aversive behavior is due to the presence of high amounts of a toxic volatile compound – phenol – specifically enriched in carnivore feces, and is detected by Or46a-expressing OSNs located on the maxillary palps (**Figure 6B**). Silencing this single neuronal population, through directed expression of TNT (Sweeney et al., 1995), abolishes aversion to phenol as well as carnivore feces (**Figure 6C, D**). I further demonstrate that phenol suppresses egg-laying in flies, even in an attractive odor background. This observation suggests that phenol itself affects both oviposition site selection and the oviposition rate in flies.



**Figure 6. Flies avoid carnivore feces and the bacterial volatile phenol**


A) Oviposition index (OI) of wild-type flies from a binary-choice test between standard cornmeal fly food and fly food mixed with mammalian feces. Error bars represent the SEM. Deviation of the response indices against zero was analyzed for significance with Student's t test ( $p < 0.05$ ). B) Expression pattern of *Or46a* in the maxillary palp. C) and D) OI of flies expressing TNT from the putative promoter of *Or46a* in a binary-choice assay between standard cornmeal fly food and fly food mixed with (C) phenol (10<sub>2</sub> dilution) or (D) lion feces.

## Phenol signals pathogenic bacteria to the fly

Detecting and avoiding pathogens before infection is naturally beneficial, because once an animal is infected, it has to pay a high metabolic price to mount immune responses and to recover (McKean and Lazzaro, 2011; Kominsky et al., 2010). *D. melanogaster* can detect bacterial molecules called lipopolysaccharides (LPS), which alerts the fly's immune system to the presence of dangerous bacteria (Abbas et al., 2014). Gr66a detects these bacterial molecules, which mediates aversive feeding and oviposition responses (Soldano et al., 2016). In **Paper II** of this thesis, I show that carnivore feces, unlike herbivore feces is enriched by a family of bacteria – Enterobacteriaceae – that contains a wide range of pathogenic taxa causing severe diseases in humans (**Figure 7A**) (Rossetto et al., 2014). Some of these bacteria are able to produce phenol under special circumstances; when these bacteria grow in substrates containing L-tyrosine, an enzyme called tyrosine phenol-lyase convert L-tyrosine to phenol, pyruvate, and ammonia (Kumagi and Yamada, 1970). Accordingly, bacteria in carnivore feces are able to produce phenol from L-tyrosine, whereas bacteria in herbivore feces, scarce in L-tyrosine,

do not (**Figure 7B**). I further demonstrate that a dung beetle, *Scarabaeus (Kheper) lamarcki*, avoids carnivore feces and phenol presumably for a similar reason (**Figure 7C**).

Why would a fruit-dwelling fly then have a receptor for a fecal chemical? Fecal matter generates a wide range of volatiles, several of which are released by fruits as well. For example, limonene is emitted from thick epicarp fruit and also from many fecal samples. Flies in the native Southeast African habitat face feces more frequently than fruit, especially during the dry season when fruit is scarce. In this situation, distinguishing feces that contains harmful bacteria from fruit is necessary and beneficial for flies.




**Figure 7. Dangerous dung**

A) Multidimensional scaling plot from a Random Forest analysis of bacterial content in herbivore and carnivore feces B) Schematic model of phenol production by bacteria presence in the carnivore feces, presenting (1) L-tyrosine, (2) bacteria, and (3) L-tyrosine-phenol lyase enzyme. C) Dung beetles avoid carnivore feces. From left to right, tracks from beetles provided with a choice between lion and giraffe feces, confronted with lion feces and sand in an open olfactory binary-choice arena.

### III. Humidity sensation in the fly


Insects must find a proper level of humidity for their survival, as they are in high risk of dehydration due to their low capacity to store water. Insects have evolved a sensory system to detect humidity changes in their environment, however, the molecular basis of humidity sensation has remained unknown. In **Paper III** of this thesis, I study humidity preference in the fly. In this paper, I present that drosophilid flies have an innate and species-specific preference towards various humidity levels in harmony with the climate of their native habitat. For instance, the cosmopolitan *D. melanogaster* prefers an intermediate humidity of 70% relative humidity (RH) at 25°C, whereas *Drosophila mojavensis*, a native to the dry Sonoran desert in southwestern United States and northern Mexico, prefers a 20% RH. *Drosophila teissieri*, native to warm and humid tropical rainforest in western Africa, prefers 85% RH (**Figure 8A-C**).

I have also searched for the hygrosensory sensilla and the molecular components of this sense on the antenna (Altner and Loftus, 1985). The hygrosensilla in *D. melanogaster* are located in the sacculus (**Figure 9A**), composed of three chambers (**Figure 9B**), each housing sensilla with distinct morphology. Those housed in chambers I and II are poreless, and show similar morphology to hygrosensilla of other insects. I show that Ir40a, and Ir93a along with the co-receptor Ir25a are expressed in neurons found in sensilla housed in chamber I and II (**Figure 9C**). Flies lacking any of these genes display deficient humidity guided behavior. Previous studies have proposed the requirement of thermosensory pathways for hygrosensation (Altner and Loftus, 1985), therefore, I test if these



**Figure 8. Innate and species-specific humidity preference**

A) *D. teissieri* and *D. mojavensis*. B) Humidity-preference indices after 4 hr of *D. melanogaster* (white), *D. teissieri* (green), and *D. mojavensis* (yellow) tested in a binary humidity-preference arena with 20% RH versus 70% RH and, C) 70% RH versus 85% RH.



**Figure 9 The humidity sensing apparatus**

A) The antenna of the fly with the sacculus highlighted. B) Maximum-intensity projection of cuticular autofluorescence showing the outline of the sacculus. C) Maximum-intensity projection of antennae from a *Ir40a-Gal4>UAS-GFP* shows *Ir40a* is expressed in neurons targeting chamber I and II.


IRs are also involved in thermosensation. Genetic inactivation of *Ir40a* (through RNAi) caused a loss of the physiological and behavioral response only to the humidity stimulus, but not to temperature, whereas, both *Ir25a* or *Ir93a* mutant flies showed abnormal thermosensory along with hygrosensory responses. *Ir25a* and *Ir93a* are also expressed in the thermosensory cells of arista (Benton et al., 2009) and in the larval dorsal organ cool cells (DOCCs) (Knecht et al., 2016).

### Early processing of hygrosensory information in the fly brain

In **Paper III**, I show that humidity information is processed in a region called the posterior antennal lobe (PAL) in the brain. The axon terminals of dry sensing neurons, expressing *Ir40a*, project to a glomerulus called “arm” while cool sensing neurons expressing the same receptor – *Ir40a* – project to “column” (**Figure 10A**). As *Ir25a* is required for both humidity and temperature sensing, neurons expressing this co-receptor projects to both “arm” and “column”, as well as “hot” and “cold” glomeruli in the PAL. Moreover, since *Ir25a* is also co-expressed with a subset of other odor sensing IRs, *Ir25a* positive neurons also project to olfactory glomeruli of the AL. *Ir93a*, necessary for humidity preference, projects to arm and column, as well as thermosensory glomeruli.

I also examine the response of “arm” and “column” to humidity and temperature changes by calcium sensors (**Figure 10B**). In imaging experiments the “arm” glomerulus showed increased calcium signal in response to stimulation with dry air, and a reduction in response to humid air, consistent with a function as the dry





**Figure 10. The arm glomerulus in the brain response to humidity stimuli**  
 A) Maximum-intensity projection from the PAL of a *Ir40a-Gal4>UAS-GFP* fly. B) The antennal lobe of a *Ir25a-Gal4>UAS-CaMPARI* fly following photoconversion during dry air stimulation.

neuron. In response to temperature stimulation, the “arm” glomerulus only responded to strong cold stimulation. The “column” on the other hand did not respond to humidity changes, but to changes in temperature, both hot and cold.


I did not succeed in identifying the moist neurons in this study, and accordingly I did not identify the corresponding glomerulus either. Subsequent studies have, however, found that these neurons express *Ir68a*, project to a glomerulus called “bean”, which is adjacent to the “arm” and “column” (Frank et al., 2017; Knecht et al., 2017). The “hot” and “cold” glomeruli processing temperature information are found adjacent to these humidity-sensing glomeruli (Gallio et al., 2011; Knecht et al., 2017, Frank et al., 2017), and together form a hygro and thermo topographic map in the fly brain.

## IV. Marula – The ancestral host of the lord of flies

As stated, *D. melanogaster* displays strong egg-laying preference towards *Citrus* spp. over other type of fruits (Dweck et al., 2013). This distinct host preference implies that *D. melanogaster* might form a close association with a specific fruit, or group of fruits, that probably share some characteristics with citrus in the native habitat. In preparation for **Paper IV** of this thesis, I searched for wild non-commensal populations of *D. melanogaster* within its native range, along with a search for its ancestral host. I searched mopane woodlands of the Matobos national park in Southwestern Zimbabwe (**Figure 11A**). This park is within the predicted ancestral range of *D. melanogaster* (Pool et al 2012), and covers 424 km<sup>2</sup>, have no human habitation and is covered in mopane and kopje woodlands (**Figure 11B**). In the park I found a population of *D. melanogaster*, which turned out to be closely

associated with marula fruit (*Sclerocarya birrea*), a seasonally abundant resource during March to May. Marula, like citrus, has a thick rind, which covers a sweet-sour juicy pulp (**Figure 11C**). Marula is well-known for its high ethanol content (Morris et al., 2006), which would allow *D. melanogaster* to exploit its greater ethanol tolerance (David et al., 1986; Montooth et al., 2006).

We demonstrate that flies have a strong oviposition preference towards marula over orange both in the lab and in field traps (Figure. ). We furthermore localized drosophilid larvae in all marula examined in the field traps, from which *D. melanogaster* adults emerged. Interestingly, we only found the wild population of *D. melanogaster* where the marula trees were present, but not in any other location of the Matobos with similar vegetation (**Figure 11D**), unlike *D. simulans*, which was present in high number at all trap locations (**Figure 11E**). This distribution pattern suggests that *D. melanogaster* has a specialized lifestyle, and is in fact a seasonal specialist, like its siblings *D. erecta* and *D. orena* in the *melanogaster* subgroup.




**Figure 11** Picturetext headline

A) Location of the Matobos National Park, Zimbabwe. B) View of the park. Photo: M. Stensmyr. C) Marula fruit has a thick rind (in my right hand) and a soft flesh (in my left hand). Photo: E. Jirle. D) Number of *D. melanogaster* (red fly) trapped at sites with and without marula. E) Number of *Drosophila simulans* (yellow fly) caught at the same sites as in D).

## Or22a detects the marula volatile ethyl isovalerate

The two main chemicals of marula, ethyl isovalerate (an ester) and beta-caryophyllene (a sesquiterpene), together make up ~55% of the marula headspace and both trigger oviposition in flies. Similar to citrus, marula emits large amounts of terpenes. Citrus headspace is typically dominated by limonene, whereas, marula primarily emits beta-caryophyllene, which also activates the same olfactory pathway as limonene, Or19a (Dweck et al., 2013). Unlike citrus though, marula emits high amounts of esters, primarily ethyl isovalerate, which activates Or22a, as I show via calcium imaging (**Figure 12A**). Silencing the *Or22a* pathway via *Or22a-Gal4>UAS-TNT*, however, did not fully abolish the marula oviposition preference, suggesting that additional pathways are involved in this behavioral preference. The primary function of this pathway might be to locate the host over distance rather than mediating egg-laying preference. *Or22a* in general is broadly tuned to detect fruity esters in *D. melanogaster* and its eight close relatives (Stensmyr et al., 2003). On the other hand, Or22a has a selective and species specific receptive range based on the species preferred breeding substrates (Pelz et al., 2006; Linz et al., 2013). For example in *D. erecta* and *D. sechellia*, Or22a is most sensitive to the key ligands of their respective host fruits. In *D. erecta*, Or22a is most selective towards the *Pandanus* volatile, 3-methyl-2-butenyl acetate (Linz et al., 2013), while in *D. sechellia* this receptor detects the morinda volatile, methyl hexanoate (Dekker et al., 2006). Moreover, this selectivity and higher sensitivity is accomplished by a numerical increase in neurons expressing Or22a in both species in comparison to the other melanogaster species subgroup members (Dekker et al., 2006, Linz et al., 2013).



**Figure 12. Picturetext headline**

A) Pseudocolored image showing ethyl isovalerate-induced fluorescence changes in the antennal lobe (AL) of a *Or22a-Gal4>UAS-GCaMP6f* fly. B) Dose-response curve of ab3A neurons from the Rwanda (RG18N) and a lab strain of North American origin (Canton-S) toward ethyl isovalerate and ethyl hexanoate. Shaded area shows standard deviation. C) Genetic differentiation among populations at *Or22a* and *Or22b* is depicted via Circos plots [Krzywinski et al., 2009] based on  $F_{ST}$  quantiles. Only connections between populations with unusually high  $F_{ST}$  values (elevated genetic differentiation) is shown. The red color, for example, indicates that between this pair of populations, less than 0.1% of windows on the same chromosome arm have an  $F_{ST}$  value this high. Abbreviations display geographic origin of examined *D. melanogaster* populations e.g. RG stands for Rwanda, FR for France.

Interestingly, electrophysiological examination of OSNs expressing *Or22a/Or22b* from Sub-Saharan African flies (collected in Rwanda), revealed stronger responses to ethyl isovalerate than to ethyl hexanoate – the primary ligand of *Or22a* (Munch and Galizia, 2016). In contrast, the corresponding OSNs from a lab strain of North American origin, displayed stronger response to ethyl hexanoate than ethyl isovalerate (**Figure 12B**).

As a consequence of the functional difference of these *Or22a/Or22b* expressing OSNs between the two examined populations, I looked for signs of local genetic adaptation of *Or22a/Or22b* between *D. melanogaster* populations utilizing other hosts than marula. The results of comparing the local genetic differentiation within the Or family between genomes of ten African populations, plus one European revealed a strong genetic differentiation at the *Or22a/Or22b* locus between almost all population pairs (**Figure 12C**), whereas, most of the other ORs exhibited little or no sign of local adaptation. Taken together, unlike most members of the OR family in *D. melanogaster* that are conserved, *Or22a/Or22b* displays strong signs of local adaptation, in line with a function associated with host specific chemistry. Curiously, we found that the Sub-Saharan flies predominantly carries a specific allele at this locus, with a chimeric receptor formed from *Or22a* and *Or22b*. Whether or not this “*Or22ab*” gene indeed is responsible for the shifted ligand tuning of these neurons remains to be investigated. In brief, Southern African population of *D. melanogaster* not only detects ethyl isovalerate, they are even more sensitive to this marula ester than flies from outside Africa.



# Concluding remarks

In this thesis, I have showed that the fly olfactory system does not detect odor molecules randomly, but capture and process specific odors associated with needs and dangers. For example, the fly has several olfactory pathways dedicated to odors coming from alcoholic fermentation and activation of these receptors and the related pathways guides the fly to suitable food sources rich in sugar and yeast (**Paper I**). Although, we outline a range of behavioral and dedicated ecological functions for a number of olfactory pathways, a number of unknowns remain to be studied, such as identifying ligands for the few orphan receptors left e.g. Or2a and Or23a whose functions are likely important given their evident extreme specificity. In addition, the ecological significance of a number of receptors with known key ligands, like Gr21a/Gr63a that detect CO<sub>2</sub> remain puzzling. I have also investigated how the olfactory system cope with toxic and harmful matters in the natural habitat and I could identify an olfactory circuit that mediates repellency towards phenol, which is produced by pathogenic bacteria, predominantly present in carnivore feces (**Paper II**). Determining ligands for the few remaining orphan ORs should be a priority.

Furthermore, I reveal that flies have an innate and species-specific ability to find suitable humidity levels, related to their native habitat. Flies can sense humidity changes in their environment through a trio of IRs expressed in the sacculus of the antenna, which projects to the “arm” glomerulus in the brain (**Paper III**). But we still do not know what the real stimulus is that activates these hygrosensitive neurons. Moreover, the hygro-sensilla expressing IRs are poreless, which brings up the question “how these receptors detect changes in water vapor in the air”.

Although vinegar flies are known as generalists, wild populations of *D. melanogaster* from a mopane forest within the potential ancestral habitat have a strong breeding preference towards marula fruit. This fruit is seasonally abundant, native to Southern Africa, and is presumably the ancestral host of the fly. I also speculate that marula was a vehicle for *D. melanogaster* to become a human commensal. Archaeological excavations from one of the caves in the Matobos uncovered 24 million marula stones (Walker, 1995). These fruit were collected by a San tribe settling in this region during the Late Pleistocene to the Early Holocene and stored for later consumption. The smell of stored marula from these caves would have invited flies to a massive source of food, making the favored host

available long after fruiting season. (**Paper IV**). The finding of a wild population of *D. melanogaster*, and the ancestral habitat, opens up for a range of interesting questions to be addressed. For example, how do these wild flies differ from their commensal relatives, i.e. which genetic factors underlie this shift in lifestyle? The finding that *D. melanogaster* has a close association with a single host fruit, will greatly facilitate studies in host specific chemosensory adaptations, which so far have had to be conducted in other insects in which the wealth of tools available in *D. melanogaster* are unavailable.

In summary, the research presented in my thesis enhances the basic science behind how the olfactory system operates and its mechanisms, the wild behavior of flies, and introduces the genetic and neural basis underlying humidity sensation in insects. These basic understandings may lead us to better strategies for controlling insect pests, as well as human disease vectors such as mosquitoes.

# References

- Abbas, A. K., Lichtman, A. H., and Pillai, S. (2014). "Cellular and molecular immunology E-book," Elsevier Health Sciences.
- Abuin, L., Bargeton, B., Ulbrich, M. H., Isacoff, E. Y., Kellenberger, S., and Benton, R. (2011). Functional architecture of olfactory ionotropic glutamate receptors. *Neuron* **69**, 44-60.
- Ache, B. W., and Young, J. M. (2005). Olfaction: diverse species, conserved principles. *Neuron* **48**, 417-430.
- Adams, M. D., Celniker, S. E., Holt, R. A., Evans, C. A., Gocayne, J. D., Amanatides, P. G., Scherer, S. E., Li, P. W., Hoskins, R. A., and Galle, R. F. (2000). The genome sequence of *Drosophila melanogaster*. *Science* **287**, 2185-2195.
- Ai, M., Min, S., Grosjean, Y., Leblanc, C., Bell, R., Benton, R., and Suh, G. S. (2010). Acid sensing by the *Drosophila* olfactory system. *Nature* **468**, 691.
- Altner, H., and Loftus, R. (1985). Ultrastructure and function of insect thermo- and hygroreceptors. *Annual review of entomology* **30**, 273-295.
- Altner, H., and Prillinger, L. (1980). Ultrastructure of Invertebrate Chemo-, Thermo-, and Hygroreceptors and Its Functional Significance. In "International Review of Cytology" (G. H. Bourne and J. F. Danielli, eds.), Vol. 67, pp. 69-139. Academic Press.
- Amendt, J., Krettek, R., and Zehner, R. (2004). Forensic entomology. *Naturwissenschaften* **91**, 51-65.
- Bargiello, T. A., Jackson, F. R., and Young, M. W. (1984). Restoration of circadian behavioural rhythms by gene transfer in *Drosophila*. *Nature* **312**, 752.
- Bargiello, T. A., and Young, M. W. (1984a). Molecular genetics of a biological clock in *Drosophila*. *Proceedings of the National Academy of Sciences* **81**, 2142-2146.
- Bargiello, T. A., and Young, M. W. (1984b). Molecular genetics of a biological clock in *Drosophila*. *Proc Natl Acad Sci U S A* **81**, 2142-6.
- Begon, M. (1982). Yeasts and *Drosophila*. *The genetics and biology of Drosophila* **3**, 345-384.
- Benton, R., Sachse, S., Michnick, S. W., and Vosshall, L. B. (2006). Atypical membrane topology and heteromeric function of *Drosophila* odorant receptors in vivo. *PLoS biology* **4**, e20.
- Benton, R., Vannice, K. S., Gomez-Diaz, C., and Vosshall, L. B. (2009a). Variant ionotropic glutamate receptors as chemosensory receptors in *Drosophila*. *Cell* **136**, 149-62.



- Benton, R., Vannice, K. S., Gomez-Diaz, C., and Vosshall, L. B. (2009b). Variant ionotropic glutamate receptors as chemosensory receptors in *Drosophila*. *Cell* **136**, 149-162.
- Bhalerao, S., Sen, A., Stocker, R., and Rodrigues, V. (2003). Olfactory neurons expressing identified receptor genes project to subsets of glomeruli within the antennal lobe of *Drosophila melanogaster*. *Developmental Neurobiology* **54**, 577-592.
- Boyle, J., and Cobb, M. (2005). Olfactory coding in *Drosophila* larvae investigated by cross-adaptation. *Journal of Experimental Biology* **208**, 3483-3491.
- Brand, A. H., and Perrimon, N. (1993). Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. *development* **118**, 401-415.
- Bridges, C. B. (1916). Non-disjunction as proof of the chromosome theory of heredity (concluded). *Genetics* **1**, 107-163.
- Bridges, C. B. (1921). Current maps of the location of the mutant genes of *Drosophila melanogaster*. *Proceedings of the National Academy of Sciences* **7**, 127-132.
- Buck, L., and Axel, R. (1991). A novel multigene family may encode odorant receptors: a molecular basis for odor recognition. *Cell* **65**, 175-87.
- Capy, P., and Gibert, P. (2004). *Drosophila melanogaster*, *Drosophila simulans*: so similar yet so different. In "Drosophila melanogaster, Drosophila simulans: So Similar, So Different", pp. 5-16. Springer.
- Carson, H. L. (1971). ecology of *Drosophila* breeding sites.
- Chapman, R. (1982a). Chemoreception: the significance of receptor numbers. In "Advances in insect physiology", Vol. 16, pp. 247-356. Elsevier.
- Chapman, R. F. (1982b). Chemoreception: The Significance of Receptor Numbers. In "Advances in Insect Physiology" (M. J. Berridge, J. E. Treherne and V. B. Wigglesworth, eds.), Vol. 16, pp. 247-356. Academic Press.
- Chu, I.-W., and Axtell, R. (1971). Fine structure of the dorsal organ of the house fly larva, *Musca domestica* L. *Zeitschrift für Zellforschung und mikroskopische Anatomie* **117**, 17-34.
- Chyb, S., Dahanukar, A., Wickens, A., and Carlson, J. R. (2003). *Drosophila* Gr5a encodes a taste receptor tuned to trehalose. *Proceedings of the National Academy of Sciences* **100**, 14526-14530.
- Clyne, P., Grant, A., O'Connell, R., and Carlson, J. R. (1997a). Odorant response of individual sensilla on the *Drosophila* antenna. *Invert Neurosci* **3**, 127-35.
- Clyne, P., Grant, A., O'Connell, R., and Carlson, J. R. (1997b). Odorant response of individual sensilla on the *Drosophila* antenna. *Invertebrate Neuroscience* **3**, 127-135.
- Cobb, M. (1999). What and how do maggots smell? *Biological reviews* **74**, 425-459.
- Cobb, M., Bruneau, S., and Jallon, J.-M. (1992). Genetic and developmental factors in the olfactory response of *Drosophila melanogaster* larvae to alcohols. *Proc. R. Soc. Lond. B* **248**, 103-109.
- Cobb, M., and Domain, I. (2000). Olfactory coding in a simple system: adaptation in *Drosophila* larvae. *Proceedings of the Royal Society of London B: Biological Sciences* **267**, 2119-2125.

- Colomb, J., Grillenzoni, N., Ramaekers, A., and Stocker, R. F. (2007). Architecture of the primary taste center of *Drosophila melanogaster* larvae. *Journal of Comparative Neurology* **502**, 834-847.
- Comeault, A. A., Serrato-Capuchina, A., Turissini, D. A., McLaughlin, P. J., David, J. R., and Matute, D. R. (2017). A nonrandom subset of olfactory genes is associated with host preference in the fruit fly *Drosophila oreana*. *Evolution Letters* **1**, 73-85.
- Couto, A., Alenius, M., and Dickson, B. J. (2005). Molecular, anatomical, and functional organization of the *Drosophila* olfactory system. *Current Biology* **15**, 1535-1547.
- Croset, V., Rytz, R., Cummins, S. F., Budd, A., Brawand, D., Kaessmann, H., Gibson, T. J., and Benton, R. (2010). Ancient protostome origin of chemosensory ionotropic glutamate receptors and the evolution of insect taste and olfaction. *PLoS genetics* **6**, e1001064.
- Dahanukar, A., Foster, K., and Carlson, J. R. (2001). A Gr receptor is required for response to the sugar trehalose in taste neurons of *Drosophila*. *Nature neuroscience* **4**, 1182.
- Dahanukar, A., Lei, Y.-T., Kwon, J. Y., and Carlson, J. R. (2007). Two Gr genes underlie sugar reception in *Drosophila*. *Neuron* **56**, 503-516.
- David, J., Mercot, H., Capy, P., McEvey, S., and Van Herrewege, J. (1986). Alcohol tolerance and Adh gene frequencies in European and African populations of *Drosophila melanogaster*. *Genetique, selection, evolution* **18**, 405.
- David, J. R., and Capy, P. (1988). Genetic variation of *Drosophila melanogaster* natural populations. *Trends in Genetics* **4**, 106-111.
- Davis, R. L. (2005). Olfactory memory formation in *Drosophila*: from molecular to systems neuroscience. *Annu. Rev. Neurosci.* **28**, 275-302.
- De Bruyne, M., Foster, K., and Carlson, J. R. (2001). Odor coding in the *Drosophila* antenna. *Neuron* **30**, 537-552.
- Dekker, T., Ibba, I., Siju, K., Stensmyr, M. C., and Hansson, B. S. (2006). Olfactory shifts parallel superspecialism for toxic fruit in *Drosophila melanogaster* sibling, *D. sechellia*. *Current Biology* **16**, 101-109.
- Depetris-Chauvin, A., Galagovsky, D., and Grosjean, Y. (2015). Chemicals and chemoreceptors: ecologically relevant signals driving behavior in *Drosophila*. *Frontiers in Ecology and Evolution* **3**, 41.
- Dietzl, G., Chen, D., Schnorrer, F., Su, K.-C., Barinova, Y., Fellner, M., Gasser, B., Kinsey, K., Oettel, S., and Scheiblaue, S. (2007). A genome-wide transgenic RNAi library for conditional gene inactivation in *Drosophila*. *Nature* **448**, 151.
- Dobritsa, A. A., van Naters, W. v. d. G., Warr, C. G., Steinbrecht, R. A., and Carlson, J. R. (2003). Integrating the molecular and cellular basis of odor coding in the *Drosophila* antenna. *Neuron* **37**, 827-841.
- Dweck, C. S. (2013). "Self-theories: Their role in motivation, personality, and development," psychology press.

- Dweck, H. K., Ebrahim, S. A., Khallaf, M. A., Koenig, C., Farhan, A., Stieber, R., Weißflog, J., Svatoš, A., Grosse-Wilde, E., and Knaden, M. (2016). Olfactory channels associated with the *Drosophila* maxillary palp mediate short-and long-range attraction. *Elife* **5**.
- Dweck, H. K., Ebrahim, S. A., Kromann, S., Bown, D., Hillbur, Y., Sachse, S., Hansson, B. S., and Stensmyr, M. C. (2013). Olfactory preference for egg laying on citrus substrates in *Drosophila*. *Current Biology* **23**, 2472-2480.
- Dweck, H. K., Ebrahim, S. A., Thoma, M., Mohamed, A. A., Keesey, I. W., Trona, F., Lavista-Llanos, S., Svatoš, A., Sachse, S., and Knaden, M. (2015). Pheromones mediating copulation and attraction in *Drosophila*. *Proceedings of the National Academy of Sciences* **112**, E2829-E2835.
- Enjin, A. (2017). Humidity sensing in insects—from ecology to neural processing. *Current opinion in insect science* **24**, 1-6.
- Enjin, A., Zaharieva, E. E., Frank, D. D., Mansourian, S., Suh, G. S., Gallio, M., and Stensmyr, M. C. (2016). Humidity sensing in *Drosophila*. *Current Biology* **26**, 1352-1358.
- Ferveur, J.-F., Cortot, J., Rihani, K., Cobb, M., and Everaerts, C. (2018). Desiccation resistance: effect of cuticular hydrocarbons and water content in *Drosophila melanogaster* adults. *PeerJ PrePrints*.
- Fishilevich, E., and Vosshall, L. B. (2005). Genetic and functional subdivision of the *Drosophila* antennal lobe. *Current Biology* **15**, 1548-1553.
- Frank, D. D., Enjin, A., Jouandet, G. C., Zaharieva, E. E., Para, A., Stensmyr, M. C., and Gallio, M. (2017). Early integration of temperature and humidity stimuli in the *Drosophila* brain. *Current Biology* **27**, 2381-2388. e4.
- Galizia, C. G. (2014). Olfactory coding in the insect brain: data and conjectures. *European Journal of Neuroscience* **39**, 1784-1795.
- Gallio, M., Ofstad, T. A., Macpherson, L. J., Wang, J. W., and Zuker, C. S. (2011). The coding of temperature in the *Drosophila* brain. *Cell* **144**, 614-624.
- Gao, Q., and Chess, A. (1999). Identification of candidate *Drosophila* olfactory receptors from genomic DNA sequence. *Genomics* **60**, 31-39.
- Gao, Q., Yuan, B., and Chess, A. (2000). Convergent projections of *Drosophila* olfactory neurons to specific glomeruli in the antennal lobe. *Nature neuroscience* **3**, 780.
- Ha, T. S., and Smith, D. P. (2006). A pheromone receptor mediates 11-cis-vaccenyl acetate-induced responses in *Drosophila*. *Journal of Neuroscience* **26**, 8727-8733.
- Hallem, E. A., and Carlson, J. R. (2006). Coding of odors by a receptor repertoire. *Cell* **125**, 143-160.
- Hallem, E. A., Ho, M. G., and Carlson, J. R. (2004). The molecular basis of odor coding in the *Drosophila* antenna. *Cell* **117**, 965-979.
- Heimbeck, G., Bugnon, V., Gendre, N., Häberlin, C., and Stocker, R. F. (1999). Smell and Taste Perception in *Drosophila melanogaster* Larva: Toxin Expression Studies in Chemosensory Neurons. *Journal of Neuroscience* **19**, 6599-6609.

- Heimbeck, G., Bugnon, V., Gendre, N., Keller, A., and Stocker, R. F. (2001). A central neural circuit for experience-independent olfactory and courtship behavior in *Drosophila melanogaster*. *Proceedings of the National Academy of Sciences* **98**, 15336-15341.
- Heisenberg, M. (2003). Mushroom body memoir: from maps to models. *Nature Reviews Neuroscience* **4**, 266.
- Hendrichs, J., Katsoyannos, B., and Prokopy, R. (1993). Bird feces in the nutrition of adult Mediterranean fruit flies *Ceratitis capitata* (Diptera: Tephritidae) in nature. *Mitteilungen der Deutschen Gesellschaft fuer Allgemeine und Angewandte Entomologie (Germany)*.
- Hussain, A., Zhang, M., Üçpunar, H. K., Svensson, T., Quillery, E., Gompel, N., Ignell, R., and Kadow, I. C. G. (2016). Ionotropic chemosensory receptors mediate the taste and smell of polyamines. *PLoS biology* **14**, e1002454.
- Joiner, W. J., Crocker, A., White, B. H., and Sehgal, A. (2006). Sleep in *Drosophila* is regulated by adult mushroom bodies. *Nature* **441**, 757.
- Joseph, R. M., Devineni, A. V., King, I. F., and Heberlein, U. (2009). Oviposition preference for and positional avoidance of acetic acid provide a model for competing behavioral drives in *Drosophila*. *Proceedings of the National Academy of Sciences* **106**, 11352-11357.
- Kaiser, M., and Cobb, M. (2008). The behaviour of *Drosophila melanogaster* maggots is affected by social, physiological and temporal factors. *Animal Behaviour* **75**, 1619-1628.
- Karlson, P., and Butenandt, A. (1959). Pheromones (ectohormones) in insects. *Annual review of entomology* **4**, 39-58.
- Keil, T. A., and Steinbrecht, R. A. (1984). Mechanosensitive and olfactory sensilla of insects. In "Insect ultrastructure", pp. 477-516. Springer.
- Knecht, Z. A., Silbering, A. F., Cruz, J., Yang, L., Croset, V., Benton, R., and Garrity, P. A. (2017). Ionotropic Receptor-dependent moist and dry cells control hygrosensation in *Drosophila*. *Elife* **6**.
- Knecht, Z. A., Silbering, A. F., Ni, L., Klein, M., Budelli, G., Bell, R., Abuin, L., Ferrer, A. J., Samuel, A. D., and Benton, R. (2016). Distinct combinations of variant ionotropic glutamate receptors mediate thermosensation and hygrosensation in *Drosophila*. *Elife* **5**.
- Koh, T.-W., He, Z., Gorur-Shandilya, S., Menuz, K., Larter, N. K., Stewart, S., and Carlson, J. R. (2014). The *Drosophila* IR20a clade of ionotropic receptors are candidate taste and pheromone receptors. *Neuron* **83**, 850-865.
- Kominsky, D. J., Campbell, E. L., and Colgan, S. P. (2010). Metabolic shifts in immunity and inflammation. *The Journal of Immunology* **184**, 4062-4068.
- Kreher, S. A., Mathew, D., Kim, J., and Carlson, J. R. (2008). Translation of sensory input into behavioral output via an olfactory system. *Neuron* **59**, 110-124.
- Kühnel, S., Brückner, A., Schmelzle, S., Heethoff, M., and Blüthgen, N. (2017). Surface area–volume ratios in insects. *Insect science* **24**, 829-841.

- Kumagai, H., Yamada, H., Matsui, H., Ohkishi, H., and Ogata, K. (1970). Tyrosine phenol lyase I. Purification, crystallization, and properties. *Journal of Biological Chemistry* **245**, 1767-1772.
- Kurtovic, A., Widmer, A., and Dickson, B. J. (2007). A single class of olfactory neurons mediates behavioural responses to a *Drosophila* sex pheromone. *Nature* **446**, 542.
- Kwon, J. Y., Dahanukar, A., Weiss, L. A., and Carlson, J. R. (2007). The molecular basis of CO2 reception in *Drosophila*. *Proceedings of the National Academy of Sciences* **104**, 3574-3578.
- Kwon, J. Y., Dahanukar, A., Weiss, L. A., and Carlson, J. R. (2011). Molecular and cellular organization of the taste system in the *Drosophila* larva. *Journal of Neuroscience* **31**, 15300-15309.
- Lacaille, F., Hiroi, M., Twele, R., Inoshita, T., Umemoto, D., Maniere, G., Marion-Poll, F., Ozaki, M., Francke, W., and Cobb, M. (2007). An inhibitory sex pheromone tastes bitter for *Drosophila* males. *PLoS one* **2**, e661.
- Lachaise, D., Cariou, M.-L., David, J. R., Lemeunier, F., Tsacas, L., and Ashburner, M. (1988). Historical biogeography of the *Drosophila melanogaster* species subgroup. In "Evolutionary biology", pp. 159-225. Springer.
- Lachaise, D., Harry, M., Solignac, M., Lemeunier, F., Benassi, V., and Cariou, M.-L. (2000). Evolutionary novelties in islands: *Drosophila santomea*, a new *melanogaster* sister species from Sao Tome. *Proceedings of the Royal Society of London B: Biological Sciences* **267**, 1487-1495.
- Lachaise, D., and Silvain, J.-F. (2004). How two Afrotropical endemics made two cosmopolitan human commensals: the *Drosophila melanogaster*-*D. simulans* palaeogeographic riddle. In "*Drosophila melanogaster*, *Drosophila simulans*: So Similar, So Different", pp. 17-39. Springer.
- Lacher, V. (1964). Elektrophysiologische untersuchungen an einzelnen rezeptoren für geruch, kohlendioxid, luftfeuchtigkeit und temperatur auf den antennen der arbeitsbiene und der drohne (*Apis mellifica* L.). *Zeitschrift für vergleichende Physiologie* **48**, 587-623.
- Larsson, M. C., Domingos, A. I., Jones, W. D., Chiappe, M. E., Amrein, H., and Vosshall, L. B. (2004). Or83b encodes a broadly expressed odorant receptor essential for *Drosophila* olfaction. *Neuron* **43**, 703-714.
- Lauzon, C. R. (2003). Symbiotic relationships of tephritids. *Insect symbiosis* **2**, 115-129.
- Lee, J., and Strausfeld, N. (1990). Structure, distribution and number of surface sensilla and their receptor cells on the olfactory appendage of the male moth *Manduca sexta*. *Journal of neurocytology* **19**, 519-538.
- Lee, Y., Moon, S. J., and Montell, C. (2009). Multiple gustatory receptors required for the caffeine response in *Drosophila*. *Proceedings of the National Academy of Sciences* **106**, 4495-4500.
- Ling, F., Dahanukar, A., Weiss, L. A., Kwon, J. Y., and Carlson, J. R. (2014). The molecular and cellular basis of taste coding in the legs of *Drosophila*. *Journal of Neuroscience* **34**, 7148-7164.

- Linz, J., Baschwitz, A., Strutz, A., Dweck, H. K., Sachse, S., Hansson, B. S., and Stensmyr, M. C. (2013). Host plant-driven sensory specialization in *Drosophila erecta*. *Proc. R. Soc. B* **280**, 20130626.
- Lu, B., and Vogel, H. (2009). *Drosophila* models of neurodegenerative diseases. *Annual Review of Pathological Mechanical Disease* **4**, 315-342.
- Ludwig, D. (1945). The effects of atmospheric humidity on animal life. *Physiological Zoology* **18**, 103-135.
- Mansourian, S., and Stensmyr, M. C. (2015). The chemical ecology of the fly. *Current opinion in neurobiology* **34**, 95-102.
- Marella, S., Fischler, W., Kong, P., Asgarian, S., Rueckert, E., and Scott, K. (2006). Imaging taste responses in the fly brain reveals a functional map of taste category and behavior. *Neuron* **49**, 285-295.
- Martin, J.-R., Ernst, R., and Heisenberg, M. (1998). Mushroom bodies suppress locomotor activity in *Drosophila melanogaster*. *Learning & Memory* **5**, 179-191.
- Mathew, D., Martelli, C., Kelley-Swift, E., Brusalis, C., Gershow, M., Samuel, A. D., Emonet, T., and Carlson, J. R. (2013). Functional diversity among sensory receptors in a *Drosophila* olfactory circuit. *Proc Natl Acad Sci U S A* **110**, E2134-43.
- McKean, K. A., and Lazzaro, B. P. (2011). The costs of immunity and the evolution of immunological defense mechanisms. pp. 299-310. Oxford University Press Oxford.
- Menuz, K., Larter, N. K., Park, J., and Carlson, J. R. (2014). An RNA-seq screen of the *Drosophila* antenna identifies a transporter necessary for ammonia detection. *PLoS Genetics* **10**, e1004810.
- Mizunami, M., Nishino, H., and Yokohari, F. (2016). Status of and future research on thermosensory processing. *Frontiers in physiology* **7**, 150.
- Montooth, K. L., Siebenthal, K. T., and Clark, A. G. (2006). Membrane lipid physiology and toxin catabolism underlie ethanol and acetic acid tolerance in *Drosophila melanogaster*. *Journal of Experimental Biology* **209**, 3837-3850.
- Moon, S. J., Lee, Y., Jiao, Y., and Montell, C. (2009). A *Drosophila* gustatory receptor essential for aversive taste and inhibiting male-to-male courtship. *Current Biology* **19**, 1623-1627.
- Morgan, T. H. (1910). Sex limited inheritance in *Drosophila*. *Science* **32**, 120-122.
- Morris, S., Humphreys, D., and Reynolds, D. (2006). Myth, marula, and elephant: an assessment of voluntary ethanol intoxication of the African elephant (*Loxodonta africana*) following feeding on the fruit of the marula tree (*Sclerocarya birrea*). *Physiological and Biochemical Zoology* **79**, 363-369.
- Muller, H. J. (1927). Artificial Transmutation of the Gene. *Science* **66**, 84-7.
- Münch, D., and Galizia, C. G. (2016). DoOR 2.0-comprehensive mapping of *Drosophila melanogaster* odorant responses. *Scientific reports* **6**, 21841.
- Nakai, J., Ohkura, M., and Imoto, K. (2001). A high signal-to-noise Ca<sup>2+</sup> probe composed of a single green fluorescent protein. *Nature biotechnology* **19**, 137.
- Ni, J.-Q., Liu, L.-P., Binari, R., Hardy, R., Shim, H.-S., Cavallaro, A., Booker, M., Pfeiffer, B. D., Markstein, M., and Wang, H. (2009). A *Drosophila* resource of transgenic RNAi lines for neurogenetics. *Genetics* **182**, 1089-1100.

- Ni, L., Klein, M., Svec, K. V., Budelli, G., Chang, E. C., Ferrer, A. J., Benton, R., Samuel, A. D., and Garrity, P. A. (2016). The ionotropic receptors IR21a and IR25a mediate cool sensing in *Drosophila*. *Elife* **5**.
- Niimura, Y., Matsui, A., and Touhara, K. (2014). Extreme expansion of the olfactory receptor gene repertoire in African elephants and evolutionary dynamics of orthologous gene groups in 13 placental mammals. *Genome research* **24**, 1485-1496.
- Niimura, Y., and Nei, M. (2007). Extensive gains and losses of olfactory receptor genes in mammalian evolution. *PLoS one* **2**, e708.
- Norris, D. M., and Chu, H.-m. (1974). Morphology and ultrastructure of the antenna of male *Periplaneta americana* as related to chemoreception. *Cell and tissue research* **150**, 1-9.
- Okada, J., Kanamaru, Y., and Toh, Y. (2002). Mechanosensory control of antennal movement by the scapal hair plate in the American cockroach. *Zoological science* **19**, 1201-1210.
- Okada, J., and Toh, Y. (2006). Active tactile sensing for localization of objects by the cockroach antenna. *Journal of Comparative Physiology A* **192**, 715-726.
- Okal, M. N., Francis, B., Herrera-Varela, M., Fillinger, U., and Lindsay, S. W. (2013). Water vapour is a pre-oviposition attractant for the malaria vector *Anopheles gambiae* sensu stricto. *Malaria journal* **12**, 365.
- Pelz, D., Roeske, T., Syed, Z., Bruyne, M. d., and Galizia, C. G. (2006). The molecular receptive range of an olfactory receptor in vivo (*Drosophila melanogaster* Or22a). *Developmental Neurobiology* **66**, 1544-1563.
- Pfeiffer, B. D., Jenett, A., Hammonds, A. S., Ngo, T.-T. B., Misra, S., Murphy, C., Scully, A., Carlson, J. W., Wan, K. H., and Laverly, T. R. (2008). Tools for neuroanatomy and neurogenetics in *Drosophila*. *Proceedings of the National Academy of Sciences* **105**, 9715-9720.
- Pitman, J. L., McGill, J. J., Keegan, K. P., and Allada, R. (2006). A dynamic role for the mushroom bodies in promoting sleep in *Drosophila*. *Nature* **441**, 753.
- Poivet, E., Gallot, A., Montagné, N., Glaser, N., Legeai, F., and Jacquin-Joly, E. (2013). A comparison of the olfactory gene repertoires of adults and larvae in the noctuid moth *Spodoptera littoralis*. *PLoS one* **8**, e60263.
- Pool, J. E., and Aquadro, C. F. (2006). History and structure of sub-Saharan populations of *Drosophila melanogaster*. *Genetics* **174**, 915-929.
- Pool, J. E., Corbett-Detig, R. B., Sugino, R. P., Stevens, K. A., Cardeno, C. M., Crepeau, M. W., Duchon, P., Emerson, J., Saelao, P., and Begun, D. J. (2012). Population genomics of sub-Saharan *Drosophila melanogaster*: African diversity and non-African admixture. *PLoS genetics* **8**, e1003080.
- Pool, J. E., Wong, A., and Aquadro, C. F. (2006). Finding of male-killing *Spiroplasma* infecting *Drosophila melanogaster* in Africa implies transatlantic migration of this endosymbiont. *Heredity* **97**, 27.
- Rong, Y. S., Titen, S. W., Xie, H. B., Golic, M. M., Bastiani, M., Bandyopadhyay, P., Olivera, B. M., Brodsky, M., Rubin, G. M., and Golic, K. G. (2002). Targeted mutagenesis by homologous recombination in *D. melanogaster*. *Genes & development* **16**, 1568-1581.

- Rossetto, O., Pirazzini, M., and Montecucco, C. (2014). Botulinum neurotoxins: genetic, structural and mechanistic insights. *Nature Reviews Microbiology* **12**, 535.
- Rouault, J., and David, J. (1982). Evolutionary biology of *Drosophila melanogaster* and *D. simulans*: a behavioral divergence in microhabitat selection [flies, sibling species, interspecific coexistence; Tunisia]. *Acta Oecologica Oecologia Generalis*.
- Rytz, R., Croset, V., and Benton, R. (2013). Ionotropic receptors (IRs): chemosensory ionotropic glutamate receptors in *Drosophila* and beyond. *Insect biochemistry and molecular biology* **43**, 888-897.
- Sakai, T., and Kitamoto, T. (2006). Differential roles of two major brain structures, mushroom bodies and central complex, for *Drosophila* male courtship behavior. *Developmental Neurobiology* **66**, 821-834.
- Schafer, R., and Sanchez, T. V. (1973). Antennal sensory system of the cockroach, *Periplaneta americana*: postembryonic development and morphology of the sense organs. *Journal of Comparative Neurology* **149**, 335-353.
- Schaller, D. (1978). Antennal sensory system of *Periplaneta americana* L. *Cell and tissue research* **191**, 121-139.
- Schawaroch, V. (2002). Phylogeny of a paradigm lineage: the *Drosophila melanogaster* species group (Diptera: Drosophilidae). *Biological Journal of the Linnean Society* **76**, 21-37.
- Schneider, D. (1964). Insect antennae. *Annual review of entomology* **9**, 103-122.
- Scott, K., Brady Jr, R., Cravchik, A., Morozov, P., Rzhetsky, A., Zuker, C., and Axel, R. (2001). A chemosensory gene family encoding candidate gustatory and olfactory receptors in *Drosophila*. *Cell* **104**, 661-673.
- Seki, Y., Rybak, J., Wicher, D., Sachse, S., and Hansson, B. S. (2010). Physiological and morphological characterization of local interneurons in the *Drosophila* antennal lobe. *Journal of neurophysiology* **104**, 1007-1019.
- Semmelhack, J. L., and Wang, J. W. (2009). Select *Drosophila* glomeruli mediate innate olfactory attraction and aversion. *Nature* **459**, 218.
- Shelford, V. (1918). A comparison of the responses of animals in gradients of environmental factors with particular reference to the method of reaction of representatives of the various groups from protozoa to mammals. *Science* **48**, 225-230.
- Silbering, A. F., Rytz, R., Grosjean, Y., Abuin, L., Ramdya, P., Jefferis, G. S., and Benton, R. (2011). Complementary function and integrated wiring of the evolutionarily distinct *Drosophila* olfactory subsystems. *Journal of Neuroscience* **31**, 13357-13375.
- Singh, R. N. (1997). Neurobiology of the gustatory systems of *Drosophila* and some terrestrial insects. *Microscopy research and technique* **39**, 547-563.
- Singh, R. N., and Singh, K. (1984). Fine structure of the sensory organs of *Drosophila melanogaster* Meigen larva (Diptera: Drosophilidae). *International Journal of Insect Morphology and Embryology* **13**, 255-273.
- Soldano, A., Alpizar, Y. A., Boonen, B., Franco, L., Lopez-Requena, A., Liu, G., Mora, N., Yaksi, E., Voets, T., and Vennekens, R. (2016). Gustatory-mediated avoidance of bacterial lipopolysaccharides via TRPA1 activation in *Drosophila*. *Elife* **5**.



- Stanley, S. M., Parsons, P. A., Spence, G., and Weber, L. (1980). Resistance of species of the *Drosophila melanogaster* subgroup to environmental extremes. *Australian Journal of Zoology* **28**, 413-421.
- Starmer, W. T., and Aberdeen, V. (1990). The nutritional importance of pure and mixed cultures of yeasts in the development of *Drosophila mulleri* larvae in *Opuntia* tissues and its relationship to host plant shifts. In "Ecological and evolutionary genetics of *Drosophila*", pp. 145-160. Springer.
- Steinbrecht, R. A. (1984). Chemo-, hygro-, and thermoreceptors. In "Biology of the integument", pp. 523-553. Springer.
- Stensmyr, M. C., Dekker, T., and Hansson, B. S. (2003). Evolution of the olfactory code in the *Drosophila melanogaster* subgroup. *Proceedings of the Royal Society of London B: Biological Sciences* **270**, 2333-2340.
- Stensmyr, M. C., Dweck, H. K., Farhan, A., Ibba, I., Strutz, A., Mukunda, L., Linz, J., Grabe, V., Steck, K., and Lavista-Llanos, S. (2012). A conserved dedicated olfactory circuit for detecting harmful microbes in *Drosophila*. *Cell* **151**, 1345-1357.
- Stocker, R., Lienhard, M., Borst, A., and Fischbach, K. (1990). Neuronal architecture of the antennal lobe in *Drosophila melanogaster*. *Cell and tissue research* **262**, 9-34.
- Stocker, R. F. (1994). The organization of the chemosensory system in *Drosophila melanogaster*: a review. *Cell and tissue research* **275**, 3-26.
- Stocker, R. F. (2001). *Drosophila* as a focus in olfactory research: mapping of olfactory sensilla by fine structure, odor specificity, odorant receptor expression, and central connectivity. *Microscopy research and technique* **55**, 284-296.
- Stöckl, J., Strutz, A., Dafni, A., Svatos, A., Doubsky, J., Knaden, M., Sachse, S., Hansson, B. S., and Stensmyr, M. C. (2010). A deceptive pollination system targeting drosophilids through olfactory mimicry of yeast. *Current Biology* **20**, 1846-1852.
- Störtkuhl, K. F., and Kettler, R. (2001). Functional analysis of an olfactory receptor in *Drosophila melanogaster*. *Proceedings of the National Academy of Sciences* **98**, 9381-9385.
- Strausfeld, N. J., and Hildebrand, J. G. (1999). Olfactory systems: common design, uncommon origins? *Current opinion in neurobiology* **9**, 634-639.
- Sturtevant, A. H. (1965). "History of genetics," Harper And Row; New York And London.
- Sweeney, S. T., Broadie, K., Keane, J., Niemann, H., and O'Kane, C. J. (1995). Targeted expression of tetanus toxin light chain in *Drosophila* specifically eliminates synaptic transmission and causes behavioral defects. *Neuron* **14**, 341-351.
- Tichy, H., and Kallina, W. (2010). Insect hygrometric responses to continuous changes in humidity and air pressure. *Journal of neurophysiology* **103**, 3274-3286.
- Tichy, H., and Loftus, R. (1996). Hygrometric receptors in insects and a spider: humidity transduction models. *Naturwissenschaften* **83**, 255-263.
- Tissot, M., Gendre, N., Hawken, A., Störtkuhl, K. F., and Stocker, R. F. (1997). Larval chemosensory projections and invasion of adult afferents in the antennal lobe of *Drosophila*. *Developmental Neurobiology* **32**, 281-297.

- Toh, Y. (1981). Fine structure of sense organs on the antennal pedicel and scape of the male cockroach, *Periplaneta americana*. *Journal of ultrastructure research* **77**, 119-132.
- Tsacas, L., and David, J. (1978). Une septième espèce appartenant au sous-groupe *Drosophila melanogaster* Meigen: *Drosophila orena* spec. nov. du Cameroun (Diptera: Drosophilidae). *Beitrage zur Entomologie*.
- Van Breugel, F., Riffell, J., Fairhall, A., and Dickinson, M. H. (2015). Mosquitoes use vision to associate odor plumes with thermal targets. *Current Biology* **25**, 2123-2129.
- Von Arx, M., Goyret, J., Davidowitz, G., and Raguso, R. A. (2012). Floral humidity as a reliable sensory cue for profitability assessment by nectar-foraging hawkmoths. *Proceedings of the National Academy of Sciences* **109**, 9471-9476.
- Vosshall, L. B., Amrein, H., Morozov, P. S., Rzhetsky, A., and Axel, R. (1999). A spatial map of olfactory receptor expression in the *Drosophila* antenna. *Cell* **96**, 725-736.
- Vosshall, L. B., and Stocker, R. F. (2007). Molecular architecture of smell and taste in *Drosophila*. *Annu. Rev. Neurosci.* **30**, 505-533.
- Vosshall, L. B., Wong, A. M., and Axel, R. (2000). An olfactory sensory map in the fly brain. *Cell* **102**, 147-159.
- Walker, N. J. (1995). Late Pleistocene and Holocene hunter-gatherers of the Matopos: an archaeological study of change and continuity in Zimbabwe.
- Wang, Z., Singhvi, A., Kong, P., and Scott, K. (2004). Taste representations in the *Drosophila* brain. *Cell* **117**, 981-991.
- Weiss, L. A., Dahanukar, A., Kwon, J. Y., Banerjee, D., and Carlson, J. R. (2011). The molecular and cellular basis of bitter taste in *Drosophila*. *Neuron* **69**, 258-272.
- Wilson, R. I. (2013). Early olfactory processing in *Drosophila*: mechanisms and principles. *Annual Review of Neuroscience* **36**, 217-241.
- Wilson, R. I., and Laurent, G. (2005). Role of GABAergic inhibition in shaping odor-evoked spatiotemporal patterns in the *Drosophila* antennal lobe. *Journal of Neuroscience* **25**, 9069-9079.
- Yao, C. A., Ignell, R., and Carlson, J. R. (2005). Chemosensory coding by neurons in the coeloconic sensilla of the *Drosophila* antenna. *Journal of Neuroscience* **25**, 8359-8367.
- Yokohari, F. (1999). Hygro- and thermoreceptors. *Atlas of Arthropod Sensory Receptors*. Springer, Berlin Heidelberg New York, 191-210.
- Zhou, X., Slone, J. D., Rokas, A., Berger, S. L., Liebig, J., Ray, A., Reinberg, D., and Zwiebel, L. J. (2012). Phylogenetic and transcriptomic analysis of chemosensory receptors in a pair of divergent ant species reveals sex-specific signatures of odor coding. *PLoS genetics* **8**, e1002930.



# Acknowledgement

Writing acknowledgment is the finishing touch on my dissertation. It has been a period of intense learning for me, not only in scientific arena, but also on a personal level. I am pleased to reflect on all my memories and thank all people who have supported and helped me throughout this period.

*Marcus Stensmyr*, my energetic and positive supervisor. Doing my PhD by your positive disposition and wonderful supervision was extremely enjoyable for me. Working in the lab was the most enjoyable for me when the lab was your office. You always spoke to me, made me laugh and encouraged me to do my best. I have learned a lot from you in addition to being creative, and to learn to express my ideas, even when they were weird. Thank you so much for taking me to the most exciting fieldwork in Zambia, which made me grow as a person. Thanks for all the happy moments and encouragement. Whenever I needed to talk to you, not only about work but also about my troubles in my life, you would make time, even if you were not in Lund, you were always reachable. After talking to you, all obstacles were gone and everything became possible. Thanks for having an open ear to my questions and worries. My warmest thanks are to you.

*Marie Dacke*, my inspiring co-supervisor. Thank you for being always encouraging and supportive. I look at you as an inspiration for women in science and I wish to become like you, a successful science communicator.

*Teun Dekker*, my kind teacher and great supervisor before starting my PhD. You have prepared me to become a competitive PhD candidate. You gave me the opportunity to work in your projects and taught me patiently many new techniques in the field. The deepest thanks for allowing me to grow as a research scientist and my appreciation for your willingness to help in different occasions.

I feel extremely lucky to have three great supervisors.

*Christer Löfstedt*, my special appreciation and thanks for accepting me to be part of your pheromone group when Stensmyr group was consisting of me only. I felt welcomed by you and your group members and I still I feel that I am member of the pheromone group. Also, thank you for taking care of all the PhD meetings and motivations.

*Erling Jirle*, a fun friend, nice neighbor and Swedish Papa. Thank you for always being there for me and kindly helped me whenever I asked you. Thanks for your fun company during our field trips to Africa. Lots of unforgettable memories, “look at them, are not they lovely?”, Thank you Winston”. Thank you Erling ☺ for giving me enjoyable company whenever I felt lonely and showing me many beautiful places in Skåne. Tusen tack för den svenska populära sammanfattningen.

*Anders*, Many thanks for teaching me how “fly pushing” works and answering my questions regarding fly genetic. Thanks for your valuable comments on my thesis.

*Nadia*, my sweet friend. Working environment became more enjoyable for me since you joined us. I am so glad to have won you as my friend to talk about everything that burns our tongues and soul.

*Nicoletta, Jothi, and Erika*, Thank you for all nice time during lunch, parties and your fun company. Simply, thanks for our friendships.

*Jacob*, Thanks for introducing HEK-cell to me and helping me with it. It made a positive impact on my paper.

Past and present members of the “Pheromone group”, *Olle, Hong-Lei, Dan Dan, Martin, Glenn, Bao-jian, , Inis, Colm, Xiaoqing, Yihan, Johan* My big thanks to all of you for being nice to me and for all of your help and group meetings.

*Ylwa, Katarina, Anne and Anne L*, my Great thanks to you for taking care of all the administrative issues.

*Camila*, Thanks for organizing fun kick-of-days.

*David, Fredrik, Agneska, Tini, Elisa*, Thanks for your smiles when meeting me in the corridor.

*Mehrnaz*, Thanks for knocking at my office door, asking me how it is going, and if I needed any help. That made my heart warm on stressful days.

Thanks to all people working in Ecology and Biology building for making such a great working environment.

I would like to thank all the funding agencies *Fysiografen, VR, and STINT* for financing my trip to the scientific meetings.

I would also like to thank all the fellow people in Alnarp, SLU for welcoming me to your group when I moved to Sweden. Your kind attitudes, your smile, your willingness to improve my skills made my new life in Sweden easier. Thank you all especially, *Aman, William Walker, Göran Birgersson, Marie Bengtsson, Paul Becher, Petter Anderson, Sebastian, Eduardo, Ian, Franklin, David, Magali, Miriam* and many more.

I also made lots of friends during three years in Alnarp. It might be along time since I met you guys, but I always count you as my cool friends and miss you all. *Shahid, Malin, Zubi, Ali, Faraz, Mikael, Gökhan, Negar, Mohamad Habibi, Santosh and Mike*, many many thanks for wonderful memories and friendship.

*Samareh*, my closest friend. You know all of my feelings. Thank you for always being by my side. We really have had shared pains and laughters in the last 7 years.

*Raha, Iren, Sudi, Arezoo and Soroush*, you are my old- gold friends. Love you all and thanks for giving me hope all the time and make me smile.

Elnazi, Thanks for the cute painting of the vinegar fly.

*Lale and Amo Saeed*, my kind-hearted aunt and uncle, you have always giving me warm welcome to your home since I moved to Sweden. Your door is the one that I can knock whenever I feel home sick. Thanks for your kindness and warm hospitality. *Alex and Asal*, I get so much energy and happiness from you. It is amazing to have a small family close by when you are far away from home.

*Sepideh*, My lovely sister, I am so lucky to have you close by! You are one of the reasons that I moved to Sweden, to be closer to you. I will never forget whatever you have done for me, specially your company in my first days in Alnarp. A big thank you *Khahari* for the unfailing support in these years. *Shahriar*, azize khale, for me, You are a bundle of happiness and love. Thanks for bringing joy to me through your visits during the summers.

*Mostafa*, my best friend, my beloved partner, you absorbed much of my frustration, stress and fears during the writing phase. Thank you so much for your great understanding and support when there was no one else to answer my queries. I am lucky enough to have had you next to me especially in the last phase of my PhD. Thanks hamsi for the Persian translation of the thesis summary.

Finally, my wonderful parents, *Mansour and Shokoufe*, my deepest gratitude is to you, who encouraged me to continue my education, supported me during my PhD study and pushed me forward in all the tough situations. You have been a constant source of strength and inspiration. Without your support, I could not have reached this success. Words cannot express how grateful I am to you. *Ehsan*, my kindest brother, you have had a great impact on my personality and my attitude since my childhood. You are my encourager to pursue my Phd. Thank you *dadashi mehrabonam* for being my mentor and thanks so much for taking care of our parents during my absence.

