



LUND UNIVERSITY

Postnatal sense of security, anxiety and risk for postnatal depression

Persson, Eva-Kristina; Kvist, Linda

Published in:
Journal of Women's Health: issues and care

DOI:
[10.4172/2325-9795.1000141](https://doi.org/10.4172/2325-9795.1000141)

2014

[Link to publication](#)

Citation for published version (APA):
Persson, E.-K., & Kvist, L. (2014). Postnatal sense of security, anxiety and risk for postnatal depression. *Journal of Women's Health: issues and care*, 3(3). <https://doi.org/10.4172/2325-9795.1000141>

Total number of authors:
2

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Postnatal sense of security, anxiety and risk for postnatal depression

Eva K. Persson, RN, RM, PhD, Senior lecturer¹
Linda J. Kvist, RN, RM, PhD, Associate professor^{1,2}

¹ Division of Nursing, Department of Health Sciences, Lund University, PO Box 157,
S-221 00 Lund, Sweden.

² Department of Obstetrics & Gynaecology, Helsingborg Hospital, S-251 87 Helsingborg,
Sweden

Correspondence: eva-kristina.persson@med.lu.se Tel: +46 46 222 1890, fax +46 46 2221808

Abstract

Background: Approximately 10% - 15% of mothers and 10% of fathers suffer from depressive symptoms during pregnancy and the postpartum period. Both maternal and paternal postnatal depression impact adversely on the family and the child's behavioural development. The Edinburgh Postnatal Depression Scale (*EPDS*) is a widely used instrument for the measurement of risk for postnatal depression and the State Trait Anxiety Inventory (*STAI-state*) instrument is well established as a measurement of present anxiety. The recently-developed Parent's Postnatal Sense of Security (*PPSS*) instrument measures parents' security in the early postnatal period. *Aim:* The aims of this paper were to determine the levels of correlation between scores on the *PPSS*, the *EPDS* and the *STAI-state* instruments and to test concurrent validity between the *EPDS* and *STAI-state* instruments. *Methods:* A questionnaire study which included socio-demographic questions, the *EPDS*, the *STAI-state* and the *PPSS* instruments was sent out to 160 mothers and 160 fathers. A total of 71% of the mothers and 63% of the fathers answered the questionnaires. Correlation tests (Spearman) were used to examine relationships between answers to the three questionnaires. *Results:* Significant correlations were shown between the *PPSS*, the *EPDS* and the *STAI-state* for both mothers and fathers. A significant correlation between the *EPDS* and the *STAI-state* instrument for both mothers and fathers indicated substantial concurrent validity. *Conclusions and implications:* A correlation between parents' postnatal sense of security, risk for postnatal anxiety and depression were indicated in this study. The *PPSS* instrument may be used to give an early indication of those at risk for postnatal distress, anxiety and depressive illness.

Key words: Anxiety, EPDS, PPSS, postnatal depression, preparation for parenthood, sense of security, STAI-state

Background

Researchers have reported that approximately 10% - 15% of mothers suffer from depressive symptoms during pregnancy and the postpartum period (1, 2) and a recent meta-analysis has shown that about 10% of fathers also suffer from these symptoms (3). Both maternal and paternal postnatal depression may impact adversely on outcomes for the child, for example psychiatric disorders and the child's behavioural development (4, 5). Infant temperament may be negatively affected and the child's biological response to stress may also be compromised (4, 5). It has been demonstrated that states of anxiety and depression are closely related (6, 7) and that more research is needed (8). Tammentie et al (9) showed that in postnatally depressed mothers, there was a discrepancy between their expectations of early parenthood and the reality they experienced. It is possible that this discrepancy might be linked to risk for postnatal depression. Parents, especially mothers, wanted the experience of parenthood to be perfect and had high expectations of their life as a family. Despite these expectations, there was a sense that the infant tied them down. Tammentie et al pointed out the need for support in the postpartum period as an emergent theme in their analysis (9).

Demands and expectations, from both parents, concerning fathers' participation during childbirth, responsibility for the child and equal participation in family life, have increased, in recent years, especially in high-income countries (10-13). It has been shown that increased involvement of fathers in maternity care, both pre- and postnatally, increases mothers', fathers' and children's wellbeing and may be protective in maternal and paternal depression (14, 15). Persson et al (16, 17) have shown that fathers' participation from early pregnancy and parents' antenatal preparation for the early postnatal period are important factors for postnatal sense of security for both parents.

The Edinburgh Postnatal Depression Scale (*EPDS*) (1) is a widely used and validated instrument for the measurement of risk for postnatal depression, which was originally developed for use in women, but is in some studies used also for fathers (18, 19). The scale measures both anxiety and depression and is commonly used at approximately two to three months postpartum. Recently Massoudi et al (20) showed that that the *EPDS* scale picked up more worry, anxiety and unhappiness than depression when used for fathers. The *STAI*-state (State Trait Anxiety Inventory) instrument was developed to measure present anxiety (21). However, this instrument is not a specific instrument for the postnatal period but may be used for evaluating anxiety in any specific situation (22).

Feelings of security in the early postnatal period are important for parents and for the wellbeing of the infant (16, 17, 23). Earlier research has identified factors important for parents' postnatal sense of security and the instrument Parents' Postnatal Sense of Security (*PPSS*) has been developed to measure these factors (23). Factor analysis identified four dimensions of importance for security during the first week after childbirth, for both parents. Three of the dimensions were common to mothers and fathers; a sense of the midwives'/nurses' empowering behaviour, a sense of affinity within the family and a sense of general well being. For the mothers, the fourth dimension was a sense that breast-feeding was manageable and for the fathers the fourth dimension was a sense of the mother's well being which included breast-feeding (23).

We hypothesise that a lack of security in the early postnatal period may be a contributing factor to anxiety and depressive illness. Thus, if correlations exist between the *PPSS*, *EPDS* and *STAI*-state, it may be possible to identify parents at risk for anxiety and depression in an earlier phase. This could help health care providers to form the provision of individually

tailored postnatal care in order to reduce the risk for depressive illness. Early identification and treatment of parental depressive symptoms may improve infant outcomes.

The aims of this paper were to determine the levels of correlation between scores on the *PPSS*, the *EPDS* and the *STAI*-state instruments and to test concurrent validity between the *EPDS* and *STAI*-state instruments.

Materials and Methods

This paper represents an analysis of material that was collected in a previous study (23). Questionnaires were sent to mothers and fathers separately, approximately ten weeks postpartum.

Participants and Setting

Mothers and fathers ($n = 160 + 160$) to every fifth child born live at term (≥ 37 weeks) at five hospitals in southern Sweden were sent a letter of invitation to participate in the study by answering questionnaires. After two reminders, 113 (71%) of the mothers and 99 (63%) of the fathers returned the questionnaires. The mean age for mothers was 29.8 years (17-40) and for fathers 32.4 years (21-52). Most of the parents were cohabitating (106 of the mothers and 94 of the fathers). For 52 (46%) of the mothers and 48 (49%) of the fathers it was their first child. Ninety-nine (88%) of the mothers had experienced a normal vaginal birth and 89 (91%) of the fathers answered that their partner had had a normal vaginal birth. All deliveries were singleton. Twelve (11%) of the mothers and 7 (7%) of the fathers had basic school education (9 years), 53 (47%) of the mothers and 58 (59%) of the fathers had high-school education (12 years) and 44 (39%) of the mothers and 31 (31%) of the fathers had a university/college education. Of the mothers 92 (81%) and of the fathers 81 (82%) were born in Sweden.

Procedures and measurements

The questionnaire included socio-demographic variables, the *EPDS* instrument (1), *STAI*–state instrument (18) and the *PPSS* instrument (23).

The *EPDS* instrument consists of 10 items in the form of statements on four point Likert-scales (0-3) to which the parents are asked if they agree. Possible scores are between 0-30. A higher score denotes a higher risk for postnatal depression.

The *STAI*-state instrument is well validated and consists of 20 items measured on four point Likert scales (1-4). Possible scores are between 20 and 80. A higher score denotes a higher level of anxiety.

The *PPSS* instrument exists in two similar versions, one for mothers (Cronbach's coefficient alpha 0.88) and one for fathers (Cronbach's coefficient alpha in the present study was 0.77) (23). The items are statements on four point Likert-scales to which the parents are asked if they agree or not, (1- strongly disagree and 4 – strongly agree). The mother's version of the *PPSS* instrument consists of 18 items (possible score 18-72) and the father's version of 13 items (possible score 13-52). A higher score denotes a higher sense of security.

Both the *EPDS* instrument and the *STAI*- state instrument measured, in this study, the situation approximately ten weeks after childbirth. Parents were also asked to re-call the first postnatal week and to answer items on the *PPSS* instrument. Their answers were used to estimate the correlation between *PPSS* the first postnatal week with risk for depression and anxiety at approximately 10 weeks after birth.

Statistics

Spearman's correlation coefficient (*rho*) was used to examine relationships between the *PPSS*, the *EPDS* and the *STAI* - state instruments. This correlation test was also used to test concurrent validity between the *EPDS* and the *STAI-state* instruments. Correlation coefficients of 0.1-0.3 were considered to indicate weak relationship, 0.3-0.5 a moderate relationship and > 0.5 a strong relationship (24). Concurrent validity measures how well an instrument correlates with another established instrument within the same area.

Ethical considerations

The project was implemented in compliance with the ethical principles of the Declaration of Helsinki. The material that this paper is based on is taken from an earlier published study (reference removed for peer review) which was granted permission by the Research Ethics Committee at (name removed for peer review) University, (number removed for peer review). The letter of invitation contained details of the study, explained that participation was completely voluntary and refusal to join the study would in no way effect present or future care. Informed consent was assumed if parents returned a completed questionnaire.

Results

Correlations between PPSS, EPDS and STAI – state instruments

Statistically significant correlations were shown between scores for postnatal sense of security one week after childbirth, anxiety and the risk for postnatal depression after childbirth for both mothers and fathers; lower scores for security correlate with increased risk for anxiety and depression (Table 1).

Insert table 1 here

Concurrent validity between EPDS and STAI state

Results of the analysis of concurrent validity showed high levels of correlation between the *EPDS* and *STAI-state* instruments for both mothers and fathers; higher levels of anxiety correlate with an increased risk for depression (Table 2).

Discussion

The results from the present study show moderate to strong correlations between the *PPSS*, *EPDS* and *STAI* – state instruments for both mothers and fathers. Correlations between *PPSS* and *EPDS* and also between *PPSS* and *STAI-state* for mothers were strong, whereas for fathers the correlation between *PPSS* and *EPDS* was weak. However, fathers' scores for *PPSS* and *STAI-state* were moderately well correlated which is a finding in line with findings from Massoudi et al (20). This suggests that our hypothesis that a lack of security in the early postnatal period might predispose parents to anxiety and postnatal depression and that anxiety and depression can be seen as indicators of insecurity. The fact that there was a strong concurrent validity between measurements of anxiety and depression indicates that a care environment aimed at reducing anxiety might also help reduce the occurrence of postnatal depression, which can be a serious problem for new families. This probably also indicates that anxiety increases when a person does not feel secure, which is not surprising. A limitation to this study is the risk for recall bias when asking parents to fill in the *PPSS* instrument 10 weeks after the event. Ideally, the *PPSS* instrument should be administered within two weeks from the birth. However, it has been demonstrated that women's memories of childbirth are generally accurate (25).

Earlier research has indicated areas of particular importance for mothers' and fathers' feelings of security in the immediate postnatal period; being well prepared for the time after birth, fathers' participation from early pregnancy and throughout the childbirth period, knowing who to turn to for advice, being met as an individual and having a planned follow-up before discharge (16, 17). The quality of antenatal preparation for parenthood in Sweden has been discussed (26) but little change has ensued since these discussions. Care providers must focus on giving a realistic picture of life with a small infant in order to reduce the discrepancy between expectations and reality felt by many parents (9). It is possible that friends, relatives and care providers, in an effort to not disrupt pre-birth bonding, avoid discussion of the difficulties of early parenthood. This may be a misguided approach.

If we can increase parents' postnatal sense of security by helping them to be prepared we may decrease the frequency of anxiety and postnatal depression and in that way improve the well-being of the child. Research has focused on the *effects* of preparation for childbirth but little is known regarding parents' views concerning their needs for preparation for parenthood. It is therefore essential to investigate, using qualitative research methods, preparation needs as expressed by parents. By also focusing on both paternal and maternal perspectives, the father's position can be consolidated, gender equality can increase and the family unit can be strengthened. It has been shown that an involved father has positive effects on the attachment to his child (27).

Conclusions, implications and further research

A correlation between parents' postnatal sense of security and risk for postnatal depression was indicated in this study. Further research will address the limitations of the present study so that data is collected prospectively and the *PPSS* instrument is administered in the early

postpartum. This research should also include measurement of *EPDS* and *STAI* during the antenatal period in order to provide baseline values. At present, in some areas of Sweden, mothers are screened at two months post-partum using the *EPDS*. Routine measurement of *PPSS* after one postpartum week may help to give an early indication of those at risk for postpartum distress. An increased understanding of both parents' preparation needs for the first weeks after childbirth is essential and should be investigated.

References

1. Cox JL, Holden JM, Sagovsky R (1987) Detection of postnatal depression. Development of the 10-item Edinburgh Postnatal Depression Scale. *Br J Psychiatry* 150:782-786.
2. Rubertsson C, Börjesson K, Berglund A, Josefsson A, Sydsjö G (2011) The Swedish validation of Edinburgh Postnatal Depression Scale (EPDS) during pregnancy. *Nord J Psychiatry* 65:414-418.
3. Paulson JF, Bazemore SD (2010) Prenatal and postpartum depression in fathers and its association with maternal depression: a meta-analysis. *JAMA* 303:1961-1969.
4. Ramchandani P G, Stein A, O'Connor TG, Heron J, Murray, Evans J (2008) Depression in Men in the Postnatal Period and Later Child Psychopathology; Population Cohort Study. *J Am Acad Child Adolesc Psychiatry* 47: 390-398.
5. Waxler E, Thelen K, Muzik M (2011) Maternal Perinatal Depression – Impact of Infant and Child Development. *European Psychiatric Review* 7: 41-47.
6. Ross LE, Gilbert Evans SE, Sellers EM, Romah MK (2003) Measurement issues in postpartum depression part 1: anxiety as a feature of postpartum depression. *Arch Womens Ment Health* 6:51-57.
7. Matthey S (2008) Using the Edinburgh Postnatal Depression Scale to screen for anxiety disorders. *Depress Anxiety* 25:926-931.
8. Matthey S, Fisher J, Rowe H (2013) Using the Edinburgh postnatal depression scale to screen for anxiety disorders: conceptual and methodological considerations. *J Affect Disord* 146:224-230.
9. Tammentie T, Paavilainen E, Åstedt-Kurki P, Tarkka MT (2004) Family dynamics of postnatally depressed mothers - discrepancy between expectations and reality. *J Clin Nurs* 13: 65-74.

10. WHO (2007) Fatherhood and Health outcomes in Europe. Denmark: World Health Organization Accessed [2014-01-28] from http://www.euro.who.int/_data/assets/pdf_file/0017/69011/E91129.pdf
11. McKellar L, Pincombe J, Henderson A (2008) Enhancing Fathers' Educational Experiences During the Early Postnatal Period. *J Perinat Educ* 17: 12-20.
12. SFOG [Swedish Association for Obstetric and Gynaecology] (2008) Maternity Health care, Sexual and Reproductive Health. Stockholm: Arbets- & Referensgrupp (ARG) :59.
13. Ellberg L, Högberg U, Lindh V (2010) We feel like one, they see us as two: new parents' discontent with postnatal care. *Midwifery* 26: 463-468.
14. de Montigny F, Lacharite C (2004) Fathers' perceptions of the immediate postpartal period. *J Obstet Gynecol Neonatal Nurs* 33: 328-339.
15. Schytt E, Waldenström U (2007) Risk factors for poor self-rated health in women at 2 months and 1 year after childbirth. *J Womens Health* 16:390-405.
16. Persson EK, Kvist LJ, Fridlund B, Dykes AK (2011) Mothers' sense of security in the first postnatal week: interview study. *J Adv Nurs* 67:105-116.
17. Persson EK, Kvist LJ, Fridlund B, Dykes AK (2012) Fathers' Sense of Security during the first postnatal week: a qualitative interview study in Sweden. *Midwifery* 28: e697-704.
18. Matthey S, Barnett B, Kavanagh DJ, Howie P (2001) Validation of the Edinburgh Postnatal Depression Scale for men and comparison of item endorsement with their partners. *J Affect Disord* 64:175-184.
19. Figueredo B, Conde A (2011) Anxiety and depression in women and men from early pregnancy to 3-months postpartum. *Arch Womens Ment Health* 14:247-255.

20. Massoudi P, Hwang CP, Wickberg B (2013) How well does the Edinburgh Postnatal Depression Scale identify depression and anxiety in fathers? A validation study in a population based Swedish sample. *J Affect Disord* 149: 67-74.
21. Spielberger CD, Gorsuch RL, Lushene R, Vagg PR, Jacobs GA (1983) Manual for the State-Trait Anxiety Inventory STAI (form Y) ("Self-Evaluation Questionnaire"). Palo Alto: Consulting Psychologists Press.
22. Britton JR (2005) Pre-discharge anxiety among mothers of well newborns: prevalence and correlates. *Acta Paediatr* 94: 1771-1776.
23. Persson EK, Fridlund B, Dykes AK (2007) Parents' postnatal sense of security (PPSS): development of the PPSS instrument. *Scand J Caring Sci* 21:118-125.
24. Burns N, Grove SK (1997) *The Practice of Nursing Research - Conduct, Critique & Utilisation*. Philadelphia: JB Saunders Company.
25. Simkin P (1992) Just another day in a woman's life? Part II: Nature and consistency of women's long-term memories of their first birth experiences. *Birth* 19:64-81
26. Andersson E, Christensson K, Hildingsson I (2012) Parents' experiences and perceptions of group-based antenatal care in four clinics in Sweden. *Midwifery* 28: 442-448.
27. Sarkadi A, Kristiansson R, Oberklaid F, Bremberg S (2008) Fathers' involvement and children's developmental outcomes: a systematic review of longitudinal studies. *Acta Paediatr* 97:153-158.

Acknowledgements

The authors wish to thank all parents who made this study possible. This research project was funded by Faculty of Medicine, (Lund university).

Disclosure of Interest

We have no interests to disclose.

Table 1. Correlations (*rho*) between the PPSS-instrument measuring postnatal sense of security one week after childbirth, the *EPDS* instrument measuring risk for postnatal depression and the STAI-state instrument, measuring present anxiety, approximately ten weeks after childbirth for mothers and fathers. Spearman's correlation test (2-tailed).

<i>Validators</i>	<i>EPDS</i>	<i>STAI</i>
Mothers (<i>n</i> = 111)		
<i>PPSS</i>	-0.575 **	-0.584 **
<i>EPDS</i>		0.731**
Fathers (<i>n</i> = 99)		
<i>PPSS</i>	-0.231 *	-0.352 **
<i>EPDS</i>		0.681**

* $p \leq 0.05$, ** $p \leq 0.01$ (2-tailed)