



LUND UNIVERSITY

Burn injuries in small children, a population-based study in Sweden.

Carlsson, Anna; Udén, Giggi; Håkansson, Anders; Karlsson, Elisabeth Dejin

Published in:
Journal of Clinical Nursing

DOI:
[10.1111/j.1365-2702.2006.01259.x](https://doi.org/10.1111/j.1365-2702.2006.01259.x)

2006

[Link to publication](#)

Citation for published version (APA):

Carlsson, A., Udén, G., Håkansson, A., & Karlsson, E. D. (2006). Burn injuries in small children, a population-based study in Sweden. *Journal of Clinical Nursing*, 15(2), 129-134. <https://doi.org/10.1111/j.1365-2702.2006.01259.x>

Total number of authors:
4

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

This is an author produced version of a paper published in Journal of Clinical Nursing. This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Citation for the published paper:

Carlsson, Anna and Udén, Giggi and Hakansson, Anders
and Karlsson, Elisabeth Dejin

“Burn injuries in small children, a population-based study in Sweden.”

J Clin Nurs. 2006 Feb;15(2):129-34.

<http://dx.doi.org/10.1111/j.1365-2702.2006.01259.x>.

Access to the published version may require journal subscription.

Published with permission from: Blackwell Synergy

Burn injuries in small children, a population-based study in Sweden

ANNA CARLSSON, MSc, RN

Doctoral student, School of Health and Society, Department of Nursing, Malmö University, Malmö, Sweden.

GIGGI UDÉN, PhD, RNM, RNT

Professor, School of Health and Society, Department of Nursing, Malmö University, Malmö, Sweden.

ANDERS HÅKANSSON, PhD, MD

Professor, Department of Community Medicine, Malmö University Hospital, Malmö, Sweden.

ELISABETH DEJIN KARLSSON, PhD, RNM, RNT

Senior Lecturer, School of Health and Society, Department of Nursing, Malmö University, Malmö, Sweden.

Number of words: 3479

Journal of Clinical Nursing ISSN 1365-2702

Summary

- * Epidemiological studies of burns in children have mostly been hospital-based and the cases that never reached hospital have been excluded.
- * The aim of this study was to describe characteristics of burn injuries in children (0–6 years old), consulting primary care and hospital-based care in Malmö, Sweden.
- * Burn-injured children consulting the University Hospital or the 21 Health Centres, during 1998 and 2002 were included.
- * The study had a retrospective design with data collected from medical records.
- * Chi-squared test was used to analyse differences in nominal data and cross-tabulation was used to analyse the proportions between the characteristics of the injuries and sex, age and nationality.
- * There were 148 burns, 80% of which were scalds caused by hot liquid (71%) or hot food (29%).
- * The majority of injuries were to boys between one and two years old.
- * Children of foreign-born parents were more frequently affected and the extent of the injuries often greater.
- * Almost all the accidents (96%) occurred in the home environment, while a family member was next to the child.
- * The Health Centres more often received children affected on hand/arm and by causes such as hot food than the University Hospital.
- * Our data demonstrate the importance of developing a programme for the prevention of paediatric scalds with education of family members to be aware of the danger.

Key words: burns; children; prevention and scalds

Correspondence to: Anna Carlsson, School of Health and Society, Department of Nursing, Malmö

University, SE-205 06 Malmö, Sweden (tel.: +46 40 665 79 39; fax: +46 40 665 81 00;

e-mail: anna.carlsson@hs.mah.se).

Introduction

Sweden has the lowest accident mortality in children in the world (SOU 2002:99; IRTAD, 2003). The mortality rate for accidents in Sweden in 1999 was 22/100 000 children (WHO, 2003).

Children get hurt by accidents, mostly in their homes, in the age range 0–4 years (March & Kendrick, 2000; den Hertog et al., 2000; Mukerji et al., 2000). The most common injuries leading to consultation of hospitals or primary care units are commotio, fractures, luxations, distortions and ruptures of muscles, all of them connected with fall accidents. Other accidents course drowning, poisoning, suffocation and burn injuries (March & Kendrick, 2000).

Burn injuries are often a serious tragedy for both parents and children caused by the pain and the trauma of dressings, procedures, hospitalisation and emotional adjustments in an immature child, the mortality risk, the long-term rehabilitation and the cosmetic disfigurement (Andersson, 1996; Muerji et al., 2001).

In Copenhagen, Denmark, epidemiological analyses of burn injuries have been carried out every tenth year. The majority of accidents occurred in the home, younger children were at risk, and scalding was the most common type of injury often caused by water from toppled electrical kettles (Sheller & Thuesen, 1998). Also, a one-year study of burn victims in four counties south-east England revealed that children under the age of five had the highest rate of admission to hospital. The admission from the district health authorities to hospital burn units was 14/100 000 population per year (Wilkinson, 1998). The incidence of burns in the Netherlands was 210/100 000, in 0–4-year-old children, and 65% of them were scalds. Among the children aged less than four, 80% of all scald injuries were caused by pushing down or dropping a hot liquid (den Hertog et al., 2000). In a French study, the typical burn-injury patient was a boy with a mean age

of 4 years, who suffered a scald in the kitchen, and the burn injuries covered less than 10% of the body surface (Mercier & Blond, 1996). In Canada, 67% of burns in children under five were caused by food preparation and consumption, 14% were bathtub scalds and 19% were due to flames. Of children under two years of age, 70% were affected (Ray, 1995).

In Malmö, Sweden, a 6-year prospective study of scald-injured children (0–6 years old), treated at the inpatient burn unit at the University Hospital (UMAS), ran 1990–1996. Scalding was the cause of 87% of the burn injured children (Freccero et al., 2000).

Many studies have estimated the epidemiology of paediatric burns but they have often yielded different results depending on the regions where the studies were conducted (WHO, 2004; den Hertog et al., 2000; Sheller & Thuesen, 1998; Mercier & Blond, 1996; Ray, 1995). Also, epidemiological studies of burns in children have mostly been hospital-based and the cases that never reached the hospital were excluded. It is important to prevent as many burns as possible and to achieve this it is necessary to be aware of the characteristics of both minor and severe burn injuries in children.

The aim of this study is to describe characteristics in burn injuries in children (0–6 years old), consulting primary care or hospital-based care in Malmö.

Material and Methods

STUDY AREA AND POPULATION

This study was performed in Malmö, a city with 257 574 inhabitants (2000), in Sweden during 1998 and 2002. There were 21 Health Centres and one University Hospital in Malmö. Among other specialities, the hospital also had a department for burn injuries

with plastic surgery. Parents of children with burn injuries could voluntarily choose to consult either the emergency unit at the hospital or the Health Centre in their neighbourhood (includes a doctor-on-duty centre).

DATA COLLECTING

The data was collected from medical records, in a retrospectively designed study. All emergency incoming patients at the University Hospital were recorded, according to age, sex and diagnosis. Approximately 10 000 incoming patients per year were registered. All the burn injuries in children found in these registers in both 1998 and 2002 were compiled through medical records, by one of the authors (AC). The internal dropout was due to the fact that not all of the questions could be answered through the medical records.

Children admitted during the study period with any type of burn-related injury were selected for review. Data on the burn-injured children consulting the Health Centres were collected through the diagnosis registration.

Patients who consulted the University Hospital and the Health Centres in Malmö are coded in accordance with the *International Classification of Diseases, 9th revision* (ICD-9, 1990). All medical records of the children were registered for sex, age and nationality of the parents.

DEFINITIONS

The nationality of the parents was defined as Swedish-born when at least one of the parents was born in Sweden and as foreign-born when both parents were born in any country outside Sweden (Landstingsförbundet, 2002). Body site, extent and depth of injuries but also causes and location of injuries were investigated. The extent of the

injuries was classified in two groups: <2% and >2% (den Hertog et al., 2000). The depth of the injuries was categorised in three degrees: degree 1 was redness in the skin, but intact, degree 2 was blisters and degree 3 was when parts of the skin need to be taken away for recovering (Freccero et al., 2000).

STATISTICAL ANALYSIS

To analyse differences in nominal data, chi-squared test was used and a difference was considered statistically significant if $p \leq 0.05$. The cross tables was used to analyse the proportions between sex, age, nationality and injuries. The statistical analyses were performed using the SPSS program for Windows 11.0 (Norusis, 2002).

The study was approved by the Ethics Committee of Lund University (LU 37-99).

Results

During the study period 1998 and 2002, 148 children were treated for burn injuries. Children aged 1–2 years suffered most frequently ($p=0.0001$). Boys were represented to the largest extent with 64% and girls with 36% ($p=0.001$). Table 1 shows the distribution of children according to their sex and age.

Children aged 0–2 years suffered from injuries on the hand, arm, foot or leg to a significantly higher level ($p=0.04$). Children with foreign-born parents suffered more often from burn injuries than children with Swedish-born parents ($p=0.001$). Children with foreign-born parents also had a larger extent of injury more often than children with Swedish parents ($p=0.04$) (Table 2).

Burns were primarily scalds 81% ($n=95$) ($p=0.001$). The non-scald burns were caused by a hand on a stove, a foot in a hot candle-wax, a hand on a lamp or by sitting down on a barbeque grill.

The scalds were mostly caused by hot liquid 71% (n=67) (p=0.001). Scalds by hot food caused the injuries in 29% of cases (n=28). Accident scenarios in which the child pulled hot food or hot liquid off the stove frequently involved the child trying to climb up and/or reach a pan on the stove or the worktop. The extent of injury was largest to the children injured by hot liquid (p=0.01). The scalds happened mostly in home environment (95%), (p=0.001) when a family member was present. The depth of injury was often greater in the accidents that occurred outside the home (p=0.01) (Table 2).

The distribution between the children consulting the Health Centres, 49% (n=73), and the University Hospital, 51% (n=75), differed according to causes of accidents. The University Hospital received more children with other causes of injuries than hot liquid/hot food (p=0.04) and the site of injury was more often the face (p=0.001) or the leg/foot (p=0.001) or with several sites involved (p=0.001). Those children also suffered more often from injuries of a greater extent (p=0.001) and depth (p=0.05). The Health Centres more often received children affected by hot food than the University Hospital (p=0.03). Arms and hands were more involved in the injuries consulting the Health Centres (p=0.03) (Table 3).

Discussion

In this study 72% (n=106) of the burn-injured children were under 3 years old. This age group is also seen as the most common age group for burn injuries in other studies (Sheller & Thuesen, 1998; den Hertog, 2000; Wilkinson, 1998). Children aged under 3 often stay near their parents while they are cooking and therefore they are more exposed to burn risks. Almost all of the children (96%) in the present study had their parent close by when the accident occurred. Many of those accidents could have been avoided as they were caused by preventable events such as when the handle of a pan was reachable

from the cooker front or overheated liquids were handled when the children were nearby.

According to earlier study and as shown in the present study, there seems to be a peak in the affected child between 1 and 2 years old, and at that age close observation of the child is a main rule. Education about preventive precautions can be added to observations, starting at around 2 years old. When children ask questions it offers opportunities for teaching about risks in daily life. From about 3 years children understand the concept of danger (Mercier & Blond, 1996).

As in other studies (den Hertog, 2000; Mercier & Blond, 1996) boys were more frequently affected by burn injuries than girls.

The frequency of burn injuries, in particular scalds with a larger extent, was higher among children whose both parents were born abroad than for children of parents of Swedish origin. These findings are supported by studies in the Netherlands (den Hertog, 2000). It is not yet known whether this is due to possible differences in cultural background or economic position. Interesting future research can be directed to the more general question of how a lower social economic position, problems in reading and understanding instructions in another language, a limited living situation, unemployment and distinctive cultural traditions influence the risk of accidents among children.

Previous studies have shown that burn injuries in children are most frequently scalds caused by hot liquid, especially hot coffee or tea (den Hertog et al., 2000; Sheller & Thuesen, 1998). The present study showed the same causes, although frequently caused by hot food (29%). Burns caused by hot food were more frequent among the children consulting the Health Centres; these burn injuries are seldom described in the literature and may be the reason for the higher extent of hot food causes in the present

study. Other studies have shown that hot baths were the most frequently observed causes of scalds (Fukunishi et al., 2000; Mercier & Blond, 1996; Ray, 1995). None of the 148 children in the present study had a scald caused by hot bath water. A major reason may be the regulation of water temperature in taps, to 55 degrees centigrade, according to Swedish law. The limit of 55 degrees is set due to the risk of infection from legionnaires disease.

Most of the injuries were defined as degree 1+2 (redness and blisters) and none of the injuries had a fatal outcome. Many other studies show that the fatal incidence of injuries is often caused by flames or hot tap water (den Hertog, 2000; Fukinishi et al., 2000). Those causes have not been seen in the present study.

During the study period of 1998 and 2002, 148 children were treated for burn injuries. In Malmö, the total numbers of reported accidents in children (0–6 years old) in 1998 and 2002 were 1300 and 1478 respectively (n=2778). About 30% of the patients consulted the Health Centres and 70% consulted the University Hospital. Eighty-five percent (n=1653) of the children consulting the University Hospital for accidents in the home environment, had parents of foreign origin. Scald injuries were not separated from other accidents in the injury registrations nor in the registrations for Malmö or for Sweden as a whole (Dept. of Social Medicine, 2004).

It is a long-term process to prevent child accidents. This long-term process cooperates to motivate the parents to take precautions to prevent accidents (McClure & Douglas, 1996; Arborelius, 1993). The responsibilities of the landlords, concerning the environmental risks in flats, are well regulated in Sweden but not always so well known by the citizens. Campaigns aimed to educate the public about environmental risks in children can be advertised on the airwave media, on billboard posters, at nursery schools and in the Child Health Services.

Some of the results in the present study are reduced in internal validity due to the internal dropout values between 8% and 29% (Rothman & Greenland, 1998). A negative effect of conducting a record study is the limitations in answering all questions, due to the documentation.

A strength of this study is that it aimed to find all children with burn injuries consulting medical care in Malmö and carefully went through the 20 000 diagnoses at the emergency unit at the University Hospital and all diagnoses in the registration at the Health Centres during 1998 and 2002. The children not reached in the present study were those with burn injuries not consulting medical care.

Conclusions

From the investigation of burn injuries in children aged less than 7 years consulting the University Hospital and the Health Centres in Malmö during 1998 and 2002, the following results were obtained:

- * Burn injuries were primarily scalds
- * The scalds were caused by hot liquid in 71% and by hot food in 29% of cases
- * Children aged 1–2 years suffered most frequently and the sites of injuries were more often on the hand, arm, foot or leg
- * More boys than girls suffered from burn injuries
- * Most of the burn accidents occurred in the home environment when the parent was close by
- * Children of foreign-born parents had burn injuries more frequently than children of Swedish-born parents and they also more often had injuries on the hands, arms, leg or feet and injuries of larger extent

- * The Health Centres more often received children affected on the hand/arm and the causes were more often hot food

Acknowledgement

This study has been supported by the Primary Health Care Research Department in Scania (Primärvården Skåne, FoUU).

References

- Andersson, I. (1996) *Fakta om skador bland barn och ungdom. (Facts about injuries among children and youth)*. Stockholm: Barnombudsmannen, I. (In Swedish)
- Arborelius, E. (1993) *Teori och praktik om att påverka människors levnadsvanor. (Theory and practice to affect patients' lifestyles)* Stockholm: Allmänmedicinskt utvecklingscentrum. (In Swedish).
- den Hertog, P C, Blankendaal, F, Ten Hag, S M. (2000) Burn injuries in the Netherlands. *Accident Analysis and Prevention*, 32, p. 355-364.
- Department of Social Medicine. (2004) *Skaderegistrering i Region Skåne. (Injury registration in Region Scania)*. University Hospital UMAS, in Malmö. (In Swedish).
- Freccero, C, Svensson, H, Kalhor, S, Larsson, A. (2000) Scalding injuries in immigrant families. *Scand. J. of Plastic and Reconstruc. Surg. and Hand Surg.* 34 (4). P. 309-13.
- Fukunishi, K, Takahashi, H, Kitagishi, H, Matsushima, T, Kanai, T, Ohsawa, H, Sakata, I. (2000) Epidemiology of childhood burns in the Critical Care Medical Center of Kinki University Hospital in Osaka, Japan. *Burns* 26, p. 465-469.
- ICD-9 (1990) *International Classification of Diseases, 9th Revision, Clinical Modification*. 3rd ed. Washington, DC: Health Care Financing Administration.
- IRTAD, OECD-International Road Traffic and Accident Database,
www.bast.de/htdocs/fachthemen/irtad/, 2003 08 25.
- Landstingsförbundet. (2002) *Statistisk årsbok. (The Federation of Swedish County Councils)*. Stockholm: Landstingsförbundet. (In Swedish)
- March, P, Kendrick, D. (2000) Near miss and minor injury information – can it be used to plan and evaluate injury prevention programmes? *Accident Analysis and*

- Prevention* 32, p. 345-354.
- McClure, R, Douglas, R M. (1996) The public health impact of minor injury. *Accident Analysis and Prevention* 28, (4), p. 443-451.
- Mercier, C. & Blond, M H. (1996) Epidemiological survey of childhood burn injuries in France. *Burns* 22, p.29-34.
- Mukerji, G, Chaminia, S, Patidar, G P, Gupta, S. (2000) Epidemiology of paediatric burns in Indore, India. *Elsevier Science*, India.
- Mukerji, G, Shobha, C, Patidar, G H, Suurabh, G. (2001) Epidemiology of paediatric burns in Indore, India. *Burns* 27, p. 33-38.
- Norusis, M. (2002) *The SPSS 11.0 guide to data analysis*. Upper Saddle River, N.J.: Prentice Hall.
- Ray, J G. (1995) Burns in young children: a study of mechanism of burns in children aged 5 years and under in the Hamilton, Ontario Burn Unit. *Burns* 21, p. 463-466.
- Rothman, K J. & Greenland, S. (1998) *Modern Epidemiology*. Second edition. Philadelphia: Lippincott-Raven Publishers, p. 119.
- Sheller, J L P. & Thuesen, B. (1998) Scalds in children caused by water from electrical kettles: effect of prevention through information. *Burns* 24, p. 420-424.
- SOU 2002:99 Statens offentliga utredningar. (State Public Inquiries). *Barns skador i Sverige. (Children's injuries in Sweden)*. Barnsäkerhetsdelegationen, Stockholm 2002. (In Swedish).
- World Health Organization (WHO).
- http://www3.who.int/whosis/mort/table1_process.cfm , 2003 09 11
- World Health Organization (WHO). <http://www.who.int/ceh/risks/cehinjuries2/en/> , 2004 07 05
- Wilkinson, E. (1998) The epidemiology of burns in secondary care, in a population of

2.6 million people. *Burns*, 24, p. 139-143.

Table 1 Distribution of children according to age and sex (number, percentage and p values).

Age	Boys			Girls			Total		
	N	%	p value	N	%	p value	N	%	p value
≤1	18		NS	11		NS	29	20	NS
1–2	50		NS	27		NS	77	52	0.0001
3–4	19		NS	10		NS	29	20	NS
5–6	8		NS	5		NS	13	8	NS
Total	95	64	0.001	53	36		148	100	NS

NS= non significant

Table 2 Differences regarding burn injuries in small children (N=148).

		Dimensions of injuries					
		Extent of injury ¹		Site of injury ²		Depth of injury ³	
Variable	%	%	p value	%	p value	%	p value
Age							
3–6 y	28	42		80		11	
0–2 y	72	58	NS	88	0.04	6	NS
Sex							
Girls	36	53		78		9	
Boys	64	54	NS	90	NS	6	NS
Parents' origin							
Swedish-born	44	44		76		7	
Foreign-born	57	63	0.04	92	0.04	7	NS
Cause of injury							
Hot food	29	36		100		7	
Hot liquid	71	67	0.01	93	NS	6	NS
Location							
Home	95	52		96		6	
Outside	5	80	NS	100	NS	33	0.01

¹ =2%² Hand, arm, foot and leg³ Degree 3

Table 3 Distribution of children according to place of consultation and occurrence of injuries, in percent (N=136).

Occurrence of injuries	Consultation		p value
	Health Centre/ D.on dc. ¹	University Hospital	
	%	%	
Cause of accident			
Hot liquid	36	64	NS
Hot food	59	41	0.03
Other	29	71	0.04
Site of injury			
Face	17	83	0.001
Trunk	42	58	NS
Arm/hand	61	39	0.03
Leg/foot	21	79	0.001
>3 of above	0	100	0.001
Extent of injury			
A) < 2%	75	25	NS
B) = 2%	41	59	0.001
Depth of injury			
Degree 1+2	55	45	NS
Degree 3	22	78	0.05
Total	51	49	

¹) D. on dc.=Doctor-on-duty centre.