



LUND UNIVERSITY

A Fallacy on Correlated White Noise Processes

Åström, Karl Johan; Hagander, Per

1973

Document Version:

Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (APA):

Åström, K. J., & Hagander, P. (1973). *A Fallacy on Correlated White Noise Processes*. (Research Reports TFRT-3105). Department of Automatic Control, Lund Institute of Technology (LTH).

Total number of authors:

2

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

A FALLACY ON CORRELATED WHITE NOISE
PROCESSES

K.J. ÅSTRÖM
P. HAGANDER

TILLHÖR REFERENSBIBLIOTEKET
UTLÄNAS EJ

Report 7322 (C) July 1973
Lund Institute of Technology
Division of Automatic Control

A FALLACY ON CORRELATED WHITE NOISE PROCESSES

K.J. Åström

P. Hagander

A FALLACY ON CORRELATED WHITE NOISE PROCESSES

Let T be the set of positive and negative integers and let $\{e(t), t \in T\}$ and $\{v(t), t \in T\}$ be white noise processes. It is tempting to believe that the covariance function $r_{ve}(\tau)$ is nonzero in one point only. It is shown that this is not the case.

Example

Consider the process v defined by

$$v(t) = a_0 e(t) + a_1 e(t-1) + \varepsilon(t) + b_1 \varepsilon(t-1)$$

with e and ε being independent white noise processes with variance 1. The covariance function for v is

$$r_v(t,s) = \begin{cases} a_0 a_1 + b_1 & |t-s| = 1 \\ a_0^2 + a_1^2 + 1 + b_1^2 & t = s \\ 0 & \text{otherwise} \end{cases}$$

and the cross covariance between v and e :

$$r_{ve}(t,s) = \begin{cases} a_1 & s = t-1 \\ a_0 & s = t \\ 0 & \text{otherwise} \end{cases}$$

If the parameters are chosen in such a way that

$$b_1 = -a_0 a_1$$

the process $\{v(t), t \in T\}$ is obviously a white noise process. We have thus given two white processes whose joint covariance function is different from zero for two arguments and a counterexample is provided.

The idea behind the example is that if three stochastic variables are given and if the correlation is known between two pairs say (x_1, x_2) and (x_2, x_3) it is not possible to say anything in general about the correlation between x_1 and x_3 .

The example can obviously be generalized and extended in many ways.