



LUND UNIVERSITY

Cardiovascular autonomic neuropathy associated with carotid atherosclerosis in Type 2 diabetic patients.

Gottsäter, Anders; Rydén Ahlgren, Åsa; Szelag, Beata; Hedblad, Bo; Persson, J; Berglund, Göran; Wroblewski, M; Sundkvist, Göran

Published in:
Diabetic Medicine

DOI:
[10.1046/j.1464-5491.2003.00956.x](https://doi.org/10.1046/j.1464-5491.2003.00956.x)

2003

[Link to publication](#)

Citation for published version (APA):

Gottsäter, A., Rydén Ahlgren, Å., Szelag, B., Hedblad, B., Persson, J., Berglund, G., Wroblewski, M., & Sundkvist, G. (2003). Cardiovascular autonomic neuropathy associated with carotid atherosclerosis in Type 2 diabetic patients. *Diabetic Medicine*, 20(6), 495-499. <https://doi.org/10.1046/j.1464-5491.2003.00956.x>

Total number of authors:
8

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Cardiovascular autonomic neuropathy associated with carotid atherosclerosis in Type 2 diabetic patients

A. Gottsäter, Å. Rydén-Ahlgren*, B. Szelag†, B. Hedblad†, J. Persson‡, G. Berglund†, M. Wroblewski† and G. Sundkvist§

Departments of Vascular Diseases, *Clinical Physiology, †Medicine, ‡Cardiology, and §Endocrinology, University of Lund, University Hospital, Malmö, Sweden

Accepted 25 February 2003

Abstract

Aims To clarify if cardiovascular autonomic neuropathy is associated with carotid artery atherosclerotic plaques in Type 2 diabetic patients.

Methods Cardiovascular autonomic nerve function was related to carotid artery ultrasound in 61 Type 2 diabetic patients 5–6 years after diagnosis of diabetes.

Results Cardiovascular autonomic neuropathy [abnormal age corrected expiration/inspiration (E/I) ratio or acceleration index (AI)] was found in 13/61 (21%) patients. Patients with cardiovascular autonomic neuropathy showed increased degree of stenosis in the common carotid artery ($24.6 \pm 13.2\%$ vs. $14.7 \pm 9.2\%$; $P = 0.014$) and a tendency towards a higher plaque score (4.0 ± 1.7 vs. 3.2 ± 1.6 ; $P = 0.064$). Controlled for age, AI correlated inversely with degree of stenosis ($r = -0.39$; $P = 0.005$), plaque score ($r = -0.39$; $P = 0.005$), and mean ($r = -0.33$; $P = 0.018$) and maximum ($r = -0.39$; $P = 0.004$) intima-media thickness in the common carotid artery. In contrast, E/I ratio correlated only slightly with mean intima-media thickness in the common carotid artery ($r = -0.28$; $P = 0.049$).

Conclusions Cardiovascular autonomic neuropathy was associated with carotid atherosclerosis in Type 2 diabetic patients. Abnormal E/I ratios reflect efferent structural damage to parasympathetic nerves whereas abnormal AI reflects afferent autonomic dysfunction possibly due to impaired baroreceptor sensitivity secondary to carotid atherosclerosis.

Diabet. Med. 20, 495–499 (2003)

Keywords Type 2 diabetes mellitus, carotid atherosclerosis, autonomic neuropathy

Abbreviations AI, acceleration index; E/I ratio, expiration/inspiration ratio; SD, standard deviation

Introduction

Autonomic neuropathy is a serious complication of diabetes [1], implicated in sudden death [2], nephropathy [3–5] and proliferative retinopathy [6,7]. Moreover, cardiac complications such as silent myocardial infarction [8], reduced pain

threshold for myocardial ischaemia [9], cardiac hypertrophy [10], impaired cardiac function during exercise [11], and disturbed coronary blood flow [12] have also been associated. We have shown connections between cardiovascular autonomic neuropathy and hyperinsulinaemia [13] and increased plasminogen activator inhibitor 1 activity [14], indicating a link between cardiovascular autonomic neuropathy and the metabolic syndrome. Others [15,16] have recently confirmed these assumptions. Hence, there is evidence that cardiovascular autonomic neuropathy may damage the vasculature in diabetic patients [17].

Correspondence to: Dr. Anders Gottsäter, Department of Vascular Diseases, University of Lund, University Hospital, S-205 02 Malmö, Sweden. E-mail: anders.gottsater@skane.se

Recently, Töyry *et al.* reported an association between stroke and cardiovascular autonomic neuropathy in Type 2 diabetic patients [18]. We therefore evaluated whether carotid artery atherosclerosis was associated with cardiovascular autonomic neuropathy as evaluated by the heart rate reactions to deep breathing [19] and tilting [20], amongst Type 2 diabetic patients in whom we previously found associations between parasympathetic neuropathy and the metabolic syndrome [14]. The aims of this study were to clarify the relationship between atherosclerotic plaques in the carotid artery and cardiovascular autonomic neuropathy.

Patients and methods

Patients

From a population-based study of the diabetes incidence in Malmö between 1990 and 1992 [21], all Type 2 diabetic patients below 60 years of age at diagnosis ($n = 91$) were invited to an assessment of cardiac autonomic nerve function and biochemical risk profile 5 years after diagnosis, and 76 accepted [14]. One year later, 61/76 (80%) patients underwent carotid ultrasound examination for quantification of atherosclerosis. Their age then ranged from 40 to 64 years, median 54. In this report, data from all 61 patients that underwent carotid ultrasound examination are reported.

Informed consent was obtained from all subjects. The study was conducted in accordance with the Declaration of Helsinki and approved by the ethics committee of the University of Lund.

Methods

Body mass index was measured as weight (kg)/height (m^2). Blood pressure was defined in the supine position with a sphygmomanometer in the right upper arm after 10 min rest.

Autonomic nerve function tests, deep breathing test (R–R interval variation)

Six maximal expirations and inspirations were performed during 1 min in the supine position during the recording of a continuous ECG, and the R–R intervals were recorded. The expiration/inspiration (E/I) ratio, a test of parasympathetic vagal nerve function, was calculated as the mean of the longest R–R interval during expiration (E) divided by the mean of the shortest R–R interval during inspiration (I) [19].

The immediate heart rate reaction to tilt

After 10 min of supine rest, the subject was rapidly tilted (< 2 s) to the upright position (head up 90°) and remained there for 8 min. The initial heart rate reaction to tilt, an immediate acceleration followed by a transient deceleration, was evaluated by continuous ECG recording and determination of the acceleration and brake indices [20]. In the formulae below A indicates mean R–R interval before tilt, B indicates the shortest R–R interval during the immediate acceleration, and C indicates the longest R–R interval during the deceleration. The acceleration index

(AI) is $(A - B)/A \times 100$ and the brake index (BI) is $(C - B)/A \times 100$. Although the immediate postural acceleration (AI) in heart rate reflects withdrawal of parasympathetic nerve activity and the later transient deceleration reinstatement of parasympathetic nerve activity, the indices also evaluate sympathetic and not only parasympathetic nerve function [22,23]. The difference between tests (repeatability) of AI is 24% and of BI 29%.

Definitions of abnormalities

The E/I ratio, AI, and BI, were expressed in age corrected values [i.e. z -scores in standard deviations (SD)] [24]. Values < -1.64 SD (95% confidence interval, one-sided test) below the age-related reference values were considered abnormal.

Ultrasound investigation

Ultrasound investigation of the right carotid artery was performed using Acuson 128/XP10 (Acuson, Mountain View, CA, USA) equipped with a linear array 7-MHz transducer with subjects supine and the head turned 45° to the left. The right carotid artery was scanned for the presence of plaques, defined as focal thickenings of the arterial wall. The extent of atherosclerosis was determined on-line according to a semi-quantitative scale as a plaque score ranging from '0' (representing a vessel without plaques or wall thickenings, the latter defined as intima media thickness ≥ 1.25 mm) to '5' (representing a vessel containing two or more plaques ≥ 10 mm² or one circumferential plaque or a plaque causing $\geq 50\%$ stenosis). The degree of stenosis was decided based on blood flow velocity at the location of maximum lumen diameter reduction. When no increase in flow velocity (change in Doppler shift) could be detected, the degree of stenosis was subjectively decided from eye-balling the plaque on-line and determining to what extent the plaque protruded into the lumen (maximum 30%, otherwise an increase in flow velocity should be possible to detect). From R-triggering in an ECG-tracing, end-diastolic images were recorded for off-line analysis of intima-media thickness and lumen diameter in the common carotid artery, and intima-media thickness in the bulb of the carotid bifurcation in a computer-assisted analysing system with the identity of the subject coded.

This method has been described previously [25,26]. It has been proven valid in measurements of atherosclerosis [27,28] and the reproducibility is acceptable [29].

Statistical analysis

Differences between groups were evaluated with the Mann–Whitney U -test. Differences in frequency were evaluated with the χ^2 test. Partial correlations were evaluated, controlled for age. Tests were two-tailed and P -values < 0.05 were considered significant. Results are presented as mean \pm SD. SPSS, version 10.0 software (Chicago, IL, USA) was used for the statistical calculations.

Results


Patient characteristics are shown in Table 1. Cardiovascular autonomic neuropathy (an abnormal age corrected E/I ratio

Table 1 Patient characteristics 5 years after diagnosis of diabetes and results of carotid ultrasound investigation 1 year later in 61 Type 2 diabetic patients

	Cardiovascular autonomic neuropathy		P-value
	With <i>n</i> = 13	Without <i>n</i> = 48	
Age (years)	53 ± 6	53 ± 7	0.685
Females [<i>n</i> (%)]	3 (23)	18 (37)	0.320
Body mass index (kg/m ²)	31.9 ± 6.7	28.6 ± 4.8	0.159
Systolic blood pressure (mmHg)	145 ± 16	144 ± 18	0.853
Diastolic blood pressure (mmHg)	89 ± 8	89 ± 10	0.930
<i>Common carotid artery</i>			
Plaque score	4.0 ± 1.7	3.2 ± 1.6	0.064
Degree of stenosis (%)	24.6 ± 13.2	14.7 ± 9.2	0.014
Mean intima-media thickness (mm)	0.994 ± 0.288	0.866 ± 0.146	0.267
Maximum intima-media thickness (mm)	1.291 ± 0.382	1.081 ± 0.250	0.120
Mean lumen diameter (mm)	7.200 ± 1.175	6.589 ± 0.980	0.074

Mean standard deviation if not stated otherwise. Significant difference shown in bold.

Figure 1 Degree of common carotid artery stenosis and plaque score vs. autonomic nervous function; acceleration index (AI), $[A - B]/A \times 100$, where *A* indicates mean R–R interval before tilt, *B* indicates the shortest R–R interval during immediate acceleration [20]) and expiration/inspiration (E/I) ratio [mean of the longest R–R interval during expiration (E) divided by the mean of the shortest R–R interval during inspiration (I)] in 61 Type 2 diabetic patients 5–6 years after diagnosis. AI correlated significantly with the degree of stenosis and plaque score ($r = -0.39$; $P = 0.005$ for both), whereas there were no such correlations for the E/I ratio ($r = -0.20$; $P = 0.14$ and $r = -0.11$; $P = 0.44$, respectively).



or AI) was found in 13/61 (21%). There were no age or sex differences between patients with and without cardiovascular autonomic neuropathy (Table 1). Patients with cardiovascular autonomic neuropathy showed significantly increased degree of stenosis at plaque sites in the common carotid artery ($24.6 \pm 13.2\%$ vs. $14.7 \pm 9.2\%$; $P = 0.014$, Table 1) and a non-significant tendency towards a higher plaque score (4.0 ± 1.7 vs. 3.2 ± 1.6 ; $P = 0.064$, Table 1). In age-controlled partial correlation, AI correlated inversely with degree of stenosis ($r = -0.39$; $P = 0.005$), plaque score ($r = -0.39$; $P = 0.005$), and mean ($r = -0.33$; $P = 0.018$) and maximum ($r = -0.39$; $P = 0.004$) intima-media thickness in the common carotid artery. In contrast, E/I ratio correlated only and slightly with mean intima-media thickness in the common carotid artery ($r = -0.28$; $P = 0.049$, Table 2 and Fig. 1).

Table 2 Age-controlled partial correlations between expiration/inspiration (E/I) ratio and acceleration index (AI) 5 years after diagnosis of diabetes vs. carotid ultrasound investigation 1 year later in 61 Type 2 diabetic patients

	E/I ratio	AI
<i>Common carotid artery</i>		
Plaque score	-0.11; 0.44	-0.39; 0.005
Degree of stenosis	-0.20; 0.14	-0.39; 0.005
Mean intima-media thickness	-0.28; 0.049	-0.33; 0.018
Maximum intima media thickness	-0.13; 0.35	-0.39; 0.004
Lumen diameter	-0.02; 0.87	-0.26; 0.07

R; P-values. Significant correlations shown in bold.

Discussion

We have previously shown in this group of patients with Type 2 diabetes [14] and another [13] that Type 2 diabetic patients with cardiovascular autonomic neuropathy, as defined by an abnormal E/I ratio (i.e. an established sign of parasympathetic neuropathy [22]) display signs of the metabolic syndrome. This suggested that cardiovascular autonomic neuropathy by mechanisms related to the metabolic syndrome may affect the vasculature in Type 2 diabetic patients, and is supported by the current study. Type 2 diabetic patients with cardiovascular autonomic neuropathy had increased degree of stenosis and plaque score in the common carotid artery compared with those without. Most interestingly AI, and not the E/I ratio, was associated with these changes as well as with carotid intima-media thickness. The weak negative relationship between AI and the lumen diameter of the common carotid artery ($r = -0.26$; $P = 0.07$) might seem puzzling at first glance. However, artery diameters increase in the presence of both atherosclerotic risk factors and atherosclerosis [30], presumably as a compensatory regulatory mechanism to preserve luminal area [30,31]. Consequently, our study shows that cardiovascular autonomic neuropathy is associated with atherosclerotic plaques in the carotid artery. However, we cannot draw any conclusions about causality from cross-sectional results in a limited cohort, especially as the time delay of 1 year between the autonomic function tests and the ultrasound examinations in the study might have affected our results.

Carotid artery atherosclerosis and increased intima-media thickness often occur together with atherosclerosis in coronary [32] and peripheral [33,34] arteries, and increased carotid intima-media thickness predicts myocardial infarction and stroke [35]. The present study of relationships between cardiovascular autonomic neuropathy and carotid intima-media thickness was performed 5–6 years after the clinical diagnosis of diabetes, a disease in which macrovascular complications are known to develop steadily over time [36]. Follow-up of this cohort will reveal whether carotid artery plaques associated with cardiovascular autonomic neuropathy and the metabolic syndrome [13–16] indicate increased risk of atherosclerotic events with increasing diabetes duration [18].

The correlation between carotid plaque variables and autonomic dysfunction was mainly confined to AI and not to the E/I ratio. AI reflects a fast inhibition of parasympathetic nerve tone when a subject is tilted [20], whereas a disturbed heart rate reaction during deep breathing, as expressed by an abnormal E/I ratio, is believed to reflect efferent parasympathetic nerve damage [37]. These assumptions are supported by the current study. The association between low AI and carotid plaques is a strong argument that afferent nerve activity secondary to postural changes in baroreceptor activity is the mechanism behind the immediate acceleration after tilt. Our study further supports the idea that the E/I ratio reflects efferent and most likely structural degeneration of the vagal nerve

[38]. Previously we suggested that AI also might reflect sympathetic nerve activity [23]. Withdrawal of parasympathetic nerve activity in combination with increased sympathetic nerve activity is probably involved in the postural heart rate reaction. We speculate that parasympathetic neuropathy as manifested by an abnormal E/I ratio through factors related to the metabolic syndrome induces an early atherosclerosis in the carotid artery and, by impairing baroreceptor function, leads to an afferent autonomic dysfunction as demonstrated by a low AI.

In conclusion, cardiovascular autonomic neuropathy in Type 2 diabetic patients was associated with atherosclerosis and narrowing of the carotid artery. This association could be due to factors related to the metabolic syndrome present, a feature of autonomic neuropathy in Type 2 diabetic patients.

Acknowledgements

We thank Mrs Helene Brandt, Ulrika Gustavsson, Ann Radelius, Christina Rosborn, Gerd Östling, and Birgitta Frid for skilful technical assistance. This study was supported by grants from the Swedish Diabetes Associations, the Ernhold Lundström Foundation, Research Funds of Malmö University Hospital, Swedish Heart-Lung Foundation, Research Funds at University Hospital MAS, the Albert Pålsson Foundation, Hulda Ahlmroth Foundation, NW Lundblad Foundation, and the Swedish Life Assurances Fund.

References

- Ewing DJ, Campbell IW, Clarke BF. The natural history of diabetic autonomic neuropathy. *Q J Med* 1980; 49: 95–108.
- Page MM, Watkins PJ. Cardiorespiratory arrest and diabetic autonomic neuropathy. *Lancet* 1978; 1: 14–16.
- Lilja B, Nosslin B, Bergström B, Sundkvist G. Glomerular filtration rate, autonomic nerve function, and orthostatic blood pressure in patients with diabetes mellitus. *Diabetes Res* 1985; 2: 179–181.
- Sundkvist G, Lilja B. Autonomic neuropathy predicts deterioration in glomerular filtration rate (GFR) in patients with IDDM. *Diabetes Care* 1993; 16: 773–779.
- Spallone V, Gambardella S, Maiello MR, Barini A, Frontoni S, Menzinger G. Relationship between autonomic neuropathy, 24-h blood pressure profile, and nephropathy in normotensive IDDM patients. *Diabetes Care* 1994; 17: 578–584.
- Krolewski AS, Barzilay J, Warram JH, Martin BC, Pfeifer M, Rand LI. Risk of early-onset proliferative retinopathy in IDDM is closely related to cardiovascular autonomic neuropathy. *Diabetes* 1992; 41: 430–437.
- Dosso A, Golay A, Morel Y, Furrara T, Assal JP, Leuenberger PM. Cardiovascular autonomic neuropathy in diabetic patients with macular oedema. *Diabet Metab* 1995; 21: 41–46.
- Niakan E, Harati Y, Rolak LA, Comstock JP, Rockey R. Silent myocardial infarction and diabetic cardiovascular autonomic neuropathy. *Arch Intern Med* 1986; 146: 2229–2230.
- Ambepityia G, Kopelman PG, Ingram D, Swash M, Mills PG, Timmis AD. Exertional myocardial ischemia in diabetes: a quantitative analysis of anginal perceptual threshold and the influence of autonomic function. *J Am Coll Cardiol* 1990; 15: 72–77.
- Gambardella S, Frontoni S, Spallone V, Rosaria Maiello M, Civetta E, Lanza G et al. Increased left ventricular mass in normotensive

- diabetic patients with autonomic neuropathy. *Am J Hypertens* 1993; **6**: 97–102.
- 11 Bottini P, Tantucci C, Scionti L, Dottorini ML, Puxeddu E, Reboldi G *et al*. Cardiovascular response to exercise in diabetic patients: influence of autonomic neuropathy of different severity. *Diabetologia* 1995; **38**: 244–250.
 - 12 Di Carli MF, Bianco-Battles D, Landa ME, Kazmers A, Groehn H, Muzik O *et al*. Effects of autonomic neuropathy on coronary blood flow in patients with diabetes mellitus. *Circulation* 1999; **100**: 813–819.
 - 13 Gottsäter A, Ahmed M, Fernlund P, Sundkvist G. Autonomic neuropathy in Type 2 diabetic patients associated with the metabolic syndrome. *Diabet Med* 1999; **16**: 49–54.
 - 14 Szelag B, Wroblewski M, Castenfors J, Henricsson M, Fernlund P, Berntorp K *et al*. Obesity, microalbuminuria, hyperinsulinaemia, and increased plasminogen activator inhibitor 1 activity associated with parasympathetic neuropathy in type 2 diabetes. *Diabetes Care* 1999; **22**: 1907–1908.
 - 15 Kerenyi Z, Stella P, Nadasdi A, Tabak AG, Tamas G. Associations between cardiovascular autonomic neuropathy and multimetabolic syndrome in a special cohort of women with prior gestational diabetes mellitus. *Diabet Med* 1999; **16**: 794–795.
 - 16 Takayama S, Sakura H, Katsumori K, Wasada T, Iwamoto Y. A possible involvement of parasympathetic neuropathy on insulin resistance in patients with type 2 diabetes. *Diabetes Care* 2001; **24**: 968–969.
 - 17 Edmonds ME, Morrison N, Laws JW, Watkins PJ. Medial arterial calcification and diabetic neuropathy. *Br Med J (Clin Res Ed)* 1982; **284**: 928–930.
 - 18 Toyry JP, Niskanen LK, Lansimies EA, Partanen KP, Uusitupa MI. Autonomic neuropathy predicts the development of stroke in patients with non-insulin-dependent diabetes mellitus. *Stroke* 1996; **27**: 1316–1318.
 - 19 Sundkvist G, Almér L-O, Lilja B. Respiratory influence on heart rate in diabetes mellitus. *Br Med J* 1979; **1**: 924–925.
 - 20 Sundkvist G, Lilja B, Almér L-O. Abnormal diastolic blood pressure and heart rate reactions to tilting in diabetes mellitus. *Diabetologia* 1980; **19**: 433–438.
 - 21 Wroblewski M, Gottsäter A, Lindgärde F, Fernlund P, Sundkvist G. Gender, autoantibodies, and obesity in newly diagnosed diabetic patients aged 40–75 years. *Diabetes Care* 1998; **21**: 250–255.
 - 22 Bergström B, Manhem P, Brammert M, Lilja B, Sundkvist G. Impaired responses of plasma catecholamines to exercise in diabetic patients with abnormal heart rate reactions to tilt. *Clin Phys* 1989; **9**: 259–267.
 - 23 Sundkvist G, Lilja B, Manhem P, Almér L-O. Responses of plasma catecholamines to tilt in patients with diabetes mellitus. *Acta Med Scand* 1984; **216**: 223–227.
 - 24 Bergström B, Lilja B, Rosberg K, Sundkvist G. Autonomic nerve function tests. Reference values in healthy subjects. *Clin Physiol* 1986; **6**: 523–528.
 - 25 Wendelhag I, Gustavsson T, Suurkula M, Berglund G, Wikstrand J. Ultrasound measurement of wall thickness in the carotid artery. Fundamental principles and description of a computerised image analysing system. *Clin Physiology* 1991; **11**: 565–577.
 - 26 Wendelhag I, Liang Q, Gustavsson T, Wikstrand J. A new automated computerised analysing system simplifies readings and reduces variability in ultrasound measurement of intima-media thickness. *Stroke* 1997; **28**: 2195–2200.
 - 27 Wong M, Edelstein J, Wollman J, Bond MG. Ultrasonic-pathological comparison of the human arterial wall. Verification of intima-media thickness. *Arterioscler Thromb* 1993; **13**: 482–486.
 - 28 Persson J, Formgren J, Israelsson B, Berglund G. Ultrasound determined intima-media thickness and atherosclerosis. A direct and indirect validation. *Arterioscler Thromb* 1994; **14**: 261–264.
 - 29 Persson J, Stavenow L, Wikstrand J, Israelsson B, Formgren J, Berglund G. Non-invasive quantification of atherosclerosis. Reproducibility of ultrasonographic measurement of arterial wall thickness and plaque size. *Arterioscler Thromb* 1992; **12**: 261–266.
 - 30 Labropoulos N, Zarge J, Mansour MA, Kang SS, Baker WH. Compensatory arterial enlargement is a common pathobiologic response in early atherosclerosis. *Am J Surg* 1998; **176**: 140–143.
 - 31 Jensen-Urstad K, Jensen-Urstad M, Johansson J. Carotid artery diameter correlates with risk factors for cardiovascular disease in a population of 55-year old subjects. *Stroke* 1999; **30**: 1572–1576.
 - 32 Mack WJ, LaBree L, Liu C, Selzer RH, Hodis HN. Correlations between measures of atherosclerosis change using carotid ultrasonography and coronary angiography. *Atherosclerosis* 2000; **150**: 371–379.
 - 33 Cheng SW, Wu LL, Ting AC, Lau H, Wong J. Screening for asymptomatic carotid stenosis in patients with peripheral vascular disease: a prospective study and risk factor analysis. *Cardiovasc Surg* 1999; **7**: 303–309.
 - 34 Simons PC, Algra A, Eikelboom BC, Grobbee DE, van der Graaf Y, SMART study group. Carotid artery stenosis in patients with peripheral arterial disease: the SMART study. *J Vasc Surg* 1999; **30**: 519–525.
 - 35 Fathi R, Marwick TH. Noninvasive tests of vascular function and structure: why and how to perform them. *Am Heart J* 2001; **141**: 694–703.
 - 36 Laakso M. Hyperglycemia and cardiovascular disease in type 2 diabetes. *Diabetes* 1999; **48**: 937–942.
 - 37 Lloyd-Mostyn RH, Watkins PJ. Total cardiac denervation in diabetic autonomic neuropathy. *Diabetes* 1976; **25**: 748–751.
 - 38 Duchon LW, Anjorin A, Watkins PJ, Mackay JD. Pathology of autonomic neuropathy in diabetes mellitus. *Ann Intern Med* 1980; **92**: 301–303.