



# LUND UNIVERSITY

## High prevalence of osteoarthritis 14 years after an anterior cruciate ligament tear in male soccer players: a study of radiographic and patient relevant outcomes.

von Porat, Anette; Roos, Ewa; Roos, Harald

*Published in:*  
Annals of the Rheumatic Diseases

*DOI:*  
[10.1136/ard.2003.008136](https://doi.org/10.1136/ard.2003.008136)

2004

[Link to publication](#)

*Citation for published version (APA):*  
von Porat, A., Roos, E., & Roos, H. (2004). High prevalence of osteoarthritis 14 years after an anterior cruciate ligament tear in male soccer players: a study of radiographic and patient relevant outcomes. *Annals of the Rheumatic Diseases*, 63(3), 269-273. <https://doi.org/10.1136/ard.2003.008136>

*Total number of authors:*  
3

### General rights

Unless other specific re-use rights are stated the following general rights apply:  
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00



## High prevalence of osteoarthritis 14 years after an anterior cruciate ligament tear in male soccer players: a study of radiographic and patient relevant outcomes

A von Porat, E M Roos and H Roos

*Ann. Rheum. Dis* 2004;63:269-273  
doi:10.1136/ard.2003.008136

---

Updated information and services can be found at:  
<http://ard.bmjournals.com/cgi/content/full/63/3/269>

---

*These include:*

### References

This article cites 37 articles, 13 of which can be accessed free at:  
<http://ard.bmjournals.com/cgi/content/full/63/3/269#BIBL>

2 online articles that cite this article can be accessed at:  
<http://ard.bmjournals.com/cgi/content/full/63/3/269#otherarticles>

### Rapid responses

You can respond to this article at:  
<http://ard.bmjournals.com/cgi/eletter-submit/63/3/269>

### Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

---

### Topic collections

Articles on similar topics can be found in the following collections

[Other Rheumatology](#) (1397 articles)  
[Orthopedic and Trauma Surgery](#) (262 articles)

---

### Notes

---

To order reprints of this article go to:  
<http://www.bmjournals.com/cgi/reprintform>

To subscribe to *Annals of the Rheumatic Diseases* go to:  
<http://www.bmjournals.com/subscriptions/>

## EXTENDED REPORT

# High prevalence of osteoarthritis 14 years after an anterior cruciate ligament tear in male soccer players: a study of radiographic and patient relevant outcomes

A von Porat, E M Roos, H Roos

*Ann Rheum Dis* 2004;**63**:269–273. doi: 10.1136/ard.2003.008136

See end of article for authors' affiliations

Correspondence to:  
Anette von Porat, Sport  
Medicine Centre, Södra  
Tvärgången 3, S-252 54  
Helsingborg, Sweden;  
[anette.vonporat@telia.com](mailto:anette.vonporat@telia.com)

Accepted 14 June 2003

**Objective:** To identify the consequences of an anterior cruciate ligament (ACL) tear in a cohort of male soccer players 14 years after the initial injury with respect to radiographic knee osteoarthritis and patient relevant outcomes.

**Methods:** Of 219 male soccer players with an ACL injury in 1986, 205 (94%) were available for follow up after 14 years; 75% of the cohort (154/205) answered mailed questionnaires (KOOS, SF-36, and Lysholm knee scoring scale) and 122 of these consented to weight bearing radiographs.

**Results:** Radiographic changes were found in 95 (78%) of the injured knees, while more advanced changes, comparable with Kellgren-Lawrence grade 2 or higher, were seen in 50 (41%). In the uninjured knees more advanced changes, comparable with Kellgren-Lawrence grade 2 or higher, were seen in five knees (4%). No differences were seen between surgically and conservatively treated players. The patient relevant outcome was affected and did not differ between subjects with and without radiographic changes. Eighty per cent reported reduced activity level.

**Conclusions:** A high prevalence of radiographic knee osteoarthritis was seen in male soccer players 14 years after an ACL disruption. The injury and the osteoarthritis, irrespective of the treatment provided to these patients, often result in knee related symptoms that severely affect the knee related quality of life by middle age.

Age and female sex are the most prominent risk factors for osteoarthritis over the age of 50 years.<sup>1,2</sup> Before 50, osteoarthritis is more common in men.<sup>3</sup> Risk factors for osteoarthritis at this young age include major knee injuries and joint loading.<sup>4,5</sup> It is estimated that 3.5% of all people between 35 and 54 years of age have radiological osteoarthritis after knee injuries.<sup>6–9</sup> Post-traumatic knee osteoarthritis is a good model for studies of the osteoarthritic process: the group is relatively homogeneous in an otherwise heterogeneous osteoarthritis population, the condition is common, and the start of the disease can be identified at the time of the trauma. Disease progression can be followed prospectively.

Anterior cruciate ligament (ACL) tears often influence knee stability.<sup>10–13</sup> Nearly 40% of all knee injuries in soccer are ACL injuries<sup>14</sup> and the affected players are often young. The overall injury rate in soccer has been estimated to be 8–30 injuries per 1000 game hours and 2–5 injuries per 1000 practice hours.<sup>15–17</sup>

A major knee injury may not only lead to an interrupted soccer career,<sup>4,12,18</sup> but also to the early development of knee osteoarthritis. Patients with osteoarthritis report handicap in six areas: functional and social activities, relationships, socioeconomic status, emotional wellbeing, and body image. Such patients also report more handicaps in each of the six areas than patients with rheumatoid arthritis.<sup>19</sup>

Our aim in this study was to assess the consequences of an ACL tear 14 years after the initial injury in a cohort of male soccer players with respect to radiographic knee osteoarthritis and patient relevant outcomes, such as activity level, symptoms, function, and quality of life.

## METHODS

### Subjects

The subjects in this nationwide study of ACL injuries were identified in 1989. In Sweden all soccer players participating

in league soccer have compulsory insurance through the same company (Folksam). A search was made in the Folksam archives in 1989 for all knee injuries that had occurred in 1986. In all, 937 knee injuries were found. The data from questionnaires and from the hospital records showed a total of 344 ACL injuries, of which 238 were in men and 106 in women. In 2000 the male players were contacted for a 14 year follow up, and 205 were available. The questionnaires were answered by 154, and of those 65 (42%) had been treated without surgery, and 89 (58%) had undergone ACL reconstruction. The most common surgical method was patellar tendon graft. The reasons behind the choice of treatment were not recorded.

In 1989, 59 (38%) of the patients were identified as having a meniscus tear combined with their ACL injury. The rupture was medial in 35 cases, lateral in 17 cases, and both in seven cases.

The mean age of the study group at 14 years after the index injury was 38 years (range 30 to 56) (table 1). The subgroup with a combined injury (mean age 38, range 31 to 54) did not differ in age from the study group as a whole ( $p = 0.4$ ), nor did the non-participants differ in age from those participating (mean age 37 years (range 29 to 48),  $p = 0.7$ ).

### Radiographic examination

Of the 154 subjects, 122 (79%) consented to have radiographs taken. The main reason for not having radiographs taken was lack of time. Posterior-anterior radiographs with the knees at 15° of flexion were taken in the weight bearing position, the weight equally distributed on both legs, for examination of

**Abbreviations:** ACL, anterior cruciate ligament; KOOS, knee injury and osteoarthritis outcome score; SF-36, medical outcomes study 36 item short form health survey

**Table 1** Characteristics of the study group and the subgroups operated, non-operated, osteoarthritis, and non-osteoarthritis

	Study group (n = 154)	Operated (n = 89)	Not operated (n = 65)	OA (n = 50)	Non-OA (n = 104)
Age (years)	38 (5.2), (30 to 56)	37 (4.2), (30 to 49)	40 (6.2), (30 to 56)	39 (4.9), (32 to 53)	38 (5.4), (30 to 56)
Weight (kg)	83 (8.5), (58 to 110)	83 (8.1), (67 to 105)	83 (9.0), (71 to 110)	85 (8.4), (68 to 108)	83 (8.4), (58 to 110)
Height (m)	181 (6.4), (165 to 205)	181 (6.8), (165 to 205)	181 (5.6), (170 to 193)	182 (6.2), (170 to 205)	180 (6.3), (165 to 199)
BMI (kg/m <sup>2</sup> )	26 (2.3), (21 to 34)	26 (2.3), (21 to 32)	26 (2.3), (21 to 34)	26 (2.3), (21 to 32)	26 (2.3), (21 to 34)

Values are mean (SD), (range).

BMI, body mass index; OA, osteoarthritis.

the tibiofemoral joint. All radiographs were taken with a Siemens basic radiological system (film focus distance 1.4 m; Siemens GmbH, Erlangen, Germany) at 70 kV and 10 mA. All were obtained with the same standardised technique.

The frontal views of the tibiofemoral joint from both knees of the subjects were classified according to the recommendations of the Osteoarthritis Research Society.<sup>20</sup> A radiographic atlas was used to evaluate the appearance of the joint space and the presence of osteophytes and to grade these features on a scale from 0 to 3.<sup>20</sup> Radiographic osteoarthritis was defined as joint space narrowing grade 1 combined with osteophytes, or joint space narrowing grade 2 or more.<sup>13</sup> This definition of osteoarthritis corresponds to Kellgren and Lawrence knee osteoarthritis grade 2.<sup>21</sup> The same reader (HR) evaluated all the radiographs. The reader was blinded with regard to the injured side in non-operated subjects. In ACL reconstructed subjects the bone tunnels were visible on x ray films. The reader in the present study had been shown in an earlier report to have high agreement with another reader. In the previous study there was agreement for medial joint space narrowing in 119 of 136 knees; for lateral narrowing in 133 of 136 knees; and for osteophytes in 243 of 272 compartments.<sup>13</sup>

### Questionnaires

Three questionnaires were used to evaluate patient relevant outcomes: the disease specific knee injury and osteoarthritis outcome score (KOOS), the generic short form 36 (SF-36), and the Lysholm knee scoring scale. In addition, data including duration of knee problems and current physical activity level at work and recreation were collected. The subjects were also asked to report current activity level (work and recreation) compared with before the knee injury on a five point Likert scale, ranging from much lower to much higher. The players noted the reasons as "knee problem," "other reasons," or "both" if a change in activity level was experienced.

### Knee injury and osteoarthritis outcome score (KOOS)

The KOOS is a 42 item self administered questionnaire based on the WOMAC osteoarthritis index,<sup>22</sup> proven valid for subjects with ACL injury and early osteoarthritis.<sup>23 24</sup> The KOOS assesses five separate dimensions: pain, symptoms, activities of daily living, sport and recreation function, and knee related quality of life. A score from 0–100 is calculated for each dimension, with 100 representing best possible score.

### SF-36

The medical outcomes study 36 item short form health survey (SF-36) is a widely used measure of general health which comprises eight subscales: physical functioning, role–physical, bodily pain, general health, vitality, social functioning, role–emotional, and mental health.<sup>25</sup> The SF-36 has previously been used in studies of subjects with ACL injury.<sup>26</sup> A score from 0 to 100 is calculated for each dimension, 100 representing the best result. The Swedish acute version 1.0 was used.<sup>27</sup>

### Lysholm knee scoring scale

The Lysholm knee scoring scale is an eight item questionnaire developed to assess symptoms and functional disabilities resulting from an ACL injury.<sup>28</sup> All eight items (pain, instability, locking, stairs, swelling, squat, limp, and support) are aggregated into one score from 0 to 100, where 100 indicates normal knee function. The Lysholm scale is intended to be observer administered and no patient instructions are provided in the original version. In this study the Lysholm knee scoring scale was patient administered and the subjects were instructed to consider the last week when filling out the questionnaire.

### Statistics

Non-parametric statistics were used. The Kruskal–Wallis test was employed to determine differences for each subscale of the KOOS between groups without or with radiographic osteoarthritis grade 1 to grade 3. The Mann–Whitney U test was used when comparing two groups. A probability (p) value of  $\leq 0.05$  was regarded as significant.

## RESULTS

### Radiography

Radiographic changes were found in 95 (78%) of the injured knees, and of these, radiographic osteoarthritis equivalent to Kellgren and Lawrence grade 2 was seen in 50 (41%). Twenty seven subjects (22%) had no radiographic changes at all. No difference in radiographic outcome was determined between those treated with or without surgery.

The subjects who filled out the questionnaires but declined to undergo a radiographic examination did not differ in age, weight, height, or functional status from those who had radiographs taken. Age and BMI did not differ between subjects with different radiographic grading, nor between subjects in the operated or non-operated group (table 1).

Subjects with a meniscus tear had significantly more radiographic changes equivalent to Kellgren and Lawrence grade 2 or worse (59% v 31%,  $p = 0.002$ ), but the severity of radiographic osteoarthritis between isolated ACL rupture and associated meniscus tear was similar.

### Questionnaires

Only limited problems were noted for the KOOS activity of daily living dimension (table 2). Eighty eight subjects (57%) had an excellent score for this dimension (defined as 95–100), while only 28 subjects (18%) scored excellent in sport/recreation, and 18 (12%) in knee specific quality of life. The most influenced items in sport/recreation were "What difficulty have you experienced in the last week when turning/twisting on your injured knee?" and "What difficulty have you experienced in the last week when kneeling?" One of four subjects reported severe problems with turning/twisting at least, and one of three reported severe problems with kneeling at least. Of those reporting problems with kneeling, 59% were from the operated group and 41% from the non-operated group. The most affected item in the subscale "knee specific quality of life" was "Have you modified your lifestyle to avoid potentially damaging

**Table 2** Values for the KOOS questionnaire, SF-36, and Lysholm score for the study group, with comparison values for different previously published reference groups

	Study group			Reference groups		
	Mean (SD)	Range	95% CI	Mean (SD)	Range	95% CI
KOOS (n = 154 + n = 50)						
Pain*	84 (16.1)	(28 to 100)	81.9 to 87.1	96 (9.9)	(50 to 100)	93.7 to 99.0
Symptoms*	76 (20.2)	(14 to 100)	72.5 to 79.0	94 (9.9)	(57 to 100)	92.3 to 96.9
ADL*	90 (13.5)	(37 to 100)	88.3 to 92.6	96 (10.0)	(47 to 100)	93.2 to 98.5
Sport/rec*	63 (28.7)	(0 to 100)	58.9 to 68.0	90 (17.9)	(25 to 100)	85.6 to 95.1
QOL*	60 (24.6)	(6 to 100)	56.2 to 63.9	92 (13.5)	(38 to 100)	88.6 to 95.7
SF-36 (n = 152 + n = 948)						
Physical functioning*	84.5 (14.5)	(35 to 100)	82.1 to 86.8	93.1 (15.4)	(0 to 100)	92.1 to 94.1
Role-physical*	81.4 (30.9)	(0 to 100)	76.5 to 86.4	88.5 (26.7)	(0 to 100)	86.7 to 90.2
Bodily pain	74.3 (22.9)	(22 to 100)	70.6 to 78.0	78.2 (25.3)	(0 to 100)	76.6 to 79.8
General health	82.1 (17.0)	(15 to 100)	79.4 to 84.8	78.3 (20.2)	(0 to 100)	77.0 to 79.6
Vitality	70.8 (20.8)	(15 to 100)	67.5 to 74.1	71.6 (21.2)	(0 to 100)	70.2 to 72.9
Social functioning*	93.6 (13.9)	(25 to 100)	91.3 to 95.8	89.5 (20.0)	(0 to 100)	88.2 to 90.8
Role-emotional	91.0 (22.4)	(0 to 100)	87.4 to 94.6	89.4 (24.3)	(0 to 100)	87.8 to 91.0
Mental health*	86.4 (12.9)	(40 to 100)	84.4 to 88.5	82.2 (18.6)	(0 to 100)	81.0 to 83.4
Lysholm score* (n = 154 + n = 131)						
	77.0 (19.1)	(9 to 100)	73.5 to 79.6	99 (2.7)	(77 to 100)	98.8 to 99.4

For the KOOS questionnaire, reference data are from 59 subjects aged 36 to 79 years obtained in a previous study.<sup>39</sup> For the SF-36 questionnaire, reference values are for the general Swedish male population aged 35 to 44 years.<sup>27</sup> For the Lysholm score, reference values are for healthy team athletes aged 13 to 25 years.<sup>40</sup>

\*Significant difference at  $p < 0.05$ .

ADL, activities of daily living; CI, confidence interval; KOOS, knee injury and osteoarthritis outcome score; QOL, knee related quality of life; SF-36, medical outcomes study 36 item short form health survey; Sport/rec, sport and recreation function.

activities to your knee?" Thirty per cent reported severe changes in lifestyle at least because of their knee injury.

The injured players reported significantly worse outcome in the SF-36 subscales "physical functioning" and "role-physical" compared with Swedish men aged 35 to 44. However, in the subscales "social functioning" and "mental health," the players scored significantly higher than the reference group<sup>27</sup> (table 2).

A mean score of 77 (range 9 to 100) was obtained in the Lysholm knee scoring scale. Sixty three subjects (41%) showed excellent or good results (100–84 points). Subjects with radiographic osteoarthritis did not have significantly different scores from subjects without osteoarthritis (table 3).

There was no difference in outcome of the questionnaire between those treated with ACL surgery and those who were

not (table 3). A surgically treated meniscus injury in addition to the ACL injury did not influence the outcome as measured by the KOOS, SF-36, or Lysholm questionnaires.

### Activity level

Eighty two subjects (53%) reported level 2–4, easy to moderate load at work. Eighty four subjects (55%) reported level 5–6, a high level of recreational activities. One hundred and twenty three subjects (80%) reported reduced activity level after the knee injury, and of these the majority (69%) reported the knee injury as the cause. Subjects with no radiographic changes reported the same workload and recreational activities as subjects with radiographic changes. Twelve (7.8%) of the 154 participants were still participating in organised soccer. Of these, nine had been surgically treated and three conservatively treated.

**Table 3** Values for the five measured outcomes in the KOOS questionnaire (n = 154), the eight measured outcomes in the SF-36 questionnaire (n = 152), and the outcomes in Lysholm score (n = 154) for subjects with and without osteoarthritis and for operated and non-operated subjects

	Study group	OA (n = 50)	No OA (n = 72)	Operated (n = 89)	Non-operated (n = 65)
KOOS (n = 154)					
Pain	84 (81.9 to 87.1)	85 (80.2 to 89.9)	84 (81.2 to 87.3)	86 (82.2 to 89.2)	83 (79.1 to 86.7)
Symptoms	76 (72.5 to 79.0)	73 (66.9 to 78.7)	77 (73.3 to 81.1)	75 (70.4 to 79.6)	77 (72.3 to 81.2)
ADL	90 (88.3 to 92.6)	91 (87.1 to 95.3)	90 (87.5 to 92.7)	91 (88.5 to 94.0)	89 (85.9 to 92.9)
Sport/rec	63 (58.9 to 68.0)	64 (55.5 to 72.0)	63 (57.7 to 68.8)	64 (57.7 to 70.1)	63 (55.9 to 69.6)
QOL	60 (56.2 to 63.9)	61 (55.1 to 67.2)	60 (54.5 to 64.6)	62 (56.6 to 67.1)	58 (51.6 to 63.6)
SF-36 (n = 152)					
Physical function	84 (82.1 to 86.8)	84 (80.0 to 88.4)	85 (81.8 to 87.5)	84 (80.4 to 86.9)	86 (82.3 to 89.1)
Role-physical	81 (76.5 to 86.4)	84 (75.1 to 91.4)	80 (74.2 to 86.6)	83 (76.4 to 88.9)	80 (71.4 to 87.9)
Bodily pain	74 (70.6 to 78.0)	76 (69.0 to 82.6)	74 (69.1 to 78.0)	75 (70.2 to 80.1)	73 (67.6 to 78.8)
General health	82 (79.4 to 84.8)	86 (82.2 to 89.8)	80 (76.6 to 83.8)	82 (78.7 to 85.5)	82 (78.7 to 86.7)
Vitality	71 (67.5 to 74.1)	71 (64.8 to 77.2)	71 (66.7 to 74.7)	71 (67.0 to 76.1)	70 (64.8 to 74.8)
Social functioning	94 (91.3 to 95.8)	96 (92.9 to 99.1)	92 (89.4 to 95.4)	93 (90.2 to 96.2)	94 (90.7 to 97.6)
Role-emotional	91 (87.4 to 94.6)	94 (88.7 to 99.3)	90 (84.4 to 94.2)	92 (87.8 to 97.1)	89 (83.3 to 94.8)
Mental health	86 (84.4 to 88.5)	88 (84.5 to 91.4)	86 (83.1 to 88.3)	87 (84.1 to 89.4)	86 (82.6 to 89.4)
Lysholm scale (n = 154)					
	77 (73.5 to 79.6)	76 (71.1 to 81.7)	77 (73.1 to 80.7)	78 (73.4 to 81.7)	75 (70.8 to 79.8)

Values are mean values (95% confidence interval).

ADL, activities of daily living; KOOS, knee injury and osteoarthritis outcome score; QOL, knee related quality of life; SF-36, medical outcomes study 36 item short form health survey; Sport/rec, sport and recreation function.

## DISCUSSION

The purpose of this report was to identify the consequences of an ACL tear in a cohort of male soccer players 14 years after the initial injury with respect to radiographic knee osteoarthritis, activity level, and patient related symptoms. The cohort was from the Swedish population of soccer players. They have all been followed for 14 years after the initial injury, with specific follow ups after three and seven years. The results reflect the severe consequences of an ACL injury in middle aged male former soccer players. Nearly 80% of the cohort had significant radiographic changes in the knee, and more than 40% fulfilled the criteria for definite radiographic osteoarthritis. The majority suffered from knee symptoms affecting their knee related quality of life.

Most studies describe only radiographic knee osteoarthritis, but in the present study the patient relevant outcome was taken into consideration. We also analysed the patient relevant outcome and activity level in relation to the radiographic knee osteoarthritis.

This is a nationwide study of ACL injuries in Swedish male soccer players. All players were injured in 1986, and the cohort was identified in 1989. They were previously followed up in 1993.<sup>12</sup>

A limitation of the study was the response rate. Of 205 male soccer players located, 75% responded to the questionnaires, but of these, 36 did not accept the invitation to radiographic examination. Of all 238 male soccer players with an ACL injury in 1986, 122 (51%) consented to have a radiograph taken in 2000. This may constitute a selection bias, with an increased interest in participating in the study if symptoms from the knee joint were present. However, the information from the questionnaires did not support the theory that the group who did not have radiographs taken had less or more knee symptoms than the group who had radiographs taken.

The treatment of the subjects with or without surgery was not randomised, limiting the interpretation of differences, or lack of differences, between these groups.

There are difficulties in comparing studies with regard to osteoarthritis prevalence, mainly because of factors such as different radiographic techniques and classifications. There can also be differences in patient material and in the time of follow up. This study is unique, as all subjects have been followed during the same span of time. The results are in agreement with other studies on osteoarthritis after an ACL injury.<sup>14 29 30</sup>

Male soccer players have an increased risk of knee osteoarthritis.<sup>29 31-33</sup> Injuries to the menisci and cruciate ligaments are believed to be the main reason for this, but also when injuries are excluded, a relation between elite soccer and osteoarthritis persists.<sup>32</sup> This means that a player who returns to soccer after an ACL injury has an even higher risk of osteoarthritis, both because of the soccer itself and because of the injury. Kohatsu and Schurman reported in a case-control study that osteoarthritis cases were almost five times more likely to have had a significant knee injury than the controls.<sup>1</sup> The natural course of an ACL injury in relation to the risk of developing osteoarthritis is not known. Further research is required to understand the different mechanisms whereby some individuals develop osteoarthritis and others do not. Sensitive instruments are needed to monitor the process, such as magnetic resonance imaging with specific sequences or biomarkers. Our study has not covered these issues.

Our study population reported similar symptoms to patients on the waiting list for ACL reconstruction—for example, pain and functional limitations.<sup>24</sup> However, our population with chronic ACL injury reported a better knee related quality of life than the group with an acute tear,

possibly because of reduced activity level and an acceptance of their reduced knee function.<sup>23 24</sup>

This study cohort was followed up in 1993, seven years after their index injury.<sup>12</sup> At this time, 49 subjects (32%) were still participating in organised soccer, compared with 12 (7.8%) in the present 14 year follow up. One reason for this is that as the subjects get older they have families and have less time for soccer.

At the 14 year follow up the subjects reported better knee function than they did at the seven year follow up: a mean score of 77 (range 9 to 100) was obtained in the Lysholm knee scoring scale in the present follow up, compared with 66 (4 to 100) in the seven year follow up. Sixty three subjects in the present study (41%) reported excellent or good results (100 to 84 points) compared with 49 (32%) at the seven year follow up. Perhaps the reason for this is that most subjects had stopped playing soccer and that they had learned to live with their knee problems.

There was no detectable difference in the prevalence of osteoarthritis or symptoms between subjects who had undergone knee surgery and those who had not. This finding was expected as there are no published data supporting a preventive effect of reconstructive ACL surgery against osteoarthritis. On the contrary, a study by Daniel *et al* in 1994 showed a trend towards a greater prevalence of osteoarthritis in surgically treated than in non-surgically treated patients.<sup>34</sup> An increased ability to return to soccer following surgical reconstruction may be disadvantageous with regard to the risk for developing knee osteoarthritis.<sup>32</sup> Subjects with an ACL rupture *and* a meniscus tear had a higher prevalence of radiographic osteoarthritis than those with an isolated ACL rupture. This was as expected and in accordance with other studies.<sup>35 36</sup>

The need for osteotomy and knee arthroplasty will arise well before the age of 50 in many of the subjects in this cohort. Considering that soccer is the most commonly played athletic game in the world, the risk of a latent and growing group of osteoarthritis patients is high. For this group of young people of working age, the indirect costs at an individual level, and also for society, will be considerable. Thus the prevention of knee injuries appears to be very important to prevent soccer players from getting early osteoarthritis and knee related symptoms. The results of the present study, and of others<sup>14 30 32 37 38</sup> that have shown significant radiographic osteoarthritis in young former soccer players, must be considered alarming.

## ACKNOWLEDGEMENTS

This study was supported by the Zoega Foundation, Swedish National Centre for Research in Sport, the Gorton Foundation, and the Medical Faculty of the University of Lund.

## Authors' affiliations

**A von Porat**, Department of Physical Therapy, Lund University Hospital, Lund, Sweden

**E M Roos**, Department of Orthopaedics, Lund University Hospital

**H Roos**, Hospital of Helsingborg, Helsingborg, Sweden

## REFERENCES

- 1 Kohatsu ND, Schurman DJ. Risk factors for the development of osteoarthritis of the knee. *Clin Orthop Rel Res* 1990;**261**:242-6.
- 2 Felson DT. The course of osteoarthritis and factors that affect it. *Osteoarthritis Cartilage* 1993;**19**:607-15.
- 3 Felson DT, Lawrence RC, Dieppe PA, Hirsch R, Helmick CG. Osteoarthritis: the disease and its prevalence and impact. *Ann Intern Med* 2000;**133**:635-46.
- 4 Sandmark H, Vingård E. Sports and risk for severe osteoarthritis of the knee. *Scand J Med Sci Sports* 1999;**9**:279-84.
- 5 Cooper C, Snow S, McAlindon TE, Kellingrey S, Stuart B, Coggon D, *et al*. Risk factors for the incidence and progression of radiographic knee osteoarthritis. *Arthritis Rheum* 2000;**43**:995-1000.

- 6 **Williams M**, Frankel S, Nanchahal K, Coast J, Donovan J. Total knee replacement. In: Stevens A, Raftery J, eds. *Health needs assessment*, vol 1. Oxford: Radcliffe, 1994:448–523.
- 7 **Tzonchev VT**, Pilosoff T, Kanev. *Prevalence of osteoarthritis in Bulgaria in population studies of the rheumatic diseases*. Amsterdam: Excerpta Medica Foundation, 1968.
- 8 **Hernborg JS**, Nilsson BE. Age and sex incidence of osteophytes in the knee joint. *Acta Orthop Belg* 1973;**44**:66–8.
- 9 **HANES**. Basic data on osteoarthritis of the knee, hip and sacroiliac joints in adults aged 25–74 years. USA 1971–75. *Natl Center Health Stat* 1979:series 11(213).
- 10 **Graham GP**, Fairclough JA. Early osteoarthritis in young sportsmen with severe anterolateral instability of the knee. *Injury* 1988;**19**:247–8.
- 11 **Kannus P**, Järvinen M. Age, overweight, sex and knee instability: their relationship to the post-traumatic osteoarthritis of the knee joint. *Injury* 1988;**19**:105–8.
- 12 **Roos H**, Ornell M, Gärdsell P, Lohmander S, Lindstrand A. Soccer after anterior cruciate ligament - an incompatible combination. *Acta Orthop Scand* 1995;**66**:107–12.
- 13 **Roos H**, Laurén M, Adalberth T, Roos EM, Jonsson K, Lohmander SL. Knee osteoarthritis after meniscectomy. Prevalence of radiographic changes after twenty-one years, compared with matched controls. *Arthritis Rheumatism* 1998;**41**:687–93.
- 14 **Roos H**, Adalberth T, Dahlberg L, Lohmander SL. Osteoarthritis of the knee after injury to the anterior cruciate ligament or meniscus: the influence of time and age. *Osteoarthritis Cartilage* 1995;**3**:261–7.
- 15 **Ekstrand J**, Gillquist J, Möller M, Öberg B, Liljedahl S-O. Incidence of soccer injuries and their relation to training and team success. *Am J Sports Med* 1983;**11**:63–7.
- 16 **Arendt E**, Randall D. Knee injury patterns among men and women in collegiate basket and soccer: NCAA data and review of literature. *Am J Sports Med* 1995;**23**:694–701.
- 17 **Árnason A**, Gudmundsson Á, Dahl HA, Jóhannsson E. Soccer injuries in Iceland. *Scand J Med Sci Sports* 1996;**6**:40–5.
- 18 **Engström B**, Johansson C, Törnkvist H. Soccer injuries among elite female players. *Am J Sports Med* 1991;**19**:372–5.
- 19 **Carr AJ**. Beyond disability: measuring the social and personal consequences of osteoarthritis. *Osteoarthritis Cartilage* 1999;**7**:230–8.
- 20 **Altman RD**, Hochberg M, Murphy WA, Wolfe F, Lequesne M. Atlas of individual radiographic features in osteoarthritis. *Osteoarthritis Cartilage* 1995;**3**(suppl A):3–70.
- 21 **Kellgren J**, Lawrence J. Radiological assessment of osteoarthrosis. *Ann Rheum Dis* 1957;**16**:494–502.
- 22 **Bellamy N**, Watson Buchanan W, Goldsmith C, Campbell J, Stitt L. Validation study of WOMAC: a health status instrument for measuring clinically important patient relevant outcomes to antirheumatic drug therapy in patient with osteoarthritis of the hip or the knee. *J Rheumatol* 1988;**15**:1833–40.
- 23 **Roos EM**, Roos H, Ekdahl C, Lohmander SL. Knee injury and osteoarthritis outcome score (KOOS) – validation of a Swedish version. *Scand J Med Sci Sports* 1998;**8**:439–48.
- 24 **Roos EM**, Roos H, Lohmander SL, Ekdahl C, Beynon BD. Knee injury and osteoarthritis outcome score (KOOS) – development of a self-administered outcome measure. *J Orthop Sports Phys Ther* 1998;**78**:88–96.
- 25 **Ware JE**, Sherbourne CD. The MOS 36-item short form health survey (SF-36). I. Conceptual framework and item selection. *Med Care* 1992;**30**:472–83.
- 26 **Shapiro ET**, Richmond JC, Rockett SE, McGrath MM, Donaldson WR. The use of a generic patient-based health assessment (SF-36) for evaluation of patients with anterior cruciate ligament injuries. *Am J Sports Med* 1996;**24**:No 2, 196–200.
- 27 **Sullivan M**, Karlsson J, Ware JE. The Swedish SF-36 health survey-I. evaluation of data quality, scaling assumptions, reliability and construct validity across general populations in Sweden. *Soc Sci Med* 1995;**41**:1349–58.
- 28 **Tegner Y**, Lysholm J. Rating systems in the evaluation of knee ligament injuries. *Clin Orthop Rel Res* 1985;**198**:43–9.
- 29 **Neyret P**, Donell ST, DeJour D, DeJour H. Partial meniscectomy and anterior cruciate ligament rupture in soccer players. A study with a minimum 20-year followup. *Am J Sports Med* 1993;**21**:455–60.
- 30 **Segawa H**, Omori G, Koga Y. Long-term results of non-operative treatment of anterior cruciate ligament injury. *Knee* 2000;**8**:5–11.
- 31 **Ferretti A**, Conteduca F, De Carli A, Fontana M, Mariani PP. Osteoarthritis of the knee after ACL reconstruction. *Int Orthop* 1991;**15**:367–71.
- 32 **Roos H**, Lindberg H, Gärdsell P, Lohmander SL, Wingstrand H. The prevalence of gonarthrosis and its relation to meniscectomy in former soccer players. *Am J Sports Med* 1994;**22**:219–22.
- 33 **Kujala UM**, Kaprio J, Sarna S. Osteoarthritis of weight bearing joints of lower limbs in former elite male athletes. *BMJ* 1994;**308**:231–4.
- 34 **Daniel DM**, Stone ML, Dobson BE, Fithian DC, Rossman DJ, Kaufman KR. Fate of the ACL injured patient: a prospective outcome study. *Am J Sports Med* 1994;**22**:632–44.
- 35 **Shelbourne KD**, Gray T. Results of anterior cruciate ligament reconstruction based on meniscus and articular cartilage status at the time of surgery. Five- to fifteen-year evaluations. *Am J Sports Med* 2000;**28**:446–52.
- 36 **Wu WH**, Hackett T, Richmond JC. Effects of meniscal and articular surface status on knee stability, function, and symptoms after anterior cruciate ligament reconstruction: a long-term prospective study. *Am J Sports Med* 2002;**30**:845–50.
- 37 **Sherman MF**, Warren RF, Marshall JL, Savatsky GJ. A clinical and radiographical analysis of 127 anterior cruciate insufficient knees. *Clin Orthop Rel Res* 1988;**227**:229–37.
- 38 **Gelber AC**, Hochberg MC, Mead LA, Wang N-Y, Wigley FM, Klag MJ. Joint injury in young adults and risk for subsequent knee and hip osteoarthritis. *Ann Intern Med* 2000;**133**:321–8.
- 39 **Englund M**, Roos EM, Lohmander LS. Impact of type of meniscal tear on radiographic and symptomatic knee osteoarthritis: a sixteen-year followup of meniscectomy with matched controls. *Arthritis Rheum* 2003;**48**:2178–87.
- 40 **Demirdjian AM**, Petrie SG, Guanache CA, Thomas KA. The outcomes of two knee scoring questionnaires in a normal population. *Am J Sports Med* 1998;**26**:46–51.