



LUND UNIVERSITY

Are Sulfonylureas Less Desirable Than DPP-4 Inhibitors as Add-on to Metformin in the Treatment of Type 2 Diabetes?

Ahrén, Bo

Published in:
Current Diabetes Reports

DOI:
[10.1007/s11892-011-0179-x](https://doi.org/10.1007/s11892-011-0179-x)

2011

[Link to publication](#)

Citation for published version (APA):

Ahrén, B. (2011). Are Sulfonylureas Less Desirable Than DPP-4 Inhibitors as Add-on to Metformin in the Treatment of Type 2 Diabetes? *Current Diabetes Reports*, 11, 83-90. <https://doi.org/10.1007/s11892-011-0179-x>

Total number of authors:
1

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00



LUND UNIVERSITY
Faculty of Medicine

LUP

Lund University Publications

Institutional Repository of Lund University

This is an author produced version of a paper published in Current Diabetes Reports. This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Citation for the published paper:
Bo Åhrén

"Are Sulfonylureas Less Desirable Than DPP-4 Inhibitors as Add-on to Metformin in the Treatment of Type 2 Diabetes?"

Current Diabetes Reports
2011 Jan 18

<http://dx.doi.org/10.1007/s11892-011-0179-x>

The original publication is available at
www.springerlink.com

Access to the published version may require journal subscription.
Published with permission from: Springer

Are Sulfonylureas Less Desirable Than DPP-4 Inhibitors as Add-on to Metformin in the Treatment of Type 2 Diabetes?

Bo Ahrén

Department of Clinical Sciences, Lund University, Lund, Sweden

Key words: Diabetes, hyperglycemia, sulfonylureas, glimepiride, glipizide, DPP-4 inhibitors, vildagliptin, sitagliptin, saxagliptin, islet function, insulin secretion, glucagon secretion, hypoglycemia, safety, durability, comparison

Address

Dr Bo Ahrén

Department of Clinical Sciences Lund

Lund University

B11 BMC

SE-221 84 LUND, Sweden

e-mail Bo.Ahren@med.lu.se

Abstract

Sulfonylureas (SUs) are commonly used as add-on to metformin in treatment of type 2 diabetes in patients who are insufficiently controlled by metformin alone. They have good efficacy and have shown to prevent microvascular complications. However, treatment with SUs is also associated with a high frequency of hypoglycemia, increased body weight, and a high risk of secondary failure. During recent years, dipeptidyl peptidase-4 (DPP-4) inhibitors have emerged as alternatives to SUs. They show similar efficacy as SUs but with lower risk of hypoglycemia, and reduction or no change in body weight, and if confirmed in humans, they may preserve islet function and thereby minimizing the risk for secondary failure. Their limitation at present is the lack of long-term (>5 years) experience on durability and safety. Overall, therefore, the conclusion emerges that SUs are less desirable than DPP-4 inhibitors in management of hyperglycemia in type 2 diabetes.

Introduction

Targeting the hyperglycemia is of key importance in management of type 2 diabetes, because lowering glycemia reduces both acute symptoms and the increased risk for retinopathy, nephropathy and neuropathy (1). Treating hyperglycemia may also improve the increased cardiovascular (CV) risk in type 2 diabetes, although this needs a multifactorial approach (2).

The key defect underlying hyperglycemia in type 2 diabetes is islet dysfunction, which has three components: 1) Impaired insulin secretion, i.e., insufficiently increased insulin secretion to match insulin resistance (3), 2) Defective suppression of glucagon secretion which results in hyperglucagonemia and increased hepatic glucose production (4), and 3) Reduced islet cell mass (5). These defects are early phenomena and are seen already several years before the diabetes diagnosis (6).

Treatment guidelines from national and international bodies have presented algorithms for sequential introduction of pharmacological agents to treat the hyperglycemia (7-10). The first-line pharmacological agent is, for most patients, metformin, which mainly improves insulin sensitivity. In many patients, however, this is insufficient to reach target for glycemic control and a second agent is then suggested to be added. Sulfonylureas (SUs) have for a long time been recommended as this second treatment. During recent years dipeptidyl peptidase-4 (DPP-4)-inhibitors have emerged as a new class of therapeutic agents (11). These agents seem to be equally effective in improving glycemia as SUs but lack some of the negative effects of SUs. It may therefore be discussed

whether DPP-4 inhibitors are more desirable than SUs as add-on to metformin in patients with inadequate glycemic control.

Sulfonylureas (SUs)

History. Sulfonylureas were developed as agents stimulating insulin secretion based on incidental finding of hypoglycemia during treatment with sulfonamides in the 1940s (12). Their introduction in the treatment during the 1950s represented the first reliable oral treatment of diabetes. Since then, several SUs have been of key and central value in the treatment. Following the first (chlorpropamide, acetohexamide and tolazamide) and second generations (tolbutamide), the third generation SUs (glibenclamide (=glyburide), gliclazide, glipizide and glimipiride) are the most commonly used SUs today (17,18).

Mechanisms. SUs stimulate insulin secretion by activating beta-cell sulfonylurea receptors 1 (SUR1), closing ATP-dependent potassium channels (Kir6.2 channels) thereby inhibiting potassium flow across the plasma membrane (14). This results in depolarization, which opens voltage sensitive calcium channels, allowing uptake of extracellular calcium, increased cytosolic calcium and exocytosis of insulin containing granules. Importantly, this effect is glucose-independent, i.e., SUs stimulate insulin secretion both at low and high glucose concentrations.

SUR1 receptors are expressed also in the glucagon producing islet alpha-cells cells and the net effect of Kir6.2 channel closure in these cells is a stimulation of glucagon secretion (15). A stimulation of glucagon secretion by SU has also been observed in

humans with insulin deficiency in type 1 diabetes (16); when insulin secretion is maintained, it may counteract the effect of SUs on alpha cells, which therefore may be difficult to observe. SUs may also reduce beta cell mass, through apoptosis (17). Therefore, SUs do not sufficiently target the pathophysiological defects of the disease: they indeed stimulate insulin secretion, but they do so in a glucose-independent manner, they stimulate rather than inhibit glucagon secretion and they tend to reduce rather than to increase beta cell mass.

Clinical effects. There is an extensive experience of antidiabetic efficacy by SUs (13). Thus, SUs result in robust and marked initial reductions in HbA_{1c}, which is most evident after 6-12 months of therapy, when HbA_{1c} usually is reduced by approximately 0.5-2%. Furthermore, SUs are in general well tolerated with few adverse events (apart from hypoglycemia and weight gain, see below) and the price for SUs is low.

SUs have also been demonstrated to reduce the risk for microangiopathy in type 2 diabetes, as evident in the UKPDS over a 10 year study period (1). In contrast, whether SUs also reduce the risk of CV disease in type 2 diabetes has been a matter of discussion. The first long-term study examining this possibility, the UGDP study, showed an increased CV risk in patients treated with SU (18). It should be emphasized that it was tolbutamide which was used in that study and therefore that the results are of less value for the present-day SUs. However, also later studies have not been able to convincingly show that SUs may improve CV outcome. For example, in the UKPDS there was a trend, since 16% reduction in nonfatal myocardial infarction was observed in patients treated

with SUs, but this was not significant (1). Similarly, intense treatment with gliclazide resulted in a non-significant reduction in major macrovascular events in another study (19). Furthermore, a retrospective cohort analyses has shown an excess of mortality and congestive heart failure in patients treated with SUs (20).

The reason why the improved glycemia by SUs does not seem to be associated with reduced CV risk, in spite of improving glycemia, is not known, but two aspects may be of importance. 1) SUs may have direct effects on sulfonylurea 2A (SUR2A) and sulfonylurea 2B (SUR2B) receptors expressed in cardiomyocytes and heart smooth muscle cells, which might have negative effects. 2) Treatment with SUs is associated with hypoglycemia, which is proarrhythmogenic and may increase CV events (22,23).

Limitations. A limitation with SUs is a high rate of hypoglycemia. This is mainly because SUs stimulate insulin secretion also when glucose levels are low. It is difficult to estimate the risk for hypoglycemia during SU treatment, because many events go undetected. In the UKPDS, the annual risk for hypoglycemia in patients treated with glibenclamide was 18% during ten years (1). A recent study based on self reporting showed that the risk for severe hypoglycemia in patients treated with SUs was 7% (24). Furthermore, an observational study found that >30% of patients treated with SUs have experienced hypoglycemia during the last six months and 4% have experienced severe hypoglycemia (25). Other studies have reported that SUs are associated with a risk of 0.24-1.23 severe hypoglycemic events per 100 person-years, with a lower risk with the third generation SUs than with the older SUs (26).

One implication of hypoglycemia is severe symptoms which may need assistance of a third party and emergency assistance in hospitals. It was recently estimated that >5 000 patients each year experience severe hypoglycemia requiring emergency assistance only in the UK (26). Furthermore, patients who experience hypoglycemia often fear a tight glycemic control of the disease which may have implications for quality of life and the treatment to target. Moreover, hypoglycemia is associated with increased risk for CV diseases because it is proarrhythmogenic (22), which might explain increased mortality during intense glucose control with compounds associated with high risk of hypoglycemia (23).

A second limitation with SUs is weight gain. This is seen in most patients and is usually within 1-3 kg interval (27). The mechanism is not known, but may be related to the hyperinsulinemia caused by SU, since insulin is a trophic and anabolic hormone. It may also occur, however, due to increased food intake to defend frequent hypoglycemia induced by SUs. Regardless the reason, the consequences of increased body weight is worsened insulin resistance and further increased CV risk.

A third limitation with SUs is that glucose control often deteriorates in spite of continuous treatment (28). This is common for several different treatments, but particularly characteristic for SUs. Secondary SU failure may be related to beta cell desensitization and/or to beta cell apoptosis induced by SUs (29). Secondary failure is associated with worsening of the glycemic control.

DPP-4 inhibitors

History. The antidiabetic action of DPP-4 inhibitors is based on the incretin hormone glucagon-like peptide-1 (GLP-1) (30). GLP-1 is a 30 amino acid peptide which is produced in the small intestine and is released after meal ingestion. Together with GIP (glucose-dependent insulintropic polypeptide), it is responsible for the incretin effect, i.e., the augmentation of glucose-stimulated insulin secretion which is seen after meal ingestion. GLP-1 activates G protein coupled receptors in the pancreatic beta cells, which raises cAMP (31). This results in a glucose-dependent stimulation of insulin secretion. GLP-1 may also increase beta cell mass through neogenesis and proliferation and inhibited apoptosis, as shown in rodents (32) and it inhibits glucagon secretion (4). GLP-1 also inhibits gastric emptying and induces satiety, which add to the effects which are of potential value in the treatment of type 2 diabetes (30).

In the early 1990s, it was suggested that GLP-1 might be a potential target in the treatment of type 2 diabetes (33). Native GLP-1 is, however, rapidly inactivated by the enzyme dipeptidyl peptidase-4 (DPP-4) (11), which makes GLP-1 unsuitable use in therapy. Two strategies to take advantage of the effects of GLP-1 were instead explored. 1)The use of GLP-1 receptor agonists which are largely resistant to the action of DPP-4, and 2)the use of inhibitors of DPP-4, which prevent the inactivation of endogenous GLP-1 (11).

The proof-of-concept study showing that DPP-4 inhibition reduces glycemia in type 2 diabetes was published in 2002 (34). Several DPP-4 inhibitors have now been developed and are in different stages in clinical development. Sitagliptin, vildagliptin and saxagliptin have been approved in many countries world-wide. Linagliptin and alogliptin are in late clinical development (alogliptin has been approved in Japan) (35-39).

Mechanisms. As shown for vildagliptin and sitagliptin, DPP-4 inhibitors stimulate insulin secretion and inhibit glucagon secretion (35,36). These effects are glucose dependent, which means that they vanish when glucose levels drop. Furthermore, a recent study also showed that vildagliptin reduces insulin secretion and sustains glucagon secretion during hypoglycemia (2.5 mmol/l), verifying the glucose dependency (40). This suggests that this treatment minimizes the risk for hypoglycemia. Furthermore, DPP-4 inhibition increases beta-cell mass, as demonstrated in animal models of diabetes (41), although no such information is available in humans. Therefore, DPP-4 inhibitors target several key islet defects in type 2 diabetes: they stimulate insulin in a glucose-dependent manner, they inhibit glucagon secretion and they have a potential of increasing beta cell mass.

Clinical effects, All DPP-4 inhibitors in clinical use or in late clinical development reduce HbA_{1c} when used in monotherapy as well as when used in combination with other treatments (11,30,35-39). Overall, the reduction in HbA_{1c} is approximately 0.6-1.1% depending on the initial baseline HbA_{1c}. Furthermore, DPP-4 inhibitors are safe with a low risk for adverse events. Indeed, the overall incidence of adverse events with DPP-4 inhibitors is not higher than in placebo (45-49). Furthermore, DPP-4 inhibitors are either

weight neutral or slightly reduce body weight, and a consistent finding is that they are associated with low risk for hypoglycaemia.

Limitations. Based on published studies lasting up to two years and of clinical experience now lasting >4 years, there seems to be no limitation to DPP-4 inhibitors in clinical use from a safety and tolerability point of view. A practical limitation is that since clinical experience is low in subjects with renal insufficiency, DPP-4 inhibitors should not be used in this condition unless, for some of the DPP-4 inhibitors, the dose is reduced.

An important limitation is, however, that long-term experience, including safety, is lacking. Long-term surveillance is therefore warranted. Also, due to lack of long-term experience, it is not known whether DPP-4 inhibitors improve complications to type 2 diabetes, including CV risk. Based on surrogate markers, however, this may be likely, because DPP-4 inhibitors have been shown to improve atherogenic markers (42). However, further studies are required in this area.

Comparisons between SUs and DPP-4 inhibitors as add-on to metformin

Five head-to-head studies have compared efficacy, safety and tolerability of SUs versus DPP-4 inhibitors when added to on-going metformin therapy in subjects with type 2 diabetes with insufficient glycemic control when treated with metformin alone.

Glimepiride versus vildagliptin. One study examined the efficacy and safety of vildagliptin (50 mg twice daily, n=1562) vs. glimepiride (up to 6 mg/day, mean dose 4.5

mg/day; n=1556) when added to metformin; baseline HbA_{1c} was 7.3%. The results were reported after one year (43) and after completing two years (44). HbA_{1c} was reduced during the initial 24-30 weeks in both groups. After one year, HbA_{1c} was reduced by 0.44% by vildagliptin and by 0.53% by glimepiride, after the two years, HbA_{1c} was reduced by 0.1% in both groups compared to baseline, showing that vildagliptin and glimepiride are equipotent in reducing glycemia. The study also examined the durability of action of vildagliptin versus glimepiride (44), and found that the increase in HbA_{1c} between week 24 and 104 was significantly lower for vildagliptin (0.4%/year) than for glimepiride (0.5%/year) and by examining sustainability of treatment revealed that patients treated with vildagliptin maintained their initial response for a longer period (309 days) than those treated with glimepiride (244 days). This would suggest a better durability effect by vildagliptin, although it must be emphasized that a two-year study is too short for allowing conclusions on long-term durability.

There were no differences in the overall number of adverse events between the two groups, with the important exception of hypoglycemia. Hypoglycemia was thus reported in 281 patients (18.2%) treated with glimepiride (838 events), whereas it occurred in 35 patients (2.3%) patients treated with vildagliptin (59 events). Of particular importance is that severe hypoglycemia occurred in 14 patients treated with glimepiride (8.0%) but only in one patient treated with vildagliptin (0.2%).

Another difference between the two compounds was that whereas vildagliptin was associated with a slight weight loss (0.3kg), glimepiride treatment resulted in weight gain

(+1.2kg). In regard to dyslipidemia, vildagliptin had a modest beneficial effect compared to glimepiride on HDL cholesterol, total cholesterol and triglycerides.

In a subgroup of patients, meal tests were undertaken before and after two years of treatment; meal-induced glucagon secretion was lowered by vildagliptin, whereas glucagon secretion was increased by glimepiride (45). This shows that the hyperglucagonemia, which is an important portion of the islet dysfunction in diabetes, was targeted over a two years time by vildagliptin but not by glimepiride.

Glipizide versus sitagliptin. Another study evaluated the efficacy and safety of adding sitagliptin (100 mg once daily; n=588) versus glipizide (5 mg/day with up-titration to 20 mg/day, mean dose 9.2 mg/day; n=584) when added to ongoing metformin; baseline HbA_{1c} was 7.3%. The results were reported after one year (46) and after completing two years (47). HbA_{1c} was reduced during the initial 30 week study period in both groups. After the two years, baseline HbA_{1c} was reduced by 0.54% by sitagliptin and by 0.51% by glipizide. It was also found that the rise in HbA_{1c} from week 24 to the end of the study was less with sitagliptin (0.16%/year) than with glipizide (0.26%/year). This would suggest a better long-term effect of sitagliptin, although, as underlined above, a study of only two years is too short to estimate durability of improved glycemia.

A few specific adverse events were more common each of the groups but these differences were minor. The only adverse event which showed a large difference between the groups was hypoglycemia. Thus, hypoglycemia was reported in 199 patients (34%)

treated with glipizide (805 events), compared to 31 patients (5%) treated with sitagliptin (57 events). Severe hypoglycemia occurred in 18 patients treated with glipizide (7.0%) and in 2 patients treated with sitagliptin (0.8%).

Also in this study, a difference in body weight was observed between the groups. Thus, sitagliptin was associated with a slight weight loss (-1.6 kg) compared with weight gain (+0.7 kg) with glipizide.

In a subgroup of patients, a meal test was undertaken before and after two years with estimation of glucose excursion and beta cell function. It was found that glucose excursion was smaller in the sitagliptin than in the glipizide group. Furthermore, beta-cell function (C-peptide data) was stable in the sitagliptin group as compared to before the study, whereas a reduction in insulin secretion versus baseline was observed in the glipizide group.

Gliclazide versus vildagliptin. A third study evaluated the efficacy and safety in a 52 week study of adding vildagliptin (50 mg twice daily; n=513) vs gliclazide (up to 320 mg/day; n=494) to metformin treatment; baseline HbA_{1c} was 8.5% (48). After 52 weeks, HbA_{1c} was reduced by 0.81% by saxagliptin and by 0.85% by gliclazide. Six of the patients treated with vildagliptin reported hypoglycemia compared to 11 of the patients on gliclazide. Body weight was not significantly changed by vildagliptin but increased (by 1.4 kg) by glipizide. Apart from hypoglycemia, adverse events were not different between the groups.

Glipizide versus saxagliptin. A fourth study evaluated the efficacy and safety in a 52 week study of adding saxagliptin (5 mg once daily; n=588) vs glipizide (5 mg/day with up-titration to 20 mg/day, mean dose 14.7 mg/day; n=584) to metformin treatment; baseline HbA_{1c} was 7.7%. The results are so far only available in abstract form (49). After 52 weeks, HbA_{1c} was reduced by 0.74% by saxagliptin and by 0.80% by glipizide. In the patients treated with saxagliptin, 3% reported hypoglycemia, whereas 36% of patients treated with glipizide reported hypoglycemia. Body weight was slightly reduced (by 1.1 kg) by saxagliptin but slightly increased (by 1.0 kg) by glipizide. Apart from hypoglycemia, adverse events were not different between the groups.

Glimepiride versus sitagliptin. A fifth study evaluated the efficacy and safety in a 30 week study of adding sitagliptin (100 mg once daily; n=516) versus glimepiride (titration up to 6 mg/day; n=519) to metformin treatment; baseline HbA_{1c} was 7.5%. The results are so far only available in abstract form (50). After 30 weeks, HbA_{1c} was reduced by 0.47% by sitagliptin and by 0.54% by glimepiride. A total of 7% of patients treated with sitagliptin reported hypoglycemia (73 events), whereas of those treated with glimepiride, 22% reported hypoglycemia (460 events). Furthermore, body weight was slightly reduced (by 0.8 kg) by sitagliptin but slightly increased (by 1.2 kg) by glimepiride. Apart from hypoglycemia, adverse events were not different between the groups.

These comparative studies show together that DPP-4 inhibition when added to on-going therapy with metformin in patients with insufficient glycemic control when treated with metformin alone is equipotent with SUs over periods up to two years. Furthermore, DPP-4 inhibition is associated with much lower incidence of hypoglycemia than SUs and with a reduction in body weight, whereas SUs increases body weight.

Sulfonylureas and DPP-4 inhibitors as add-on to metformin – which is most desirable?

Several aspects need to be considered when choosing between sulfonylureas versus DPP-4 inhibitors as add-on to metformin in patients with insufficient glycemic control when treated with metformin alone:

- 1) The therapy should *target pathophysiological defects in type 2 diabetes*. This is important for being able to prevent deterioration of the disease. This is a great advantage of DPP-4 inhibitors over SUs. Thus, DPP-4 inhibitors a)stimulate insulin secretion and b)inhibit glucagon secretion, both effects being in glucose-dependent manner, and c)in rodent studies they increase beta cell mass. In contrast, SUs a)stimulate insulin secretion in a glucose-independent manner, b)stimulate rather than inhibit glucagon secretion, and 3)reduce beta cell mass through an augmented apoptosis, as shown in rodents.
- 2) The treatment should result in *clinically meaningful reduction in HbA_{1c}*. This is undertaken by both DPP-4 inhibitors and SUs, since they reduce HbA_{1c} and in

head-to-head studies over thirty weeks to two years, they do so in an equipotent manner.

- 3) The reduction in HbA_{1c} should show *sustained durability*. Mechanistically, DPP-4 inhibitors have been shown to increase beta cell mass in rodents and GLP-1 reduces rather than increases beta cell apoptosis, which may suggest potential for long-term beta cell preservation and durability. Clinically, DPP-4 inhibitors show durability over the two years they have been studied (44,47). In contrast, SUs may induce beta cell apoptosis and have in general been associated with poor long-term clinical durability with secondary failures (28). However, more long-term studies with DPP-4 inhibitors vs SUs are required, and so far no conclusion regarding this criterion is possible.
- 4) The treatment should *be safe with no hypoglycemia, weight gain or other adverse events*. Number of adverse events, apart from hypoglycemia and weight gain, does not seem to differ between SUs and DPP-4 inhibitors in head-to-head studies. However, hypoglycemia, which is rarely observed during treatment with DPP-4 inhibitors, is commonly observed during treatment with SU, as was clearly demonstrated in the head-to-head studies (44,45,47-50). Similarly, whereas DPP-4 inhibitors are weight neutral or reduce body weight, there is an increase in body weight during treatment with SUs.
- 5) The treatment should *prevent retinopathy, nephropathy and neuropathy and reduce the risk for CV events*. SUs have been shown to prevent microangiopathic complications to diabetes (1), whereas no such evidence exists for the DPP-4 inhibitors. In regard to reducing CV risk, no clear evidence for such an effect of

SUs exist (20). For DPP-4 inhibitors, no long term studies with hard end point data have been performed; such studies are under way.

- 6) The treatment should be *acceptable for the patient, the health care providers and the community*, including a favorable cost-benefit profile. There is a difference in how the treatment is initiated, in that SUs need up-titration, whereas DPP-4 inhibitors are given at the same dose in all patients, except in those with renal impairment. This may indicate that DPP-4 inhibitors are more user friendly, to which is added that self glucose monitoring is not required in patients treated with DPP-4 inhibitors, due to the low risk for hypoglycemia. Another advantage for DPP-4 inhibitors is that patients given these compounds do not fear hypoglycemia or increased body weight, which are limitations to SUs. On the contrary, pricing is an advantage for SUs, since these are cheaper. However, in this context it needs to be emphasized that full health economy analysis need to be undertaken including additional costs due to hypoglycemic events and long-term failure.

The Jury.

The development and introduction of SUs in the treatment of type 2 diabetes had a tremendous importance for the patients, since they were offered efficient oral treatment to combat the disease. SUs have also been in the center of treatment for several decades. However, after more than 50 years of use, researchers have provided the patients and clinicians with a class of compounds that may challenge the SUs as central compounds for management of hyperglycemia in type 2 diabetes. As is evident from this article,

several different aspects can be raised when comparing the two groups. Fig. 1 illustrates the different aspects.

It is clear from the comparisons reported here and illustrated in Fig. 1 that there are several advantages of DPP-4 inhibitors over SUs. Hence, the important limitations with SUs, i.e., the high risk for hypoglycemia and the increase in body weight are not seen with DPP-4 inhibitors. Furthermore, in terms of efficacy, the two approaches seem equipotent over the two year periods they have been compared, but based on animal studies, DPP-4 inhibitors might be more favorable in terms of potential for long-term durability. However, it is very important with long-term surveillance to explore the long-term safety of DPP-4 inhibitors, and, also, to establish the long-term consequences for secondary complications to diabetes. SUs have a good record for microvascular complications, but less convincing for macrovascular complications. Any long-term benefits of any treatment of type 2 diabetes will be of more importance than the cost of the compound; the prize issue may therefore become less important as times go by. The conclusion of the comparison between SUs and DPP-4 inhibitors is therefore that SUs have been the number one agent to add-on to metformin for many decades, but now they are less desirable than the new DPP-4 inhibitors, although long-term efficacy and safety (>4 years) are still needed for DPP-4 inhibitors.

Acknowledgements and disclosure. The author acknowledges the Swedish Research Council (grant nr 6834), Region Skåne, Faculty of Medicine, Lund University, and Novartis AG for financial support in own studies reported here. The author discloses

receipt of honoraria for lectures from AstraZeneca, GSK, Merck, Novartis, Novo Nordisk, Roche, Sanofi Aventis and Servier, and receipt of research grants from AstraZeneca, Merck and Novartis.

References

1. U.K. Prospective Diabetes Study (UKPDS) Group: Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *Lancet* 1998, 352: 837-853.
2. Gaede P, Vedel P, Larsen N, et al.: Multifactorial intervention and cardiovascular disease in patients with type 2 diabetes. *N Engl J Med* 2003, 348:383-393.
3. Kahn SE: The relative contributions of insulin resistance and beta-cell dysfunction to the pathophysiology of type 2 diabetes. *Diabetologia* 2003, 46:3-19.
4. Dunning BE, Foley JE, Ahrén B: Alpha cell function in health and disease: influence of glucagon-like peptide-1. *Diabetologia* 2005, 48:1700-1713.
5. Butler AE, Janson J, Bonner-Weir S, Ritzel R, Rizza RA, Butler PC: Beta-cell deficit and increased beta-cell apoptosis in humans with type 2 diabetes. *Diabetes* 2003, 52:102-110
6. Ahrén B: Beta- and alpha-cell dysfunction in subjects developing impaired glucose tolerance: outcome of a 12-year prospective study in postmenopausal Caucasian women. *Diabetes* 2009, 58:726-731.
7. Adler AI, Shaw EJ, Stokes T, Ruiz F: Newer agents for blood glucose control in type 2 diabetes: summary of NICE guidance. *Brit Med J* 2009, 338:b1668

8. IDF Clinical guidelines Task Force: Global guideline for type 2 diabetes: recommendations for standard, comprehensive, and minimal care. *Diabet Med.* 2006,23:579–593.
9. Rodbard HW, Jellinger PS, Davidson JA, et al.: Statement by an American Association of Clinical Endocrinologists/American College of Endocrinology consensus panel on type 2 diabetes mellitus: an algorithm for glycemic control. *Endocr Pract* 2009, 15:540-559.
10. Nathan DM, Buse JB, Davidson MB, et al.: Medical management of hyperglycaemia in type 2 diabetes mellitus: a consensus algorithm for the initiation and adjustment of therapy. A consensus statement from the American Diabetes Association and the European Association for the Study of Diabetes. *Diabetologia* 2009, 52:17-30.
11. Ahrén B: Emerging dipeptidyl peptidase-4 inhibitors for the treatment of diabetes. *Expert Opin Emerg Drugs* 2008, 13: 593-507.
12. Levine R: Sulfonylureas: background and development of the field. *Diabetes Care* 1984, 7: suppl 1:3-7.
13. Rendell M: The role of sulphonylureas in the management of type 2 diabetes mellitus. *Drugs* 2004, 64:1339-1358.
14. Gribble FM, Reimann F: Sulphonylurea action revisited: the post-cloning area. *Diabetologia* 2003, 46:875-891.
15. Bokvist K, Olsen HL, Hoy M, et al.: Characterisation of sulphonylurea and ATP-regulated K⁺ channels in rat pancreatic A-cells. *Pfl Arch* 1999, 438:428–436.

16. Østergård T, Degn KB, Gall MA, et al.: The insulin secretagogues glibenclamide and repaglinide do not influence growth hormone secretion in humans but stimulate glucagon secretion during profound insulin deficiency. *J Clin Endocrinol Metab* 2004, 89:297-302.
17. Lupi R, Del Prato S: Beta-cell apoptosis in type 2 diabetes: quantitative and functional consequences. *Diabetes Metab* 2008, 34: suppl 2:S56-64.
18. Meinert CL, Knatterud GL, Prout TE, Klimt CR: A study of the effects of hypoglycemic agents on vascular complications in patients with adult-onset diabetes. II. Mortality results. *Diabetes* 1970, 19: suppl.: 789-830.
19. ADVANCE Collaborative Group: Intensive blood glucose control and vascular outcomes in patients with type 2 diabetes. *N Engl J Med* 2008, 358:2560-2572.
20. Tzoulaki I, Molokhia M, Curcin V, et al.: Risk of cardiovascular disease and all cause mortality among patients with type 2 diabetes prescribed oral antidiabetes drugs: retrospective cohort study using UK general practice research base. *Brit Med J* 2009, 339:b4731
21. Nordin C: The case for hypoglycaemia as a proarrhythmic event: basic and clinical evidence. *Diabetologia* 2010, 53:1552-1561.
22. Zhou M, He HJ, Suzuki R, et al.: Localization of sulfonylurea receptor subunits, SUR2A and SUR2B, in rat heart. *J Histochem Cytochem* 2007, 55:795-804.
23. Mannucci E, Monami M, Lamanna C, Gori F, Marchionni N: Prevention of cardiovascular disease through glycemic control in type 2 diabetes: a meta-analysis of randomized clinical trials. *Nutr Metab Cardiovasc Dis* 2009, 19:604-612

24. Belcher G, Lambert C, Goh KL, Edwards G, Valbuena M: Cardiovascular effects of treatment of type 2 diabetes with pioglitazone, metformin and gliclazide. *Int J Clin Pract* 2004, 58: 833– 837.
25. Amiel SA, Dixon T, Mann R, Jameson K: Hypoglycaemia in type 2 diabetes. *Diabet Med* 25:245-254, 2008
26. Gerstein HC, Miller ME, Byington RP, et al.: Effects of intensive glucose lowering in type 2 diabetes. *N Engl J Med* 2008, 358:2545-2559.
27. Campbell IW: Comparing the actions of older and newer therapies on body weight: to what extent should these effects guide the selection of antidiabetic therapy? *Int J Clin Pract* 2010, 64:791-801.
28. Harrower AD: Comparison of efficacy, secondary failure rate, and complications of sulfonylureas. *J Diabetes Complications* 1994, 8:201-203.
29. Del Guerra S, Marselli L, Lupi R, et al.: Effects of prolonged in vitro exposure to sulphonylureas on the function and survival of human islets. *J Diabetes Complications* 2005, 19:60-64.
30. Drucker DJ, Nauck MA: The incretin system: glucagon-like peptide-1 receptor agonists and dipeptidyl peptidase-4 inhibitors in type 2 diabetes. *Lancet* 2006, 368:1696-1705.
31. Ahrén B: Islet G-protein-coupled receptors as potential targets for treatment of type 2 diabetes. *Nat Rev Drug Discov* 2009, 8: 369-385.
32. Perfetti R, Hui H: The role of GLP-1 in the life and death of pancreatic beta cells. *Horm Metab Res* 2004, 36: 804-810.

33. Gutniak M, Ørskov C, Holst JJ, et al.: Antidiabetic effect of glucagon-like peptide-1 (7-36) amide in normal subjects and patients with diabetes mellitus. *N Engl J Med* 1992, 326: 1316-1322.
34. Ahrén B, Simonsson E, Larsson H, et al.: Inhibition of dipeptidyl peptidase IV improves metabolic control over a 4 week study period in type 2 diabetes. *Diabetes Care* 2002, 25: 869-875.
35. Ahrén B: Vildagliptin: an inhibitor of dipeptidyl peptidase-4 with antidiabetic properties. *Exp Opin Invest Drugs* 2006, 15:431-442.
36. Dhillon S: Sitagliptin: a review of its use in the management of type 2 diabetes mellitus. *Drugs* 2010, 70:489-512
37. Deacon CF, Holst JJ: Saxagliptin: a new dipeptidyl peptidase-4 inhibitor for the treatment of type 2 diabetes. *Adv Ther* 2009; 26: 488-499.
38. Deacon CF, Holst JJ: Linagliptin, a xanthine-based dipeptidyl peptidase-4 inhibitor with an unusual profile for the treatment of type 2 diabetes. *Exp Opin Invest Drugs* 210, 19: 133-140.
39. Pratley RE: Alogliptin: a new, highly selective dipeptidyl peptidase-4 inhibitor for the treatment of type 2 diabetes. *Exp Opin Pharmacother* 2009, 20:503-512.
40. ● Ahren B, Schweizer A, Dejager S, et al. Vildagliptin enhances islet responsiveness to both hyper- and hypoglycemia in patients with type 2 diabetes. *J Clin Endocrinol Metab* 2009, 94:1236-1243. This study shows that the glucagon counterregulation to hypoglycemia is preserved during treatment with vildagliptin.

41. Mu J, Woods J, Zhou YP, et al.: Chronic inhibition of dipeptidyl peptidase-4 with a sitagliptin analog preserves pancreatic beta-cell mass and function in a rodent model of type 2 diabetes. *Diabetes* 2006, 55:1695-1704
42. Rizzo M, Rizvi AA, Spinas GA, Rini GB, Berneis K: Glucose lowering and anti-atherogenic effects of incretin-based therapies: GLP1 analogues and DPP-4 inhibitors. *Exp Opin Investig Drug* 2009, 18:1495-1503
43. • Ferrannini E, Fonseca V, Zinman B, et al.: Fifty-two-week efficacy and safety of vildagliptin vs. glimepiride in patients with type 2 diabetes mellitus inadequately controlled on metformin monotherapy. *Diabetes Obes Metab* 2009, 11:157-166. This study demonstrates that vildagliptin and glimepiride have the same reduction in HbA_{1c} but with marked difference in that glimepiride is associated with much more hypoglycemic events and body weight gain.
44. Matthews DR, Dejager S, Ahrén B, et al.: Vildagliptin add-on to metformin produces similar efficacy and reduced hypoglycaemic risk compared with glimepiride, with no weight gain: results from a 2-year study. *Diabetes Obes Metab* 2010, 12:780-789.
45. Ahrén B, Foley JE, Ferrannini E, et al.: Changes in prandial glucagon levels after a 2-year treatment with vildagliptin or glimepiride in patients with type 2 diabetes inadequately controlled with metformin monotherapy. *Diabetes Care* 2010, 33:730-732.
46. • Nauck MA, Maininger G, Sheng D, et al.: Efficacy and safety of the dipeptidyl peptidase-4 inhibitor, sitagliptin, compared with the sulfonylurea, glipizide, in patients with type 2 diabetes inadequately controlled on metformin alone: a

- randomized, double-blind, non-inferiority trial. *Diabetes Obes Metab* 2007, 9:194-205. This study demonstrates that sitagliptin and glipizide have the same reduction in HbA_{1c} but with marked difference in that glipizide is associated with much more hypoglycemic events and body weight gain.
- 47.
48. Seck T, Nauck M, Sheng D, et al.: Safety and efficacy of treatment with sitagliptin or glipizide in patients with type 2 diabetes inadequately controlled on metformin: a 2-year study. *Int J Clin Pract* 2010, 64:562-576.
49. Filozof C, Gautier JF: A comparison of efficacy and safety of vildagliptin and glimepiride in combination with metformin in patients with type 2 diabetes inadequately controlled with metformin alone: a 52-week, randomized study. *Diabetic Med* 2010, 27:318-326.
50. Göke B, Gallwitz B, Eriksson J, et al.: Saxagliptin is non-inferior to glipizide when added to metformin in patients with type 2 diabetes mellitus. *Diabetes* 2010, 59: suppl 2: A158.
51. Goldstein BJ, Seck T, Chen Y, et al.: Sitagliptin compared with glimepiride provides similar efficacy with weight loss and less hypoglycemia when added to metformin therapy in patients with type 2 diabetes mellitus. *Diabetologia* 2010, 53: suppl 1: S325

Legends to Figure 1

Illustration of the balance between advantages and disadvantages for SUs vs DPP-4 inhibitors in the management of hyperglycemia in 2 diabetes. Management with SU is placed on the left at the balance, whereas management with DPP-4 inhibition (DPP-4i) is placed on the right. Advantages of each of the two approaches are listed above the scale, which make up the force of each strategy to be more desirable over the other, whereas disadvantages are listed below the scale, working against desirability.




Fig. 1