



LUND UNIVERSITY

Autistic-like traits and their association with mental health problems in two nationwide twin cohorts of children and adults.

Lundström, Sebastian; Chang, Z; Kerekes, Nora; Gumpert, C H; Råstam, Maria; Gillberg, C; Lichtenstein, P; Anckarsäter, Henrik

Published in:
Psychological Medicine

DOI:
[10.1017/S0033291711000377](https://doi.org/10.1017/S0033291711000377)

2011

[Link to publication](#)

Citation for published version (APA):

Lundström, S., Chang, Z., Kerekes, N., Gumpert, C. H., Råstam, M., Gillberg, C., Lichtenstein, P., & Anckarsäter, H. (2011). Autistic-like traits and their association with mental health problems in two nationwide twin cohorts of children and adults. *Psychological Medicine*, 41(11), 2423-2433.
<https://doi.org/10.1017/S0033291711000377>

Total number of authors:
8

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Autistic-like traits and their association with mental health problems in two nationwide twin cohorts of children and adults

S. Lundström^{1,2*}, Z. Chang³, N. Kerekes^{2,4}, C. H. Gumpert⁵, M. Råstam^{6,7}, C. Gillberg⁸, P. Lichtenstein³ and H. Anckarsäter^{1,4}

¹ Institute of Clinical Sciences, Lund University, Malmö, Sweden

² Swedish Prison and Probation Service, R&D Unit, Gothenburg, Sweden

³ Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden

⁴ Institute of Neuroscience and Physiology, Forensic Psychiatry, University of Gothenburg, Sweden

⁵ Division of Social and Forensic Psychiatry, Department of Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden

⁶ Institute of Clinical Sciences, Child and Adolescent Psychiatry, Lund University, Lund, Sweden

⁷ Institute of Neuroscience and Physiology, Child and Adolescent Psychiatry, University of Gothenburg, Sweden

⁸ Institute of Neuroscience and Physiology, Gillberg Neuropsychiatry Center, University of Gothenburg, Sweden

Background. Autistic-like traits (ALTs), that is restrictions in intuitive social interaction, communication and flexibility of interests and behaviors, were studied in two population-based Swedish twin studies, one in children and one in adults: (1) to examine whether the variability in ALTs is a meaningful risk factor for concomitant attention deficit hyperactivity disorder (ADHD), anxiety, conduct problems, depression and substance abuse, and (2) to assess whether common genetic and environmental susceptibilities can help to explain co-existence of ALTs and traits associated with such concomitant problems.

Method. Two nationwide twin cohorts from Sweden (consisting of 11 222 children and 18 349 adults) were assessed by DSM-based symptom algorithms for autism. The twins were divided into six groups based on their degree of ALTs and the risk for concomitant mental health problems was calculated for each group. Genetic and environmental susceptibilities common to ALTs and the other problem types were examined using bivariate twin modeling.

Results. In both cohorts, even the lowest degree of ALTs increased the risk for all other types of mental health problems, and these risk estimates increased monotonically with the number of ALTs. For all conditions, common genetic and environmental factors could be discerned. Overall, the phenotypic correlation between ALTs and the traits examined were less pronounced in adulthood than in childhood and less affected by genetic compared with environmental factors.

Conclusions. Even low-grade ALTs are relevant to clinical psychiatry as they increase the risk for several heterotypical mental health problems. The association is influenced partly by common genetic and environmental susceptibilities. Attention to co-existing ALTs is warranted in research on a wide range of mental disorders.

Received 29 October 2010; Revised 10 February 2011; Accepted 12 February 2011

Key words: Autism, autistic traits, epidemiology, genetics, twin.

Introduction

Autism spectrum disorders (ASDs), which correspond to the pervasive developmental disorders in DSM-IV (APA, 1994), include autistic disorder, Asperger's disorder and pervasive developmental disorder not otherwise specified (sometimes referred to as atypical autism). They affect about 1% of the population (as shown in several different European countries,

e.g. Gillberg *et al.* 2006; Baird *et al.* 2006) and are characterized by deficits in social interaction, communication and behavioral flexibility. 'Classic' autism is defined as a severe neurodevelopmental disorder with pronounced socio-communicative deficits noted before 3 years of age, often accompanied by mental retardation, whereas Asperger's disorder is defined by an overall normal level of verbal functioning but significantly decreased intuitive social communication skills and ability to adapt behaviors and interests to contexts and circumstances (APA, 1994). It is not clear whether these diagnoses represent true categories among the ASDs or whether they are clinically defined

* Address for correspondence: Mr S. Lundström, Forensic Psychiatry, Lillhagsparken 3, 422 50 Hisings Backa, Sweden.
(Email: sebastian.lundstrom@med.lu.se)

types characterized by uneven intelligence profiles, cognitive dysfunctions, age at onset and gender-specific variation.

Other mental disorders, such as attention deficit hyperactivity disorder (ADHD), mood disorders, schizophrenia and personality disorders, seem to be present in one form or another in the majority of cases diagnosed with an ASD (e.g. in children: Simonoff *et al.* 2008; in adults: Hofvander *et al.* 2009). Clinical (Lee & Ousley, 2006), epidemiological (Kadesjö *et al.* 1999), behavioral genetic (Lichtenstein *et al.* 2010) and molecular genetic (Roohi *et al.* 2009) studies have reported commonalities in ASDs and ADHD, and clinical studies have shown overlaps between ASDs and anxiety (Pine *et al.* 2008), depression (Ghaziuddin & Greden, 1998) and conduct disorder (reviewed by Anckarsäter *et al.* 2008).

ASDs can be construed as the phenotypical expression of an extreme lowermost end of a continuum of socio-communicative abilities that are distributed dimensionally in the general population (Gillberg, 1992). Autistic-like traits (ALTs), that is significant problems or peculiarities in social style, perception of self and others, pragmatic communication and adaptation to the environment, which do not meet formal criteria for a mental disorder, have also been found in the normal distribution of social responsiveness (Constantino & Todd, 2003). These traits are similar to the 'broader autism phenotype' described by Folstein & Rutter (1977) in unaffected siblings of children identified with autism, but the nosological link between the disorder, broader phenotype and ALTs has not yet been established. Neither have the three different symptom clusters described in ASDs been established for the ALTs, although two twin studies, one British and one Swedish, have recently demonstrated etiological heterogeneity within the autism-related symptomatology (Ronald *et al.* 2006, 2010). The current state of knowledge on ASDs has not been paralleled by systematic research on ALTs, although such traits have been a noted feature in clinical studies of ADHD (Reiersen *et al.* 2008; Rommelse *et al.* 2009), internalizing problems/depression (Kanne *et al.* 2009), conduct disorder (Gilmour *et al.* 2004) and behavioral problems (Hoekstra *et al.* 2007). Recent twin studies have also documented substantial shared genetic influences operating over both ALTs and traits associated with ADHD and anxiety in children (Ronald *et al.* 2008, 2010; Hallett *et al.* 2009). It has not yet been established, however, whether interindividual differences in subthreshold ALTs are associated with the risk for ADHD, anxiety, conduct problems, depression and substance abuse, nor whether there are similar associations between ALTs and other mental health problems both in children and in adults.

We have used data from two large-scale, population-based twin cohorts, one of children and one of adults, (1) to test whether varying degrees of ALTs are meaningful risk factors for concomitant ADHD, anxiety disorder, conduct problems, depression and substance abuse both in childhood and in adulthood, and (2) to assess whether there are common genetic and environmental susceptibilities explaining the co-existence of ALTs and traits associated with ADHD, anxiety, conduct problems, depression and substance abuse both in childhood and in adulthood.

Method

Sample

The present study is based on two prospective, nationally representative twin studies from Sweden: the Child and Adolescent Twin Study in Sweden (CATSS), previously used to investigate the etiology of ASDs and their triad structure (Lichtenstein *et al.* 2010; Lundström *et al.* 2010), and the Study of Twin Adults: Genes and Environment (STAGE), previously used to study addiction and psychiatric disorders (Furberg *et al.* 2008; Frisell *et al.* 2010). Both cohorts are described in Table 1.

Measures in CATSS

Parents of twins were interviewed with the Autism – Tics, ADHD, and other Co-morbidities inventory (A-TAC; Hansson *et al.* 2005), a psychiatric telephone interview validated for administration by lay interviewers. The A-TAC has good test–retest and excellent inter-rater reliability and construct validity according to two validation studies (Hansson *et al.* 2005; Larson *et al.* 2010), and convergent validity with the Child Behavior Checklist (CBCL; Halleröd *et al.* 2010). Items in the A-TAC are scored 1, 0.5 or 0 for 'yes', 'yes, to some extent' or 'no' respectively in a lifetime perspective and are organized in modules assessing different psychiatric diagnoses. Psychometric data for the dimensional and categorical assessments of the targeted conditions are based on the final validation study (Larson *et al.* 2010) and detailed below.

ASDs

To correspond with the STAGE data in this paper, the DSM-IV score from the A-TAC was chosen (Larson *et al.* 2010), containing 12 items assessing the DSM-IV algorithm for the identification of autism. This score has an excellent validity for identifying ASDs [area under a receiver operating characteristics (ROC) curve, AUC=0.96; Larson *et al.* 2010] and a high internal consistency (Cronbach's $\alpha=0.86$). A cut-off

Table 1. Description of the two cohorts of twins: CATSS and STAGE

Cohort	Participants ^a	ALT measure	Assessment method	<i>n</i>	Zygosity ^b	Sex (% male)
CATSS (Anckarsäter <i>et al.</i> unpublished observations)	Nationwide cohort of all 9- and 12-year-old twins born from July 1992 to June 1998, assessed at ages 9 or 12	A-TAC (Hanson <i>et al.</i> 2005; Larson <i>et al.</i> 2010)	Parental telephone interviews	11 222	29% MZ (50% male) 29% DZ same-sex (54% male) 33% DZ opposite sex 9% unknown	51
STAGE (Furberg <i>et al.</i> 2008)	Nationwide cohort of all twins born 1959–1985	DSM-IV-based questions on autistic disorder, modified for self-reporting purposes	Web questionnaires and telephone interviews	18 349	32% MZ (36% male) 21% DZ same-sex (37% male) 30% DZ opposite sex 17% unknown	40

CATSS, Child and Adolescent Twin Study in Sweden; STAGE, Study of Twin Adults: Genes and Environment; A-TAC, Autism – Tics, ADHD, and other Co-morbidities inventory; ADHD, attention deficit hyperactivity disorder; MZ, monozygotic; DZ, dizygotic.

^a The baseline data consisted of 11 122 individuals in CATSS and 18 349 individuals in STAGE. Only twin pairs with no more than one item missing on the autistic-like trait (ALT) variable were included, giving 10 773 individuals in CATSS and 16 695 in STAGE. Excluding individuals for whom data were missing on the response variables yielded seven final populations: in CATSS 10 760 individuals for ADHD, 10 770 for anxiety, and 10 768 for conduct disorder, and in STAGE of 16 647 individuals for ADHD, 14 438 for anxiety, 16 354 for depression, and 14 670 for substance abuse.

^b In both cohorts, zygosity determination was based on a validated algorithm with >95% predictive value (Lichtenstein *et al.* 2002).

score of 4.5 yields a sensitivity of 0.86 and a specificity of 0.94 for the identification of ASDs (according to re-analyses of the data from the paper by Larson *et al.* 2010), and a prevalence of 1.5% in the CATSS cohorts used here. A total of 10 773 subjects in the CATSS were classified into six groups according to their ASD DSM scores and arranged to be comparable with respect to percentage proportions across CATSS and STAGE: group 1 (0 points) $n=7499$ (69.6%); group 2 (0.5–1 point) $n=2339$ (21.7%); group 3 (1.5–2 points) $n=497$ (4.6%); group 4 (2.5–3 points) $n=191$ (1.7%); group 5 (3.5–4 points) $n=88$ (0.8%); and group 6 (≥ 4.5 points) $n=159$ (1.5%), the latter thus corresponding to probable ASD cases according to the validation study (referred to as the ALT 1–5 and ASD groups). The same scale, ranging from 0 to 12, was used as a dimensional measure of ALTs. All three ASD subdomains in the A-TAC contribute to the ALTs with some overall differences in population frequency [social interaction problems (mean=0.09), communication problems (mean=0.14) and lack of flexibility (mean=0.17)].

ADHD

The A-TAC DSM-IV score for ADHD consists of 18 items and was used to assess ADHD. This scale has

excellent validity for identifying ADHD (AUC=0.94; Larson *et al.* 2010), with an optimal cut-off score at 8 yielding a sensitivity of 0.87 and a specificity of 0.88 (derived from the validation study as for the ASD score), with an excellent internal consistency (Cronbach's $\alpha=0.90$) and a prevalence of 4.8% in the CATSS. The same scale, ranging from 0 to 18, was used as a dimensional measure of ADHD traits.

Anxiety

Six items in the A-TAC were used to assess anxiety. This scale had good validity for identifying anxiety disorders in the second validation study (AUC=0.78, unpublished data) and good internal consistency (Cronbach's $\alpha=0.65$), yielding a prevalence of 6% in the present study. The same scale, ranging from 0 to 6, was used as a dimensional measure of anxiety traits.

Conduct problems

Five DSM-IV-based items in the A-TAC were used to assess and identify conduct problems. Using the second validation study enriched by A-TAC results from 66 institutionalized adolescents with a higher prevalence of conduct problems (from the study described by Ståhlberg *et al.* 2010), the AUC for identifying

conduct disorder was 0.95. Cronbach's α for the conduct scale was 0.65. As our subjects were only 9 or 12 years old, the prevalence of conduct disorder was low ($n=48$), and we chose to categorize subjects who scored ≥ 1.5 as cases, yielding a prevalence of conduct problems at 1.7%. The same scale, ranging from 0 to 5, was used as a dimensional measure of conduct problems.

Measures in STAGE

Participants were interviewed with a web questionnaire including self-rated DSM-IV-based questions for ASDs and ADHD, scored as 1 for 'yes', 0.5 for 'yes, to some extent' and 0 for 'no'. Depressive traits, anxiety traits and substance abuse were scored as 1 for 'yes' and 0 for 'no'. All questions explicitly referred to a lifetime perspective.

ASDs

By analogy with the A-TAC algorithm, 12 DSM-IV-based items (Cronbach's $\alpha=0.63$) were used to measure ALTs. The 16 695 subjects were divided into six groups based on their scoring on these items: group 1 (0–2 points) $n=12\,409$ (74.3%); group 2 (2.5–3 points) $n=2509$ (15%); group 3 (3.5–4.5 points) $n=1301$ (7.8%); group 4 (5–5.5 points) $n=265$ (1.6%); group 5 (6–6.5 points) $n=131$ (0.8%); and group 6 (≥ 7 points) $n=80$ (0.50%) (referred to as ALT groups 1–5 and ASD group).

ADHD

Eighteen items were used for the identification of ADHD (similar to the A-TAC algorithm and described previously by Frisell *et al.* 2010). Cronbach's α for the whole scale was 0.84. ADHD was considered to be probable in those who responded 'yes to some extent' or 'yes' to six or more questions in the hyperactivity dimension or in the inattentive dimension according to the cut-off used in the DSM-IV, yielding a prevalence of 2.2%. The same scale, ranging from 0 to 18, was used as a dimensional measure of ADHD traits.

Anxiety

Six items corresponding to generalized anxiety disorder (GAD) in the DSM-IV, preceded by a question about whether the subject had ever experienced symptoms of anxiety and worrying lasting for one month or more, were used for the identification of anxiety disorder (Frisell *et al.* 2010). The cut-off for a possible GAD was 3 plus an affirmative answer to the question of duration, yielding a prevalence of 4.5%.

The same scale, ranging from 0 to 6, was used as a dimensional measure of anxiety traits.

Depression

Nine DSM-IV-based items were used to identify possible major depression. The cut-off for a possible lifetime major depression was 5 plus an affirmative answer to the C criterion (i.e. that the symptoms were causing clinically significant suffering) and the E criterion (i.e. that the symptoms were not better accounted for by bereavement) in DSM-IV (APA, 1994), yielding a prevalence of 13.4%. Depressive traits were measured by an 11-item version of the Iowa version of the Center for Epidemiologic Studies Depression Scale (CES-D; Carpenter *et al.* 1998), a measure widely used in epidemiological settings to screen for depression.

Substance abuse

One item was used to capture substance and alcohol abuse: 'Do you have or have you ever had problems with alcohol or drugs?' The item is scored dichotomously, categorizing 2.5% of the sample as having experienced problems with alcohol or drugs at some time in life.

Covariates

Covariates included in all analyses were zygosity, sex and socio-economic status (SES) estimated by parental education (CATSS) or the education level of the respondent (STAGE). In STAGE, the age when the web questionnaire was answered was also used in the analyses. To control for the effects of low IQ, we also used the validated A-TAC algorithm for learning disorder (in the validation studies defined as mental retardation diagnosed clinically on the basis of neuropsychological testing, generally by the Wechsler scales; Hansson *et al.* 2005; Larson *et al.* 2010).

Statistical analyses

Association between ALTs and co-morbidity

The association between autistic traits and psychiatric co-morbidity was examined using generalized estimating equation (GEE) regression models with the PROC GENMOD procedure in SAS 9.2 (SAS Institute Inc., USA). Odds ratios (ORs) with 95% confidence intervals (CIs) were first computed, and in a subsequent step adjusted for possible confounding by the covariates specified above. The GEE models account for correlations within twin pairs and are robust to deviations from a normal distribution. Cross-twin, cross-trait correlations were conducted with the PROC CORR procedure in SAS 9.2.

Genetic analyses

Twin modeling is based on the variance in traits and covariance between traits in monozygotic (MZ) and dizygotic (DZ) twins and is decomposed into genetic (A, the part of phenotypical variance that is attributable to genetic factors), shared environment (C, factors that make the twins similar) and unique environment/non-shared environment (E, factors that make the twins different) factors. As MZ twins share all their genes but DZ twins share only half of their segregating alleles, the factors attributable to A can be traced to the difference between MZ and DZ twins. For instance, if a phenotype is solely affected by genes, the twin-pair correlation for MZ twins would be 1.00 whereas the correlation for DZ twins would be 0.50. If, however, the phenotype is solely affected by C or E, a twin correlation of 1.00, or 0.00 for both MZ and DZ twins, would be expected. Most psychiatric disorders and traits are, however, affected by both genes and environment.

Intra-class correlations and standard continuous univariate analyses, with only same-sex MZ and DZ twins, were used to obtain heritability estimates. Cross-twin, cross-trait correlations were used to assess whether common genetic and environmental susceptibilities are important for the association between ALTs and ADHD or anxiety traits: the ALT score of twin 1 is correlated with the ADHD trait score in the co-twin. Higher correlations for MZ twins than for DZ twins indicate genetic covariation. We then used bivariate twin modeling, including only MZ twins and DZ same-sex twins. Eligible for the analyses were 6624 twins (3172 MZ and 3092 DZ) in CATSS and 8170 twins (4968 MZ and 3202 DZ same-sex) in STAGE. Because of power issues we used continuous measures in the bivariate models. We found mean differences between sexes for ALTs and ADHD traits (>0.001) in CATSS, and for ALTs, anxiety traits and depressive traits in STAGE (>0.001), when sex, SES and interview order were controlled for in a regression model. Thus, residual scores, controlling for sex, SES and interview order, were used in the twin analyses. We also found mean differences between zygosity for ALTs, ADHD traits, anxiety traits and conduct problems (>0.001) in CATSS. This was probably because of the sibling interaction effect (most probably due to parents unwittingly exaggerating the difference between DZ twins), as reported previously in another study using the CATSS sample (Ronald *et al.* 2010). To resolve this, a contrast model considering the sibling interaction was used for the model-fitting analyses in CATSS. As the relationship between ALTs and ADHD traits, anxiety traits, conduct problems or depressive traits in the general population might be reciprocal, a

correlated factors solution was chosen. As the substance abuse module was dichotomous, a liability threshold model using the same six categories as in the risk analyses was used. The twin analyses were performed in Mx (Neale *et al.* 2003). To avoid oversimplification and thus bias of the estimates of A, C and E, we did not try to fit reduced models (Sullivan & Eaves, 2002).

Results

Association between ALT and mental health problems

The ASD group showed a large increase in risk for ADHD (adjusted OR 100 in children, 69 among adults, Table 2). The risk for ADHD then decreased monotonically (a function that is consistently increasing and never decreasing or a function that is consistently decreasing and never increasing) by the ALT groups in both CATSS and STAGE. Furthermore, the risk of anxiety was 22 times greater in the ASD group in CATSS and 21 times greater in STAGE compared to subjects without ALTs. Again, the risk decreased monotonically with the number of ALTs. The risk for conduct problems in the ASD group was 35 times higher than in subjects without ALTs. The corresponding figures for depression and substance abuse were 12 and 6 respectively. Both adjusted and unadjusted ORs are given in Table 2. Adjusting the ORs left the magnitude of the associations almost unchanged, with the exception of the relationship between the two highest groups (ALT 5 and ASD) and ADHD, which was strongly confounded by the presence of learning disorder in both studies, as ORs adjusted only to zygosity, sex and SES remained similar to the unadjusted ORs; data not shown).

Heritability and genetic correlation

Table 3 presents the intra-class correlations for all mental health problems in CATSS and STAGE, all of which were higher for MZ than for DZ twins, indicating genetic influences. This was confirmed in the univariate analysis, where, for instance, ALTs in childhood showed a high heritability ($h^2=0.70$). Similar effects were detected among adults but with more modest heritability estimates ($h^2=0.32$). All cross-twin, cross-trait correlations were higher for MZ than for DZ twins, indicating genetic covariance (Table 3). The full statistics from the univariate and bivariate models between ALTs and the co-occurring conditions are given in the online Appendix (Supplementary Tables S1 and S2), and Fig. 1 provides an overview of the main results. The phenotypical correlation (i.e. the degree to which two traits covary

Table 2. Associations between ALTs and risk of ADHD, anxiety, depression, substance abuse and conduct problems in CATSS and STAGE

	CATSS			STAGE		
	<i>n</i>	Unadjusted OR (95% CI)	Adjusted OR (95% CI) ^a	<i>n</i>	Unadjusted OR (95% CI)	Adjusted OR (95% CI) ^a
ADHD						
ASD	156	194.1 (128.5–293.3)	99.7 (62.1–160.1)	79	78.5 (47.9–128.4)	68.9 (40.5–117.1)
ALT 5	88	70.8 (43.7–114.6)	40.6 (23.7–69.6)	128	37.2 (23.6–58.5)	35.9 (21.9–57.0)
ALT 4	190	38.4 (26.5–55.6)	25.4 (16.8–38.4)	263	22.4 (15.2–33.0)	21.3 (14.3–31.7)
ALT 3	497	12.2 (8.7–17.1)	8.3 (5.8–12.0)	1298	8.9 (6.7–11.9)	8.7 (6.45–11.8)
ALT 2	2335	5.2 (4.0–6.8)	4.4 (3.3–5.8)	2501	3.9 (2.9–5.2)	3.7 (2.7–5.1)
ALT 1 ^b	7494	1.0	1.0	12 378	1.0	1.0
Anxiety						
ASD	159	23.4 (16.5–33.2)	21.6 (14.7–31.7)	55	21.6 (12.1–36.8)	21.2 (11.2–39.7)
ALT 5	88	11.5 (7.1–18.3)	11.1 (6.7–18.4)	103	9.3 (5.7–15.1)	11.7 (7.0–19.3)
ALT 4	190	7.4 (4.9–11.1)	7.5 (4.9–11.3)	212	8.0 (5.5–11.4)	8.9 (6.0–12.9)
ALT 3	496	5.2 (3.9–6.9)	5.2 (3.9–6.9)	1059	4.1 (3.2–5.0)	4.3 (3.3–5.4)
ALT 2	2339	2.5 (2.0–3.0)	2.6 (2.1–3.3)	2111	2.4 (1.9–2.9)	2.4 (1.9–2.9)
ALT 1 ^b	7498	1.0	1.0	10 898	1.00	1.00
Conduct problems						
ASD	158	49.1 (27.6–87.2)	35.3 (18.3–68.3)			
ALT 5	87	48.2 (25.8–90.0)	36.7 (17.9–75.1)			
ALT 4	191	19.6 (10.7–36.1)	15.7 (8.2–30.3)			
ALT 3	496	10.6 (6.1–18.5)	9.2 (5.1–16.4)			
ALT 2	2339	2.9 (1.8–4.5)	2.7 (1.6–4.3)			
ALT 1 ^b	7497	1.0	1.0			
Depression						
ASD				76	9.4 (5.9–14.8)	11.7 (7.1–19.3)
ALT 5				131	3.8 (2.7–5.6)	4.7 (3.1–7.1)
ALT 4				259	3.1 (2.4–4.2)	3.4 (2.5–4.6)
ALT 3				1267	2.3 (2.1–2.7)	2.4 (2.1–2.8)
ALT 2				2458	1.8 (1.6–2.1)	1.8 (1.6–2.0)
ALT 1 ^b				12 163	1.0	1.0
Substance abuse						
ASD				68	7.4 (3.5–15.7)	5.7 (2.4–13.3)
ALT 5				111	8.2 (1.5–2.7)	7.9 (4.4–14.2)
ALT 4				236	4.1 (2.4–6.7)	3.0 (1.6–5.5)
ALT 3				1128	3.7 (2.8–4.9)	3.4 (2.5–4.6)
ALT 2				2200	1.9 (1.5–2.5)	1.9 (1.4–2.5)
ALT 1 ^b				10 927	1.0	1.0

CATSS, Child and Adolescent Twin Study in Sweden; STAGE, Study of Twin Adults: Genes and Environment; ALT, autistic-like trait; ADHD, attention deficit hyperactivity disorder; ASD, autism spectrum disorder; OR, odds ratio; CI, confidence interval.

In CATSS: ALT 1 = 0 points, ALT 2 = 0.5–1 point, ALT 3 = 1.5–2 points, ALT 4 = 2.5–3 points, ALT 5 = 3.5–4 points and ASD \geq 4.5 points.

In STAGE: ALT 1 = 0–2 points, ALT 2 = 2.5–3 points, ALT 3 = 3.5–4.5 points, ALT 4 = 5–5.5 points, ALT 5 = 6–6.5 points and ASD \geq 7 points.

^a Adjusted for interview order, zygosity, age, sex, socio-economic status (SES) and, in CATSS, learning disorder.

^b Reference category.

in the population) between ALTs and ADHD traits was 0.55 in CATSS and 0.44 in STAGE. Of these correlations, 71% in CATSS and 54% in STAGE were

accounted for by common genetic effects. The phenotypic correlations were more modest for ALTs and anxiety traits in both CATSS and STAGE whereas

Table 3. Intra-class correlations, within-individual (phenotypic) correlations and cross-twin, cross-trait correlations

	Intra-class correlation				Within-individual (phenotypic) correlation				Cross-twin, cross-trait correlation			
	CATSS		STAGE		CATSS		STAGE		CATSS		STAGE	
	MZ	DZ	MZ	DZ	MZ	DZ	MZ	DZ	MZ	DZ	MZ	DZ
ALTs	0.77 (0.74–0.81)	0.38 (0.32–0.45)	0.38 (0.34–0.41)	0.16 (0.10–0.21)					0.46	0.21	0.22	0.11
ADHD traits	0.62 (0.59–0.66)	0.19 (0.15–0.25)	0.44 (0.41–0.48)	0.20 (0.15–0.25)	0.55		0.46		0.29	0.13	0.11	0.05
Anxiety traits	0.72 (0.64–0.81)	0.38 (0.25–0.42)	0.37 (0.26–0.49)	0.16 (–0.02 to 0.34)	0.33		0.17		0.32	0.21		
Conduct problems	0.77 (0.71–0.83)	0.48 (0.38–0.58)			0.36							
Depressive traits			0.38 (0.35–0.42)	0.17 (0.12–0.22)			0.31				0.18	0.07
Substance abuse			0.66 (0.53–0.80)	0.41 (0.13–0.69)			0.09				0.05	0.02

CATSS, Child and Adolescent Twin Study in Sweden; STAGE, Study of Twin Adults: Genes and Environment; ALT, autistic-like trait; ADHD, attention deficit hyperactivity disorder; MZ, monozygotic; DZ, dizygotic.

common genetic effects again accounted for 92% and 55%. For ALTs and depressive traits, roughly half of the correlation was accounted for by common non-shared environmental effects. Sixty-eight per cent of the correlation between ALTs and substance abuse was accounted for by common genetic effects, and 81% of the correlation between ALTs and conduct problems.

Discussion

The findings presented here add to what is known about ALTs in the general population in four respects. First, even if increasing symptoms of ALTs monotonically increase the risk for ADHD, anxiety, conduct problems, depression, and substance and alcohol abuse, the group with only one or two ALTs also had considerably increased risks for all other types of mental disorders compared to the majority of individuals who do not have such traits. Second, consistent with previous studies in children (Ronald *et al.* 2008; Hallett *et al.* 2009), we found evidence for substantial genetic influences on the correlations between ALTs and ADHD also in adulthood. Third, the phenotypic correlation between ALTs, depressive and anxiety traits in adults was almost as strongly affected by genetic as by non-shared environmental factors. Fourth, the largest part of the phenotypic correlation between ALTs and conduct problems and ALTs and substance abuse was affected by genes.

The study has several strengths. Both samples are nationwide birth cohorts, representative of children and adults. Although psychiatric symptoms were assessed by different informants, methods and instruments in the two samples (parental interviews *versus* self-report; telephone interviews and web questionnaires), overall the results were consistent.

The results also have to be interpreted in the light of important limitations. Asking adults direct questions based on DSM-IV criteria has not been formally validated as a measure of any of the targeted problem types, including ALTs, but has been used previously in epidemiological studies as a proxy for clinical instruments (Oliveira *et al.* 2007; Frisell *et al.* 2010). Until validated self-report measures for ALTs in adults are available for epidemiological studies, caution is warranted when interpreting the results. Socio-communicative restrictions may be less suited for self-rate assessment than other types of problems, and a core deficit in ASDs is the lack of understanding of emotions. For this reason, however, self-rate questionnaires for neuropsychiatric problems, just as for personality traits, are based on assessments of behavior patterns (such as avoiding to engage in small talk) rather than of emotional states (such as feeling left




Fig. 1. Phenotypical correlations with autistic-like traits (ALTs), and the portions of correlations accounted for by genetic and environmental effects. Unique environment means unique to each twin but common to ALTs and the other types of mental problems. CATSS, Child and Adolescent Twin Study in Sweden; STAGE, Study of Twin Adults: Genes and Environment; ADHD, attention deficit hyperactivity disorder.

behind). Several studies have nevertheless shown valid results from self-rating instruments assessing ALTs among individuals with known ASDs (e.g. Baron-Cohen *et al.* 2001). The cut-offs used for anxiety and depression have not been formally validated, but our prevalence estimates (4.5% for anxiety and 13.4% for depression) are compatible with previous estimates of the general population prevalence (Doris *et al.* 1999; Tyrer & Baldwin, 2006). STAGE has a comparatively low response rate compared to CATSS, which may explain the lower ASD prevalence and the lower prevalence of males in STAGE than in CATSS. Thus, the adult results might not be generalizable to the most extreme cases because non-responders might be more likely to have some type of problem. As ASDs are defined as developmental disorders, epidemiological studies without access to developmental assessments are, by definition, suboptimal. At the same time it is probably not feasible to conduct developmental assessments of rare conditions in a population-based cohort. It should also be kept in mind that the number of possible confounding factors was uncontrolled for or included by crude measures; IQ, for instance, was reflected merely by the presence of learning disability, and systematic differences in measures and sources of information could give rise to some of the noted differences between children and adults. Finally, it was not possible to formally test whether the ALTs reported here are really ‘autistic’ or just similar phenotypic expressions with different etiology. In our opinion, however, there are more arguments for than against a continuity from ASDs to ALTs created by underlying dimensionally distributed traits or abilities. First, ALTs are assessed by the same DSM-IV symptom criteria used for autistic disorder, making it likely that at least the phenotypical expression

corresponding to a specific item is similar. Second, pathogenetic processes that affect the risk for ASDs have also been shown to influence ALTs (e.g. increased paternal age, Lundström *et al.* 2010; genetic aberrations on 16p11, St Pourcain *et al.* 2010). Third, studies of relatives of probands with ASDs consistently show marked elevation of ALTs regardless of whether they are conceptually defined as ‘broader phenotypes’ (Folstein & Rutter, 1977) or traits (Szatmari *et al.* 2000). These cautionary notes should be kept in mind in the following discussion on the implications of our findings.

It has been suggested that subthreshold ALTs might entail cognitive features useful for careers in, for instance, science or engineering (Baron-Cohen *et al.* 2001; Focquaert *et al.* 2007). Our results do not contradict such possible advantages but demonstrate that, even if ‘systemizing’ abilities may be an asset in some respects, there is still an increased risk of mental health problems, such as ADHD, anxiety, and depression, substance abuse and conduct disorder. Furthermore, the presence of ALTs in subgroups of individuals with various forms of psychopathology may influence the response to pharmacological or psychotherapeutic interventions, as shown, for example, in anorexia nervosa, where the presence of ALTs was associated with a poorer psychosocial outcome (Wentz *et al.* 2009).

Clinical descriptions of Asperger’s syndrome and other types of autism-related personality profiles have long emphasized sensitivity to stress and social interaction as a basis for the development of concomitant anxiety and depression (Wing, 1981). Our results indicate that, in addition to such reactions, there are common underlying genetic susceptibilities. The observed phenotypic correlation between ALTs and traits related to ADHD and anxiety in children is in

line with previous findings (Ronald *et al.* 2008; Hallett *et al.* 2009), although our results indicate a stronger genetic component for the overlap with anxiety. This may be due to our larger sample size, but the findings are in need of corroboration. Moreover, our results suggest a modest genetic overlap between ALTs and ADHD traits among adults and between ALTs and anxiety and depressive traits, which might be expected, considering the well-known association between anxiety and depression (Kendler *et al.* 2007). The phenotypical correlation observed between ALTs and conduct problems is not surprising considering that social communication deficits are a common feature in children with conduct disorder (Gilmour *et al.* 2004; Donno *et al.* 2010). The large CIs observed for the genetic and non-shared correlations between ALTs and substance abuse preclude firm conclusions on the etiology of this particular association. However, substance abuse has previously been reported in combination with ASDs (e.g. Sizoo *et al.* 2010; Ståhlberg *et al.* 2010).

The differences between the heritability estimates and the phenotypical correlations in CATSS and in STAGE may reflect age-related effects. For instance, an increasing number of stressful life events increase the risk for anxiety disorders (Blazer *et al.* 1987), which may deflate the phenotypical correlation and increase the non-shared environmental factors. The heritability of ADHD has indeed been found to be substantially lower in adulthood than in childhood in other studies (Boomsma *et al.* 2010), indicating that unique environmental influences act in concert with the genotype. Of note, the proportion of unique environmental factors operating across ALTs and ADHD traits and across ALTs and anxiety traits was higher in adults than in children. On a speculative note, it may be argued that children experience a more homogeneous environment where genetics may play a greater role, whereas adults are exposed to a more versatile environment where environmental hazards that may affect both ALTs and ADHD traits and anxiety traits are more prevalent.

Recently, molecular genetic studies have suggested similarities in the etiologies of ASDs and ADHD. For example, a study of ADHD by Williams *et al.* (2010) detected increased rare copy number variants at locus 16p13.11, an area where microdeletions and microduplications have been reported in association with autism (Weiss *et al.* 2008). Our results call for attention to a continuum of traits related to mental disorders that are less specific than previously thought in creating susceptibility for mental disorders. As suggested by Plomin *et al.* (2009), this will have implications on genome-wide association studies, where it may be more fruitful to use continuous measures to study the

whole range of variation so as to explore genetic overlaps between quantitative traits and also polygenic factors behind these traits.

The accumulated evidence thus suggests a dimensional association between ALTs and ADHD, conduct problems, substance abuse, anxiety and depression. Such dimensional associations will challenge future versions of the DSM and ICD systems as it is clear that symptomatic and etiological commonalities exist behind disorders treated as distinct in diagnostics and that these etiological factors influence phenotypes both above and below diagnostic thresholds. In the future, clinicians need to be aware of the possibility of co-existing and complicating ALTs when assessing or treating patients with conduct problems, substance abuse, ADHD, anxiety and depression, and in clinical contacts with patients who are relatives of individuals with ASDs and thereby may be at an elevated risk for ALTs complicating other, more classic, mental disorders. New studies addressing whether the continuum of ALTs is etiologically similar or different from the ASDs are emerging (Ronald *et al.* 2010) and will influence whether to clinically define caseness from the normal variation by symptoms or by other variables, such as global mental or specific cognitive functioning.

Note

Supplementary material accompanies this paper on the Journal's website (<http://journals.cambridge.org/psm>).

Acknowledgments

STAGE was supported by CA085739 from the U.S. National Cancer Institute of the National Institutes of Health, and by CA118412. CATSS was supported by the Swedish Council for Working Life, the Research Council of the Swedish National Alcohol Monopoly, funds under the ALF agreement, and the Swedish Research Council.

Declaration of Interest

None.

References

- Anckarsäter H, Nilsson T, Saury JM, Rastam M, Gillberg C (2008). Autism spectrum disorders in institutionalized subjects. *Nordic Journal of Psychiatry* **62**, 160–167.
- APA (1994). *Diagnostic and Statistical Manual of Mental Disorders: DSM-IV*, 4th edn. American Psychiatric Association: Washington, DC.

- Baird G, Simonoff E, Pickles A, Chandler S, Loucas T, Meldrum D, Charman T** (2006). Prevalence of disorders of the autism spectrum in a population cohort of children in South Thames: the Special Needs and Autism Project (SNAP). *Lancet* **368**, 210–215.
- Baron-Cohen S, Wheelwright S, Skinner R, Martin J, Clubley E** (2001). The autism-spectrum quotient (AQ): evidence from Asperger syndrome/high-functioning autism, males and females, scientists and mathematicians. *Journal of Autism and Developmental Disorders* **31**, 5–17.
- Blazer D, Hughes D, George LK** (1987). Stressful life events and the onset of a generalized anxiety syndrome. *American Journal of Psychiatry* **144**, 1178–1183.
- Boomsma DI, Saviouk V, Hottenga JJ, Distel MA, de Moor MH, Vink JM, Geels LM, van Beek, JH, Bartels M, de Geus EJ, Willemsen G** (2010). Genetic epidemiology of attention deficit hyperactivity disorder (ADHD index) in adults. *PLoS One* **5**, e10621.
- Carpenter JS, Andrykowski MA, Wilson J, Hall LA, Rayens MK, Sachs B, Cunningham LLC** (1998). Psychometrics for two short forms of the Center for Epidemiologic Studies-Depression Scale. *Issues in Mental Health Nursing* **19**, 481–494.
- Constantino JN, Todd RD** (2003). Autistic traits in the general population: a twin study. *Archives of General Psychiatry* **60**, 524–530.
- Donno R, Parker G, Gilmour J, Skuse DH** (2010). Social communication deficits in disruptive primary-school children. *British Journal of Psychiatry* **196**, 282–298.
- Doris A, Ebmeier K, Shajahan P** (1999). Depressive illness. *Lancet* **354**, 1369–1375.
- Focquaert F, Steven MS, Wolford GL, Colden A, Gazzaniga MS** (2007). Empathizing and systemizing cognitive traits in the sciences and humanities. *Personality and Individual Differences* **43**, 619–625.
- Folstein S, Rutter M** (1977). Infantile autism: a genetic study of 21 twin pairs. *Journal of Child Psychology and Psychiatry* **18**, 297–321.
- Frisell T, Lichtenstein P, Rahman Q, Langstrom N** (2010). Psychiatric morbidity associated with same-sex sexual behaviour: influence of minority stress and familial factors. *Psychological Medicine* **40**, 315–324.
- Furberg H, Lichtenstein P, Pedersen NL, Thornton L, Bulik CM, Lerman C, Sullivan PF** (2008). The STAGE cohort: a prospective study of tobacco use among Swedish twins. *Nicotine and Tobacco Research* **10**, 1727–1735.
- Ghaziuddin M, Greden J** (1998). Depression in children with autism/pervasive developmental disorders: a case-control family history study. *Journal of Autism and Developmental Disorders* **28**, 111–115.
- Gillberg C** (1992). The Emanuel Miller Memorial Lecture 1991. *Autism and autistic-like conditions: subclasses among disorders of empathy*. *Journal of Child Psychology and Psychiatry* **33**, 813–842.
- Gillberg C, Cederlund M, Lamberg K, Zeijlon L** (2006). Brief report: 'the autism epidemic'. The registered prevalence of autism in a Swedish urban area. *Journal of Autism and Developmental Disorders* **36**, 429–435.
- Gilmour J, Hill B, Place M, Skuse DH** (2004). Social communication deficits in conduct disorder: a clinical and community survey. *Journal of Child Psychology and Psychiatry* **45**, 967–978.
- Halleröd SL, Larson T, Ståhlberg O, Carlström E, Gillberg C, Anckarsäter H, Råstam M, Lichtenstein P, Gillberg C** (2010). The Autism – Tics, ADHD and other Comorbidities (A-TAC) telephone interview: convergence with the Child Behavior Checklist (CBCL). *Nordic Journal of Psychiatry* **64**, 218–224.
- Hallett V, Ronald A, Happe F** (2009). Investigating the association between autistic-like and internalizing traits in a community-based twin sample. *Journal of the American Academy of Child and Adolescent Psychiatry* **48**, 618–627.
- Hansson SL, Svanstrom Rojvall A, Rastam M, Gillberg C, Anckarsäter H** (2005). Psychiatric telephone interview with parents for screening of childhood autism – tics, attention-deficit hyperactivity disorder and other comorbidities (A-TAC): preliminary reliability and validity. *British Journal of Psychiatry* **187**, 262–267.
- Hoekstra RA, Bartels M, Hudziak JJ, Van Beijsterveldt TC, Boomsma DI** (2007). Genetic and environmental covariation between autistic traits and behavioral problems. *Twin Research and Human Genetics* **10**, 853–860.
- Hofvander B, Delorme R, Chaste P, Nyden A, Wentz E, Stahlberg O, Herbrecht E, Stopin A, Anckarsäter H, Gillberg C, Råstam M, Leboyer M** (2009). Psychiatric and psychosocial problems in adults with normal-intelligence autism spectrum disorders. *BMC Psychiatry* **9**, 35.
- Kadesjö B, Gillberg C, Hagberg B** (1999). Brief report: autism and Asperger syndrome in seven-year-old children: a total population study. *Journal of Autism and Developmental Disorders* **29**, 327–331.
- Kanne SM, Christ SE, Reiersen AM** (2009). Psychiatric symptoms and psychosocial difficulties in young adults with autistic traits. *Journal of Autism and Developmental Disorders* **39**, 827–833.
- Kendler KS, Gardner CO, Gatz M, Pedersen NL** (2007). The sources of co-morbidity between major depression and generalized anxiety disorder in a Swedish national twin sample. *Psychological Medicine* **37**, 453–462.
- Larson T, Anckarsäter H, Gillberg C, Stahlberg O, Carlstrom E, Kadesjo B, Råstam M, Lichtenstein P, Gillberg C** (2010). The Autism – Tics, ADHD and other Comorbidities inventory (A-TAC): further validation of a telephone interview for epidemiological research. *BMC Psychiatry* **10**, 1–11.
- Lee DO, Ousley OY** (2006). Attention-deficit hyperactivity disorder symptoms in a clinic sample of children and adolescents with pervasive developmental disorders. *Journal of Child and Adolescent Psychopharmacology* **16**, 737–746.
- Lichtenstein P, Carlstrom E, Rastam M, Gillberg C, Anckarsäter H** (2010). The genetics of autism spectrum disorders and related neuropsychiatric disorders in childhood. *American Journal of Psychiatry*. Published online: 4 August 2010. doi:10.1176/appi.ajp.2010.10020223.
- Lichtenstein P, De Faire U, Floderus B, Svartengren M, Svedberg P, Pedersen NL** (2002). The Swedish Twin Registry: a unique resource for clinical, epidemiological and genetic studies. *Journal of Internal Medicine* **252**, 184–205.

- Lundström S, Haworth CM, Carlström E, Gillberg C, Mill J, Råstam M, Hultman CM, Ronald A, Anckarsäter H, Plomin R, Lichtenstein P, Reichenberg A (2010). Trajectories leading to autism spectrum disorders are affected by paternal age: findings from two nationally representative twin studies. *Journal of Child Psychology and Psychiatry, and Allied Disciplines* **51**, 850–856.
- Neale MC, Boker SM, Xie G, Maes HH (2003). *Mx Statistical Modeling*, 6th edn. VCU Box 900126, Richmond, VA 23298, Department of Psychiatry.
- Oliveira G, Ataíde A, Marques C, Miguel TS, Coutinho AM, Mota-Vieira L, Gonçalves E, Lopes NM, Rodrigues V, Carmona da Mota H, Vicente AM (2007). Epidemiology of autism spectrum disorder in Portugal: prevalence, clinical characterization, and medical conditions. *Developmental Medicine and Child Neurology* **49**, 726–733.
- Pine DS, Guyer AE, Goldwin M, Towbin KA, Leibenluft E (2008). Autism spectrum disorder scale scores in pediatric mood and anxiety disorders. *Journal of the American Academy of Child and Adolescent Psychiatry* **47**, 652–661.
- Plomin R, Haworth CM, Davis OS (2009). Common disorders are quantitative traits. *Nature Reviews Genetics* **10**, 872–878.
- Reiersen AM, Constantino JN, Grimmer M, Martin NG, Todd RD (2008). Evidence for shared genetic influences on self-reported ADHD and autistic symptoms in young adult Australian twins. *Twin Research and Human Genetics* **11**, 579–585.
- Rommelse NN, Altink ME, Fliers EA, Martin NC, Buschgens CJ, Hartman CA, Buitelaar JK, Faraone SV, Sergeant JA, Oosterlaan J (2009). Comorbid problems in ADHD: degree of association, shared endophenotypes, and formation of distinct subtypes. Implications for a future DSM. *Journal of Abnormal Child Psychology* **37**, 793–804.
- Ronald A, Edelson LR, Asherson P, Saudino KJ (2010). Exploring the relationship between autistic-like traits and ADHD behaviors in early childhood: findings from a community twin study of 2-year-olds. *Journal of Abnormal Child Psychology* **38**, 185–196.
- Ronald A, Happé F, Bolton P, Butcher LM, Price TS, Wheelwright S, Baron-Cohen S, Plomin R (2006). Genetic heterogeneity between the three components of the autism spectrum: a twin study. *Journal of the American Academy of Child and Adolescent Psychiatry* **45**, 691–699.
- Ronald A, Simonoff E, Kuntsi J, Asherson P, Plomin R (2008). Evidence for overlapping genetic influences on autistic and ADHD behaviours in a community twin sample. *Journal of Child Psychology and Psychiatry, and Allied Disciplines* **49**, 535–542.
- Roohi J, DeVincent CJ, Hatchwell E, Gadow KD (2009). Association of a monoamine oxidase-a gene promoter polymorphism with ADHD and anxiety in boys with autism spectrum disorder. *Journal of Autism and Developmental Disorders* **39**, 67–74.
- Simonoff E, Pickles A, Charman T, Chandler S, Loucas T, Baird G (2008). Psychiatric disorders in children with autism spectrum disorders: prevalence, comorbidity, and associated factors in a population-derived sample. *Journal of the American Academy of Child and Adolescent Psychiatry* **47**, 921–929.
- Sizoo B, van den Brink W, Koeter M, Gorissen van Eenige M, van Wijngaarden-Cremers P, van der Gaag RJ (2010). Treatment seeking adults with autism or ADHD and co-morbid substance use disorder: prevalence, risk factors and functional disability. *Drug and Alcohol Dependence* **107**, 44–50.
- Ståhlberg O, Anckarsäter H, Nilsson T (2010). Mental health problems in youths committed to juvenile institutions: prevalences and treatment needs. *European Child and Adolescent Psychiatry* **19**, 893–903.
- St Pourcain B, Wang K, Glessner JT, Golding J, Steer C, Ring SM, Skuse DH, Grant SF, Hakonarson H, Davey Smith G (2010). Association between a high-risk autism locus on 5p14 and social communication spectrum phenotypes in the general population. *American Journal of Psychiatry* **167**, 1364–1372.
- Sullivan PF, Eaves LJ (2002). Evaluation of analyses of univariate discrete twin data. *Behavior Genetics* **32**, 221–227.
- Szatmari P, MacLean JE, Jones MB, Bryson SE, Zwaigenbaum L, Bartolucci G, Mahoney WJ, Tuff L (2000). The familial aggregation of the lesser variant in biological and nonbiological relatives of PDD probands: a family history study. *Journal of Child Psychology and Psychiatry, and Allied Disciplines* **41**, 579–586.
- Tyrer P, Baldwin D (2006). Generalised anxiety disorder. *Lancet* **368**, 2156–2166.
- Weiss LA, Shen Y, Korn JM, Arking DE, Miller DT, Fossdal R, Saemundsen E, Stefansson H, Ferreira MA, Green T, Platt OS, Ruderfer DM, Walsh CA, Altshuler D, Chakravarti A, Tanzi RE, Stefansson K, Santangelo SL, Gusella JF, Sklar P, Wu BL, Daly MJ; Autism Consortium (2008). Association between microdeletion and microduplication at 16p11.2 and autism. *New England Journal of Medicine* **358**, 667–675.
- Wentz E, Gillberg IC, Anckarsäter H, Gillberg C, Råstam M (2009). Reproduction and offspring status 18 years after teenage-onset anorexia nervosa – a controlled community-based study. *International Journal of Eating Disorders* **42**, 483–491.
- Williams NM, Zaharieva I, Martin A, Langley K, Mantripragada K, Fossdal R, Stefansson H, Stefansson K, Magnusson P, Gudmundsson OO, Gustafsson O, Holmans P, Owen MJ, O'Donovan M, Thapar A (2010). Rare chromosomal deletions and duplications in attention-deficit hyperactivity disorder: a genome-wide analysis. *Lancet* **376**, 1401–1408.
- Wing L (1981). Asperger's syndrome: a clinical account. *Psychological Medicine* **11**, 115–129.