



LUND UNIVERSITY

The low-energy spectrum of ^{29}Si and the existence of large oblate distortions associated with certain intrinsic orbitals

Ragnarsson, Ingemar; Nilsson, Sven Gösta

Published in:
Nuclear Physics, Section A

1970

[Link to publication](#)

Citation for published version (APA):
Ragnarsson, I., & Nilsson, S. G. (1970). The low-energy spectrum of ^{29}Si and the existence of large oblate distortions associated with certain intrinsic orbitals. *Nuclear Physics, Section A*, 158(1), 155-160.

Total number of authors:
2

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

THE LOW-ENERGY SPECTRUM OF ^{29}Si AND THE EXISTENCE OF LARGE OBLATE DISTORTIONS ASSOCIATED WITH CERTAIN INTRINSIC ORBITALS

INGEMAR RAGNARSSON and SVEN GÖSTA NILSSON

Department of Mathematical Physics, Lund Institute of Technology, Lund, Sweden

Received 26 June 1970

Abstract: Equilibrium calculations involving different odd-particle excitations for ^{29}Si and neighbouring nuclei are reported. Minima corresponding to very large oblate distortions are confirmed to be associated with a recently found rotational band in ^{29}Si as suggested in ref. ¹).

1. Introduction

In a recent publication ¹) experimental evidence was given for a very strongly oblate rotational band based on the orbital $[303 \frac{7}{2}]$. The existence of strong oblate distortions for at least some intrinsic configurations of ^{29}Si is reasonable in view of the fact that the ground state band of ^{28}Si is associated with a strong intrinsic oblate quadrupole moment ^{2,3}). An analysis of the ^{29}Si spectrum was made by Bromley *et al.* ⁴) in 1957 suggesting oblate shapes for the lowest states of that nucleus. Renewed theoretical studies of the total potential-energy surface based on a generalized harmonic-oscillator potential ⁵) and a subsequent normalization of the computed total energy (so as on the average to reproduce the liquid-drop behaviour) by the Strutinsky shell correction method ⁶) is presently being undertaken for large regions of the periodic table. One aim of this study is to find the A and Z dependence of the nuclear potential parameters with important application to the problem of the superheavy elements ⁷). The theoretical equilibrium configurations are found to explain remarkably well several dominant features of the ^{29}Si spectrum.

2. Single-particle energies and the total nuclear energy

The following nuclear potential ⁵), involving essentially quadrupole (ϵ) and hexadecapole (ϵ_4) degrees of freedom has been considered (particularly in the beginning of the $A = 16$ –40 region the latter degree of freedom is of great importance)

$$V = \frac{1}{2} M \omega_0^2 (\epsilon, \epsilon_4) (1 - \frac{2}{3} \epsilon P_2 + 2 \epsilon_4 P_4) - \kappa \hbar \omega_0 [2 I_t \cdot s + \mu (I_t^2 - \langle I_t^2 \rangle_N)].$$

Fysik- & astronomibiblioteket
Lunds universitet

The parameters κ and μ are assumed to vary with A in a simple way over the periodic

table. For the $A \approx 25$ region we have (as previously) assumed ⁸⁾ $\kappa_n = \kappa_p = -0.08$, $\mu_n = \mu_p = 0$. We have furthermore assumed that in this particular region pairing is negligible ($G_n = G_p = 0$). The Coulomb energy is calculated by a numerical method due to Nix, as in ref. ⁷⁾. The total nuclear energy is subsequently evaluated as a function of ϵ and ϵ_4 by employment of the Strutinsky shell correction method ⁶⁾. In the




Fig. 1. The potential energy surface in the (ϵ, ϵ_4) plane for $^{28}_{14}\text{Si}$.


odd case the computer program places the odd particle in the orbital of specific Ω and parity that is available nearest to the Fermi surface.

The $^{28}_{14}\text{Si}$ potential energy surface for the 0^+ state is exhibited in fig. 1. Note the strongly oblate minimum corresponding to $\epsilon = -0.43$, $\epsilon_4 = 0.00$, which under the assumption of homogeneous charge correspond to an intrinsic quadrupole moment $Q_0 = -54 \text{ fm}^2$. The apparent secondary prolate minimum situated roughly 2 MeV over the ground state oblate minimum may or may not be a proper minimum as the barrier in the gamma plane as encountered in the calculations by Cusson ⁹⁾ may not be sufficient to sustain the zero-point vibration.

3. Discussion of the ^{29}Si spectrum

Let us now consider the theoretical results for the ^{29}Si case, see figs. 2 and 3. (Actually fig. 3 refers specifically to ^{29}P whose potential energy surface differs from that of its mirror nucleus only with respect to the Coulomb term, whose difference in this case is negligible.) In fig. 2 the potential energy surface corresponding to $\Omega = \frac{7}{2}$ is illustrated. It is found to exhibit only one well-developed minimum corresponding to $\epsilon = -0.49$, $\epsilon_4 = 0.00$. In fig. 3 we give cuts through the two-dimensional energy surfaces corresponding to specified Ω and parity in such a way that for each ϵ -value we let ϵ_4 assume the value that minimizes the energy. In this way fig. 3 is obtained,

assumed ⁸) $\kappa_n = \kappa_p = -0.08$,
 this particular region pairing is
 calculated by a numerical method
 subsequently evaluated as a func-
 tion correction method ⁶). In the



plane for ²⁸Si.

in the orbital of specific Ω

exhibited in fig. 1. Note the
 $\epsilon_4 = 0.00$, which under the
 intrinsic quadrupole moment
 situated roughly 2 MeV
 be a proper minimum as the
 calculations by Cusson ⁹) may not

m

case, see figs. 2 and 3. (Actu-
 surface differs from that of
 m, whose difference in this
 corresponding to $\Omega = \frac{7}{2}$ is
 d minimum corresponding
 the two-dimensional energy
 a way that for each ϵ -value
 this way fig. 3 is obtained,




Fig. 2. Same as fig. 1 valid for ²⁹Si with the odd particle placed in an orbital with $\Omega = \frac{7}{2}$ and negative parity.




Fig. 3. The total energy for ²⁹P as a function of distortion for the odd particle associated with alternative orbitals of given Ω and π . Except for the difference in Coulomb energy this diagram should be equal to that of ²⁹Si. The diagram represents cuts through surfaces of the type given in fig. 2. For each ϵ the lowest energy with respect to ϵ_4 is plotted.

representing the lowest configurations of given Ω and π for each distortion. From this figure one should expect an oblate $\frac{1}{2}^+$ configuration ($\varepsilon = -0.34$, $\varepsilon_4 = 0.01$) and an oblate ($\varepsilon = -0.39$, $\varepsilon_4 = 0.02$) $\frac{3}{2}^+$ configuration at about the same energy and a $\frac{5}{2}^+$ prolate one ($\varepsilon = 0.35$, $\varepsilon_4 = 0.11$) about 2 MeV higher. The strongly deformed oblate $[303 \frac{7}{2}]$ configuration (corresponding to $\varepsilon = -0.49$, $\varepsilon_4 = 0.00$) occurs theo-




Fig. 4. Experimental and theoretical intrinsic spectra of ^{29}Si .

retically at about 2.5 MeV of excitation. The next excited oblate odd state is $[301 \frac{3}{2}]$ with $\varepsilon = -0.48$, $\varepsilon_4 = -0.01$ predicted about 2 MeV higher in energy. The low-lying even-parity levels have earlier been identified by Bromley *et al.* ⁴⁾ as associated with weakly oblate ($\varepsilon = -0.15$), Coriolis coupled orbitals of $\Omega = \frac{1}{2}^+$ and $\Omega = \frac{3}{2}^+$. This interpretation is roughly in agreement with the theoretical results mentioned above apart from the larger theoretical distortions (the energy surface is, however, relatively flat and no pairing is included). The $[202 \frac{5}{2}^+]$ orbital is probably involved at some place in the spectrum above a few MeV of excitation. As to other higher excited even-parity states we are aware that other degrees of freedom, not directly represented by minima in these potential energy surfaces, are expected to occur, e.g. in the form of particle-hole excitations.

(On the other hand when all these kinds of excitations are introduced, there is, of course, a clear redundancy of states, as for each distortion there exists a complete set of states once all particle-hole excitations are considered. Therefore, for each Ω and parity, usually only the lowest-lying excitations can be reliably obtained by the method employed here.) The emphasis in this discussion is placed on the odd-parity states

and π for each distortion. From
ation ($\varepsilon = -0.34$, $\varepsilon_4 = 0.01$) and
at about the same energy and a
V higher. The strongly deformed
 $\varepsilon = -0.49$, $\varepsilon_4 = 0.00$) occurs theo-

EXPERIMENTAL



c spectra of ²⁹Si.

ited oblate odd state is $[301 \frac{3}{2}]$
higher in energy. The low-lying
ley *et al.* ⁴⁾ as associated with
of $\Omega = \frac{1}{2}^+$ and $\Omega = \frac{3}{2}^+$. This
tical results mentioned above
surface is, however, relatively
is probably involved at some
. As to other higher excited
dom, not directly represented
ted to occur, e.g. in the form

as are introduced, there is, of
on there exists a complete set
d. Therefore, for each Ω and
ably obtained by the method
ced on the odd-parity states

observed in ref. ¹⁾. In view of the discussion above, the assignment of the odd-parity states should be comparatively reliable, as the minima are very deep both for the $[303 \frac{7}{2}]$ state and the $[301 \frac{3}{2}]$ state, and as they indeed are the lowest odd-parity excitations of the respective Ω and parity. Both these orbitals originate from the spherical $f_{7/2}$ orbital. One may note that both theoretically and experimentally the orbital associated with the $\frac{5}{2}^-$ orbital occurs higher in energy than the $\frac{3}{2}^-$ orbital. This reordering of the $\frac{7}{2}, \frac{5}{2}, \frac{3}{2}, \frac{1}{2}$ sequence is a non-diagonal effect of the P_2 deformation. The experimental and theoretical intrinsic spectra are given in fig. 4.

Two members, $I = \frac{7}{2}$ and $\frac{5}{2}$, of the $\frac{7}{2}^-$ band are observed ¹⁾, the band head at 3.62 MeV, or about 1 MeV higher than theoretically expected. The discrepancy is even smaller for the $\frac{3}{2}^-$ excitation which experimentally occurs at 4.94 MeV. The E2 matrix element for the interband transition of the $\frac{7}{2}^-$ band corresponds to an intrinsic quadrupole moment of $|Q_0| = 66 \pm 9 \text{ fm}^2$. The theoretical equilibrium distortions would correspond to a $Q_0 = -62 \text{ fm}^2$, in excellent agreement with experiments. There remains a problem of the empirical identification of the $[330 \frac{1}{2}]$ rotational band, which is seen at an excitation of 3.09 MeV in ²⁵Al. According to the calculations the $[303 \frac{7}{2}]$ orbital is never found to occur as a ground state in this region of nuclei.

It is tempting to conjecture that similar oblate configurations associated with $[404 \frac{9}{2}]$ may be expected to occur for Z and N about 28, provided the single-particle preference for oblate distortions is strong enough to overcome the prolate tendency associated with the $N = 28$ shell ⁹⁾. No theoretical calculations corresponding to the ones presented here are presently available, however, in the relevant regions of nuclei. Results pertaining to the entire $A = 16-40$ region as obtained by the method described will be published by Ragnarsson.

The comparison of experimental and theoretical level energies (see fig. 4), shows that the agreement is much less impressive than in heavier regions exhibiting stable equilibrium distortions. Some of the partial failure in this light region is probably due to the fact that the neglected gamma-degree of freedom here plays a role important enough to invert the predicted level order. In addition the single-particle orbital occupied by the odd particle may by itself radically effect the equilibrium shape, as is exemplified by the present case. This leads to some orthogonality problems for the equilibrium configurations calculated, as was pointed out above.

Hartree-Fock calculations in fairly large numbers are available for the ²⁸Si case, indicating oblate equilibrium shapes for that nucleus (for a review and further references see Ripka ¹⁰⁾). Unfortunately, for the odd ²⁹Si case no such calculations are available for comparison.

We are grateful to Profs. B. R. Mottelson and R. Y. Cusson for valuable comments on the manuscript. Dr. Cusson also kindly communicated to us results of his unpublished calculations in the $A = 16-40$ region.

References

- 1) T. T. Bardin, J. A. Becher, T. R. Fischer and A. D. Jones, Phys. Rev. Lett. **24** (1970) 772
- 2) K. Nakai, J. L. Québert, F. S. Stephens and R. M. Diamond, Phys. Rev. Lett. **24** (1970) 903
- 3) O. Hausser, T. K. Alexander, D. Pelte, B. W. Hooton and H. C. Evans, Phys. Rev. Lett. **23** (1969) 320
- 4) D. A. Bromley, H. E. Gove and A. E. Litherland, Can. J. Phys. **35** (1957) 1057
- 5) C. Gustafsson, I-L. Lamm, B. Nilsson and S. G. Nilsson, Ark. Fys. **36** (1967) 613
- 6) V. M. Strutinsky, Nucl. Phys. **A95** (1967) 420
- 7) S. G. Nilsson, C. F. Tsang, A. Sobiczewski, Z. Szymanski, S. Wycech, C. Gustafsson, I-L. Lamm, P. Möller and B. Nilsson, Nucl. Phys. **A131** (1969) 1
- 8) B. R. Mottelson and S. G. Nilsson, Mat. Fys. Skr. Dan. Vid. Selsk. **1**, no. 8 (1969)
- 9) R. Y. Cusson, private communication
- 10) G. Ripka, Advances in nuclear physics, **1** (1968) 183