



LUND UNIVERSITY

Stabilins are expressed in bone marrow sinusoidal endothelial cells and mediate scavenging and cell adhesive functions.

Qian, Hong; Johansson, Sophie; McCourt, Peter; Smedsrød, Bård; Ekblom, Marja; Johansson, Staffan

Published in:
Biochemical and Biophysical Research Communications

DOI:
[10.1016/j.bbrc.2009.10.068](https://doi.org/10.1016/j.bbrc.2009.10.068)

2009

[Link to publication](#)

Citation for published version (APA):

Qian, H., Johansson, S., McCourt, P., Smedsrød, B., Ekblom, M., & Johansson, S. (2009). Stabilins are expressed in bone marrow sinusoidal endothelial cells and mediate scavenging and cell adhesive functions. *Biochemical and Biophysical Research Communications*, 390, 883-886.
<https://doi.org/10.1016/j.bbrc.2009.10.068>

Total number of authors:
6

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00



LUND UNIVERSITY
Faculty of Medicine

LUP

Lund University Publications

Institutional Repository of Lund University

This is an author produced version of a paper published in Biochemical and biophysical research communications. This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Citation for the published paper:

Hong Qian, Sophie Johansson, Peter McCourt, Bård Smedsrød, Marja Ekblom, Staffan Johansson

“Stabilins are expressed in bone marrow sinusoidal endothelial cells and mediate scavenging and cell adhesive functions.”

Biochemical and biophysical research communications, 2009, Issue: Oct 20

<http://dx.doi.org/10.1016/j.bbrc.2009.10.068>

Access to the published version may require journal subscription.

Published with permission from: Elsevier

Stabilins are expressed in bone marrow sinusoidal endothelial cells and mediate scavenging and cell adhesive functions

Hong Qian,^{1a} Sophie Johansson,^{2b} Peter McCourt,³ Bård Smedsrød,³ Marja Ekblom^{1,4c} and Staffan Johansson^{2,c}

¹ Hematopoietic Stem Cell Laboratory; Dept. Laboratory Medicine, Lund University, Lund, Sweden; ² Dept. of Medical Biochemistry and Microbiology, Uppsala University, Uppsala, Sweden; ³ Dept. of Cell Biology and Histology, University of Tromsø, Tromsø, Norway; and ⁴ Department of Hematology, Lund University Hospital, Lund, Sweden

^{a)} Present address: Dept. of Clinical and Experimental Medicine, Linköping University, Sweden; ^{b)} Present address: Q-Med AB, Uppsala, Sweden

Correspondence: Marja Ekblom, Hematopoietic Stem Cell Laboratory, BMC B10, Lund University, 221 84 Lund, Sweden. Phone: +4646 222 0804. Fax: +4646 222 3600. E-mail: marja.ekblom@med.lu.se

M.E. and St.J. contributed equally to this work.

Funding:

This work was supported by SSMF, the Swedish Research Council Grants No. 7147, K 2009-64X-21033-01-3 and 521-2008-2303. Swedish Cancer Foundation Grants No. 05 0175 and 2007/772., the Norwegian Research Council Grant No. 149091, the Simon Fougner Hartmanns Family Fund, Royal Physiographic Society in Lund, University Hospital of Lund Foundation, Crafoord Foundation and Georg Danielsson's Foundation.

ABSTRACT

Bone marrow sinusoidal endothelial cells have a specific function as a site of transmigration of hematopoietic stem and progenitor cells and mature blood cells between bone marrow and blood stream. However, the specific characteristics of bone marrow sinusoidal endothelial cells are still largely unclear. We here report that these cells express stabilin-1 and stabilin-2, which in liver sinusoidal endothelial cells have been identified as endocytic scavenger receptors for several ligands, including SPARC and hyaluronan. We show here that intravenously injected formaldehyde-treated serum albumin, advanced glycation end-products, and collagen I α -chains were taken up by bone marrow sinusoidal endothelial cells, showing that these cells have a scavenging function and thereby may modulate bone marrow vascular stem cell niches. Importantly, we show that hyaluronan mediated adhesion of hematopoietic stem and progenitor cells to stabilin-2-transfected cells, suggesting that stabilin-2 has a role in adhering circulating stem and progenitor cells to bone marrow.

INTRODUCTION

Endothelial cells are specialized according to the tissue-specific functions. A distinct type of endothelial cell makes up the sinusoids of liver, spleen, and bone marrow. Distinguishing features of these sinusoidal endothelial cells (SECs) are sieve plates of fenestrae without diaphragms and a non-continuous or missing basement membrane(1).

In the liver, SECs have a key role in the endocytic removal of waste products, like connective tissue molecules, atherogenic lipids, and advanced glycation end products (AGEs) from the circulation(2). In BM, SECs function as a transit site for newly produced mature blood cells and for transmigration of hematopoietic stem and progenitor cells (HSPCs) between blood stream and bone marrow (BM) intersinusoidal spaces(3). In contrast to most endothelia, BM SECs constitutively express VCAM-1 and SDF-1(4) which promote migration of HSPCs to BM. However, the molecules mediating specific BM SEC functions, including extra- and intravasation, are still incompletely characterized.

Stabilin-1 and -2 are endocytic receptors expressed in non-continuous sinusoidal endothelium. We therefore studied whether BM SECs may express these receptors and have scavenging functions similar to SECs in other tissues. We also studied the possibility that

stabilin-2 through its hyaluronan-binding ability might have a role in cell adhesive interactions with HSPCs.

DESIGN AND METHODS

Tail vein injections and immunofluorescence microscopy.

The animal experiments were approved by the animal ethics committees in Uppsala (C-192/3) and Lund (M215-05). TRITC-labelled formaldehyde-treated BSA (FSA) and DTAF-labelled collagen I α -chains were produced according to standard protocols (5, 6). FITC-labelled AGE-modified BSA (AGE-BSA) was a kind gift from Dr. Berit Hansen, Heidelberg, Germany. C57Bl6 female mice (8-10 week old) were given single intravenous injections of the ligands in PBS (140 μ g/200 μ l for DTAF-labelled collagen I and 100 μ g/200 μ l for other ligands).

Frozen sections (7-8 μ m) of BM were prepared as described (7). The sections were fixed 5 minutes in -20°C methanol, blocked in PBS, 10% FCS (Gibco), and incubated 1 hour with antibodies against stabilin-1 and -2 (8, 9) and 30 min with Alexa-Fluor 488-goat anti-rabbit antibodies (Molecular Probes, Invitrogen). The sections were photographed using a Zeiss Axioplan microscope equipped with Zeiss Plan-APOCHROMAT 20x/0.75 and 40x/0.95 lenses, ORCA-1394ER digital camera and Openlab-3 software.

RT-PCR.

RNA was purified from mouse liver and BM using the Qiagen RNeasy Mini-Kit and treated with DNase I. cDNA was produced using Superscript II Reverse Transcriptase kit (Invitrogen). PCR primers were: stabilin-1 forward (exon 67) 5'-GTGCTGGGATCTGAGCCTCC, stabilin-1 reverse (exon 69) 5'-GTGTCGGGGAAGTCCTCCTC, stabilin-2 forward (exon 68) 5'-CTTTGGCAAGCAGCAGCCTG, and stabilin-2 reverse (exon 69) 5'-GTTCTCCAGCTCCCGTTCTC. The protocol was: denaturation at 94°C for 5 min, followed by 30 cycles of 94°C for 1 minute / 68°C for 2 minutes / 72°C for 5 minutes, and a final elongation for 7 minutes at 72°C.

Cell adhesion assay.

BM cells from 8-12 week old β -actin GFP transgenic mice(10) were enriched for c-kit by using anti-CD117 magnetic beads (Miltenyi Biotech, Bergisch Gladbach, Germany). Treatment with streptomyces hyaluronidase (H1136, Sigma-Aldrich) (3U/ml in PBS, 0,5% BSA) was performed at 37°C, 5% CO₂ for 30-45 min.

HEK293 cells stably expressing human stabilin-1 or mouse stabilin-2 (8) and nontransfected cells were grown in DMEM with 4,5g/L glucose, 10% FCS, L-glutamine, β -ME (10^{-2} M) at 37°C with 5% CO₂. For cell adhesion assay, the cells were plated in 24-well tissue culture plates (1×10^5 cells/well) and grown overnight. The medium was changed to IMDM. After 4-6 hours, c-kit-enriched cells (2×10^5 cells/well) were added and incubated for 2 hours. The non-adherent cells were harvested by collecting the medium and washing the wells twice with PBS, 5% FCS. The lineage negative, kit^{hi} Sca-1^{hi} (LSK) HSPCs (11) in c-kit enriched input and nonadherent cells were analyzed by BD FACSLSR II (BD Biosciences, San Jose, CA) using FowJo software (TreeStar Inc. San Carlos, CA). Antibodies used (from BD, Biosciences, unless otherwise stated) were rat anti-mouse CD16/32 (clone 2.4G2) , pacific blue- or FITC-Sca-1 (D7) and APC-CY7- or APC-c-Kit (2B8) (E-Bioscience, San Diego, CA), PE-Cy5-rat anti-CD4 (H129.9), -CD3 (17A2), -220/CD45 (RA3-6B2, Biolegend, San Diego, CA), -Mac-1 (M1/70, Biolegend), -TER119 (TER-119, Biolegend), -GR1 (RB6-8C5, Biolegend) and -CD8a (Ly2/53-67). 7-amino-actinomycin (7-AAD; Sigma-Aldrich Co, St Louis, MO) was used to exclude dead cells. Expression of hyaluronan was analyzed using biotinylated hyaluronan-binding protein (Sigma, H9910) and streptavidin-PE.

RESULTS AND DISCUSSION

RNA expression of mouse BM stabilins was studied by RT-PCR. PCR products obtained from BM and liver, used as a positive control (Figure 1A), matched the sizes expected for the selected primers (289 and 118 base pairs for stabilin-1 and -2, respectively). By Western blotting of BM lysates, stabilin-1 was detected as a band of 280 kDa and stabilin-2 as two bands of 300 and 175 kDa (not shown), similar to their appearance in liver (9). Immunofluorescent staining showed ubiquitous expression of both stabilin-1 and stabilin-2 in BM SECs (Figure 1B, D). Control staining with the anti-rabbit antibody gave no specific staining (Figure 1H).

Stabilin-1 was originally identified in non-continuous SECs in spleen(12) and was subsequently found in liver and lymph node SECs. In addition, its expression is induced in angiogenic endothelium associated with wound healing, tumour vascularisation and chronic

inflammation, and in macrophages after “alternative activation”(13). In contrast, expression of stabilin-2 in endothelium has been found restricted to SECs(14), in agreement with the expression in SECs in BM.

To investigate the scavenging function of the bone marrow SECs, the uptake of selected scavenger receptor ligands was analyzed after intravenous administration. Accumulation of TRITC-FSA in the BM SECs was observed within one hour (Figure 1C, E, F). There was little or no fluorescence in the intersinusoidal spaces. Injection of unlabelled FSA together with TRITC-FSA reduced the fluorescence in the SECs (Figure 1G), indicating that the uptake was due to specific recognition of FSA. Likewise, labeled AGE-BSA accumulated in BM SECs (Figure 1I). These two ligands are primarily cleared by the stabilins. Uptake of DTAF- collagen I α -chains, which are eliminated by the receptor for non-native collagens,(6) was also seen in BM SECs (Figure 1I).

Several extracellular matrix components have been identified as ligands for stabilins. One example is SPARC, which is expressed by hematopoietic progenitors, megakaryocytes, osteoblasts, adipocytes, and endothelial cells(15). SPARC mediates diverse functions, including inhibition of cell proliferation and modulation of cell-matrix interactions(15), and its clearance by stabilin-1 may thus influence proliferation and migration of BM cells(16). Stabilin-2 has been identified as a receptor for hyaluronan, collagen propeptides(9), chondroitin sulphate(17) and heparin. Hyaluronan is synthesized and deposited on the cell surface by HSPCs and has been shown to mediate their mobilization and homing into BM(18). Our FACs analysis showed expression of hyaluronan in one third of LSK cells (Fig 2A), in agreement with published results(19). Importantly, we found significantly increased adhesion of c-kit enriched cells and LSK cells to cells transfected with stabilin-2 compared to cells expressing stabilin-1 or non-transfected cells. Pretreatment with hyaluronidase resulted in significantly reduced adhesion of both c-kit enriched cells and LSK cells to stabilin-2 expressing cells. These results suggest that stabilin-2 may be an important adhesion molecule for initial recruitment of circulating HSPCs to SECs, promoting their homing into BM. Similarly, binding of tumour cell-associated hyaluronan by stabilin-2 might cause tumour cells in blood to halt in bone marrow, thereby increasing the likelihood for metastasis at this site. In support of this, high level of hyaluronan synthesis in prostate cancer cells has been correlated with increased metastasis to the bone marrow(20).

In conclusion, this study shows a novel function for BM SECs as a scavenging endothelium with implications for the molecular environment in BM stroma. Further, the specific cell adhesive interactions mediated by hyaluronan on HSPCs and stabilin-2 expressing cells in vitro suggest a new role for this receptor in HSPC homing to BM.

AUTHORSHIP AND DISCLOSURES

Author's contributions: HQ. was the principal investigator and takes primary responsibility for the paper. HQ, SoJ, and ME performed laboratory work for this study. HQ, SoJ, BS, PM, StJ and ME designed research, analyzed data and wrote the paper.

Declaration of commercial interest: All authors declare that they have no conflicting interests.

Running heads:

H Qian et al. Stabilins in bone marrow sinusoids

REFERENCES

1. Wisse E. An electron microscopic study of the fenestrated endothelial lining of rat liver sinusoids. *J Ultrastruct Res.* 1970 Apr;31(1):125-50.
2. Smedsrod B. Clearance function of scavenger endothelial cells. *Comp Hepatol.* 2004 Jan 14;3 Suppl 1:S22.
3. Lapidot T, Dar A, Kollet O. How do stem cells find their way home? *Blood.* 2005 Sep 15;106(6):1901-10.
4. Sipkins DA, Wei X, Wu JW, Runnels JM, Cote D, Means TK, et al. In vivo imaging of specialized bone marrow endothelial microdomains for tumour engraftment. *Nature.* 2005 Jun 16;435(7044):969-73.
5. Elvevold KH, Nedredal GI, Revhaug A, Smedsrod B. Scavenger properties of cultivated pig liver endothelial cells. *Comp Hepatol.* 2004 Aug 12;3(1):4.
6. Malovic I, Sorensen KK, Elvevold KH, Nedredal GI, Paulsen S, Erofeev AV, et al. The mannose receptor on murine liver sinusoidal endothelial cells is the main denatured collagen clearance receptor. *Hepatology (Baltimore, Md.)* 2007 Jun;45(6):1454-61.
7. Gu Y, Sorokin L, Durbeej M, Hjalt T, Jonsson JI, Ekblom M. Characterization of bone marrow laminins and identification of alpha5-containing laminins as adhesive proteins for multipotent hematopoietic FDCCP-Mix cells. *Blood.* 1999 Apr 15;93(8):2533-42.
8. Hansen B, Longati P, Elvevold K, Nedredal GI, Schledzewski K, Olsen R, et al. Stabilin-1 and stabilin-2 are both directed into the early endocytic pathway in hepatic sinusoidal endothelium via interactions with clathrin/AP-2, independent of ligand binding. *Experimental cell research.* 2005 Feb 1;303(1):160-73.
9. McCourt PA, Smedsrod BH, Melkko J, Johansson S. Characterization of a hyaluronan receptor on rat sinusoidal liver endothelial cells and its functional relationship to scavenger receptors. *Hepatology (Baltimore, Md.)* 1999 Nov;30(5):1276-86.
10. Wagers AJ, Allsopp RC, Weissman IL. Changes in integrin expression are associated with altered homing properties of Lin(-/lo)Thy1.1(lo)Sca-1(+)/c-kit(+) hematopoietic stem cells following mobilization by cyclophosphamide/granulocyte colony-stimulating factor. *Experimental hematology.* 2002 Feb;30(2):176-85.
11. Qian H, Tryggvason K, Jacobsen SE, Ekblom M. Contribution of alpha6 integrins to hematopoietic stem and progenitor cell homing to bone marrow and collaboration with alpha4 integrins. *Blood.* 2006 May 1;107(9):3503-10.
12. Goerdt S, Walsh LJ, Murphy GF, Pober JS. Identification of a novel high molecular weight protein preferentially expressed by sinusoidal endothelial cells in normal human tissues. *J Cell Biol.* 1991 Jun;113(6):1425-37.
13. Kzhyshkowska J, Gratchev A, Goerdt S. Stabilin-1, a homeostatic scavenger receptor with multiple functions. *Journal of cellular and molecular medicine.* 2006 Jul-Sep;10(3):635-49.
14. Falkowski M, Schledzewski K, Hansen B, Goerdt S. Expression of stabilin-2, a novel fasciclin-like hyaluronan receptor protein, in murine sinusoidal endothelia, avascular tissues, and at solid/liquid interfaces. *Histochemistry and cell biology.* 2003 Nov;120(5):361-9.
15. Framson PE, Sage EH. SPARC and tumor growth: where the seed meets the soil? *Journal of cellular biochemistry.* 2004 Jul 1;92(4):679-90.
16. Kzhyshkowska J, Workman G, Cardo-Vila M, Arap W, Pasqualini R, Gratchev A, et al. Novel function of alternatively activated macrophages: stabilin-1-mediated clearance of SPARC. *J Immunol.* 2006 May 15;176(10):5825-32.

17. Weigel JA, Weigel PH. Characterization of the recombinant rat 175-kDa hyaluronan receptor for endocytosis (HARE). *J Biol Chem*. 2003 Oct 31;278(44):42802-11.
18. Avigdor A, Goichberg P, Shivtiel S, Dar A, Peled A, Samira S, et al. CD44 and hyaluronic acid cooperate with SDF-1 in the trafficking of human CD34+ stem/progenitor cells to bone marrow. *Blood*. 2004 Apr 15;103(8):2981-9.
19. Nilsson SK, Haylock DN, Johnston HM, Occhiodoro T, Brown TJ, Simmons PJ. Hyaluronan is synthesized by primitive hemopoietic cells, participates in their lodgment at the endosteum following transplantation, and is involved in the regulation of their proliferation and differentiation in vitro. *Blood*. 2003 Feb 1;101(3):856-62.
20. Draffin JE, McFarlane S, Hill A, Johnston PG, Waugh DJ. CD44 potentiates the adherence of metastatic prostate and breast cancer cells to bone marrow endothelial cells. *Cancer research*. 2004 Aug 15;64(16):5702-11.

FIGURE LEGENDS

Figure 1. Expression of stabilin-1 and stabilin-2 in bone marrow (BM) sinusoidal endothelial cells (SECs). (A). cDNA fragments generated from liver and bone marrow (BM) using PCR primers for stabilin-1 (lanes 1 and 4) and stabilin-2 (lanes 2 and 3) were separated on 1.5% agarose gel. Bp: 100-300 base pair molecular weight markers. (B-E) Immunostaining of stabilin-1 (B) and stabilin-2 (D) of in mouse bone marrow SECs and uptake of TRITC-FSA by SECs in the corresponding tissue sections (C, E). The bone marrow was isolated 60 minutes after the injection of TRITC-conjugated FSA in the tail vein and the frozen sections were stained by antibodies against stabilin-1 and stabilin-2. (H) Control staining with the anti-rabbit antibody. (F, G). Competition of uptake of TRITC-FSA by unlabelled FSA. TRITC-FSA (0.5 mg/ml) was injected in the tail vein alone (F) or together with 6.4 mg/mL of unlabelled FSA (G), and the accumulation of fluorescence in BM after 60 minutes was analyzed. (I). Localization in mouse BM of intravenously injected AGE-BSA. The bone marrow was collected 60 minutes after injection of the FITC-conjugated protein. (J, K), Localization in BM (J) and liver (K) of intravenously injected DTAF-collagen α -chains. The tissues were collected 30 minutes after injection of the protein. The sections were photographed using a Zeiss Axioplan microscope equipped with Zeiss Plan-APOCHROMAT 20x/0.75 and 40x/0.95 lenses, ORCA-1394ER digital camera and Openlab-3 software. Original magnification was (B-H, J, K) x40, (I) x20.

Figure 2. Expression of hyaluronan (HA) on bone marrow (BM) stem and progenitor cells and adhesion of BM c-kit-enriched and lineage negative, c-kit^{hi} Sca-1^{hi} (LSK) cells to HEK293 cells and HEK293 cells transfected with stabilin-1 or stabilin-2. (A). HA expression in LSK cells. Left panel: FACS profile showing gating of SCA-1^{high}c-Kit^{high} of lineage negative (not shown) cells. Right panel: HA expression (dashed line) and control staining with the secondary streptavidin-PE antibody (solid line) in the gated LSK cells. (B). The number of nonadherent c-kit enriched BM cells (KIT) (left) and LSK cells (right) after incubation in wells plated with HEK293 cells (293) or cells transfected with stabilin-1 (ST1) or stabilin-2 (ST2). ST2+HASE; adhesion of cells treated with hyaluronidase to HEK cells expressing stabilin-2. ** p<0.01; *** p<0.001. Data are from 2 experiments performed in 3-6 replicates; horizontal bars show mean values. Each dot represents nonadherent cells form one well. Except for two samples pooled in

(293) group, the frequency of LSK cells was analyzed separately from each well. The mean number of LSK cells within 2×10^5 c-kit enriched cells was $13.8-15.5 \times 10^3$. (C). FACS profiles showing the frequency of LSK cells in the corresponding nonadherent cell populations.

Figure 1




Figure 2.

