



LUND UNIVERSITY

The IAU 2000 Resolutions for Astrometry, Celestial Mechanics, and Metrology in the Relativistic Framework: Explanatory Supplement

Soffel, M.; Klioner, S. A.; Petit, G.; Wolf, P.; Kopeikin, S. M.; Bretagnon, P.; Brumberg, V. A.; Capitaine, N.; Damour, T.; Fukushima, T.; Guinot, B.; Huang, T.-Y.; Lindegren, Lennart; Ma, C.; Nordtvedt, K.; Ries, J. C.; Seidelmann, P. K.; Vokrouhlický, D.; Will, C. M.; Xu, C.

Published in:
Astrophysical Journal

DOI:
[10.1086/378162](https://doi.org/10.1086/378162)

2003

[Link to publication](#)

Citation for published version (APA):

Soffel, M., Klioner, S. A., Petit, G., Wolf, P., Kopeikin, S. M., Bretagnon, P., Brumberg, V. A., Capitaine, N., Damour, T., Fukushima, T., Guinot, B., Huang, T.-Y., Lindegren, L., Ma, C., Nordtvedt, K., Ries, J. C., Seidelmann, P. K., Vokrouhlický, D., Will, C. M., & Xu, C. (2003). The IAU 2000 Resolutions for Astrometry, Celestial Mechanics, and Metrology in the Relativistic Framework: Explanatory Supplement. *Astrophysical Journal*, 126(6), 2687-2706. <https://doi.org/10.1086/378162>

Total number of authors:
20

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

THE IAU 2000 RESOLUTIONS FOR ASTROMETRY, CELESTIAL MECHANICS, AND METROLOGY IN THE RELATIVISTIC FRAMEWORK: EXPLANATORY SUPPLEMENT

M. SOFFEL,¹ S. A. KLIONER,¹ G. PETIT,² P. WOLF,² S. M. KOPEIKIN,³ P. BRETAGNON,⁴ V. A. BRUMBERG,⁵ N. CAPITAINE,⁶ T. DAMOUR,⁷ T. FUKUSHIMA,⁸ B. GUINOT,⁶ T.-Y. HUANG,⁹ L. LINDEGREN,¹⁰ C. MA,¹¹ K. NORDTVEDT,¹² J. C. RIES,¹³ P. K. SEIDELMANN,¹⁴ D. VOKROUHLICKÝ,¹⁵ C. M. WILL,¹⁶ AND C. XU¹⁷

Received 2002 August 9; accepted 2003 July 2

ABSTRACT

We discuss the IAU resolutions B1.3, B1.4, B1.5, and B1.9 that were adopted during the 24th General Assembly in Manchester, 2000, and provides details on and explanations for these resolutions. It is explained why they present significant progress over the corresponding IAU 1991 resolutions and why they are necessary in the light of present accuracies in astrometry, celestial mechanics, and metrology. In fact, most of these resolutions are consistent with astronomical models and software already in use. The metric tensors and gravitational potentials of both the Barycentric Celestial Reference System and the Geocentric Celestial Reference System are defined and discussed. The necessity and relevance of the two celestial reference systems are explained. The transformations of coordinates and gravitational potentials are discussed. Potential coefficients parameterizing the post-Newtonian gravitational potentials are expounded. Simplified versions of the time transformations suitable for modern clock accuracies are elucidated. Various approximations used in the resolutions are explicated and justified. Some models (e.g., for higher spin moments) that serve the purpose of estimating orders of magnitude have actually never been published before.

Key words: astrometry — celestial mechanics — reference systems — time

1. INTRODUCTION

It is clear that, beyond some threshold of accuracy, any astronomical problem has to be formulated within the framework of Einstein's theory of gravity (general relativity theory, or GRT). Many high-precision astronomical techniques have already passed this threshold. For example,

lunar laser ranging measures the distance to the Moon with a precision of a few centimeters, thereby operating at the 10^{-10} level. At this level, several relativistic effects are significant and observable. Relativistic effects related to the motion of the Earth-Moon system about the Sun are of order $(v_{\text{orbital}}/c)^2 \simeq 10^{-8}$. The Lorentz contraction of the lunar orbit about Earth that appears in barycentric coordinates has an amplitude of about 100 cm, whereas in some suitably chosen (local) coordinate system that moves with the Earth-Moon barycenter, the dominant relativistic range oscillation reduces to only a few centimeters (Mashhoon 1985; Soffel, Ruder, & Schneider 1986).

¹ Lohrmann-Observatorium, Institut für Planetare Geodäsie, Technische Universität Dresden, Mommsenstrasse 13, D-01062 Dresden, Germany.

² Bureau International des Poids et Mesures, Pavillon de Breteuil, F-92312 Sèvres, France.

³ Department of Physics and Astronomy, 322 Physics Building, University of Missouri–Columbia, Columbia, MO 65211.

⁴ Bureau des Longitudes, 77 Avenue Denfert-Rechereau, F-75014 Paris, France.

⁵ Institute of Applied Astronomy, Russian Academy of Sciences, Naberezhnaya Kutuzova 10, St. Petersburg 191187, Russia.

⁶ Observatoire de Paris, 61 Avenue de l'Observatoire, F-75014 Paris, France.

⁷ Institut des Hautes Etudes Scientifiques, 35 Route de Chartres, F-91440 Bures-sur-Yvette, France.

⁸ National Astronomical Observatory, 2-21-1 Osawa, Mitaka, Tokyo 181, Japan.

⁹ Department of Astronomy, Nanjing University, 210093 Nanjing, China; and National Astronomical Observatories, Chinese Academy of Sciences.

¹⁰ Institutionen för Astronomi, Lunds Universitet, Box 43, SE-221 00 Lund, Sweden.

¹¹ Space Geodesy Branch, Code 926, Laboratory for Terrestrial Physics, NASA Goddard Space Flight Center, Greenbelt, MD 20771.

¹² Northwest Analysis, 118 Sourdough Ridge Road, Bozeman, MT 59715.

¹³ Center for Space Research, University of Texas at Austin, 1 University Station R1000, Austin, TX 78712.

¹⁴ Department of Astronomy, University of Virginia, P.O. Box 3818, Charlottesville, VA 22903.

¹⁵ Astronomický Ústav, Universita Karlova, V. Holešovičkách 2, CZ-180 00 Praha 8, Czech Republic.

¹⁶ McDonnell Center for the Space Sciences and Department of Physics, Washington University, 1 Brookings Drive, St. Louis, MO 63130.

¹⁷ Department of Physics, Nanjing Normal University, 210097 Nanjing, China.

The situation is even more critical in the field of astrometry. It is well known that the gravitational light deflection at the limb of the Sun amounts to $1''.75$ and decreases only as $1/r$ with increasing impact parameter r of a light ray to the solar center. Thus, for light rays incident at about 90° from the Sun the angle of light deflection still amounts to 4 mas. To describe the accuracy of astrometric measurements, it is useful to make use of the parameter γ of the parameterized post-Newtonian (PPN) formalism. We would like to emphasize that this paper deals solely with Einstein's theory of gravity, where $\gamma = 1$, and not with the PPN formalism. Nevertheless, the introduction of γ is useful if one wishes to talk about measurement accuracies. In the PPN formalism, the angle of light deflection is proportional to $(\gamma + 1)/2$, so that astrometric measurements might be used for a precise determination of the parameter γ . Meanwhile, very long baseline interferometry (VLBI) has achieved accuracies of better than 0.1 mas, and regular geodetic VLBI measurements have frequently been used to determine the space curvature parameter. A detailed analysis of VLBI data from the projects MERIT and POLARIS/IRIS yielded $\gamma = 1.000 \pm 0.003$ (Robertson & Carter 1984; Carter, Robertson, & MacKay 1985), where a formal standard error is given. Recently an advanced processing of VLBI data provided the best current estimates, $\gamma = 0.9996 \pm 0.0017$

(Lebach et al. 1995) and $\gamma = 0.99994 \pm 0.00031$ (Eubanks et al. 1997). The current accuracy of modern optical astrometry, as represented by the *Hipparcos* Catalogue, is about 1 mas, which gave a determination of γ at the level of 0.997 ± 0.003 (Froeschlé, Mignard, & Arenou 1997). Future astrometric missions such as *SIM* and especially *GAI*A will push the accuracy to the level of a few microarcseconds, and the expected accuracy of determinations of γ will be 10^{-6} to 10^{-7} . The accuracy of $1 \mu\text{as}$ should be compared with the maximal possible light deflection due to various parts of the gravitational field: the post-Newtonian effect of $1''.75$ due to the mass of the Sun, $240 \mu\text{as}$ caused by the oblateness of Jupiter, J_2 ($10 \mu\text{as}$ due to Jupiter's J_4), the post-post-Newtonian effect of $11 \mu\text{as}$ due to the Sun, etc. This illustrates how complicated the relativistic modeling of future astrometric observations will be. It is clear that for such high accuracy the corresponding model must be formulated in a self-consistent relativistic framework.

Another problem worth mentioning is that of time measurement. The realization of the SI (*Système international*) second (the unit of proper time) has improved by 1 order of magnitude in the last few years with the advent of laser-cooled atomic clocks (Lemondé et al. 2001; Weyers et al. 2001; references therein) and is now below 2 parts in 10^{15} . This should be compared with the dimensionless quantity $U_E/c^2 \simeq 7 \times 10^{-10}$, which gives the order of magnitude of relativistic effects produced by the gravity field of Earth itself in the vicinity of its surface. In the near future, laser-cooled atomic clocks in microgravity are expected to lead to a further improvement by at least 1 order of magnitude. At present, several clock experiments in terrestrial orbit are planned, such as the Atomic Clock Ensemble in Space project (Lemondé et al. 2001). These in turn are likely to lead to clock experiments in solar orbits, such as the *Solar Orbit Relativity Test* project. All of these experiments require a detailed account of many subtle relativistic effects.

Finally, we would like to mention the problem of geodetic precession and nutation (a relativistic effect that is discussed in more detail below; Misner, Thorne, & Wheeler 1973; Soffel 1989) and the description of Earth's rotation in a suitably chosen geocentric celestial reference system. Geodetic precession amounts to $1''.9$ per century, and geodetic nutation is dominated by an annual term with amplitude 0.15 mas. Since the geocentric reference system is chosen to be kinematically nonrotating, geodetic precession and nutation should be contained in the model describing the relation between the geocentric system and the International Terrestrial Reference System (ITRS). According to IAU Resolution B1.6 (2000), this relativistic precession-nutation is indeed contained in the present IAU precession-nutation model.

These examples show clearly that high-precision modern astronomical observations can no longer be described by Newtonian theory but require Einstein's theory of gravity. The consequences of this are profound for the basic formalism to be used, since one often tends to express it in terms of "small relativistic corrections" to Newtonian theory. This can lead to misconceptions and mistakes. One central point is that in Newton's theory, globally preferred coordinate systems exist that have a direct physical meaning. In the Newtonian framework, idealized clocks show absolute time everywhere in the universe at all times, and global spatial inertial coordinates exist in which dynamical equations of motion show no inertial forces. This is no longer true in

GRT. Usually, spacetime coordinates have no direct physical meaning and it is essential to construct the observables as coordinate-independent quantities, that is, scalars, in mathematical language. This construction usually occurs in two steps: first one formulates a coordinate picture of the measurement procedure, and then one derives the observable out of it. This leads us to the problem of defining useful and adequate coordinate systems in astronomy. The underlying concept in relativistic modeling of astronomical observations is a relativistic four-dimensional *reference system*. By reference system, we mean a purely mathematical construction (a chart or a coordinate system) giving "names" to spacetime events. In contrast to this, a *reference frame* is some materialization of a reference system. In astronomy, the materialization is usually realized by means of a catalog or ephemeris, containing positions of some celestial objects relative to the reference system under consideration. Hence it is very important to understand that any reference frame is defined only through a well-defined reference system, which has been used to construct physical models of observations.

In the following, a four-dimensional spacetime reference system will be described by four coordinates $x^\alpha = (x^0, x^i) = (x^0, x^1, x^2, x^3)$. Here and below, the Greek indices (e.g., α) take the values 0, 1, 2, and 3, and the Latin ones (e.g., i) take the values 1, 2, and 3. The index 0 refers to the time variable and the indices 1, 2, and 3 refer to the three spatial coordinates. For dimensional reasons, one usually writes $x^0 = ct$, where c is the speed of light and t is a time variable. According to the mathematical formalism of general relativity, a particular reference system is fixed by the specific form of the metric tensor $g_{\alpha\beta}(t, x^i)$, which allows one to compute the 4-distance ds between any two events x^α and $x^\alpha + dx^\alpha$ according to the rule

$$ds^2 = g_{\alpha\beta}(t, x^i) dx^\alpha dx^\beta \\ \equiv g_{00} c^2 dt^2 + 2g_{0i} c dt dx^i + g_{ij} dx^i dx^j, \quad (1)$$

where Einstein's summation convention (summation over repeated indices) is implied. The metric tensor allows one to derive translational and rotational equations of motion of bodies, to describe the propagation of light, and to model the process of observation. Examples of such modeling include relating the observed (proper) time of an observer to the coordinate time t , and relating the angles between two incident light rays as observed by that observer to the corresponding coordinate directions. All of these components can be combined into a single relativistic model for a particular kind of observation. Such a model contains a certain set of parameters describing various properties of the objects participating in the process of observation. These parameters should be determined from observations. Many of these parameters crucially depend upon the reference system used to formulate the model of observations (e.g., the initial positions and velocities of certain bodies). Some other parameters might not depend at all upon the reference system (e.g., the speed of light in vacuum). On the other hand, according to the principle of covariance, different reference systems covering the region of spacetime under consideration are mathematically equivalent in the sense that any such system can be used to model the observations. This freedom to choose the reference system can be used to simplify the models or to make the resulting parameters more physically adequate.

It is widely accepted that in order to adequately describe modern astronomical observations, one has to use several relativistic reference systems. The solar system Barycentric Celestial Reference System (BCRS) can be used to model light propagation from distant celestial objects, as well as the motion of bodies within the solar system. The Geocentric Celestial Reference System (GCRS) is physically adequate to describe processes occurring in the vicinity of Earth (Earth's rotation, the motion of Earth's satellites, etc.). The introduction of further local systems (selenocentric, martian, etc.) might be of relevance for specific applications, where physical phenomena in the vicinity of the corresponding body play a role.

The necessity of using several reference systems can be understood from the following example: If we were to characterize terrestrial observers by the difference between their BCRS coordinates and the BCRS coordinates of the geocenter, the positions of the observers relative to the geocenter would be altered by time-dependent, relativistic coordinate effects (such as Lorentz contraction) that have nothing to do with Earth's rotation or with geophysical factors and which would vanish if one employed suitable GCRS coordinates instead. On the other hand, the coordinate positions derived with VLBI observations are used to investigate local geophysical processes, and some adequate geocentric reference system allows one to simplify their description.

The basic idea is to construct a special local reference system for each material subsystem, in which the relativistic equations of motion for a test body inside the subsystem under consideration take a particularly simple form. In such a local reference system the influence of external matter, in

accordance with the equivalence principle, should be given by tidal potentials only, that is, by potentials whose expansions in powers of local spatial coordinates in the vicinity of the origin of the corresponding reference system start with the second order (the linear terms representing inertial forces may also exist, but they can be eliminated if desired by a suitable choice of the origin of the local coordinates).

Two advanced relativistic formalisms have been worked out to tackle this problem in the first post-Newtonian approximation of general relativity. One formalism is due to Brumberg and Kopeikin (Brumberg & Kopeikin 1989; Kopeikin 1988, 1990; Brumberg 1991; Klioner & Voinov 1993), and another is due to Damour, Soffel, and Xu (Damour, Soffel, & Xu 1991, 1992, 1993, 1994). The IAU 2000 Resolutions B1.3–B1.5 are based upon these approaches. These resolutions extend corresponding older ones that are reconsidered in the next section. From a mathematical point of view, Resolution B1.3 recommends the use of certain coordinates and the way of writing the metric tensor. Clearly, one might use any coordinate system that is well adapted to a specific problem of interest. Nevertheless, because of the high risk of possible confusion, the strategy of recommending special coordinate systems (to fix the gauge completely, in mathematical language) has significant advantages. If different coordinates are employed to derive certain results, this should be stressed explicitly so that they can be transformed into the reference systems recommended by the IAU and can be compared with the results derived in the IAU framework.

The organization of the present paper is as follows (see also Fig. 1): In § 2, the principal content of the IAU 1991 recommendations on relativity (§ 2.1) and the further




FIG. 1.—Notations used for quantities of the Barycentric and Geocentric Celestial Reference Systems (coordinates, metric, potentials, and multipole moments) with references to the sections and resolutions where they appear.

related IAU and International Union of Geodesy and Geophysics (IUGG) resolutions (§ 2.2) are repeated. The IAU 2000 resolutions on relativity (Resolutions B1.3, B1.4, B1.5, and B1.9) are discussed in § 3. The full text of the IAU 2000 resolutions on relativity is given in Appendix A. Section 3.1 briefly clarifies the necessity and the role of the two celestial reference systems defined by the IAU resolutions. The Barycentric Celestial Reference System, defined by Resolution B1.3, is discussed in § 3.2. Section 3.3 is devoted to a discussion of the Geocentric Celestial Reference System, as well as the definition of the geocentric gravitational potentials also defined by Resolution B1.3. The coordinate transformations between the BCRS and GCRS, also fixed by Resolution B1.3, are explained in § 3.4. Potential coefficients that can be used to represent in a meaningful way the post-Newtonian geocentric gravitational potential of Earth in its immediate vicinity are fixed by Resolution B1.4 and explained in § 3.5. As an illustration, the gravitational potentials of the BCRS are calculated in § 3.6 for the simplified case in which all gravitating bodies of the solar system can be characterized by their masses only (no further structure of the gravitational field of the bodies is considered). A similar form of the barycentric gravitational potentials is used in Resolution B1.5, where a practical relativistic framework for time and frequency applications in the solar system is formulated. This practical relativistic framework is discussed in § 4. The practical transformation between the coordinate times of the BCRS and GCRS is explained in § 4.1, while the transformations between the various kinds of time scales appropriate for Earth's vicinity are discussed in § 4.2. Appendix B contains an explicit proof that the BCRS metric coincides with well-known results from the literature.

2. THE IAU 1991 FRAMEWORK AND PREVIOUS RECOMMENDATIONS

2.1. *The IAU 1991 Recommendations*

IAU Resolution A4 (1991) contains nine recommendations, the first five of which are directly relevant to our discussion.

In the first recommendation, the metric tensor for spacetime coordinate systems (t, \mathbf{x}) centered at the barycenter of an ensemble of masses is recommended in the form

$$\begin{aligned} g_{00} &= -1 + \frac{2U(t, \mathbf{x})}{c^2} + O(c^{-4}), \\ g_{0i} &= O(c^{-3}), \\ g_{ij} &= \delta_{ij} \left(1 + \frac{2U(t, \mathbf{x})}{c^2} \right) + O(c^{-4}), \end{aligned} \quad (2)$$

where c is the speed of light in vacuum ($c = 299,792,458 \text{ m s}^{-1}$) and U is the Newtonian gravitational potential (here a sum of the gravitational potentials of the ensemble of masses and of an external potential generated by bodies external to the ensemble, the latter potential vanishing at the origin). The algebraic sign of U is taken to be positive, and it satisfies Poisson's equation,

$$\nabla^2 U = -4\pi G\rho. \quad (3)$$

Here G is the gravitational constant, ρ is the matter density, and ∇^2 is the usual Laplacian $\nabla^2 \equiv \partial^2/\partial x^2 + \partial^2/\partial y^2 +$

$\partial^2/\partial z^2$, where $\mathbf{x} = (x, y, z)$. This recommendation also recognizes that spacetime cannot be described by a single coordinate system. The recommended form of the metric tensor can be used not only to describe the barycentric reference system of the whole solar system, but also to define the geocentric reference system centered at the center of mass of Earth with a suitable function instead of U , now depending upon geocentric coordinates. In analogy to the geocentric reference system, a corresponding reference system can be constructed for any other body of the solar system.

In the second recommendation, the origin and orientation of the spatial coordinate grids for the barycentric and geocentric reference systems are defined. Notably, it is specified that the spatial coordinates of these systems should show no global rotation with respect to a set of distant extragalactic objects. It also specifies that the SI second and meter should be the units of time and length in all coordinate systems. It states in addition that the time coordinates should be derived from an Earth atomic time scale.

The third recommendation defines TCB (Barycentric Coordinate Time) and TCG (Geocentric Coordinate Time) as the time coordinates of the BCRS and GCRS, respectively. Here we write $(t = \text{TCB}, x^i)$ and $(T = \text{TCG}, X^i)$ for the respective coordinates. The recommendation also defines the origin of the time scales in terms of International Atomic Time (TAI). The reading of the coordinate time scales on 1977 January 1, 00:00:00 TAI (JD = 2,443,144.5 TAI) must be 1977 January 1, 00:00:32.184. Finally, the recommendation declares that the units of measurement of the coordinate times of all reference systems should be chosen so that they are consistent with the SI second. The relationship between TCB and TCG is then given by the time part of the full four-dimensional transformation between the barycentric and geocentric reference systems,

TCB – TCG

$$= \frac{1}{c^2} \left[\int_{t_0}^t \left(\frac{v_E^2}{2} + U_{\text{ext}}(\mathbf{x}_E) \right) dt + v_E^i r_E^i \right] + O(c^{-4}), \quad (4)$$

where x_E^i and v_E^i are the barycentric coordinate position and velocity of the geocenter, $r_E^i = x^i - x_E^i$ with x^i the barycentric position of some observer, and $U_{\text{ext}}(\mathbf{x}_E)$ is the Newtonian potential of all solar system bodies apart from Earth evaluated at the geocenter.

In the fourth recommendation, another coordinate time, Terrestrial Time (TT), is defined. It differs from TCG by a constant rate only,

$$\begin{aligned} \text{TCG} - \text{TT} &= L_G \times (\text{JD} - 2,443,144.5) \times 86,400, \\ L_G &\approx 6.969291 \times 10^{-10}, \end{aligned} \quad (5)$$

where JD is TAI measured in Julian days, so that the mean rate of TT agrees with that of the proper time of an observer situated on the geoid up to a certain accuracy limit. Up to a constant shift of 32.184 s, TT represents an ideal form of TAI, the divergence between them being a consequence of the physical defects of atomic clocks. It is also recognized that TT is nothing but a rescaling of the geocentric time TCG.

The fifth recommendation states that the old barycentric time TDB may still be used where discontinuity with previous work is deemed to be undesirable. Let us note, however, that TDB was never defined in a self-consistent and exact manner. For that reason it cannot be used in

theoretical considerations. In the notes to the third recommendation, the relation of TCB to TDB is given as

$$\begin{aligned} \text{TCB} - \text{TDB} &= L_B \times (\text{JD} - 2,443,144.5) \times 86,400, \\ L_B &\approx 1.550505 \times 10^{-8}. \end{aligned} \quad (6)$$

Note, however, that according to IAU Resolution C7 (see § 2.2), JD is defined in Terrestrial Time, which makes this formula problematic.

2.2. Further Resolutions

Resolution 2 (1991) of the IUGG defined the Conventional Terrestrial Reference System (CTRS) as a reference system resulting from a (time dependent) spatial rotation of the geocentric reference system defined by the 1991 IAU recommendations, the spatial rotation being chosen such that the CTRS has no global residual rotation with respect to horizontal motions at Earth's surface. The coordinate time of the CTRS coincides with TCG.

IAU Resolution C7 (1994) recommends that the epoch J2000, as well as the Julian (ephemeris) day, be defined in TT. IAU Resolution B6 (1997) has supplemented this framework by one more recommendation stating that no scaling of spatial axes should be applied in any reference system (even if a scaled time coordinate such as TT is used). Note, however, that this resolution has been ignored in the construction of the International Terrestrial Reference Frame, which is defined not with the GCRS spatial coordinates \mathbf{X} but with scaled coordinates $\mathbf{X}_{\text{TT}} = (1 - L_G)\mathbf{X}$.

3. THE IAU 2000 RESOLUTIONS ON RELATIVITY

The IAU 1991 framework is unsatisfactory from many points of view. The Einstein-Infeld-Hoffmann equations of motion, which have been used since the 1970s to construct the JPL numerical ephemerides of planetary motion, cannot be derived from the metric of equation (2). In other words, for the motion of massive solar system bodies this metric is not the post-Newtonian metric of Einstein's theory of gravity. In the years prior to the 23d General Assembly in Kyoto (1997), it became obvious that the IAU 1991 set of recommendations concerning relativity in astrometry, celestial mechanics, and metrology was not sufficient for the accuracies that were achievable. Especially with respect to planned astrometric missions with microarcsecond accuracy, extended and improved resolutions had become indispensable. For that reason, the IAU Working Group on Relativity in Celestial Mechanics and Astrometry, together with a joint committee of the Bureau International des Poids et Mesures and the IAU on relativity for spacetime reference systems and metrology, suggested such an extended set of resolutions (B1.3–B1.5 and B1.9), which was finally adopted at the IAU General Assembly in Manchester in the year 2000. The relevant resolutions can be found in Appendix A. It is clear that because of their brevity they need additional explanation, and there is also a need to show how they work in practice. This paper now presents a detailed explanatory supplement for these IAU 2000 resolutions.

3.1. The Role of the Two Celestial Reference Systems, BCRS and GCRS

Some of the reasons why two different celestial astronomical reference systems have to be introduced have already

been mentioned in § 1. Here we would like to deepen this discussion in several respects. It is clear that for many applications in the fields of astrometry, celestial mechanics, geodynamics, geodesy, etc., some quasi-inertial or “space-fixed” reference system has to be introduced. Resolution B1.3 actually defines *two* different celestial reference systems: the Barycentric Celestial Reference System and the Geocentric Celestial Reference System.

In *Newtonian theory*, one can easily introduce inertial spacetime coordinates that cover the entire universe. Such inertial coordinates in Newton's theory are unique up to the choice of origin, scales, and orientation of the spatial axes and up to a constant velocity of origin. In astronomy, *conceptually* we may talk about two different relevant celestial systems: a barycentric one and a geocentric one, which basically serve different purposes. The barycentric celestial system is considered to be inertial (external Galactic and extragalactic matter normally being neglected) and is used for solar system ephemerides, for concepts such as an ecliptic, for interplanetary spacecraft navigation, etc. The positions of remote objects can be defined in that system. The barycentric celestial system presents the fundamental astrometric system, in which concepts such as “proper motion” and “radial velocity” can be defined.

On the other hand, the geocentric celestial system might be called quasi-inertial, since the spatial axes are non-rotating in the Newtonian absolute sense, whereas the geocenter is accelerated. It is employed for the description of physical processes in the vicinity of Earth, for satellite theory, the dynamics of Earth (including Earth's rotation), etc. It is also used for the introduction of concepts such as the equator and the ITRS. Let us denote the time and space coordinates of the barycentric celestial system by (t, \mathbf{x}) , and those of the geocentric celestial system by (T, \mathbf{X}) . In Newton's framework the relation between these two sets of coordinates is trivial:

$$T = t, \quad \mathbf{X} = \mathbf{x} - \mathbf{x}_E(t),$$

where $\mathbf{x}_E(t)$ denotes the barycentric position of the geocenter. Because these relations are so trivial, for some purposes the barycentric and the geocentric celestial systems are not always clearly distinguished in the Newtonian framework.

Of course, for astrometric problems one always distinguishes between the two celestial systems and apparent places of stars from true (barycentric) places. However, annual parallax and aberration were merely understood as correction terms that have to be applied to get the “true” positions for the realization of the astronomical quasi-inertial, space-fixed celestial system. Note that the definition of the classical astronomical (α, δ) system uses concepts from both systems: some ecliptic from the barycentric celestial system, and some Earth rotation pole, the Celestial Ephemeris Pole or Celestial Intermediate Pole, and its corresponding equator from the geocentric celestial system.

In *relativity theory*, the situation is more complicated. Even in the absence of gravitational fields and a uniformly moving geocenter, the two coordinate systems are related by a four-dimensional Lorentz transformation from special relativity. In our solar system, BCRS and GCRS coordinates are related by a complicated four-dimensional spacetime transformation (a generalized Lorentz transformation) that also contains acceleration terms and gravitational potentials. This implies that the two astronomical reference

systems, the BCRS and the GCRS, are actually quite different. This has profound consequences for many classical astronomical concepts.

The BCRS is the basic astrometric celestial reference system. Usually one considers the solar system to be isolated—that is, one ignores all matter and fields outside the system and assumes that the gravitational potentials vanish far from the system. It is obvious that ignoring the Galaxy and extragalactic objects is an unphysical idealization for several specific questions (which, however, will not be touched upon here). If the solar system is considered to be isolated, we might follow light rays from some very remote source back in time to the region $|\mathbf{x}| \rightarrow \infty$, which might be called the celestial sphere. In the vicinity of the celestial sphere, a certain light ray defines spherical angles that might appear as catalog values. Actually, for reference stars the physical distance from Earth usually plays a role. In that case we might associate any star with a corresponding BCRS coordinate position \mathbf{x}_* , which will be a function of TCB. From this position, vector spherical angles (α_*, δ_*) can be introduced in a very simple manner by

$$\frac{\mathbf{x}_*}{|\mathbf{x}_*|} = \begin{pmatrix} \cos \alpha_* \cos \delta_* \\ \sin \alpha_* \cos \delta_* \\ \sin \delta_* \end{pmatrix}, \quad (7)$$

which can be considered as catalog values. If the coordinate distance of some source tends to infinity, the two constructions for an astrometric position will coincide. From $x_*(t)$, quantities such as “proper motion” and “radial velocity” can be defined as coordinate quantities in the BCRS. Note that the problem of “radial velocity” has exhaustively been discussed by Lindegren & Dravins (2003; see also IAU 2000 Resolutions C1 and C2 [Rickman 2001]). Other fields of application of the BCRS are solar system ephemerides, interplanetary navigation, etc.

The definition of the BCRS given by IAU Resolution B1.3 (2000) does not fix the orientation of the spatial axes uniquely but only up to some constant, time-independent rotation matrix about the origin. One natural choice of orientation is provided by the International Celestial Reference System (ICRS). Actually, for the construction of the International Celestial Reference Frame (ICRF) and its optical counterpart, the *Hipparcos* Catalogue, the recommended form of the barycentric metric tensor has already been used explicitly in the underlying models. This implies that a set of definitions that completely fix the ICRS contains the BCRS definitions.

There might be other useful possibilities for the orientation of barycentric spatial coordinates. One possibility is an orientation according to some ecliptic \mathcal{E}_0 at a certain epoch t_0 defined by corresponding solar system ephemerides. Such an ecliptic would coincide with the x - y plane of a BCRS(\mathcal{E}_0), which might be useful for reasons of historical continuity.

On the other hand, quantities and concepts related to the physics in the immediate vicinity of Earth should be formulated in the GCRS. This concerns the gravity field of Earth itself and satellite theory and especially applies to theories of Earth’s rotation and their parameters. Clearly, the spatial GCRS coordinates \mathbf{X} can be used to define corresponding unit vectors at the geocenter, which might be employed to compute spherical angles $(\alpha_{\text{GCRS}}, \delta_{\text{GCRS}})$ that might be

called “geocentric places.” Note, however, that the coordinates of the remote astronomical sources are defined in the BCRS only. The calculated GCRS places $(\alpha_{\text{GCRS}}, \delta_{\text{GCRS}})$ are determined by incident light rays at the geocenter. They differ from corresponding ICRS (α, δ) -values because of annual aberration, annual parallax, and gravitational light deflection due to the gravitational fields of the solar system bodies (apart from Earth) and are independent of Earth’s rotation. Whether these GCRS places, however, will ever play a role in practice is not clear.

In the past, apparent places of stars that were annually published, for example, in the “Apparent Places of Fundamental Stars” played a role for certain problems. These places are related to the old, traditional astronomical reference system, that is, with some equator and equinox of date. Now with the ICRS, we have a highly precise astronomical reference system that is basically independent of Earth’s rotation parameters and their determination. For several applications, however, the introduction of quantities such as apparent places might still be useful, especially if there is a reference to the local plumb line—that is, to the zenith—and the astronomical (or nautical) triangle can be employed. In that case Resolutions B1.7 and B1.8 (IAU 2001; Rickman 2001) come into play. These two resolutions define some intermediate system that can be used for the definition of an intermediate position $(\alpha_{\text{inter}}, \delta_{\text{inter}})$ by the Celestial Intermediate Pole and the Celestial Ephemeris Origin. Such an intermediate position can be considered a modern version of the apparent place, defined in the GCRS.

For astrometry at microarcsecond accuracies, neither GCRS places nor intermediate places likely will play a role. To avoid problems related to nonlinearities, it is simpler to use an overall BCRS picture to describe not only the light rays and the motion of gravitating bodies, but also the trajectory of an observer. In that case, only catalog and observed positions will be of importance.

3.2. The Barycentric Celestial Reference System

Resolution B1.3 concerns the definition of the Barycentric Celestial Reference System and the Geocentric Celestial Reference System. The BCRS is defined with coordinates $(ct, \mathbf{x}^i) = x^\mu$, where $t = \text{TCB}$. The BCRS is a particular version of the barycentric reference system of the solar system. The resolution recommends that the metric tensor of the BCRS be written in the form

$$\begin{aligned} g_{00} &= -1 + \frac{2w}{c^2} - \frac{2w^2}{c^4} + O(c^{-5}), \\ g_{0i} &= -\frac{4}{c^3} w^i + O(c^{-5}), \\ g_{ij} &= \delta_{ij} \left(1 + \frac{2}{c^2} w \right) + O(c^{-4}). \end{aligned} \quad (8)$$

A comparison reveals that this form of the metric presents an extension of equation (2). Whereas the old form contains only the Newtonian potential U , the new one contains a scalar potential w and a vector potential w^i .

Actually, the equations for g_{00} and g_{0i} from equation (8) without the order symbols $O(c^{-5})$ are always correct and can simply be considered definitions of w and w^i in terms of g_{00} and g_{0i} . In contrast to the concrete form of the resolution, we have added order symbols in equation (8). For example, for g_{00} the order symbol indicates that terms of order c^{-5} will

systematically be neglected, as stated in the notes to the resolution. With these forms for g_{00} and g_{0i} , one finds that spatially isotropic coordinates x^i exist such that g_{ij} from equation (8) with the potential w from g_{00} solves Einstein's field equations to first post-Newtonian order. Note that the form of equation (8) implies that the barycentric spatial coordinates x^i satisfy the harmonic gauge condition (see, e.g., Brumberg & Kopeikin 1989; Damour et al. 1991). At this point, because of the freedom in the time coordinate, many different "time gauge conditions" are still possible. The resolution proceeds by recommending a specific kind of space and time *harmonic gauge*. One argument in favor of the harmonic gauge is that tremendous work on general relativity has been done with the harmonic gauge, which was found to be a useful and simplifying gauge for many kinds of applications. Moreover, the harmonic gauge condition,

$$g^{\mu\nu}\Gamma_{\mu\nu}^{\lambda} = 0 \quad (9)$$

(e.g., Weinberg 1972; Fok 1959), where $\Gamma_{\mu\nu}^{\lambda}$ are the Christoffel symbols of the metric tensor, is not restricted to some post-Newtonian approximation but can be defined in Einstein's theory of gravity without any approximations. This may be important for future refinements of the IAU framework. With the harmonic gauge condition, the post-Newtonian Einstein field equations take the form

$$\left(-\frac{1}{c^2}\frac{\partial^2}{\partial t^2} + \nabla^2\right)w = -4\pi G\sigma + O(c^{-4}), \quad (10)$$

$$\nabla^2 w^i = -4\pi G\sigma^i + O(c^{-2}). \quad (11)$$

Here σ and σ^i are the gravitational mass and mass current density, respectively. Mathematically they are related to the *energy-momentum tensor* $T^{\mu\nu}$ by

$$\sigma = \frac{1}{c^2}(T^{00} + T^{ss}), \quad \sigma^i = \frac{1}{c}T^{0i}. \quad (12)$$

The energy-momentum tensor $T^{\mu\nu}$ generalizes the density ρ of the Poisson equation (eq. [3]). In relativity, energy density, pressure, and stresses all act as sources of the gravitational field. This implies that different kinds of energy contribute to the gravitational sources—kinetic energy, gravitational potential energy, energy of deformation, etc. Since the kinetic energy depends upon the state of motion of the matter, the energy-momentum tensor, which really acts as the gravitational source, exhibits nontrivial transformation behavior if we go from one reference system to another. In practice, however, the energy-momentum tensor will usually not appear explicitly. This is because the gravitational potentials w and w^i from equations (10)–(11) are completely determined by σ and σ^i , which can be considered primary quantities. If we deal with problems in which gravitational fields play a role only *outside* of astronomical bodies and admit a useful convergent expansion in terms of multipole moments (potential coefficients), only corresponding integral characteristics of the bodies such as masses and quadrupole moments show up explicitly, which are defined in terms of σ and σ^i and whose numerical values will be fixed by observations. Because of equation (11), w^i is sometimes called the gravitomagnetic potential, since it results from mass currents (moving or rotating masses) just as the electromagnetic vector potential results from electric currents in Maxwell's theory of electromagnetism.

Equation (10) generalizes the Poisson equation (eq. [3]), and hence the scalar potential w presents a relativistic generalization of the Newtonian potential U . Because of problems related to homogeneous solutions and boundary conditions, mathematically it is clear that these differential equations do not fix the harmonic solutions uniquely. Assuming spacetime to be asymptotically flat (no gravitational fields far from the system), that is,

$$\lim_{\substack{r \rightarrow \infty \\ t = \text{const}}} g_{\mu\nu} = \text{diag}(-1, +1, +1, +1), \quad (13)$$

the recommended solution reads

$$w(t, \mathbf{x}) = G \int d^3x' \frac{\sigma(t, \mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|} + \frac{1}{2c^2} G \frac{\partial^2}{\partial t^2} \int d^3x' \sigma(t, \mathbf{x}') |\mathbf{x} - \mathbf{x}'|, \quad (14)$$

$$w^i(t, \mathbf{x}) = G \int d^3x' \frac{\sigma^i(t, \mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|}. \quad (15)$$

It is obvious that the second time derivative term in equation (14) results from the corresponding operator in the field equation (eq. [10]). This operator modifies the Laplacian from the Newtonian Poisson equation to the d'Alembertian, and the similarity between the harmonic post-Newtonian field equations and Maxwell's equations of electromagnetism in the Lorentz gauge becomes obvious (actually, one might replace the Laplacian by the d'Alembertian in eq. [11] to post-Newtonian accuracy). From Maxwell's theory, it is well known that the retarded potential solves the corresponding field equation:

$$w_{\text{ret}}(t, x^i) = G \int d^3x' \frac{\sigma(t_{\text{ret}}, \mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|} \quad (16)$$

with

$$t_{\text{ret}} = t - |\mathbf{x} - \mathbf{x}'|/c. \quad (17)$$

One might then expand the retarded potential in terms of $1/c$. Note that such an expansion also yields a term proportional to $1/c$. If we stay within the first post-Newtonian approximation, these $1/c$ terms vanish as a consequence of the Newtonian mass conservation law. Such odd powers of $1/c$ indicate time-asymmetric terms, that is, they break time-reversal symmetry. It is well known that such time-asymmetric terms appear only to higher post-Newtonian order, and they will not be considered here. For that reason, the retarded potential (eq. [16]) leads to the recommended solution above.

Comparing the form of the metric tensor in equation (8) with other forms that can be found in the literature (e.g., Will 1993), one might get the erroneous impression that something is missing in equation (8), which is not the case. If matter is described by some fluid model, then formally (w, w^i) might be split into various pieces resulting from kinetic energy, gravitational potential energy, specific internal energy density, pressure, etc., and the equivalence of our form of the metric tensor, for example, to that given by Will (1993) can be shown. This is explicitly demonstrated in Appendix B.

The point, however, is that a split of (σ, σ^i) of our metric potentials (w, w^i) or of the metric tensor itself into various pieces is usually unnecessary. If only gravitational

fields outside the relevant bodies play a role (as is typically the case in celestial mechanics and astrometry), then it is advantageous to keep such pieces together, since it will be the sum that determines the observables. One might argue that U is the “Newtonian potential” and the rest can be identified as “relativistic corrections.” This way of thinking, however, can be very misleading and presents a source of errors. As has been shown in the literature (e.g., Damour et al. 1991, 1993), suitably defined potential coefficients based upon w (not U) and w^i can be introduced that can be determined from satellite data. From a more theoretical point of view, the introduction of (w, w^i) has the advantage that the field equations (eqs. [10]–[11]) are formally *linear*, although the corresponding metric is not (because of the w^2 term). We used the word “formally” because σ depends upon w implicitly. This nonlinearity has been explicitly treated, for example, by Brumberg & Kopeikin (1989), but this dependence becomes irrelevant in practice if the fields outside of some matter distribution are parameterized by means of potential coefficients. This linearity implies that for an ensemble of N bodies,

$$w(t, \mathbf{x}) = \sum_{A=1}^N w_A(t, \mathbf{x}), \quad w^i(t, \mathbf{x}) = \sum_{A=1}^N w_A^i(t, \mathbf{x}), \quad (18)$$

with the index A indicating the contribution related to body A , where the integrals have to be taken over the support of body A only. This linearity, however, does not imply that body-body interaction terms have been neglected. If written explicitly, w_A will in general contain contributions from bodies $B \neq A$ (see, e.g., eq. [54]).

The BCRS metric tensor from IAU Resolution B1.3 (2000) extends the form of the metric tensor given in the IAU 1991 Resolutions such that its accuracy is sufficient for most applications in the coming years. Note that an extension of the old metric (eq. [2]) is necessary (and has been in use for decades) for the derivation of the relativistic equations that form the basis of any modern solar system ephemeris (such as the JPL DE ephemerides). Resolution B1.3 formalizes this extension.

3.3. The Geocentric Celestial Reference System

Resolution B1.3 goes on to define the GCRS, which represents a particular version of the local geocentric reference system for Earth. Its spatial coordinates X^a are kinematically nonrotating with respect to the barycentric ones (see, e.g., Brumberg & Kopeikin 1989; Klioner & Soffel 1998). The geocentric coordinates are denoted by (T, \mathbf{X}) , where $T = \text{TCG}$. In the relation between x^i and X^a from Resolution B1.3, let us replace the unit matrix δ_{ai} by a general rotation matrix R_{ai} :

$$X^a = R_{ai} \left[r_E^i + \frac{1}{c^2} (\dots) \right] + O(c^{-4}),$$

where $\mathbf{r}_E = \mathbf{x} - \mathbf{x}_E$. If the two sets of spatial coordinates are aligned for all times, that is, if $R_{ai} = \delta_{ai}$ as is the case for the GCRS spatial coordinates, then X^a is defined to be kinematically nonrotating with respect to the barycentric spatial coordinates x^i . The resolution recommends writing the metric tensor of the GCRS in the same form as the barycentric

one but with potentials $W(T, \mathbf{X})$ and $W^a(T, \mathbf{X})$. Explicitly,

$$\begin{aligned} G_{00} &= -1 + \frac{2W}{c^2} - \frac{2W^2}{c^4} + O(c^{-5}), \\ G_{0a} &= -\frac{4}{c^3} W^a + O(c^{-5}), \\ G_{ab} &= \delta_{ab} \left(1 + \frac{2}{c^2} W \right) + O(c^{-4}), \end{aligned} \quad (19)$$

and the geocentric field equations formally look the same as the barycentric ones (eqs. [10]–[11]) but with all variables referred to the GCRS. Again, one decisive advantage of this recommendation is the formal linearity of the field equations. This linearity admits a unique split of the geocentric metric into a part coming from Earth itself and a remaining part resulting from inertial and tidal forces. Therefore, it is recommended to split the potentials W and W^a according to

$$\begin{aligned} W(T, \mathbf{X}) &= W_E(T, \mathbf{X}) + W_{\text{ext}}(T, \mathbf{X}), \\ W^a(T, \mathbf{X}) &= W_E^a(T, \mathbf{X}) + W_{\text{ext}}^a(T, \mathbf{X}). \end{aligned} \quad (20)$$

Earth’s potentials W_E and W_E^a are defined in the same way as w_E and w_E^a (i.e., eqs. [14]–[15] with integrals taken over the volume of the whole Earth) but with quantities calculated in the GCRS. Outside Earth the potentials (W, W^a) admit a power series expansion in terms of $R \equiv |\mathbf{X}|$, and all negative powers of R are contained in W_E and W_E^a . For that reason, Earth’s potentials admit multipole expansions that look very similar to the Newtonian ones. This point will be discussed below in more detail.

It is useful to split the external potentials W_{ext} and W_{ext}^a further. They can be written in the form

$$W_{\text{ext}} = W_{\text{tidal}} + W_{\text{iner}}, \quad W_{\text{ext}}^a = W_{\text{tidal}}^a + W_{\text{iner}}^a, \quad (21)$$

where the tidal terms are at least quadratic in X^a and the inertial contributions W_{iner} and W_{iner}^a are just linear in X^a . Explicitly,

$$W_{\text{iner}} = Q_a X^a, \quad W_{\text{iner}}^a = -\frac{1}{4} c^2 \epsilon_{abc} \Omega_{\text{iner}}^b X^c. \quad (22)$$

Mathematically, the Q_a term is related to the 4-acceleration of the geocenter in the external gravitational field, a quantity that vanishes for a purely spherical and nonrotating Earth (for a mass monopole, more precisely) that moves along a geodesic in the external gravitational field. The Q_a term therefore results from the coupling of higher order multipole moments of Earth to the external tidal gravitational fields (to the external curvature tensor of spacetime, in mathematical language). The quantity Q_a characterizes the deviation of the actual worldline of the origin of the GCRS from a geodesic in the external gravitational field. With

$$w_{\text{ext}}(t, \mathbf{x}) = \sum_{A \neq E} w_A(t, \mathbf{x}), \quad w_{\text{ext}}^i(t, \mathbf{x}) = \sum_{A \neq E} w_A^i(t, \mathbf{x}),$$

to Newtonian order Q_a is given by

$$Q_a = \delta_{ai} \left[\frac{\partial}{\partial x^i} w_{\text{ext}}(\mathbf{x}_E) - a_E^i \right]. \quad (23)$$

Here $x_E^i(t)$, $v_E^i(t) = dx_E^i/dt$, and $a_E^i = dv_E^i/dt$ are respectively the barycentric coordinate position, velocity, and acceleration of the origin of the GCRS (geocenter). The appearance

of δ_{ai} results from the fact that the GCRS is defined as kinematically nonrotating with respect to the BCRS. We retain δ_{ai} in the transformations here and below because of a desire to distinguish between BCRS quantities (spatial indices taken from the second part of the Latin alphabet, starting with the letter i) and GCRS quantities (spatial indices taken from the first part of the alphabet).

The full post-Newtonian expression for Q_a [denoted $G_a(T)$ in the Damour-Soffel-Xu papers] can be derived from equation (6.30a) of Damour et al. (1991). To get an idea about orders of magnitude, the absolute value of Q_a due to the action of the Moon is on the order of $4 \times 10^{-11} \text{ m s}^{-2}$ (Kopeikin 1991).

The term W_{iner}^a describes a relativistic Coriolis force due to the rotation of the GCRS with respect to a dynamically nonrotating geocentric reference system. Such a rotation has several components, often referred to as geodetic, Lense-Thirring, and Thomas precessions:

$$\mathbf{\Omega}_{\text{iner}} = \mathbf{\Omega}_{\text{GP}} + \mathbf{\Omega}_{\text{LTP}} + \mathbf{\Omega}_{\text{TP}} \quad (24)$$

with

$$\begin{aligned} \mathbf{\Omega}_{\text{GP}} &= -\frac{3}{2c^2} \mathbf{v}_E \times \nabla w_{\text{ext}}(\mathbf{x}_E), \\ \mathbf{\Omega}_{\text{LTP}} &= -\frac{2}{c^2} \nabla \times \mathbf{w}_{\text{ext}}(\mathbf{x}_E), \\ \mathbf{\Omega}_{\text{TP}} &= -\frac{1}{2c^2} \mathbf{v}_E \times \mathbf{Q}, \end{aligned} \quad (25)$$

in obvious notation. As a relativistic precession, the geodetic precession $\mathbf{\Omega}_{\text{GP}}$ is proportional to $1/c^2$. It is also proportional to the barycentric coordinate velocity v_E and the gradient of the external gravitational scalar potential w_{ext} at the geocenter (the barycentric coordinate acceleration of the geocenter, to sufficient accuracy). The order of magnitude is given by

$$\begin{aligned} |\mathbf{\Omega}_{\text{GP}}| &\sim 1.5 \left(\frac{v_E}{c} \right) \left(\frac{GM_\odot}{c^2 \text{ AU}} \right) \left(\frac{c}{1 \text{ AU}} \right) \\ &\sim 3 \times 10^{-15} \text{ s}^{-1} \sim 2'' \text{ per century}. \end{aligned}$$

Thomas precession is also proportional to $1/c^2$ and the barycentric coordinate velocity of the geocenter, but to the geodetic deviation term Q_a as well. The order of magnitude of Thomas precession is $|\mathbf{\Omega}_{\text{TP}}| \sim 0.5(v_E/c)|\mathbf{Q}|/c \sim 7 \times 10^{-24} \text{ s}^{-1} \sim 4 \times 10^{-9} \text{ arcseconds per century}$, that is, negligible with respect to geodetic precession.

Finally, the Lense-Thirring precession results from the gradient of the external gravitomagnetic potential at the geocenter. If we consider some spherically symmetric solar system body A , then its gravitomagnetic potential W_A^a is given by

$$W_A^a = \frac{G}{2} \frac{(\mathbf{S}_A \times \mathbf{X})^a}{R^3}$$

in its own local rest frame (see eq. [49] below). Transformation into the BCRS according to the rule indicated below in equation (31) leads to

$$w_A^i(t, \mathbf{x}) = G \left[\frac{(\mathbf{S}_A \times \mathbf{r}_A)^i}{2r_A^3} + \frac{M_A}{r_A} v_A^i \right],$$

where $\mathbf{r}_A \equiv \mathbf{x} - \mathbf{x}_A$ and \mathbf{v}_A is the barycentric velocity of

body A . In our case, the spin and motion of the Sun and Moon will provide the dominant contributions to $\mathbf{\Omega}_{\text{LTP}}$: $|\mathbf{\Omega}_{\text{LTP}}| \sim 2 \times 10^{-3} \text{ arcseconds per century}$.

The definition of the GCRS implies that the spatial GCRS coordinates \mathbf{X} are kinematically nonrotating with respect to the BCRS ones, \mathbf{x} (as indicated by the δ_{ai} term in Resolution B1.3). Because of geodetic precession, locally inertial coordinates precess with respect to the GCRS by an amount $|\mathbf{\Omega}_{\text{iner}}| = 1''.9198$ per century (Brumberg, Bretagnon, & Francou 1991). Let us forget about the mass of Earth and imagine a torque-free gyroscope at the geocenter, moving along the actual trajectory of the geocenter. It will precess by this amount in our GCRS. Since the GCRS does not present a locally inertial reference system, Coriolis forces caused by geodetic precession and nutation appear in every GCRS dynamical equation of motion, for example, that of Earth's satellites. As recommended in the IERS Conventions (2003) (McCarthy & Petit 2003), these additional forces should be taken into account. Moreover, geodetic precession-nutation has to be considered in the precession-nutation model formulated in the GCRS. For example, the basic post-Newtonian equation of Earth's intrinsic angular momentum \mathbf{S} reads

$$\frac{d\mathbf{S}}{dT} + \mathbf{\Omega}_{\text{iner}} \times \mathbf{S} = \mathbf{D}, \quad (26)$$

where \mathbf{D} is the external torque (Damour et al. 1993). As long as observations of Earth's orientation parameters are referred to the GCRS, they will contain geodetic precession-nutation automatically.

Because of the eccentricity of Earth's orbit, the leading term in $\mathbf{\Omega}_{\text{GP}}$ has an annual and a semiannual part that lead to geodetic nutation in longitude with

$$\Delta\psi_{\text{GP}} = 0.153 \sin l' + 0.002 \sin 2l', \quad (27)$$

where the amplitudes are in milliarcseconds and l is the mean anomaly of the Earth-Moon barycenter (Fukushima 1991; Brumberg et al. 1991; Bois & Vokrouhlický 1995).

The quantity W_{tidal} is a generalization of the Newtonian tidal potential

$$W_{\text{tidal}}^{\text{Newton}}(T, \mathbf{X}) = w_{\text{ext}}(\mathbf{x}_E + \mathbf{X}) - w_{\text{ext}}(\mathbf{x}_E) - \mathbf{X} \cdot \nabla w_{\text{ext}}(\mathbf{x}_E). \quad (28)$$

Full post-Newtonian expressions for W_{tidal} and W_{tidal}^a can be found in Damour et al. (1992). There W_{ext} is denoted \bar{W} , and a tidal expansion in powers of local spatial coordinates by means of suitably defined tidal moments is given in equation (4.15) of that paper. Expressions for W_{tidal} and W_{tidal}^a in closed form are given in Klioner & Voinov (1993). The quadratic term, which is the dominant term in the expansion of W_{tidal} , reads

$$W_{\text{tidal}} \Big|_{l=2} = \frac{1}{2} G_{ab}^{\text{tidal}} X^a X^b. \quad (29)$$

If the external bodies are taken to be mass monopoles, the explicit expression for G_{ab}^{tidal} (not to be confused with the GCRS metric tensor) is given by equation (3.23) of Damour et al. (1994). Higher order terms in this approximation can be found in Klioner et al. (2003).

Finally, the local gravitational potentials W_E and W_E^a of Earth are related to the barycentric gravitational potentials

w_E and w_E^i by

$$W_E(T, \mathbf{X}) = w_E(t, \mathbf{x}) \left(1 + \frac{2}{c^2} v_E^2 \right) - \frac{4}{c^2} v_E^i w_E^i(t, \mathbf{x}) + O(c^{-4}),$$

$$W_E^a(T, \mathbf{X}) = \delta_{ia}^a [w_E^i(t, \mathbf{x}) - v_E^i w_E(t, \mathbf{x})] + O(c^{-2}), \quad (30)$$

where $\delta_i^a = \delta_a^i = \delta_{ai}$, or by the inverse transformation

$$w_E(t, \mathbf{x}) = W_E(T, \mathbf{X}) \left(1 + \frac{2}{c^2} v_E^2 \right) + \frac{4}{c^2} \delta_{ia} v_E^i W_E^a(T, \mathbf{X}) + O(c^{-4}),$$

$$w_E^i(t, \mathbf{x}) = \delta_a^i W_E^a(T, \mathbf{X}) + v_E^i W_E(T, \mathbf{X}) + O(c^{-2}). \quad (31)$$

The relations between the geocentric gravitational potentials W and W^a and the barycentric ones w and w^i follow from the coordinate transformations between the BCRS and GCRS discussed below.

3.4. Coordinate Transformations

The metric tensors in the BCRS and GCRS allow one to derive the rules for the transformations between the BCRS coordinates x^μ and the GCRS ones X^α from the tensorial transformation rules. It is obvious that these transformations can be written in two equivalent forms: as $x^\mu(T, X^a)$ or as $X^\alpha(t, x^i)$. Whereas the first form was used in the Damour-Soffel-Xu formalism (Damour et al. 1991, 1992, 1993, 1994), the second one was presented in the Brumberg-Kopeikin formalism (Brumberg & Kopeikin 1989; Kopeikin 1988; Brumberg 1991; Klioner & Voinov 1993). It should be pointed out that the transformation from one version to the other is not trivial, because of the barycentric coordinate position of the geocenter, which appears in the first form as function of TCG and as function of TCB in the second one. In Resolution B1.3, $T = \text{TCG}$ and X^a are presented as functions of $t = \text{TCB}$ and x^i . The explicit form of the transformations is given in the text of Resolution B1.3 (see Appendix A). Apart from the terms of order $|\mathbf{X}|^3$ that appear in the $O(c^{-4})$ time transformation, all terms can be obtained from the results derived by Kopeikin (1988) and Damour et al. (1991). The cubic and higher order terms in $|\mathbf{X}|$ as represented by the function C in Resolution B1.3 have been derived by Kopeikin (1988) and are analyzed in full detail in Klioner & Voinov (1993). As is also clear from Klioner & Voinov (1993), the expression for C is not unique but only constrained by the gauge and field equations, so that the simplest possibility is an expression for C containing cubic terms only. It is this simplest expression that is recommended in Resolution B1.3.

The full four-dimensional coordinate transformation is just an extension of the usual Lorentz transformation. Indeed, if we neglect all gravitational fields and acceleration terms, then the coordinate transformation in Resolution B1.3 can be written in the form

$$T = t \left(1 - \frac{\beta^2}{2} - \frac{\beta^4}{8} \right) - \left(1 + \frac{\beta^2}{2} \right) \frac{\mathbf{v} \cdot \mathbf{r}}{c^2} + O(c^{-6}),$$

$$\mathbf{X} = \mathbf{r} + \frac{1}{2} (\mathbf{v} \cdot \mathbf{r}) \frac{\mathbf{v}}{c^2} + O(c^{-4}), \quad (32)$$

where $\mathbf{r} = \mathbf{x} - \mathbf{x}_E(t)$ and $\beta = v/c = \text{const}$. If we now write

$\mathbf{x}_E(t) = \mathbf{v}t$, we obtain

$$T = t \left(1 + \frac{\beta^2}{2} + \frac{3\beta^4}{8} \right) - \left(1 + \frac{\beta^2}{2} \right) \frac{\mathbf{v} \cdot \mathbf{x}}{c^2} + O(c^{-6}),$$

$$\mathbf{X} = \mathbf{x} - \left(1 + \frac{\beta^2}{2} \right) \mathbf{v}t + \frac{1}{2} (\mathbf{v} \cdot \mathbf{x}) \frac{\mathbf{v}}{c^2} + O(c^{-4}), \quad (33)$$

which is nothing but a Lorentz transformation from special relativity theory,

$$T = \gamma \left(t - \frac{\mathbf{v} \cdot \mathbf{x}}{c^2} \right), \quad \mathbf{X} = \mathbf{x} - \gamma \mathbf{v}t + \frac{\gamma - 1}{v^2} (\mathbf{v} \cdot \mathbf{x}) \mathbf{v}$$

in the corresponding approximation, since

$$\gamma \equiv (1 - \beta^2)^{-1/2} = 1 + \frac{1}{2} \beta^2 + \frac{3}{8} \beta^4 + O(c^{-6}).$$

Note that the inverse transformations are obtained simply by replacing (t, \mathbf{x}) with (T, \mathbf{X}) and the velocity \mathbf{v} with $-\mathbf{v}$.

Neglecting the $1/c^4$ terms in the T - t relation given in Resolution B1.3, one obtains

$$T = t - \frac{1}{c^2} \left\{ \int_{t_0}^t \left[\frac{v_E^2}{2} + w_{\text{ext}}(\mathbf{x}_E) \right] dt + v_E^i x_E^i \right\} + O(c^{-4}), \quad (34)$$

which reduces to the old recommendation (eq. [4]), since $t = \text{TCB}$, $T = \text{TCG}$, and $w_{\text{ext}}(\mathbf{x}_E(t))$ reduces to $U_{\text{ext}}(t, \mathbf{x}_E(t))$ in the Newtonian limit. A more accurate version of this transformation will be discussed below.

Let us also note that the BCRS, the GCRS, and the transformation between them have been discussed by Klioner & Soffel (2000) in the framework of the PPN formalism, with parameters β and γ . For the limit of general relativity, $\beta = \gamma = 1$, all of the formulae given in that publication become equal to those derived in the framework of the new IAU resolutions, which refer solely to Einstein's theory of gravity.

3.5. Potential Coefficients

3.5.1. General Post-Newtonian Multipole Moments

For many problems it is advantageous to present the local gravitational potentials of Earth as multipole series that usually converge outside Earth. To this end one, has to introduce a certain set of multipole moments or *potential coefficients* for Earth. A certain set of potential coefficients, called Blanchet-Damour (B-D) moments (Blanchet & Damour 1989; Damour et al. 1991), defined to first post-Newtonian order, have especially attractive features. Moreover, by using such B-D moments we obtain a very simple form for the multipole expansion of the post-Newtonian potentials (these expansions have almost Newtonian form). Basically, two sets of B-D moments occur in the formalism: mass multipole moments and spin multipole moments. Theoretically, these moments can be derived from the distribution of mass and matter currents inside the body, but for an observer they simply present parameters that can be directly estimated from observations.

Expressed in terms of symmetric and trace-free Cartesian tensors, the B-D moments are denoted \mathcal{M}_L and \mathcal{S}_L . Here L is a multi-index of l different indices all taking the values 1, 2, and 3; that is, $L = i_1 i_2 \dots i_l$ and every index $i = (1, 2, 3)$. Explicit expressions for \mathcal{M}_L and \mathcal{S}_L as integrals over Earth can be found in, for example, Blanchet & Damour (1989)

and Damour et al. (1991):

$$\begin{aligned} \mathcal{M}_L(T) \equiv & \int d^3X \hat{X}^L \Sigma \\ & + \frac{1}{2(2l+3)c^2} \frac{d^2}{dT^2} \left(\int d^3X \hat{X}^L X^2 \Sigma \right) \\ & - \frac{4(2l+1)}{(l+1)(2l+3)c^2} \frac{d}{dT} \left(\int d^3X \hat{X}^{aL} \Sigma^a \right) \end{aligned} \quad (l \geq 0), \quad (35)$$

$$\mathcal{S}_L(T) \equiv \int d^3X \epsilon^{abc} \hat{X}^{L-1a} \Sigma^b \quad (l \geq 1), \quad (36)$$

where the integrations extend over the body under consideration and

$$\Sigma(T, \mathbf{X}) = \frac{1}{c^2} (\mathcal{T}^{00} + \mathcal{T}^{ss}), \quad \Sigma^a(T, \mathbf{X}) = \frac{1}{c} \mathcal{T}^{0a}. \quad (37)$$

Here the $\mathcal{T}^{\mu\nu} = \mathcal{T}^{\mu\nu}(T, X^a)$ are components of the energy-momentum tensor in the GCRS. Both the caret and the angle brackets indicate the symmetric and trace-free (STF) part of the object or of the indices enclosed by the brackets (see, e.g., Damour et al. Xu 1991, p. 3277, for an explicit definition of the STF part of an object). Some basic information on the operations with STF objects can be found in, for example, Blanchet & Damour (1989) and Damour et al. (1992).

For practical applications, however, the explicit form of \mathcal{M}_L and \mathcal{S}_L will not be needed, since these quantities are parameters characterizing the gravitational field of the corresponding body that are fitted to observations. The set \mathcal{M}_L is equivalent to the set of potential coefficients C_{lm} and S_{lm} that appear in the much more familiar spherical harmonic expansion of W_E . The first nonvanishing spin moment (the spin dipole) of a body agrees with its spin vector (total intrinsic angular momentum). The multipole expansion of W_E and W_E^a reads

$$\begin{aligned} W_E = & G \sum_{l=0}^{\infty} \frac{(-1)^l}{l!} \left(\mathcal{M}_L \partial_L \frac{1}{|\mathbf{X}|} + \frac{1}{2c^2} \ddot{\mathcal{M}}_L \partial_L |\mathbf{X}| \right) + \frac{4}{c^2} \Lambda_{,T} \\ & + O(c^{-4}), \end{aligned} \quad (38)$$

$$\begin{aligned} W_E^a = & -G \sum_{l=1}^{\infty} \frac{(-1)^l}{l!} \left(\dot{\mathcal{M}}_{aL-1} \partial_{L-1} \frac{1}{|\mathbf{X}|} \right. \\ & \left. + \frac{l}{l+1} \epsilon_{abc} \mathcal{S}_{cL-1} \partial_{bL-1} \frac{1}{|\mathbf{X}|} \right) \\ & - \Lambda_{,a} + O(c^{-2}), \end{aligned} \quad (39)$$

where

$$\Lambda = G \sum_{l=0}^{\infty} \frac{(-1)^l}{(l+1)!} \frac{2l+1}{2l+3} \mathcal{P}_L \partial_L \frac{1}{|\mathbf{X}|}, \quad (40)$$

$$\mathcal{P}_L = \int_V \Sigma^a \hat{X}^{aL} d^3X. \quad (41)$$

Here an overdot stands for $\partial/\partial T$, and ∂_L stands for $\partial^l/\partial x^{i_1} \dots \partial x^{i_l}$. The subscripted commas denote partial differentiation: $\Lambda_{,T} \equiv \partial\Lambda/\partial T$ and $\Lambda_{,a} \equiv \partial\Lambda/\partial X^a$.

The gauge function Λ does not enter the post-Newtonian equations of motion. The latter contains only the B-D

multipole moments \mathcal{M}_L and \mathcal{S}_L . The only place where the function Λ should be accounted for is in the transformation between the various time scales. However, these gauge terms are of order c^{-4} in the metric tensor, so for the problem of clock rates they are basically of second post-Newtonian order. These terms are much less than 10^{-18} in the geocentric metric tensor and will be neglected. For that reason the Λ -terms are not mentioned in Resolution B1.4.

3.5.2. Approximate Expansion of the Scalar Gravitational Potential

A spherical harmonic expansion of W_E equivalent to equation (38) without the Λ -term reads

$$\begin{aligned} W_E(T, \mathbf{X}) &= \frac{GM_E}{R} \left\{ 1 + \sum_{l=2}^{\infty} \sum_{m=0}^{+l} \left(\frac{R_E}{R} \right)^l P_{lm}(\cos\theta) \right. \\ &\quad \left. \times [\mathcal{C}_{lm}(T, R) \cos m\phi + \mathcal{S}_{lm}(T, R) \sin m\phi] \right\} \\ &\quad + O(c^{-4}) \end{aligned} \quad (42)$$

with $R = |\mathbf{X}|$ and

$$\mathcal{C}_{lm}^E(T, R) = C_{lm}^E(T) - \frac{1}{2(2l-1)} \frac{R^2}{c^2} \frac{d^2}{dT^2} C_{lm}^E(T), \quad (43)$$

$$\mathcal{S}_{lm}^E(T, R) = S_{lm}^E(T) - \frac{1}{2(2l-1)} \frac{R^2}{c^2} \frac{d^2}{dT^2} S_{lm}^E(T). \quad (44)$$

Let us stress that as stated in Resolution B1.4, $C_{lm}^E(T)$ and $S_{lm}^E(T)$ refer to the GCRS coordinates and are related approximately constant potential coefficients in a terrestrial system that is rotating with Earth (i.e., those from an Earth model) by time-dependent transformations. For a rigid, axially symmetric body rotating about its symmetry axis with angular velocity Ω_E , the second time derivative terms will vanish. Let us estimate these terms for Earth. From the order of magnitude of the $l = m = 2$ terms in the reference system corotating with Earth, one finds C_{22}^E and S_{22}^E of order 10^{-6} . The expected order of magnitude of the second time derivative terms is $(\Omega_E R_E/c)^2 \simeq 10^{-12}$ times smaller than the corresponding ‘‘Newtonian terms’’ from C_{22}^E or S_{22}^E . The Newtonian terms lead to contributions in G_{00} of order 10^{-15} , and hence the second time derivative terms lead to contributions of order 10^{-27} . This is about 9 orders of magnitude less than the $2W^2/c^4$ term in G_{00} , which is of order 10^{-18} . For that reason, these second time derivative terms in Earth’s metric can safely be neglected at present. They are not mentioned in Resolution B1.4.

3.5.3. Approximate Expansion of the Vector Gravitational Potential

Let us now come to the gravitomagnetic vector potential of Earth, W_E^a . As can be seen from equation (39), this potential is determined by the set of spin moments and the first time derivatives of the mass moments. As already mentioned, to characterize the gravitational field outside of some matter distribution in GRT, *two* independent sets of multipole moments have to be used, which in principle should be determined from observational data. So far, the spin moments of some astronomical bodies have not been studied, and more work is needed here. Formally, the spin moments of Earth are given by equation (36) above. Since for the post-Newtonian metric we need these spin moments only to Newtonian order, we might proceed with a simple

Newtonian model of a rigidly rotating Earth, with

$$\Sigma = \Sigma(\mathbf{\Omega} \times \mathbf{X}) ,$$

where Σ is the gravitational mass-energy density in the GCRS and $\mathbf{\Omega}$ is the angular velocity of rotation, which at this point has to be defined only to Newtonian order. Under this assumption, all spin moments are proportional to the angular velocity, and one might define a set of Cartesian tensors C_{Ld} such that

$$\mathcal{S}_L = C_{Ld}\Omega^d . \tag{45}$$

These tensors C_{Ld} obey the following Newtonian relations:

$$C_{Ld} = -M_{Ld} + \frac{l+1}{2l+1} \delta_{d(a_l} N_{L-1)} , \tag{46}$$

where

$$M_L \equiv \int_E \Sigma \hat{X}^L d^3X , \quad N_L \equiv \int_E \Sigma \mathbf{X}^2 \hat{X}^L d^3X . \tag{47}$$

Note that C_{Ld} is symmetric and trace-free only in the first L indices. Moreover, for the Newtonian mass moments M_L one has

$$M_L = -C_{(L)} . \tag{48}$$

For a homogeneous ($\Sigma = \text{const}$) and spherical Earth with radius R_E , one finds for $l = 1$ the usual expression for the moment-of-inertia tensor:

$$C_{ab} = \delta_{ab} \left(\frac{2}{5} MR_E^2 \right) ,$$

which yields the total intrinsic angular momentum (spin) vector of Earth according to $\mathcal{S}_a = C_{ab}\Omega^b$. For a spherically symmetric and mass-centered Earth, all mass moments M_L with $l \geq 1$ vanish, as do all quantities N_L with $l > 0$. Hence, in such a simple model only the spin vector is different from zero and all higher spin moments vanish. For this reason, we also considered a rigidly rotating, homogeneous, oblate spheroid with equatorial radius A and polar radius C . For such a model all even spin moments vanish, since they are proportional to C_L with odd l . On the other hand, odd spin moments proportional to C_L with even l are nonzero. For the spin dipole, the usual result $C_{XX} = C_{YY} = M(A^2 + C^2)/5$ and $C_{ZZ} = 2MA^2/5$ for the moment-of-inertia tensor is found. By means of computer algebra, all components C_L can be found for any value of l . Let $\eta = (4MA^4/525)\epsilon^2$ with $\epsilon^2 = (A^2 - C^2)/A^2 \simeq 2f$, where f is the usual flattening. Assuming $\Omega^d = (0, 0, \Omega)$ we find all nonvanishing $l = 3$ terms, up to symmetries and terms of order f^2 : $S_{XXZ} = S_{YYZ} = 3\eta\Omega$ and $S_{ZZZ} = -6\eta\Omega$. This implies that the metric term resulting from the spin octupole of Earth near the surface is about 10^4 times smaller than the term from the spin dipole. In the following, the contributions of higher spin moments will be neglected.

Besides the spin moments, the first time derivatives of the mass moments contribute to the gravitomagnetic field of Earth. For $l = 0$, we encounter an \mathcal{M}_a term that vanishes if the post-Newtonian center-of-mass condition $\mathcal{M}_a = 0$ is imposed. The next term is given by \mathcal{M}_{ab} , which is of order $|C_{22}^E|MR_E^2\Omega$ and would vanish for an axially symmetric rigid body rotating about its symmetry axis, as would the time derivative of all higher mass moments. For Earth the \mathcal{M}_{ab} term is smaller than the spin term (which is of order $2MR_E^2\Omega/5$) by a factor determined by $C_{22}^E \simeq 1.6 \times 10^{-6}$ and hence negligible. On the other hand, the vector potential $W_E^a(T, \mathbf{X})$ is employed only in the calculation of small

relativistic effects (e.g., Lense-Thirring effects and higher order relativistic effects in the time transformations). This implies that the expansion in equation (39) for $W_E^a(T, \mathbf{X})$ can be truncated to the approximate expression

$$W_E^a(T, \mathbf{X}) = -\frac{G}{2} \frac{(\mathbf{X} \times \mathbf{S}_E)^a}{R^3} , \tag{49}$$

where \mathbf{S}_E is a vector with components \mathcal{S}_a . This expression can be found in many standard textbooks on GRT (Weinberg 1972; Will 1993) and is usually related to Lense-Thirring effects resulting from Earth's rotational motion.

The reason for characterizing W_E^a by the spin vector and not by the angular velocity vector of Earth is a conceptual one, since it is usually advantageous to characterize the gravitational field of Earth in the outside region by multipole moments. To obtain W_E^a , Earth's spin vector is needed only to Newtonian order and can be taken from current precession-nutation models. Although one might use Newtonian concepts to relate the gravitomagnetic field of Earth to some Earth angular velocity, we prefer to employ the well-defined concept of multipole moments here, which are independent of any theoretical assumptions about the rotational motion of Earth.

3.6. The Barycentric Metric in the Mass-Monopole Approximation

In the gravitational N -body problem, the potential coefficients of a body A are defined in its corresponding local reference system (analogous to the GCRS for Earth). For many applications it is sufficient to keep only the mass monopoles of the solar system bodies, that is, to set

$$\mathcal{M}_L = 0 \text{ for } l \geq 1 , \quad \mathcal{S}_L = 0 \text{ for } l \geq 1 \tag{50}$$

for all bodies and to keep the masses only, that is, each body A is characterized by the value for its post-Newtonian mass \mathcal{M}_A (we also set $\mathcal{P}_L = 0$). In the following, we will use the notation M_A instead of \mathcal{M}_A to be consistent with the text of the IAU 2000 resolutions.

From the transformation rules for the metric potentials (eq. [31]), the expansions in equations (38)–(39), and equation (18), one derives the metric in the barycentric coordinate system in the form of equation (8) with

$$w = w_0 - \Delta/c^2 , \tag{51}$$

where

$$w_0(t, \mathbf{x}) \equiv \sum_A \frac{GM_A}{r_A} , \tag{52}$$

$$\Delta(t, \mathbf{x}) = \sum_A \Delta_A(t, \mathbf{x}) \tag{53}$$

with

$$\begin{aligned} \Delta_A(t, \mathbf{x}) &= \frac{GM_A}{r_A} \left(-\frac{3}{2} v_A^2 + \sum_{B \neq A} \frac{GM_B}{r_{BA}} \right) - \frac{1}{2} GM_{A,r_A,t} \\ &= \frac{GM_A}{r_A} \left\{ -2v_A^2 + \sum_{B \neq A} \frac{GM_B}{r_{BA}} \right. \\ &\quad \left. + \frac{1}{2} \left[\frac{(r_A^k v_A^k)^2}{r_A^2} + r_A^k a_A^k \right] \right\} \end{aligned} \tag{54}$$

and $\mathbf{r}_{BA} = \mathbf{x}_B - \mathbf{x}_A$ and $\mathbf{a}_A = d\mathbf{v}_A/dt$. Furthermore, in our approximation

$$w^i(t, \mathbf{x}) = \sum_A \frac{GM_A}{r_A} v_A^i. \quad (55)$$

Note that we have chosen a minus sign in front of Δ in order to have a plus sign in the $1/c^4$ part of g_{00} (see Resolution B1.5). Note furthermore that the post-Newtonian Einstein-Infeld-Hoffmann equations of motion for a system of mass monopoles, which form the basis of modern solar system ephemerides, can be derived from that form of the barycentric metric (for details, see Damour et al. 1991). Thus, the barycentric mass-monopole metric given above is already in use for the description of solar system dynamics.

One improvement to this simple mass-monopole model is to consider the spin dipoles of the various bodies as well (i.e., to also consider \mathbf{S}_A to be nonzero). In fact, Resolution B1.5 is based upon such a mass-monopole spin-dipole model, where modifications from the simple mass-monopole model are indicated explicitly.

4. TIME AND FREQUENCY APPLICATIONS IN THE SOLAR SYSTEM

For practical applications concerning time and frequency measurements in the solar system, it is necessary to consider a conventional model for the realization of time coordinates and time transformations. This model should be chosen so that (1) its accuracy is significantly better than the expected performance of clocks and time transfer techniques, (2) it is consistent with the general framework of § 3, and (3) it may readily be used with existing astrometric quantities, for example, solar system ephemerides.

Regarding item 1, we may derive reasonable limits on accuracy for such a model in a straightforward way. At present, the best accuracies are reached by cesium-fountain clocks operating at less than 2 parts in 10^{15} in fractional frequency (Lemondé et al. 2001; Weyers et al. 2001). Their frequency stability for time spans up to a few days characterized by a standard Allan deviation is on the order of $\sigma_y(\tau) = 4 \times 10^{-14} \tau^{-1/2}$, for an integration time τ in seconds. In the near future, high-accuracy laser-cooled rubidium clocks (Bize et al. 1999) and spaceborne cesium clocks (Lemondé et al. 2001) are expected to reach accuracies of a few parts in 10^{17} in fractional frequency and stabilities on the order of $\sigma_y(\tau) = 1 \times 10^{-14} \tau^{-1/2}$. The uncertainty in the time transformations should induce errors that are always lower than the expected performance of these future clocks. Including a factor of 2 as safety margin, we therefore conclude that time coordinates and time transformations should be realized with an uncertainty not larger than 5×10^{-18} in rate or, for quasi-periodic terms, not larger than 5×10^{-18} in rate and 0.2 ps in amplitude.

For the spatial domain of validity of the transformations, we note that projects such as the *Solar Orbit Relativity Test* plan to fly highly accurate clocks to within 0.25 AU of the Sun, which is therefore the lower limit for the distance to the barycenter that we will consider. In the geocentric system, we will consider locations from Earth's surface up to geostationary orbits ($|\mathbf{X}| < 50,000$ km).

To comply with item 2, we render the developments following the general framework outlined in § 3, and we show (item 3) how the time transformations, for example,

TCB-TCG, may be performed with existing astrometric quantities and tools.

4.1. Barycentric Reference System

Let us write the barycentric metric potential $w(t, \mathbf{x})$ in the form

$$w = w_0 + w_L - \Delta/c^2, \quad (56)$$

where w_L contains contributions from higher potential coefficients with $l \geq 1$ and can be determined from equation (38) and the transformation rules for the metric potentials. Evaluating the Δ_A terms from Resolution B1.5 (eq. [54] plus spin terms) for all bodies of the solar system, we find that in the metric tensor, $|\Delta_A(t, \mathbf{x})|/c^4$ may reach at most a few parts in 10^{17} in the vicinity of Jupiter and about 10^{-17} close to Earth. Presently, however, for all planets except Earth, the magnitude of $\Delta_A(t, \mathbf{x})/c^4$ in the vicinity of the planet is smaller than the uncertainty in w_0/c^2 or w_L/c^2 originating from the uncertainties in its mass multipole moments, so that in practice it is not necessary to account for these terms. Nevertheless, when new astrometric observations allow derivation of the moments with sufficient accuracy, it will be necessary to do so. In any case, in the vicinity of a given body A , only the effect of $\Delta_A(t, \mathbf{x})$ is needed in practice, that is, the effect of $\sum_{B \neq A} \Delta_B(t, \mathbf{x})$ is smaller than our accuracy specifications. For a comparison of the proper time of a clock in the vicinity of Earth with that of other clocks in the solar system or with TCB, it may thus be necessary to account for $\Delta_E(t, \mathbf{x})/c^4$.

From equations (8) and (56), the transformation between the proper time of some observer and TCB may be derived within our accuracy limit:

$$\frac{d\tau}{d\text{TCB}} = 1 - \frac{1}{c^2} \left(w_0 + w_L + \frac{v^2}{2} \right) + \frac{1}{c^4} \left(-\frac{v^4}{8} - \frac{3}{2} v^2 w_0 + 4v^i w^i + \frac{w_0^2}{2} + \Delta \right), \quad (57)$$

where v^i is the BCRS coordinate velocity of the observer. Similarly, the transformation between TCB and TCG in the immediate vicinity of Earth, accurate to the limits specified above, can be derived from the general post-Newtonian TCB-TCG transformation from Resolution B1.3 as

$$\begin{aligned} \text{TCB} - \text{TCG} = c^{-2} & \left\{ \int_{t_0}^t \left[\frac{v_E^2}{2} + w_{0,\text{ext}}(\mathbf{x}_E) \right] dt + v_E^i r_E^i \right\} \\ & - c^{-4} \left\{ \int_{t_0}^t \left[-\frac{1}{8} v_E^4 - \frac{3}{2} v_E^2 w_{0,\text{ext}}(\mathbf{x}_E) \right. \right. \\ & \quad \left. \left. + 4v_E^i w_{\text{ext}}^i(\mathbf{x}_E) + \frac{1}{2} w_{0,\text{ext}}^2(\mathbf{x}_E) \right] dt \right. \\ & \quad \left. - \left[3w_{0,\text{ext}}(\mathbf{x}_E) + \frac{v_E^2}{2} \right] v_E^i r_E^i \right\}, \quad (58) \end{aligned}$$

where t is TCB. Here $w_{0,\text{ext}}$ is defined by equation (52) with the summation over all solar system bodies except Earth. Note that t_0 was not explicitly defined in Resolution B1.5 (2000). It is the origin of TCB and TCG, defined in Resolution A4 (1991; see § 2.1). The external contributions to w_L and Δ are beyond our accuracy limit and can be neglected here.

TABLE 1
CONSTANTS RELATING THE MEAN RATES OF DIFFERENT RELATIVISTIC TIME SCALES

Constant	IAU 1991 (s s ⁻¹)	IAU 2000 (s s ⁻¹)	IAU 2000 (ms yr ⁻¹)
L_C	1.480813×10^{-8}	$1.48082686741 \times 10^{-8}$	467.313
L_G	6.969291×10^{-10}	$6.969290134 \times 10^{-10}$	21.993
$L_B \equiv L_C + L_G - L_C L_G$	1.550505×10^{-8}	$1.55051976772 \times 10^{-8}$	489.307

NOTE.—Both the values adopted by the IAU 1991 recommendations and the IAU 2000 resolutions are given. As an illustration, the IAU 2000 values are also given in milliseconds per Julian year.

This equation is composed of terms evaluated at the geocenter (the two integrals) and of position-dependent terms linear in $|r_E|$, terms with higher powers of $|r_E|$ having been found to be negligible. The integrals may be computed from existing planetary ephemerides (Fukushima 1995; Irwin & Fukushima 1999). Since, in general, the planetary ephemerides are expressed in terms of a time argument $T_{\text{eph}} = (1 - L_B)\text{TCB} + T_{\text{eph}_0}$ (Standish 1998; Irwin & Fukushima 1999), the first integral will be computed as

$$\int_{t_0}^t \left[\frac{v_E^2}{2} + w_{0,\text{ext}}(\mathbf{x}_E) \right] dt = \left\{ \int_{T_{\text{eph}_0}}^{T_{\text{eph}}} \left[\frac{v_E^2}{2} + w_{0,\text{ext}}(\mathbf{x}_E) \right] dT_{\text{eph}} \right\} / (1 - L_B). \quad (59)$$

Terms in the second integral of equation (58) are secular and quasi-periodic. They amount to $\sim 1.1 \times 10^{-16}$ in rate ($d\text{TCB}/d\text{TCG}$) and, primarily, a yearly term of ~ 30 ps in amplitude (i.e., corresponding to periodic rate variations of amplitude $\sim 6 \times 10^{-18}$). Position-dependent terms in c^{-4} are not negligible and reach, for example, an amplitude of 0.4 ps ($\sim 3 \times 10^{-17}$ in rate) in geostationary orbit.

4.2. Geocentric Reference System

Evaluating the contributions of the different terms in the metric tensor of the GCRS given in Resolution B1.3 to the $d\tau/d\text{TCG}$ transformation on Earth's surface and up to geostationary orbit, we find that terms of order c^{-2} reach 7 parts in 10^{10} , while the contributions from W^2 and W^a do not exceed 5 parts in 10^{19} . Also, the terms from W_{iner} in W remain below 2×10^{-20} . Therefore, the terms given in the IAU 1991 framework with the metric of the form of equation (2) are sufficient for time and frequency applications in the GCRS in the region $|X| < 50,000$ km for present and foreseeable future clock accuracies. Note that some care needs to be taken when evaluating the potential W at the location of the clock, which is not trivial when accuracies of order 10^{-18} are required (Klioner 1992; Petit & Wolf 1994; Wolf & Petit 1995).

Presently, the time scale of reference for all practical issues on Earth is Terrestrial Time or one of the scales realizing it and differing by some time offset (e.g., TAI, UTC, GPS time). TT was defined in IAU Resolution A4 (1991) as a time scale differing from the Geocentric Coordinate Time TCG by a constant rate, the unit of measurement of TT being chosen so that it agrees with the SI second on the geoid. According to the transformation between proper and coordinate time, this constant rate is given by $d\text{TT}/d\text{TCG} = 1 - U_G/c^2 = 1 - L_G$, where U_G is the gravity (gravitational plus rotational) potential on the geoid (this

notation is used instead of the usual “ W_0 ” to avoid confusion with the GCRS gravitational potential W used throughout the paper).

Some shortcomings appear in this definition of TT when considering accuracies below 10^{-17} . First, the uncertainty in the determination of U_G is on the order of $1 \text{ m}^2 \text{ s}^{-2}$ or slightly better (Burša 1995; Groten 1999). Second, even if it is expected that the uncertainty in U_G will improve with time, the surface of the geoid is difficult to realize (so that it is difficult to determine the potential difference between the geoid and the location of a clock). Third, the geoid is, in principle, variable with time. Therefore it was decided to dissociate the definition of TT from the geoid while maintaining continuity with the previous definition. The constant L_G was turned into a defining constant, with its value fixed to $6.969290134 \times 10^{-10}$ (Resolution B1.9; see Appendix A). This removes the limitations mentioned above when realizing TT from clocks on board terrestrial satellites (such as in the Global Positioning System). In Table 1, we present numerical values for the constants L_C , L_G , and L_B relating the mean rates of the different relativistic time scales.

5. FINAL REMARKS

The IAU resolutions on relativity represent a post-Newtonian framework allowing one to model any kind of astronomical observation in a rigorous, self-consistent manner with accuracies that are sufficient for the coming decades. They replace the old IAU relativistic framework, which was insufficient for many reasons discussed above. These new resolutions, however, are not expected to lead to dramatic changes. In fact, in many fields of application the models presently in use are already fully compatible with the new IAU resolutions, and in this sense the IAU resolutions officially fix the status quo. Let us give some examples of this.

The metric tensor of the BCRS allows one to derive the Einstein-Infeld-Hoffman equations, which have been used since the 1970s to construct the JPL numerical ephemerides of planetary motion (Newhall, Standish, & Williams 1983). The BCRS is the basic astrometric reference system, in which concepts such as “radial velocity” and “proper motion” are defined (Lindgren & Dravins 2003). The metric tensors of both the GCRS and BCRS and the transformation between corresponding coordinates were used to formulate the VLBI model that has been employed by the International Earth Rotation Service since 1992. In addition, the equations of motion for Earth's satellites recommended by the IERS are compatible with the new IAU framework and can be derived from the given metric tensor of the GCRS (McCarthy 1992, 1996; McCarthy & Petit 2003).

The models used for constructing the *Hipparcos* Catalogue make it clear that this catalog represents a materialization of the BCRS. The full power of the new IAU theoretical framework will be needed to construct a model for astrometric positional observations with an accuracy of 1 μ as, which will be necessary for future astrometric missions. Work in this direction has already started (Klioner 2003).

It is obvious that this explanatory supplement presents only a first step to show how the new IAU resolutions concerning relativity should be employed in practice. Much more work will be necessary to reach that goal.

The anonymous referee is thanked for valuable suggestions that improved the text and made it more readable.

APPENDIX A

IAU RESOLUTIONS CONCERNING RELATIVITY ADOPTED AT THE 24TH GENERAL ASSEMBLY¹⁸

A1. RESOLUTION B1.3: DEFINITION OF BARYCENTRIC CELESTIAL REFERENCE SYSTEM AND GEOCENTRIC CELESTIAL REFERENCE SYSTEM

The XXVIth International Astronomical Union General Assembly,

CONSIDERING,

1. That the Resolution A4 of the XXIst General Assembly (1991) has defined a system of space-time coordinates for (a) the solar system (now called the Barycentric Celestial Reference System [BCRS]) and (b) the Earth (now called the Geocentric Celestial Reference System [GCRS]), within the framework of General Relativity;

2. The desire to write the metric tensors both in the BCRS and in the GCRS in a compact and self-consistent form;

3. The fact that considerable work in General Relativity has been done using the harmonic gauge that was found to be a useful and simplifying gauge for many kinds of applications;

Recommends,

1. The choice of harmonic coordinates both for the barycentric and for the geocentric reference systems;

2. Writing the time-time component and the space-space component of the barycentric metric $g_{\mu\nu}$ with barycentric coordinates (t, \mathbf{x}) (t = Barycentric Coordinate Time [TCB]) with a single scalar potential $w(t, \mathbf{x})$ that generalises the Newtonian potential, and the space-time component with a vector potential $w^i(t, \mathbf{x})$; as a boundary condition it is assumed that these two potentials vanish far from the solar system;

Explicitly,

$$g_{00} = -1 + \frac{2w}{c^2} - \frac{2w^2}{c^4}, \quad g_{0i} = -\frac{4}{c^3}w^i, \quad g_{ij} = \delta_{ij} \left(1 + \frac{2}{c^2}w\right),$$

with

$$w(t, \mathbf{x}) = G \int d^3x' \frac{\sigma(t, \mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|} + \frac{1}{2c^2} G \frac{\partial^2}{\partial t^2} \int d^3x' \sigma(t, \mathbf{x}') |\mathbf{x} - \mathbf{x}'|, \quad w^i(t, \mathbf{x}) = G \int d^3x' \frac{\sigma^i(t, \mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|};$$

here, σ and σ^i are the gravitational mass and current densities respectively;

3. Writing the geocentric metric tensor $G_{\alpha\beta}$ with geocentric coordinates (T, \mathbf{X}) (T = Geocentric Coordinate Time [TCG]) in the same form as the barycentric one but with potentials $W(T, \mathbf{X})$ and $W^a(T, \mathbf{X})$; these geocentric potentials should be split into two parts—potentials W_E and W_E^a arising from the gravitational action of the Earth and external parts W_{ext} and W_{ext}^a due to tidal and inertial effects; the external parts of the metric potentials are assumed to vanish at the geocenter and admit an expansion into positive powers of \mathbf{X} ;

Explicitly,

$$G_{00} = -1 + \frac{2W}{c^2} - \frac{2W^2}{c^4}, \quad G_{0a} = -\frac{4}{c^3}W^a, \quad G_{ab} = \delta_{ab} \left(1 + \frac{2}{c^2}W\right);$$

the potentials W and W^a should be split according to

$$W(T, \mathbf{X}) = W_E(T, \mathbf{X}) + W_{\text{ext}}(T, \mathbf{X}), \quad W^a(T, \mathbf{X}) = W_E^a(T, \mathbf{X}) + W_{\text{ext}}^a(T, \mathbf{X});$$

the Earth's potentials W_E and W_E^a are defined in the same way as w and w^a but with quantities calculated in the GCRS with integrals taken over the whole Earth;

4. Using, if accuracy requires, the full post-Newtonian coordinate transformation between the BCRS and the GCRS as induced by the form of the corresponding metric tensors;

¹⁸ Reprinted with minor changes from the Transactions of the IAU, Vol. 24B, pp. 37–49 and 56–57 (San Francisco: ASP [2001]). Courtesy of the International Astronomical Union.

Explicitly, for the kinematically non-rotating GCRS [$T = \text{TCG}$, $t = \text{TCB}$, $r_E^i = x^i - x_E^i(t)$, and a summation from 1 to 3 over equal indices is implied],

$$T = t - \frac{1}{c^2} [A(t) + v_E^i r_E^i] + \frac{1}{c^4} [B(t) + B^i(t) r_E^i + B^{ij}(t) r_E^i r_E^j + C(t, \mathbf{x})] + O(c^{-5}),$$

$$X^a = \delta_{ai} \left\{ r_E^i + \frac{1}{c^2} \left[\frac{1}{2} v_E^i v_E^j r_E^j + w_{\text{ext}}(\mathbf{x}_E) r_E^i + r_E^i a_E^j r_E^j - \frac{1}{2} a_E^i r_E^2 \right] \right\} + O(c^{-4}),$$

where

$$\frac{d}{dt} A(t) = \frac{1}{2} v_E^2 + w_{\text{ext}}(\mathbf{x}_E), \quad \frac{d}{dt} B(t) = -\frac{1}{8} v_E^4 - \frac{3}{2} v_E^2 w_{\text{ext}}(\mathbf{x}_E) + 4v_E^i w_{\text{ext}}^i(\mathbf{x}_E) + \frac{1}{2} w_{\text{ext}}^2(\mathbf{x}_E),$$

$$B^i(t) = -\frac{1}{2} v_E^2 v_E^i + 4w_{\text{ext}}^i(\mathbf{x}_E) - 3v_E^i w_{\text{ext}}(\mathbf{x}_E),$$

$$B^{ij}(t) = -v_E^i \delta_{aj} Q^a + 2 \frac{\partial}{\partial x^j} w_{\text{ext}}^i(\mathbf{x}_E) - v_E^i \frac{\partial}{\partial x^j} w_{\text{ext}}(\mathbf{x}_E) + \frac{1}{2} \delta^{ij} \dot{w}_{\text{ext}}(\mathbf{x}_E),$$

$$C(t, \mathbf{x}) = -\frac{1}{10} r_E^2 (\dot{a}_E^i r_E^i);$$

here x_E^i , v_E^i , and a_E^i are the components of the barycentric position, velocity and acceleration vectors of the Earth, the dot stands for the total derivative with respect to t , and

$$Q^a = \delta_{ai} \left[\frac{\partial}{\partial x_i} w_{\text{ext}}(\mathbf{x}_E) - a_E^i \right];$$

the external potentials, w_{ext} and w_{ext}^i , are given by

$$w_{\text{ext}} = \sum_{A \neq E} w_A, \quad w_{\text{ext}}^i = \sum_{A \neq E} w_A^i,$$

where E stands for the Earth and w_A and w_A^i are determined by the expressions for w and w^i with integrals taken over body A only.

NOTES

It is to be understood that these expressions for w and w^i give g_{00} correct up to $O(c^{-5})$, g_{0i} up to $O(c^{-5})$, and g_{ij} up to $O(c^{-4})$. The densities σ and σ^i are determined by the components of the energy momentum tensor of the matter composing the solar system bodies as given in the references. Accuracies for $G_{\alpha\beta}$ in terms of c^{-n} correspond to those of $g_{\mu\nu}$.

The external potentials W_{ext} and W_{ext}^a can be written in the form

$$W_{\text{ext}} = W_{\text{tidal}} + W_{\text{iner}}, \quad W_{\text{ext}}^a = W_{\text{tidal}}^a + W_{\text{iner}}^a.$$

W_{tidal} generalises the Newtonian expression for the tidal potential. Post-Newtonian expressions for W_{tidal} and W_{tidal}^a can be found in the references. The potentials W_{iner} , W_{iner}^a are inertial contributions that are linear in X^a . The former is determined mainly by the coupling of the Earth's nonsphericity to the external potential. In the kinematically non-rotating Geocentric Celestial Reference System, W_{iner}^a describes the Coriolis force induced mainly by geodetic precession.

Finally, the local gravitational potentials W_E and W_E^a of the Earth are related to the barycentric gravitational potentials w_E and w_E^i by

$$W_E(T, \mathbf{X}) = w_E(t, \mathbf{x}) \left(1 + \frac{2}{c^2} v_E^2 \right) - \frac{4}{c^2} v_E^i w_E^i(t, \mathbf{x}) + O(c^{-4}), \quad W_E^a(T, \mathbf{X}) = \delta_{ai} [w_E^i(t, \mathbf{x}) - v_E^i w_E(t, \mathbf{x})] + O(c^{-2}).$$

REFERENCES

- Brumberg, V. A., Kopeikin, S. M., 1989, *Nuovo Cimento B*, **103**, 63.
 Brumberg, V. A., 1991, *Essential Relativistic Celestial Mechanics*, Hilger, Bristol.
 Damour, T., Soffel, M., Xu, C., *Phys. Rev. D*, **43**, 3273 (1991); **45**, 1017 (1992); **47**, 3124 (1993); **49**, 618 (1994).
 Klioner, S. A., Voinov, A.V., 1993, *Phys. Rev. D*, **48**, 1451.
 Kopeikin, S. M., 1988, *Celest. Mech.*, **44**, 87.

A2. RESOLUTION B1.4: POST-NEWTONIAN POTENTIAL COEFFICIENTS

The XXVIth International Astronomical Union General Assembly,

CONSIDERING,

1. That for many applications in the fields of celestial mechanics and astrometry a suitable parametrization of the metric potentials (or multipole moments) outside the massive solar system bodies in the form of expansions in terms of potential coefficients are extremely useful; and

2. That physically meaningful post-Newtonian potential coefficients can be derived from the literature;

Recommends,

1. Expansion of the post-Newtonian potential of the Earth in the Geocentric Celestial Reference System (GCRS) outside the Earth in the form

$$W_E(T, \mathbf{X}) = \frac{GM_E}{R} \left\{ 1 + \sum_{l=2}^{\infty} \sum_{m=0}^{+l} \left(\frac{R_E}{R} \right)^l P_{lm}(\cos \theta) [C_{lm}^E(T) \cos m\phi + S_{lm}^E(T) \sin m\phi] \right\};$$

here C_{lm}^E and S_{lm}^E are, to sufficient accuracy, equivalent to the post-Newtonian multipole moments introduced by Damour et al. (Damour et al., 1991, *Phys. Rev. D*, **43**, 3273); θ and ϕ are the polar angles corresponding to the spatial coordinates X^a of the GCRS and $R = |\mathbf{X}|$; and

2. Expression of the vector potential outside the Earth, leading to the well-known Lense-Thirring effect, in terms of the Earth's total angular momentum vector \mathbf{S}_E in the form

$$W_E^a(T, \mathbf{X}) = -\frac{G}{2} \frac{(\mathbf{X} \times \mathbf{S}_E)^a}{R^3}.$$

A3. RESOLUTION B1.5: EXTENDED RELATIVISTIC FRAMEWORK FOR TIME TRANSFORMATIONS AND REALISATION OF COORDINATE TIMES IN THE SOLAR SYSTEM

The XXVIth International Astronomical Union General Assembly,

CONSIDERING,

1. That the Resolution A4 of the XXIst General Assembly (1991) has defined systems of space-time coordinates for the solar system (Barycentric Reference System) and for the Earth (Geocentric Reference System), within the framework of General Relativity;

2. That Resolution B1.3 entitled "Definition of Barycentric Celestial Reference System and Geocentric Celestial Reference System" has renamed these systems the Barycentric Celestial Reference System (BCRS) and the Geocentric Celestial Reference System (GCRS), respectively, and has specified a general framework for expressing their metric tensor and defining coordinate transformations at the first post-Newtonian level;

3. That, based on the anticipated performance of atomic clocks, future time and frequency measurements will require practical application of this framework in the BCRS; and

4. That theoretical work requiring such expansions has already been performed;

Recommends, That for applications that concern time transformations and realisation of coordinate times within the solar system, Resolution B1.3 be applied as follows:

1. The metric tensor be expressed as

$$g_{00} = -\left\{ 1 - \frac{2}{c^2} [w_0(t, \mathbf{x}) + w_L(t, \mathbf{x})] + \frac{2}{c^4} [w_0^2(t, \mathbf{x}) + \Delta(t, \mathbf{x})] \right\},$$

$$g_{0i} = -\frac{4}{c^3} w^i(t, \mathbf{x}), \quad g_{ij} = \left[1 + \frac{2w_0(t, \mathbf{x})}{c^2} \right] \delta_{ij},$$

where ($t \equiv$ Barycentric Coordinate Time [TCB], \mathbf{x}) are the barycentric coordinates, $w_0 = G \sum_A M_A / r_A$, with the summation carried out over all solar system bodies A , $\mathbf{r}_A = \mathbf{x} - \mathbf{x}_A$, \mathbf{x}_A are the coordinates of the center of mass of body A , $r_A = |\mathbf{r}_A|$, and where w_L contains the expansion in terms of multipole moments [see their definition in the Resolution B1.4 entitled "Post-Newtonian Potential Coefficients"] required for each body [the vector potential $w^i(t, \mathbf{x}) = \sum_A w_A^i(t, \mathbf{x})$ and the function $\Delta(t, \mathbf{x}) = \sum_A \Delta_A(t, \mathbf{x})$ are given in note 2];

2. The relation between TCB and Geocentric Coordinate Time (TCG) can be expressed to sufficient accuracy by

$$\begin{aligned} \text{TCB} - \text{TCG} = & c^{-2} \left\{ \int_{t_0}^t \left[\frac{v_E^2}{2} + w_{0,\text{ext}}(\mathbf{x}_E) \right] dt + v_E^i r_E^i \right\} \\ & - c^{-4} \left\{ \int_{t_0}^t \left[-\frac{1}{8} v_E^4 - \frac{3}{2} v_E^2 w_{0,\text{ext}}(\mathbf{x}_E) + 4v_E^i w_{\text{ext}}^i(\mathbf{x}_E) + \frac{1}{2} w_{0,\text{ext}}^2(\mathbf{x}_E) \right] dt - \left[3w_{0,\text{ext}}(\mathbf{x}_E) + \frac{v_E^2}{2} \right] v_E^i r_E^i \right\}, \end{aligned}$$

where v_E is the barycentric velocity of the Earth and where the index ext refers to summation over all bodies except the Earth.

NOTES

1. This formulation will provide an uncertainty not larger than 5×10^{-18} in rate and, for quasi-periodic terms, not larger than 5×10^{-18} in rate amplitude and 0.2 ps in phase amplitude, for locations farther than a few solar radii from the Sun. The same uncertainty also applies to the transformation between TCB and TCG for locations within 50 000 km of the Earth. Uncertainties in the values of astronomical quantities may induce larger errors in the formulas.

2. Within the above mentioned uncertainties, it is sufficient to express the vector potential $w_A^i(t, \mathbf{x})$ of body A as

$$w_A^i(t, \mathbf{x}) = G \left[\frac{-(\mathbf{r}_A \times \mathbf{S}_A)^i}{2r_A^3} + \frac{M_A v_A^i}{r_A} \right],$$

where \mathbf{S}_A is the total angular momentum of body A and v_A^i are the components of the barycentric coordinate velocity of body A . As for the function $\Delta_A(t, \mathbf{x})$ it is sufficient to express it as

$$\Delta_A(t, \mathbf{x}) = \frac{GM_A}{r_A} \left\{ -2v_a^2 + \sum_{B \neq A} \frac{GM_B}{r_{BA}} + \frac{1}{2} \left[\frac{(v_A^k v_A^k)^2}{r_A^2} + r_A^k a_A^k \right] \right\} + \frac{2Gv_A^k (\mathbf{r}_A \times \mathbf{S}_A)^k}{r_A^3},$$

where $r_{BA} = |\mathbf{x}_B - \mathbf{x}_A|$ and a_A^k is the barycentric coordinate acceleration of body A . In these formulas, the terms in \mathbf{S}_A are needed only for Jupiter ($S \approx 6.9 \times 10^{38} \text{ m}^2 \text{ s}^{-1} \text{ kg}$) and Saturn ($S \approx 1.4 \times 10^{38} \text{ m}^2 \text{ s}^{-1} \text{ kg}$), in the immediate vicinity of these planets.

3. Because the present Recommendation provides an extension of the IAU 1991 recommendations valid at the full first post-Newtonian level, the constants L_C and L_B that were introduced in the IAU 1991 recommendations should be defined as $\langle \text{TCG}/\text{TCB} \rangle = 1 - L_C$ and $\langle \text{TT}/\text{TCB} \rangle = 1 - L_B$, where TT refers to Terrestrial Time and $\langle \rangle$ refers to a sufficiently long average taken at the geocenter. The most recent estimate of L_C is (Irwin, A., and Fukushima, T., 1999, *Astron. Astroph.*, **348**, 642–652)

$$L_C = 1.48082686741 \times 10^{-8} \pm 2 \times 10^{-17}.$$

From the Resolution B1.9 on “Redefinition of Terrestrial Time TT,” one infers $L_B = 1.55051976772 \times 10^{-8} \pm 2 \times 10^{-17}$ by using the relation $1 - L_B = (1 - L_C)(1 - L_G)$. L_G is defined in Resolution B1.9.

Because no unambiguous definition may be provided for L_B and L_C , these constants should not be used in formulating time transformations when it would require knowing their value with an uncertainty of order 1×10^{-16} or less.

4. If $\text{TCB} - \text{TCG}$ is computed using planetary ephemerides which are expressed in terms of a time argument (noted T_{eph}) which is close to Barycentric Dynamical Time (TDB), rather than in terms of TCB, the first integral in Recommendation 2 above may be computed as

$$\int_{t_0}^t \left[\frac{v_E^2}{2} + w_{0,\text{ext}}(\mathbf{x}_E) \right] dt = \left\{ \int_{T_{\text{eph}_0}}^{T_{\text{eph}}} \left[\frac{v_E^2}{2} + w_{0,\text{ext}}(\mathbf{x}_E) \right] dt \right\} / (1 - L_B).$$

A4. RESOLUTION B1.9: RE-DEFINITION OF TERRESTRIAL TIME TT

The XXVIth International Astronomical Union General Assembly,

CONSIDERING,

1. That IAU Resolution A4 (1991) has defined Terrestrial Time (TT) in its Recommendation 4; and
2. That the intricacy and temporal changes inherent to the definition and realisation of the geoid are a source of uncertainty in the definition and realisation of TT, which may become, in the near future, the dominant source of uncertainty in realising TT from atomic clocks;

Recommends, That TT be a time scale differing from TCG by a constant rate: $d\text{TT}/d\text{TCG} = 1 - L_G$, where $L_G = 6.969290134 \times 10^{-10}$ is a defining constant.

NOTE.— L_G was defined by the IAU Resolution A4 (1991) in its Recommendation 4 as equal to U_G/c^2 where U_G is the geopotential at the geoid. L_G is now used as a defining constant.

APPENDIX B

COMPARISON OF THE IAU METRIC WITH VERSIONS GIVEN IN THE LITERATURE

In this appendix, we will compare the metric of equation (8) with well-known results from the literature. However, for this purpose we will consider the material composing the various bodies of the system to behave like an ideal fluid (for the IAU 2000 resolutions this is not assumed). In the ideal-fluid case, the energy-momentum tensor can be written in the form

$$T^{00} = \rho c^2 \left[1 + \frac{1}{c^2} (\Pi + \mathbf{v}^2 + 2U) \right] + O(c^{-2}), \quad T^{0i} = \rho c v^i + O(c^{-1}), \quad T^{ij} = \rho v^i v^j + p \delta_{ij} + O(c^{-2}). \quad (\text{B1})$$

Here ρ denotes the rest-mass density, p is the pressure, Π is the specific internal energy (see, e.g., Will 1993), $v^i(t, \mathbf{x})$ is the velocity of the corresponding material element, and

$$U(t, \mathbf{x}) \equiv G \int \frac{\rho(t, \mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|} d^3 x'. \quad (\text{B2})$$

From equations (12) and (B1), we derive

$$\sigma = \rho \left[1 + \frac{1}{c^2} (\Pi + 2\mathbf{v}^2 + 2U) \right] + 3 \frac{p}{c^2} + O(c^{-4}), \quad \sigma^i = \rho v^i + O(c^{-2}). \quad (\text{B3})$$

Introducing the metric potentials

$$\Phi_1 \equiv \int \frac{\rho' v'^2}{|\mathbf{x} - \mathbf{x}'|} d^3 x', \quad \Phi_2 \equiv \int \frac{\rho' U'}{|\mathbf{x} - \mathbf{x}'|} d^3 x', \quad \Phi_3 \equiv \int \frac{\rho' \Pi'}{|\mathbf{x} - \mathbf{x}'|} d^3 x', \quad \Phi_4 \equiv \int \frac{p'}{|\mathbf{x} - \mathbf{x}'|} d^3 x', \quad (\text{B4})$$

we obtain from equations (14) and (15)

$$w = U + 2\Phi_1 + 2\Phi_2 + \Phi_3 + 3\Phi_4 - \frac{1}{2c^2} \chi_{,tt} + O(c^{-4}), \quad w^i = V_i + O(c^{-2}) \quad (\text{B5})$$

with

$$V_i \equiv \int \frac{\rho' v'_i}{|\mathbf{x} - \mathbf{x}'|} d^3 x', \quad \chi \equiv -G \int \rho' |\mathbf{x} - \mathbf{x}'| d^3 x', \quad (\text{B6})$$

and the subscript comma denoting partial differentiation, as in

$$\chi_{,tt} \equiv \frac{\partial^2 \chi}{\partial t^2}. \quad (\text{B7})$$

The post-Newtonian metric (in harmonic gauge) can then be written as

$$g_{00} = -1 + \frac{1}{c^2} (2U + 4\Phi_1 + 4\Phi_2 + 2\Phi_3 + 6\Phi_4) - \frac{2}{c^4} U^2 + \frac{1}{c^4} \chi_{,tt}, \quad g_{0i} = -\frac{4}{c^3} V_i, \quad g_{ij} = \delta_{ij} \left(1 + \frac{2U}{c^2} \right). \quad (\text{B8})$$

To compare, for example, with the metric in Will (1993), we transform from harmonic coordinates, used in the present paper and recommended by the IAU, to standard post-Newtonian coordinates, used by several authors including Will. This is achieved by a gauge transformation of the form

$$w_{\text{SPN}} = w - \lambda_{,t}/c^2, \quad w^i_{\text{SPN}} = w^i + \lambda_{,i}/4 \quad (\text{B9})$$

with

$$\lambda = -\chi_{,t}/2 \quad (\text{B10})$$

(see, e.g., eq. [3.12] of Damour et al. 1991).

This implies that the χ -term disappears from w and hence from g_{00} when standard post-Newtonian coordinates are employed, but the g_{0i} term is affected by this transformation. Using the relation

$$\chi_{,ti} = V_i - W_i \quad (\text{B11})$$

with

$$W_i \equiv \int \frac{\rho' [\mathbf{v}' \cdot (\mathbf{x} - \mathbf{x}')](x^i - x'^i)}{|\mathbf{x} - \mathbf{x}'|^3} d^3 x', \quad (\text{B12})$$

one verifies that the metric induced by the potentials in equations (14) and (15) agrees in general relativity with equation (5.28) of Will (1993).

REFERENCES

- Bize, S., Sortais, Y., Santos, M. S., Mandache, C., Clairon, A., & Salomon, C. 1999, *Europhys. Lett.*, 45, 558
- Blanchet, L., & Damour, T. 1989, *Ann. Inst. Henri Poincaré A*, 50, 377
- Bois, E., & Vokrouhlický, D. 1995, *A&A*, 300, 559
- Brumberg, V. A. 1991, *Essential Relativistic Celestial Mechanics* (Bristol: Hilger)
- Brumberg, V. A., Bretagnon, P., & Francou, G. 1991, in *Systèmes de référence spatio-temporels: Journées 1991*, ed. N. Capitaine (Paris: Obs. Paris), 141
- Brumberg, V. A., & Kopeikin, S. M. 1989, in *Reference Frames in Astronomy and Geophysics*, ed. J. Kovalevsky, I. I. Mueller, & B. Kolaczek (Dordrecht: Kluwer), 115
- Burša, M. 1995, *Travaux Assoc. Int. Geod.*, 30, 370
- Carter, W. E., Robertson, D. S., & MacKay, J. R. 1985, *J. Geophys. Res.*, 90, 4577
- Damour, T., Soffel, M., & Xu, C. 1991, *Phys. Rev. D*, 43, 3273
- . 1992, *Phys. Rev. D*, 45, 1017
- . 1993, *Phys. Rev. D*, 47, 3124
- . 1994, *Phys. Rev. D*, 49, 618
- Eubanks, T. M., Matsakis, D. N., Martin, J. O., Archinal, B. A., McCarthy, D. D., Klioner, S. A., Shapiro, S., & Shapiro, I. I. 1997, *Bull. Am. Phys. Soc.*, 42(4), abstr. No. K11.05
- Fok, V. A. 1959, *The Theory of Space, Time, and Gravitation* (New York: Pergamon)
- Froeschlé, M., Mignard, F., & Arenou, F. 1997, in *Hipparcos Venice '97*, ed. B. Battick (ESA SP-402) (Noordwijk: ESA), 49
- Fukushima, T. 1991, *A&A*, 244, L11
- . 1995, *A&A*, 294, 895
- Groten, E. 1999, *Travaux Assoc. Int. Geod.*, 31, 272
- IAU. 2001, *IAU Inf. Bull.*, No. 88, 29 (erratum No. 89, 4 [2002])
- Irwin, A. W., & Fukushima, T. 1999, *A&A*, 348, 642
- Klioner, S. A. 1992, *Celest. Mech. Dyn. Astron.*, 53, 81
- . 2003, *AJ*, 125, 1580
- Klioner, S. A., & Soffel, M. 1998, *A&A*, 334, 1123
- Klioner, S. A., Soffel, M., Xu, C., & Wu, X. 2003, in *Systèmes de référence spatio-temporels: Journées 2001* (Paris: Obs. Paris), in press (astro-ph/0303377)
- Klioner, S. A., & Soffel, M. H. 2000, *Phys. Rev. D*, 62, No. 024019
- Klioner, S. A., & Voinov, A. V. 1993, *Phys. Rev. D*, 48, 1451
- Kopeikin, S. M. 1988, *Celest. Mech.*, 44, 87
- . 1990, *AZh*, 67, 10 (English transl. *Soviet Astron.*, 34, 5)
- . 1991, *Manuscripta Geod.*, 16, 301
- Lebach, D. E., Corey, B. E., Shapiro, I. I., Ratner, M. I., Webber, J. C., Rogers, A. E. E., Davis, J. L., & Herring, T. A. 1995, *Phys. Rev. Lett.*, 75, 1439
- Lemonde, P., et al. 2001, in *Frequency Measurement and Control*, ed. A. N. Luiten (Berlin: Springer), 131
- Lindegren, L., & Dravins, D. 2003, *A&A*, 401, 1185
- Mashhoon, B. 1985, *Found. Phys.*, 15, 497
- McCarthy, D. D., ed. 1992, *IERS Standards (1992)* (IERS Tech. Note 13) (Paris: Obs. Paris)
- , ed. 1996, *IERS Conventions (1996)* (IERS Tech. Note 21) (Paris: Obs. Paris)
- McCarthy, D. D., & Petit, G., eds. 2003, *IERS Conventions (2003)* (draft; Paris: Obs. Paris)
- Misner, C. W., Thorne, K. S., & Wheeler, J. A. 1973, *Gravitation* (San Francisco: Freeman)
- Newhall, X. X., Standish, E. M., Jr., & Williams, J. G. 1983, *A&A*, 125, 150
- Petit, G., & Wolf, P. 1994, *A&A*, 286, 971
- Rickman, H., ed. 2001, *Proceedings of the Twenty-Fourth General Assembly: Manchester 2000* (Trans. IAU, Vol. 24B) (San Francisco: ASP)
- Robertson, D. S., & Carter, W. E. 1984, *Nature*, 310, 572
- Soffel, M., Ruder, H., & Schneider, M. 1986, *A&A*, 157, 357
- Soffel, M. H. 1989, *Relativity in Astrometry, Celestial Mechanics, and Geodesy* (Berlin: Springer)
- Standish, E. M. 1998, *A&A*, 336, 381
- Weinberg, S. 1972, *Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity* (New York: Wiley)
- Weyers, S., Hübner, U., Schröder, R., Tamm, C., & Bauch, A. 2001, *Metrologia*, 38, 343
- Will, C. M. 1993, *Theory and Experiment in Gravitational Physics* (rev. ed.; Cambridge: Cambridge Univ. Press)
- Wolf, P., & Petit, G. 1995, *A&A*, 304, 653