



# LUND UNIVERSITY

## Regulation of photosynthesis -Cytochrome b6f in redox regulation -Two novel proteases acting on an N-terminal peptide of LHCII

Ström, Jörgen

2005

[Link to publication](#)

*Citation for published version (APA):*

Ström, J. (2005). *Regulation of photosynthesis -Cytochrome b6f in redox regulation -Two novel proteases acting on an N-terminal peptide of LHCII*. [Doctoral Thesis (compilation), Biochemistry and Structural Biology]. Department of Plant Biochemistry, Lund University.

*Total number of authors:*

1

### General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00

# CYTOCHROME $b_6$ IN REDOX REGULATION OF PHOTOSYNTHESIS AND CHLOROPLAST GENE EXPRESSION

Jörgen Ström<sup>1</sup>, Sujith Puthiyaveetil<sup>1</sup>, Jens Forsberg<sup>2</sup>, John F. Allen<sup>1</sup>. <sup>1</sup>Plant Biochemistry, Lund University Box 124, Lund 221 00, Sweden. <sup>2</sup>Medical Biochemistry and Microbiology Uppsala Biomedical Center, Uppsala University Box 582, Uppsala 751 23, Sweden

**Keywords:** Cytochrome  $b_6f$ , redox regulation, protein phosphorylation, transcriptional control, state transitions

## INTRODUCTION

Cytochrome  $b_6$  is a component of the cytochrome  $b_6f$  complex of the chloroplast thylakoid membrane. The recently described structures of cytochrome  $b_6f$  (Kurusu et al 2003, Strobel et al 2003) have provided new insight to the function of the complex (Allen 2004). We suggest that cytochrome  $b_6$ , apart from being important to both linear and cyclic electron transport, holds another key function, namely signaling in redox regulation of chloroplast function. This suggestion is supported by the observation that cytochrome  $b_6$  is phosphorylated (Gal et al 1992) and by work of other laboratories showing that the occupancy of the  $Q_o$  quinol-binding site of the cytochrome  $b_6f$  complex is a factor governing the activation of a thylakoid protein kinase. Sequence comparisons reveal homology of an N-terminal segment of cytochrome  $b_6$  with highly conserved phosphorylatable histidine residue, which sensor histidine kinases and receptor phytochromes could act upon (Hwang et al 2002, Grebe & Stock 1999). Recent phylogenetic analysis of Xiong and Bauer (2002) suggests that cytochrome  $b_6$  is homologous with photosynthetic reaction centres, where the non-haem iron of type II centres (e.g., photosystem II) is coordinated by amino-acid side chains originally ligating haem. Electron transfer events in photosynthesis have been of over-riding importance in cell evolution, and in the function of the chloroplast genetic system.

## MATERIALS AND METHODS

**Chloroplast isolation and blue-native polyacrylamide gel electrophoresis.** Leaves were collected from 12 day-old peas (*Pisum sativum*). The leaves were homogenised in grinding buffer (0.33 M sorbitol, 50 mM HEPES pH 7.6, 5 mM  $MgCl_2$ , 10 mM NaCl, 2 mM EDTA) and the homogenate was filtered through 2 layers, and then 8 layers, of cheese cloth. The filtrate was then layered onto a Percoll cushion and centrifuged at 600g for 7 min. Intact chloroplasts collected as a green pellet in the bottom of the tube were then resuspended in grinding buffer. The chloroplast suspension (0.33 mg Chl/ml) was incubated for 5 min in light with 400  $\mu Ci$   $^{32}P$  (Amersham) present. To the  $^{32}P$  incubated chloroplast suspension was ferricyanide or dithiothreitol DTT (final concentration 2 mM), added, giving the oxidized and reduced chloroplast samples, respectively. These samples were transferred to darkness for 15 min. The remainder of the preparation was carried out under faint green light. To the chemically redox-treated chloroplasts NaF was added to give a final conc of 10 mM. This concentration of NaF was maintained for the remainder of the preparation. Chloroplasts were collected twice with Eppendorf microcentri-

fuge at 4000 rpm for 5 min, being resuspended and washed once with grinding buffer. The proteins in the final chloroplast preparation were separated using blue-native polyacrylamide gel electrophoresis essentially according to Kügler et al (1997).

**Mass spectrometry analysis.** Tryptic digests of excised protein spots were analysed with MALDI-TOF MS (Ultraflex, Bruker Daltonics). Protein identification was achieved by peptide mass fingerprinting.

**Sequence comparison.** The following databases and sequencing programs were used to establish the conserved N-terminal sequence of cyt  $b_6$ . Sequences aligned were downloaded from Swissprot Database. Multiple Alignment program used was ClustalW (<http://www.ebi.ac.uk/clustalw/>).


## RESULTS

**The Thylakoid Protein Complexes Show Different Levels of Phosphorylation.** Figure 1A and 1B show blue-native polyacrylamide gels of thylakoid membranes reduced (A) and oxidized (B). Figure 1C and 1D shows the same gels, visualizing the  $^{32}P$  labeled complexes by autoradiography.

The blue-native polyacrylamide gels are gradient gels (6–12% acrylamide) run without markers. The different protein complexes were identified using MALDI-TOF MS.

**Phosphorylation of thylakoid complexes under oxidizing conditions.** Figure 1D shows phosphorylation under oxidizing conditions. The major band in the autoradiograph of blue-native polyacrylamide gel representing the Rubisco (1) complex is strongly phosphorylated. The second most conspicuous band on the gel representing a mixture of PSII and ATPsynthase (2) is also strongly phosphorylated. The region of the gel corresponding to the  $b_6f$ -complex (3) has very low levels of phosphorylation, and the same is true for the second band, which represents PSII (4) proteins (CP47, D1, D2, cytochrome  $b_{559}$ ). Finally the band containing LHCII protein (band 5, (containing a mixture of proteins, mostly Lhcb1, Lhcb2 and CP29)) show high levels of phosphorylations.

**Phosphorylation of thylakoid complexes under reducing conditions.** Figure 1C shows phosphorylation under reducing conditions,



**Figure 1:** Coomassie stain and Phosphorimage of  $^{32}P$ -labeled protein complexes from a blue native gel of redox treated chloroplasts. (A) and (B) are Coomassie-stained gels, reduced and oxidized. (C) and (D) are phosphorimages, reduced and oxidized. The numbered complex is Rubisco (1), ATPase/PSII (2), Cytochrome  $b_6f$  (3), PSII (4), LHCII (5).




Figure 2: Sequence alignment of cytochromes *b<sub>6</sub>* from 20 different species (*Chlamydomonas*, *Arabidopsis*, *Pisum*, *Physcomitrella*, *Nostoc*, *Synechocystis*, *Pinus* and more) showing the conserved histidine near the N-terminus.

the major band in the blue-native polyacrylamide gel representing the Rubisco complex is phosphorylated to lesser extent than in Fig. 1D. The second band on the gel, representing a mixture of PS II and ATPsynthase, shows almost no phosphorylation. In contrast, the region of the gel corresponding to the *b<sub>6</sub>f*-complex is now strongly phosphorylated. The second band, of PSII proteins (CP47, D1, D2, cytochrome *b<sub>559</sub>*) shows the same low level of phosphorylation as under oxidizing conditions. Finally, the band containing LHCII proteins shows decreased phosphorylation under reducing conditions, but were still strongly phosphorylated.

**Sequence Comparison of the N-terminal Region of *b<sub>6</sub>*.** We have aligned predicted gene products from *petB* (Fig. 2). The sequences aligned are N-terminal stretch of cyt *b<sub>6</sub>*, length approximately 120 amino acids. A conserved histidine residue lies at the N-terminal part of the alignment (highlighted with a arrow). In the alignment, two of the four histidines, which ligate haem groups) are also seen. The other two haem-ligating histidines lie in the C-terminal part (not included in the alignment).

## DISCUSSION

**Phosphorylation of the Cytochrome *b<sub>6</sub>f*-Complex.** Cytochrome *b<sub>6</sub>f* is involved in redox regulation in the chloroplast, being sensitive to changes in electron flow through the thylakoid membranes. We find

that the cytochrome *b<sub>6</sub>f* complex is strongly phosphorylated under reducing conditions. It is known that subunit V of the cytochrome *b<sub>6</sub>f*-complex undergoes reversible phosphorylation (Hamel et al 2000) but subunit V seems to be present only in stroma lamella and not grana (Romanowska & Albertsson 1994). The redox state of the *b<sub>6</sub>f*-complex acts as activator of the redox dependent LHCII kinase under reducing conditions (Aro & Ohad 2003). We think that cytochrome *b<sub>6</sub>f* is closely involved in both activation of kinases and in regulation of RNA-synthesis. We suggest that the *b<sub>6</sub>* protein in the complex is phosphorylated (Gal et al 1992) under reducing conditions. This phosphorylation might be the signal initiating changes in rates of RNA synthesis. How this is done is still unclear, but the conserved histidine at the N-terminus of *b<sub>6</sub>* is a candidate for starting this signal.

## ACKNOWLEDGMENTS

This work was supported by a Swedish Research Council grant to JFA and by Wallenberg Consortium North (WCN) through the Proteomics Resource Centre at Uppsala University.

## REFERENCES

Allen, J. F. (2003a) *Philos. Trans. R. Soc.Lond. B* 358: 19–38.  
 Allen, J. F. (2003b) *Comp. Funct. Genomics* vol. 4: 31–36.  
 Allen, J. F. (2004) *Trends in Plant Sci.* vol. 3: 130–137.

- Aro, E. & Ohad, I. (2003) *Antioxidants & Redox Signaling* vol. 5 (nr 1): 55–67.
- Gal, A., Herrman, R. G., Lottspeich, F. & Ohad, I. (1992) *FEBS Lett.* 298: 33–35.
- Grebe, T. W. & Stock, J. B. (1999) *Advances in Micro. Physiol.* 41: 139–214.
- Hamel, P., Olive, J., Pierre, Y., Wollman, F. A. & de Vitry, C. A. (2000) *J. Biol. Chem.* 275: 17072–17079.
- Hwang, I., Chen, H. C. & Sheen, J. (2002) *Plant Physio.* 129: 500–515.
- Kurisu, G., Zang, H., Smith, J. L. & Cramer, A. W. (2003) *Science* 302: 1009–1014.
- Kügler, M., Jänsch, L., Krufit, V., Schmitz, U. K. & Braun, H. (1997) *Photosynthesis Res.* 53: 35–44.
- Romanowska, E. & Albertsson, P. (1994) *Plant Cell Physiol.* 35: 557–568.
- Strobel, D., Choquet, Y., Popot, J. & Picot, D. (2003) *Nature* 426: 413–418.
- Xiong, J. & Bauer, C. E. (2002) *J. Mol. Biol.* 322: 1025–1037.