



LUND UNIVERSITY

ALBEGA: A Decay Spectroscopy Setup for Chemically Separated Samples

Di Nitto, A.; Yakushev, A.; Düllmann, C. E.; Khuyagbaatar, J.; Krier, J.; Ballof, J.; Bar, J.; Budzynski, T.; Cox, D. M.; Derkx, X.; Dormand, J.; Despotopulos, J. D.; Eberhardt, K.; Even, J.; Grabiec, P.; Harkness-Brennan, L.; Herzberg, R. D.; Hübner, A.; Jäger, E.; Judson, D.; Kindler, B.; Klos, H.; Kratz, J. V.; Kulawik, J.; Kurz, N.; Lens, L.; Lommel, B.; Moody, K.; Panas, A.; Prokaryn, P.; Rudolph, Dirk; Runke, J.; Rusanov, I.; Scharrer, P.; Schausten, B.; Shaughnessy, D.; Szmigiel, D.; Ward, A. J.; Wegrzecki, M.

Published in:
GSI Report

DOI:
[10.15120/GR-2015-1-MU-NUSTAR-SHE-C-06](https://doi.org/10.15120/GR-2015-1-MU-NUSTAR-SHE-C-06)

2015

[Link to publication](#)

Citation for published version (APA):

Di Nitto, A., Yakushev, A., Düllmann, C. E., Khuyagbaatar, J., Krier, J., Ballof, J., Bar, J., Budzynski, T., Cox, D. M., Derkx, X., Dormand, J., Despotopulos, J. D., Eberhardt, K., Even, J., Grabiec, P., Harkness-Brennan, L., Herzberg, R. D., Hübner, A., Jäger, E., ... Wegrzecki, M. (2015). ALBEGA: A Decay Spectroscopy Setup for Chemically Separated Samples. In *GSI Report* (Vol. 2015-1, pp. 184). GSI Helmholtzzentrum für Schwerionenforschung. <https://doi.org/10.15120/GR-2015-1-MU-NUSTAR-SHE-C-06>

Total number of authors:
39

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Download date: 28. Apr. 2024

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

ALBEGA: A decay spectroscopy setup for chemically separated samples *

A. Di Nitto^{†1}, A. Yakushev², Ch.E. Düllmann^{1,2,3}, J. Khuyagbaatar^{2,3}, J. Krier², J. Ballof⁴, J. Bar⁷, T. Budzyński⁷, D.M. Cox⁴, X. Derkx^{1,3}, J. Dormand⁴, J.D. Despotopulos⁵, K. Eberhardt^{1,3}, J. Even³, P. Grabiec⁷, L. Harkness-Brennan⁴, R.D. Herzberg⁴, A. Huebner², E. Jäger², D. Judson⁴, B. Kindler², H. Kłos⁷, J.V. Kratz¹, J. Kulawik⁷, N. Kurz², L. Lens¹, B. Lommel², K. Moody⁵, Ch. Mokry¹, A. Panas⁷, P. Prokaryn⁷, D. Rudolph⁶, J. Runke², I. Rusanov², P. Scharrer¹, B. Schausten², D. Shaughnessy⁵, D. Szmigiel⁷, A.J. Ward⁵, and M. Wegrzecki⁷

¹Mainz University, Germany; ²GSI Darmstadt, Germany; ³HIM Mainz, Germany; ⁴Liverpool University, UK; ⁵LLNL, Livermore, USA; ⁶Lund University, Sweden; ⁷ITE Warsaw, Poland

There are many on-going programmes dedicated to elucidate the nuclear structure of SuperHeavy Elements (SHE atomic number $Z \geq 104$) based on different methods [1]. The SHE are accessed in heavy-ion induced fusion-evaporation reactions, with the nuclear spectroscopy experiments typically performed at in-flight recoil separators.

An alternative approach exploits a chemical isolation system either directly or after separator [2], this was adopted in different experiments [3-5]. A significant improvement of the background conditions was observed applying this method.

A next generation setup for measurements of ALpha-BETA-Gamma decays (ALBEGA) after chemical isolation was recently built. ALBEGA is dedicated to simultaneous measurements of α particles, electrons, photons and fission fragments. Volatile chemical species adsorb on a cooled segmented Si detector. Radiation emitted in the decay is measured with surrounding detectors, cf. Fig.1. The

the detectors were assembled in a packed configuration to maximize the geometrical efficiency.




Figure 2: The α particles are emitted isotropically at various angles in the narrow channel (a). Depending on their incident angle, they pass through different effective thicknesses of detector dead layer and gas, thus undergoing energy loss to a different degree, before entering the active detector area. Accordingly, α peaks show characteristic low-energy tailing, e.g. the $^{140}\text{Ce}(^{50}\text{Ti}, 6-7n)$ reaction energy spectrum (b). The measured energies corresponding to the two particles α_1 and α_2 are indicated, they have the same initial energy but different path. The distribution pattern observed in the experiment (c) is due to the Hg retention on the inner channel Au coating, and the maximum is observed on the segments at the channel entrance.




Figure 1: A schematic of ALBEGA, where the arrows indicate the gas flow inside the tight channel (a). A picture of the α detector mounted on the cooling support and connected to the capillaries for gas flushing.

core is a “sandwich” of two segmented Si detectors with a gas channel inside, sensitive to charged particles (“ α detector”). A round channel is etched in both wafers, glued together to form a tight channel. At both sides and parallel to the α detector two thicker, segmented Si detectors are mounted (cf. Ref. [6]). The Si detectors are backed by two 3-cm-thick broad energy Ge detectors for X and γ rays. All

* We thank the ion-source and accelerator staff at GSI. Work supported by BMBF contract-No. 06MZ7164. The development of α -detector was funded by the NCBiR (Poland) under the PBS1/A9/8/2012 Project: SiDET (ID 179409).

[†] a.dinitto@gsi.de

References

- [1] M. Leino and F.-P. Hessberger, *Ann. Rev. Nucl. Part. Sci.* 54, 175 (2004).
- [2] Ch.E. Düllmann et al., *Nucl. Instrum. Met.* A551, 528 (2005).
- [3] A. Yakushev et al., *Inorg. Chem.* 53, 1624-1629 (2014).
- [4] J. Dvorak et al., *Phys. Rev. Lett.* 97, 242501 (2006).
- [5] J. Even et al., *J. Radioanal. Nucl. Chem.* 303, 2457 (2015).
- [6] M. Wegrzecki et al., *SPIE Proc.* 9291 (2014).