



LUND UNIVERSITY

Applicability of a cognitive model and treatment to worry in children and adolescents

Perrin, Sean; Bolton, Derek; Fialko, Laura

Published in:
[Host publication title missing]

2012

[Link to publication](#)

Citation for published version (APA):
Perrin, S., Bolton, D., & Fialko, L. (in press). Applicability of a cognitive model and treatment to worry in children and adolescents. In J. Rogers (Ed.), [Host publication title missing] British Association for Behavioural and Cognitive Psychotherapy. http://www.babcpconference.com/programme/symposia_2012.htm

Total number of authors:
3

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Applicability of a cognitive model and treatment to worry in children and adolescents

Sean Perrin, Lund University & King's College London; Derek Bolton, King's College London; Laura Fialko, East London Mental Health Trust; Susanna Payne, South London & Maudsley NHS Foundation Trust.

Studies have found that the cognitive model of worry developed by Dugas, Gagnon, Ladouceur and Freeston (1998) involving Intolerance of Uncertainty (IU), Cognitive Avoidance (CA), Positive Beliefs about Worry (PB), and Negative Problem Orientation (NPO) accounts for approximately 40-50% of the variance in worry frequency in adults. A recent trial of the treatment based on this model (Dugas, Brillon, Savard et al., 2010) found it to be superior to no treatment and marginally superior to applied relaxation in adults with Generalized Anxiety Disorder (GAD). In this presentation the authors discuss the results of two studies testing the applicability of this cognitive model to children and adolescents. In the first study measures of IU, CA, PB, worry and anxiety were administered to 515 British primary and secondary school students (aged 7-19 years). A path analysis was undertaken to test whether IU acted as a higher order variable for CA and PB and whether the relationship between IU/CA/PB and anxiety was mediated by worry frequency. Significant (bivariate) correlations were observed between the measures of IU, PB, CA, worry and anxiety across the age range. However separate path models had to be fitted for children (aged 7-12 years) and adolescents (aged 13-19) with PB being dropped from the child model. CA was related to anxiety only through worry in children while IU showed direct paths to worry and anxiety in both children and adolescents. In the second study we present results from a case series where a modified version of the treatment based on the Dugas et al (1998) led to an 80% reduction in GAD diagnoses in 16 youth (aged 7-17 years) clinically referred for anxiety. Further testing and refinement of this cognitive model and its treatment may lead to further improvements in our understanding of worry in youth and its treatment. Possible modifications and refinements are discussed.