



LUND UNIVERSITY

High bacterial respiratory quotients in net heterotrophic lakes

Berggren, Martin; del Giorgio, Paul A.

2011

[Link to publication](#)

Citation for published version (APA):

Berggren, M., & del Giorgio, P. A. (2011). *High bacterial respiratory quotients in net heterotrophic lakes*. Abstract from 2011 ASLO Aquatic Sciences Meeting: Limnology and Oceanography in a Changing World, San Juan, Puerto Rico, United States.

Total number of authors:

2

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Berggren M (oral) and PA del Giorgio. 2011. High bacterial respiratory quotients in net heterotrophic lakes. 2011 ASLO Aquatic Sciences Meeting: Limnology and Oceanography in a Changing World, presented on February 18, 2011, in San Juan, Puerto Rico, USA

ABSTRACT

In most studies of bacterioplankton carbon metabolism, the need to assume a respiratory quotient (RQ; mole of CO₂ produced per mole of O₂ consumed) introduces a fundamental source of uncertainty. Many studies have assumed a fixed value, often close to 1, but this assumption has little empirical support, and ignores potential variability in RQ caused by physiological mechanisms and by varying oxygen contents of respired substrates. Here we present over 50 direct measurements of bacterioplankton RQ that we have carried out in epilimnetic samples of lakes distributed across the temperate and boreal regions of Québec, using O₂ and CO₂ sensitive mini sensors attached to the inside of closed incubation flasks. RQ was mostly in the range of 0.5-2, with the lowest values in net autotrophic and the highest in net heterotrophic systems. A plausible explanation is that bacteria were utilizing oxygen-rich organic acids in the net heterotrophic lakes (e.g., formed by photo-chemical processes), and relatively oxygen-poor phytoplankton exudates in the net autotrophic lakes. The results suggest that respiration measured with the O₂ consumption method assuming RQ=1 may underestimate the role of bacterial respiration as cause of lake net heterotrophy.