



LUND UNIVERSITY

Determination of hyperfine structure and g j factors in the sequences of 2 D states in alkali atoms using a tunable dye laser

Svanberg, Sune; Belin, G

Published in:
Journal of Physics B: Atomic and Molecular Physics

DOI:
[10.1088/0022-3700/7/3/024](https://doi.org/10.1088/0022-3700/7/3/024)

1974

[Link to publication](#)

Citation for published version (APA):
Svanberg, S., & Belin, G. (1974). Determination of hyperfine structure and g j factors in the sequences of 2 D states in alkali atoms using a tunable dye laser. *Journal of Physics B: Atomic and Molecular Physics*, 7(3).
<https://doi.org/10.1088/0022-3700/7/3/024>

Total number of authors:
2

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Determination of hyperfine structure and g_j factors in the sequences of 2D states in alkali atoms using a tunable dye laser

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

1974 J. Phys. B: At. Mol. Phys. 7 L82

(<http://iopscience.iop.org/0022-3700/7/3/024>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 130.235.188.41

The article was downloaded on 25/07/2011 at 08:28

Please note that [terms and conditions apply](#).

LETTER TO THE EDITOR

Determination of hyperfine structure and g_J factors in the sequences of 2D states in alkali atoms using a tunable dye laser

S Svanberg and G Belin

Department of Physics, Chalmers University of Technology, Fack, S-402 20
Göteborg, Sweden

Received 28 December 1973

Abstract. We have extended recently reported hyperfine structure studies of 2D states in alkali atoms to several new states. The 2D states were populated in a two-step excitation scheme, involving a conventional RF lamp and a cw tunable dye laser. Using level crossing and optical double resonance techniques we have determined the hyperfine structure of the 8 $^2D_{3/2,5/2}$ states in ^{87}Rb and the 11, 12, 13, 14 $^2D_{3/2,5/2}$ states in ^{133}Cs . Further, g_J factors for several of these states were determined.

The alkali atoms have a comparatively simple structure and are particularly suitable for spectroscopic investigations as well as for theoretical calculations. These atoms provide natural test cases for modern many-body perturbation calculations. The advent of such detailed computational schemes has renewed the interest for the experimental investigation of the fine and hyperfine structure of excited states. In the alkali atoms only the sequences of 2P states are directly optically accessible in a single-step excitation process of the electric dipole type. Consequently, for a long time only these excited alkali states were studied with high resolution resonance methods. Excited non-P states can be populated in the cascade decay of highly excited 2P levels. Several lower 2S and 2D states have recently been investigated at Columbia Radiation Laboratory using cascade decoupling (Gupta *et al* 1972a) and cascade RF spectroscopy techniques (Gupta *et al* 1972b, Gupta *et al* 1973).

In order to get access to more highly excited non-P states, the feasibility of a two-step excitation scheme, taking advantage of the recent developments in the technology of tunable dye lasers, has been demonstrated (Svanberg *et al* 1973a, b). In a first excitation step the strong D_1 or D_2 line from a powerful RF discharge lamp is used to populate one of the levels in the first excited P doublet. From this level transitions are induced to an excited S or D level using a cw dye laser, tuned to the corresponding transition. The hyperfine structure (HFS) of several 2D levels excited in this way, could be measured by the level crossing (LC) and optical double resonance (ODR) methods (Svanberg 1973, Svanberg *et al* 1974). We have now extended these measurements to several other 2D states in Rb and Cs. These states have become accessible using new dyes, extending the wavelength region covered by rhodamine 6G, which was used in the previous investigations. In this way we have now obtained HFS information on the 8 $^2D_{3/2,5/2}$ states in ^{87}Rb and on the 11, 12, 13, 14 $^2D_{3/2,5/2}$ states in ^{133}Cs . Furthermore, the Landé g_J factor has been determined for several of these states.

Our experimental arrangement is based on the set-up described by Svanberg and Rydberg (1969) and is similar to the arrangement used in the earlier laser investigations (Svanberg *et al* 1973b, 1974). An alkali vapour scattering cell is placed in a stabilized

oven (70–100 °C) and is surrounded by Helmholtz coil systems, generating a well defined external magnetic field. The strong D_1 and D_2 lines from an alkali RF lamp are isolated by a Schott colour glass filter and the light is incident upon the scattering cell in the direction of the magnetic field. In this way the first excited 2P levels are populated in a first excitation step. In the second step atoms are transferred to a highly excited 2D level by means of an appropriately tuned dye laser, the intense light beam of which traverses the scattering cell at right angles to the external field. We used a Coherent Radiation Model 490 jet stream dye laser, pumped by a Coherent Radiation Model CR8 argon ion laser. In the present measurements the dye rhodamine 110, dissolved in ethylene glycol, was used. The multi-mode bandwidth of the dye laser was about 0.5 Å. Fluorescent light following the two-step excitation was detected in the direction of the magnetic field. A Bausch and Lomb 500 mm grating monochromator with the slit parallel to the laser beam was used and the wavelength selected light was detected with a photomultiplier tube. The lock-in technique was employed for the registration of LC and ODR signals. In the LC measurements a rotating polarizer was used in the detected beam, whereas the RF field, generated in a two-turn coil around the scattering cell, was modulated in the ODR experiments. The lock-in output signal was stored in a Laben Correlation multichannel analyser, which also controlled a repetitive magnetic field sweep.

In the sequence of 2D states in ^{133}Cs , the rhodamines 6G and 110 cover the members with principal quantum number n , ranging from 8 to 14. Out of these the states




Figure 1. Level crossing signals for the 11, 12, 13, and 14 $^2D_{3/2}$ states in ^{133}Cs . The sampling time for each of the curves is 0.5–1 h.

with $n = 8, 9, 10$ were investigated in the previous works, leaving the levels with $n = 11, 12, 13, 14$ to be studied here. For ^{87}Rb the $8\ ^2\text{D}$ doublet is investigated in this work extending the previous experiments on the 6 and 7 ^2D states. All the $^2\text{D}_{3/2}$ states were studied with the LC method. In ^{133}Cs ($I = \frac{7}{2}$) as well as in ^{87}Rb ($I = \frac{3}{2}$) there are three detectable level crossings besides the zero field crossing (Hanle effect). However, in ^{133}Cs the second and third crossings are strongly overlapping, resulting in an unresolved signal structure. Experimental curves are shown in figures 1 and 2. Here the excitation and detection lines are also indicated. For an accurate evaluation of the HFS, the regions around the individual crossing signals were covered with short magnetic field sweeps. Care was taken to eliminate influences of asymmetries, time




Figure 2. Optical double resonance curve for the $8\ ^2\text{D}_{5/2}$ state, and level crossing curve for the $8\ ^2\text{D}_{3/2}$ state of ^{87}Rb .

Table 1. Absolute values of hyperfine interaction constants, and g_I values for ^2D states in ^{87}Rb and ^{133}Cs

Atom	State	$ a $ MHz	$ b $ MHz	g_I
^{87}Rb	$8\ ^2\text{D}_{3/2}$	2.85(3)	0.16(5) $b/a > 0$	
	$8\ ^2\text{D}_{5/2}$	1.2(2)		1.1997(15)
^{133}Cs	$11\ ^2\text{D}_{3/2}$	1.055(15)		
	$11\ ^2\text{D}_{5/2}$	0.24(6)		1.1994(10)
	$12\ ^2\text{D}_{3/2}$	0.758(12)		0.8001(10)
	$12\ ^2\text{D}_{5/2}$	0.19(5)		1.1996(10)
	$13\ ^2\text{D}_{3/2}$	0.556(8)		
	$13\ ^2\text{D}_{5/2}$	0.14(4)		1.1998(10)
	$14\ ^2\text{D}_{3/2}$	0.425(15)		

constants, and the earth's field. Thus the magnetic dipole interaction constant a , and for ^{87}Rb also the electric quadrupole interaction constant b , could be determined. The results are shown in table 1.

The $^2\text{D}_{5/2}$ states were all investigated with ODR. In all states the HFS turned out to be very small, and at a resonance frequency of about 110 MHz the Paschen-Back region for the HFS is well reached. Thus, $2I+1$ resonances of equal width and intensity are expected symmetrically spread around a centre of gravity, corresponding to the particular g_J value of the state. In all cases only an unresolved structure was obtained. The halfwidth of this structure was determined, extrapolated to zero RF amplitude and low vapour pressure. From the measured halfwidth and theoretically estimated widths for the individual signal contributions, only very rough values for the a factor could be obtained. From the position of the centre of gravity the g_J value was determined. The magnetic field calibration, especially critical in these last measurements, was performed using optical pumping. An experimental ODR curve is included in figure 2. ODR experiments were also performed in the $12\ ^2\text{D}_{3/2}$ state for checking the g_J value. The results of the ODR measurements are recorded in table 1. It should be noted, that these experiments only give the absolute values of the coupling constants. The signs, which are not necessarily positive due to the influence of core polarization (Gupta *et al* 1972b), will be determined in subsequent experiments. The g_J values of the studied $^2\text{D}_{5/2}$ and $^2\text{D}_{3/2}$ states are found to agree well with the values predicted by the Landé formula: 1.2005 and 0.7995, respectively.

The authors are very grateful to Professor I Lindgren for his kind interest and support. This work was financially supported by the Swedish Natural Science Research Council.

References

- Gupta R, Chang S and Happer W 1972a *Phys. Rev. A* **6** 529-44
Gupta R, Chang S, Tai C and Happer W 1972b *Phys. Rev. Lett.* **29** 695-8
Gupta R, Happer W, Lam L K and Svanberg S 1973 *Phys. Rev. A* **8** 2792-810
Svanberg S 1974 *Proceedings of the Laser Spectroscopy Conference, Vail, Colorado, 1973*, eds R G Brewer and A Mooradian (New York: Plenum)
Svanberg S and Rydberg S 1969 *Z. Phys.* **227** 216-33
Svanberg S, Tsekeris P and Happer W 1973a *Bull. Amer. Phys. Soc.* **18** 611
——— 1973b *Phys. Rev. Lett.* **30** 817-20
——— 1974 *Phys. Rev. A*, to be published