



LUND UNIVERSITY

Characteristics for Airspace Dimension Test (ADT) – A novel technique for lung diagnosis with nanoparticles

Löndahl, Jakob; Jakobsson, Jonas; Andersson, Tobias; Wollmer, Per

2014

[Link to publication](#)

Citation for published version (APA):

Löndahl, J., Jakobsson, J., Andersson, T., & Wollmer, P. (in press). *Characteristics for Airspace Dimension Test (ADT) – A novel technique for lung diagnosis with nanoparticles*. Abstract from International Aerosol Conference, 2014, Busan, Korea, Republic of.

Total number of authors:

4

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Characteristics for Airspace Dimension Test (ADT) – A novel technique for lung diagnosis with nanoparticles

Jakob Löndahl^{1*}, Jonas Jakobsson¹, Tobias Andersson², Per Wollmer²

¹Div. of Ergonomics and Aerosol Technology (EAT), Lund University, 221 00, Lund, Sweden

²Dept. of Clinical Sciences Malmö, Lund University, 205 02, Malmö, Sweden

*Corresponding author: jakob.londahl@design.lth.se

Keywords: Nanoparticles, Lung-particle interaction, Emphysema, Respiratory diagnosis

Chronic obstructive pulmonary disease (COPD) is the fourth most common cause of death globally. A general manifestation of COPD is emphysema, i.e. breakdown of lung tissue. Emphysema leads to enlargements of the peripheral airspaces that are difficult to observe without advanced and expensive tools such as X-ray computed tomography (CT) or magnetic resonance imaging with hyperpolarized ³He. This work describes a novel technique, Airspace Dimension Test (ADT), for diagnosis of emphysema with aerosolized nanoparticles.

Nanoparticles in the size range < 300 nm are known to deposit in the airways almost exclusively by diffusion, a process determined by residence time and distance to nearby surfaces. Emphysema corresponds to enlarged alveoli and therefore increased mean diffusion distances in the peripheral lung, with decreased deposition of inhaled nanoparticles as a consequence (Löndahl et al., 2012).

The ADT-instrument, shown in Figure 1, measures the deposition fraction of monodisperse nanoparticles during a well-defined single breath procedure including a 3-20 s breath-hold. It consists of three main parts: 1) aerosol generation and conditioning, 2) inhalation system and 3) particle detection and analysis. The instrument was characterized and tested for sensitivity with respect to particle size, diffusion time and inter-subject variability on a group consisting of seven young, healthy volunteers.




Figure 1. Instrumental set-up.

It was found that the ADT-instrument had a sensitivity of less than 0.3% in the measured lung deposition fractions (Figure 2). The deposition fraction increased with increased breath-hold time and decreased particle size as is expected according to diffusion theory. The inter-subject variability was significantly larger than the measurement uncertainty. The measured values were in accordance with values calculated with the multiple path particle dosimetry (MPPD) lung deposition model.




Figure 2. Deposition fractions for 7 young healthy individuals inhaling 50, 75 and 100 nm particles. Data from the simulation with the MPPD model are included.

From the theory of Brownian diffusion it is expected that the measured deposition fractions will correlate with the dimensions of the peripheral airspaces. ADT could then provide a fast, safe and cost-effective alternative for diagnosis of COPD.

Acknowledgements

We gratefully acknowledge The Swedish Research Council (project 621-2011-3560), the Crafoord foundation, Vinnova and the Swedish heart and lung foundation for financial support.

References

Löndahl J, Swietlicki W, Rissler J, Bengtsson A, Boman C, Blomberg A, Sandström T. (2012), *Particle and Fibre Toxicology*, 2012, 9:30