

# LUND UNIVERSITY

High m(T) pion andproton correlations in central Pb Pb collisions at s(NN)\*\*(1/2) = 17-GeV

Bearden, I.G.; Lörstad, Bengt

Published in: nucl-ex/0305014

2003

Link to publication

Citation for published version (APA): Bearden, I. G., & Lörstad, B. (2003). High m(T) pion andproton correlations in central Pb Pb collisions at  $s(NN)^{**}(1/2) = 17$ -GeV. Unpublished.

Total number of authors: 2

#### **General rights**

Unless other specific re-use rights are stated the following general rights apply:

- Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the
- legal requirements associated with these rights · Users may download and print one copy of any publication from the public portal for the purpose of private study
- You may not further distribute the material or use it for any profit-making activity or commercial gain
  You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: https://creativecommons.org/licenses/

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

### LUND UNIVERSITY

**PO Box 117** 221 00 Lund +46 46-222 00 00

## High $m_T$ pion and proton correlations in central PbPb collisions at $\sqrt{s_{NN}} = 17 \text{GeV}$

I.G. Bearden,<sup>1</sup> H. Bøggild,<sup>1</sup> J. Boissevain,<sup>2</sup> P.H.L. Christiansen,<sup>1</sup> L. Conin,<sup>3</sup> J. Dodd,<sup>4</sup> B. Erazmus,<sup>3</sup>

S. Esumi,<sup>5</sup> C.W. Fabjan,<sup>6</sup> D. Ferenc,<sup>7</sup> A. Franz,<sup>6</sup> J.J. Gaardhøje,<sup>1</sup> A.G. Hansen,<sup>1</sup> O. Hansen,<sup>1</sup> D. Hardtke,<sup>8</sup>

H. van Hecke,<sup>2</sup> E.B. Holzer,<sup>6</sup> T.J. Humanic,<sup>8</sup> P. Hummel,<sup>6</sup> B.V. Jacak,<sup>9</sup> K. Kaimi,<sup>5, \*</sup> M. Kaneta,<sup>5</sup>

T. Kohama,<sup>5</sup> M. Kopytine,<sup>9</sup> M. Leltchouk,<sup>4</sup> A. Ljubičić Jr.,<sup>7</sup> B. Lörstad,<sup>10</sup> N. Maeda,<sup>5</sup> L. Martin,<sup>3</sup>

A. Medvedev,<sup>4</sup> M. Murray,<sup>11</sup> H. Ohnishi,<sup>5</sup> G. Paić,<sup>6</sup> S.U. Pandey,<sup>8</sup> F. Piuz,<sup>6</sup> J. Pluta,<sup>3</sup> V. Polychronakos,<sup>12</sup>

M. Potekhin,<sup>4</sup> G. Poulard,<sup>6</sup> D. Reichhold,<sup>8</sup> A. Sakaguchi,<sup>5</sup> J. Schmidt-Sørensen,<sup>10</sup> J. Simon-Gillo,<sup>2</sup>

W. Sondheim,<sup>2</sup> T. Sugitate,<sup>5</sup> J.P. Sullivan,<sup>2</sup> Y. Sumi,<sup>5</sup> W.J. Willis,<sup>4</sup> K. Wolf,<sup>11, \*</sup> N. Xu,<sup>2</sup> and D.S. Zachary<sup>8</sup>

(The NA44 Collaboration, December 28, 2007)

<sup>1</sup>Niels Bohr Institute, University of Copenhagen, Denmark

<sup>2</sup>Los Alamos National Laboratory, Los Alamos, NM 87545

<sup>3</sup>Nuclear Physics Laboratory of Nantes, 44072 Nantes, France

<sup>4</sup>Department of Physics, Columbia University, New York, NY 10027

<sup>5</sup>Hiroshima University, Higashi-Hiroshima 739-8526, Japan

<sup>6</sup>CERN, CH-1211 Geneva 23, Switzerland

 $^{\gamma}Rudjer$ Bošković Institute, Zagreb, Croatia

<sup>8</sup>Department of Physics, The Ohio State University, Columbus, OH 43210

<sup>9</sup>State University of New York, Stony Brook, NY 11794

<sup>10</sup>Department of Physics, University of Lund, S-22362 Lund, Sweden

<sup>11</sup>Texas A&M University, College Station, Texas 77843-3366

<sup>12</sup>Brookhaven National Laboratory, Upton, New York 11973

For central PbPb collisions at  $\sqrt{S_{nn}} = 17.3$  GeV we have made the first two-dimensional measurement of the pp correlation function. These data extend the range of previous studies of HBT radii by a factor of two in  $m_T$ . They are consistent with a hydrodynamic interpretation and microscopic models that include hadronic rescattering and transverse expansion. We also report new data on pion correlations. The two particle correlations of negative pions at  $m_T = 0.92$  GeV imply source radii that are smaller than typical hydrodynamic fits and transport model simulations. It is possible that these fast pions may have left the source before the build up of hydrodynamic flow.

PACS numbers: 25.75.Gz, 25.75.Ld

Experimental studies of high-energy nuclear collisions at the AGS, SPS and RHIC accelerators (with  $\sqrt{S_{nn}} =$  $2 \rightarrow 200 \text{ GeV}$ ) have revealed many interesting features of hot and dense nuclear matter, and some characteristic signatures of a quark-gluon-plasma phase have been reported [1]. If such a high density state were formed in the initial stages of the reaction the high initial presure would result in a significant transverse flow of the hadrons that originate from the participant zone. Depending on the time scale and the amount of rescattering the system might be expected to be in local thermodynamical equilibrium. This conjecture is supported by the linear increase of the single particle inverse slopes with mass [2, 3].

The Hanbury-Brown Twiss, (HBT), technique uses the interference of particle wavefunctions to infer the angular size of stars or the length scales of nuclear systems from the two particle correlation function [4]. The "radii" measured by HBT can be thought of as "lengths of homogeneity" of the source which depend upon velocity and/or temperature gradients [5, 6, 7]. Heavy Ion HBT measurements over a wide range of energies [8, 9, 10, 11, 12, 13, 14, 15, 16] show a drop of radii with  $m_T \equiv \sqrt{p_T^2 + m^2}$  consistent with a hydrodynamical interpretation [17, 18, 19, 20].

Faster particles are more likely to come from initial collisions of the incoming baryons, so at some momentum we might expect the hydrodynamic approach to break down. This may have already been seen at RHIC ( $\sqrt{s_{NN}} =$ 200GeV) where the strength of elliptic flow falls below the hydrodynamic prediction for  $p_T > 2\text{GeV}/c$  [21]. One might expect this to occur at lower  $p_T$  at  $\sqrt{s_{NN}} = 17\text{GeV}$ since the multiplicity is lower.

NA44 is a focusing spectrometer [13]. The dipole and quadrupole magnets produce a magnified image of the target in the tracking detectors. This has the advantage that particles that are close in momentum are not necessarily close in position. Since all tracking is done after the magnets we only have to reconstruct straight lines. A high resolution pad chamber sits in the focal plane and gives the magnitude of the momentum. Downstream of the focus the direction of tracks is measured by strip chambers and scintillating hodoscopes. The momentum resolution, including effects of multiple scattering in the target, is  $\approx 11 \text{MeV}/c$  for  $p_x$ ,  $p_y$ , and  $p_z$ .

A beam counter selected events for which single lead ions hit the 3.8mm lead target. Behind the target, two scintillator bars were used to trigger on high multiplicity events. An annular silicon detector with 512 pads which was not part of the trigger measured  $dN^{\pm}/d\eta$  ( $\eta = \ln \tan \theta/2$ ) in the pseudorapidity range  $1.5 \leq \eta \leq 3.3$ . We used events in the top 18% of the multiplicity distribution as we did for our earlier PbPb results [13, 14]. For all data sets except the lower  $m_T$  protons the spectrometer was set to accept particles of momentum 6-9 GeV/cand was positioned at 131 mrad with respect to the beam axis. The setting gave a  $p_T$  window of 0.7–1.4 GeV/c. For the lower  $m_T$  proton sample the momentum window was 5.2-8.0 GeV/c and the spectrometer angle was 44 mrad. The resulting  $p_T$  range was 0.18-0.50 GeV/c. The rapidity range of the data was 2.4-2.9 and the system is symmetric about y=2.9 the center of mass rapidity. The  $p_T$  and y acceptance can be found in [22].

Particles were identified by combining time of flight measurements from the hodoscopes with velocity information from three threshold-type gas Cherenkov counters (C1, C2, TIC). C2 was set to fire only on pions. For the higher momentum proton data C1 was used to reject kaons. For the lower  $m_T$  proton data kaons were rejected exclusively by their flight time. For the  $\pi^-$  sample the TIC was used to confirm that each track was a pion [23]. Contamination of the samples by other particles was less than 2%. However there is significant feed-down of hyperons into the proton sample, ranging from 28% for the lower  $m_T$  sample to 13% for  $\langle m_T \rangle = 1.45$  GeV [22]. This tends to smear the correlation function and was accounted for by including a fraction of "fake" protons in the Monte Carlo [24]. The spectrum of hyperons was taken from RQMD with the yield scaled to match NA49 and WA97 results [25, 26].

The two-particle correlation function is defined by

$$C_2(\vec{p}_1, \vec{p}_2) = \frac{P_2(\vec{p}_1, \vec{p}_2)}{P_1(\vec{p}_1)P_1(\vec{p}_2)} \approx \frac{Real(\vec{p}_1, \vec{p}_2)}{Back(\vec{p}_1, \vec{p}_2)}, \quad (1)$$

where the numerator is the joint probability of detecting two particles with momenta  $\vec{p_1}$  and  $\vec{p_2}$ , while the denominator is the product of the probabilities of detecting the single particles. The denominator was obtained by mixing tracks from two randomly selected different events. For protons, the attractive effect of the strong interaction competes with the negative correlation due to Fermi-Dirac statistics and Coulomb repulsion. This results in characteristic "dip-peak" structure in the twoproton correlation function. The height of the correlation peak decreases as the source size increases [27, 28, 29, 30].

The true correlation function is distorted by the momentum resolution, Coulomb repulsion and by residual two particle correlations in the background. The magnitude of these effects depends on the size of the source. For the pions we we evaluated them using an iterative Monte Carlo (MC) simulation. A trial correlation function was used to generate events that we tracked through the spectrometer and analyzed in exactly the same way as the data. The Coulomb wave method was employed to simulate Coulomb effects [31]. This method was used to derive correction factors to the correlation function and the process was repeated until the changes in the corrections from step to step were insignificant [10, 32]. The theoretical correlation function for the pp data was generated by selecting proton pairs from a given source model, calculating the weight due to quantum statistics and final state interactions (strong and Coulomb) and propagating the particles through the Monte Caro. We searched for a source distribution that produced the minimum  $\chi^2$  with the data after it was fed through the Monte Carlo [24].

 $C_2(\vec{p_1}, \vec{p_2})$  is a function of 6 variables; 3 for the total momentum  $\vec{P} = \vec{p_1} + \vec{p_2}$ , and 3 for the momentum difference  $\vec{Q} = \vec{p_1} - \vec{p_2}$ . We decompose  $\vec{Q}$  into  $Q_L$ , the projection of the momentum difference onto the beam axis,  $Q_{out}$  which is parallel to  $\vec{P}$  and  $Q_{side}$  which is perpendicular to  $Q_{out}$  and  $Q_L$ . The data are analysed in the Longitudinal Co-Moving System in which  $p_{z1} = -p_{z2}$ .

To measure correlation functions efficiently we trigger on events with pairs of particles that have a small momentum difference in one direction while allowing large momentum differences in the other two dimensions. In our "vertical setting" the quadrupoles produce a small momentum acceptance in the horizontal space  $p_x$ , and a wide acceptance in the vertical direction  $p_y$ . Since the spectrometer lies in the horizontal plane this setting allows only a small range in  $Q_{side}$  but a large range in  $Q_{out}$ . In the "horizontal setting" we have a wide momentum acceptance in  $p_x$  and a small acceptance in  $p_y$ . This setting allows only a small range in  $Q_{side}$  but a large range in  $Q_{out}$ . Both settings have a large acceptance in  $Q_{Long}$ . The lower  $m_T$  proton data were taken in the vertical setting while all other data were taken in the horizontal setting.

For pions we used the maximum likelihood method to fit C2 with

$$C_2(Q_T, Q_L) = D(1 + \lambda e^{-Q_T^2 R_T^2 - Q_L^2 R_L^2}), \qquad (2)$$

where  $R_T$  and  $R_L$  parametrize the size of the source in the transverse and beam directions and D is a free parameter used for normalization. The  $\lambda$  factor is a measure of the chaoticity of quantum states of the source; the fraction of pions from resonances and experimental factors that decrease the correlation function. For these data  $Q_T \approx Q_{out}$ . Figure 1 shows projections of the correlation function and the fit onto the  $Q_T$  and  $Q_L$  axis. The  $\lambda$  parameter is larger than for our lower  $p_T$  pion measurements. This is expected since the effect of resonances, which tend to wash out  $C_2$  and reduce  $\lambda$ , should be less important as  $p_T$  increases. The systematic uncertainties in the fit parameters reflect the effect of (i) cut parameters to define a track, (ii) cut parameters to select pairs, (iii) momentum resolution, (iv) two-track resolution, (v) momentum distribution of particle production in MC, and (vi) fitting to finite bins.



FIG. 1: The  $\pi^-$  correlation function for  $\langle m_T \rangle = 0.92 GeV$  projected onto the (a)  $Q_T$  and (b) the  $Q_L$  axies. Only statistical errors are shown. The lines are projections of the fit.



FIG. 2: Projections of the pp correlation function onto the  $Q_{inv}$  axis for samples with different  $\langle m_T \rangle$ . Only statistical errors are shown. The lines show the projections of the fit.

Figure 2 shows the pp correlation function as a function of  $Q_{inv} \equiv \sqrt{Q^2 - \Delta E^2}$  for four bins in  $\langle m_T \rangle$ . At  $\langle m_T \rangle =$ 1 GeV our result is consistent with NA49 [33].  $R_{inv}$ increases from pPb to SPb and finally to PbPb, [24]. We have also analysed the pp correlation in two dimensions to extract  $R_T$  and  $R_L$ . Table I summarizes our data and compares them to two models.

RQMD 2.04 is a string model which includes reinteractions between produced hadrons [34]. It gives a good description of the proton data, as it does for our low  $m_T$ 

TABLE I: PbPb source radii versus  $m_T$  from data and models.

	$m_T$	$R_T$ (fm)			$R_L \ (\mathrm{fm})$		
	$(\mathrm{GeV})$	Data	Rqmd	Nexus	Data	Rqmd	Nexus
р	1.00	$3.6^{+0.7}_{-0.6}$	3.7	3.1	$3.5^{+1.2}_{-0.9}$	4.6	1.7
	1.21	$4.0^{+0.4}_{-0.3}$	3.6	3.2	$3.5^{+0.6}_{-0.5}$	3.8	1.9
	1.29	$3.2^{+0.2}_{-0.2}$	3.5	3.2	$3.3^{+0.6}_{-0.4}$	3.5	2.0
	1.43	$3.4^{+0.3}_{-0.2}$	3.4	2.8	$2.9^{+0.4}_{-0.3}$	3.1	2.0
$\pi^{-}$	0.92	$2.2\pm0.3$	3.9	4.5	$1.9\pm0.3$	3.1	2.0

pion results [13, 35]. The decrease of the radii with  $m_T$  has already been noticed in the model for pions and results from correlations between position and momentum caused by flow [36]. This decrease is more rapid for  $R_L$  than for  $R_T$  since the longitudinal flow is stronger. Nexus 2.00 uses pertubative QCD to treat hard nucleon-nucleon interactions and pomeron exchange for soft interactions [37]. It produces proton radii that are too small. Turning on hadronic reinteractions and resonance production results in larger radii closer to the data. RQMD fails to reproduce both pion radii at  $m_T = 0.92$ GeV. Nexus reproduces  $R_L$  but overestimates  $R_T$  by a factor of two.

Under certain conditions [17] one can derive analytic expressions for the radii:

$$\frac{1}{R_T^2} \approx \frac{1}{R^2} \left( 1 + \eta_f^2 \frac{m_T}{T} \right); \tag{3}$$

$$\frac{1}{R_L^2} \approx \frac{1}{\tau_0^2 \Delta \eta^2} \left( 1 + \Delta \eta^2 \frac{m_T}{T} \right); \tag{4}$$

where R is the geometrical size of the source,  $\tau_0$  and  $T_0$ are the time and temperature of freeze-out,  $\delta\eta$  represents the width in rapidity and  $\eta_f$  is the transverse rapidity. The radii also depend on the particle's mass but this effect is very small for reasonable values of  $\eta_f$ . Since  $\eta_f < \Delta\eta$ ,  $R_L$  should drop more rapidly with  $m_T$  than  $R_T$  because the longitudinal flow is stronger than the transverse flow.

Figure 3 shows the  $m_T$  dependence of  $R_T$  and  $R_L$  for all NA44 correlation and coalescence measurements [13, 14, 38]. The proton coalescence analysis gives only one radius which is mainly sensitive to the transverse size, (see Eqn. 6.3 of [39]). These radii are in good agreement with the pp correlation data.  $R_L$  decreases faster with  $m_T$  than does  $R_T$  as expected from RQMD simulations and Eqns 3 and 4. The lines in Fig. 3 are the result of fitting our systematics to the form  $1/R^2 = p0 + p1 \cdot M_T$ . The bands show the errors on the fits. The  $\chi^2$ /ndf is poor because the  $\pi^-$  radii at  $\langle m_T \rangle \approx 0.92$  GeV are smaller than the trend expected from our other data. Removing the high  $m_T \pi^-$  radii from the fit gives  $\chi^2/\text{ndf} \approx 1$  but has a negligible effect on the fit parameters.



FIG. 3: NA44 pion, kaon and proton source sizes versus  $m_T$  [13, 14, 38]. Both radii are fit to the form  $1/R^2 = p0+p1 \cdot M_T$ . The lines and bands show the results and errors of the fits.

At  $m_T = 0.22 \text{GeV}$  the negative pions have source radii that are 9-11% smaller than for the positive pions. However the  $\pi^-$  and  $\pi^+$  correlation functions themselves are consistent and non Gaussian in shape [13]. RQMD simulations suggest that this is due to resonance decays, particularly of the  $\omega$ . Resonances tend to reduce the  $\lambda$  parameter when fitting  $C_2$  to Eqn. 2. This effect decreases rapidly with  $p_T$  [36]. It is possible that interaction with the residual nuclear charge could reduce the  $\pi^-$  radius as compared to the  $\pi^+$  radius. Again this effect should be stronger at low  $p_T$ .

In summary, we have made the first two-dimensional measurement of the pp correlation function in relativistic nucleus-nucleus collisions. NA44 has measured HBT radii over a significantly larger  $m_T$  range than any other experiment. The pp data, and all our previous HBT results, are consistent with a hydrodynamical interpretation and microscopic models that include hadronic rescattering and transverse expansion.  $R_L$  drops more rapidly with  $m_T$  than  $R_T$  because the longitudinal flow is stronger than the transverse flow. At  $m_T = 0.92 \text{GeV}$ we find that the  $\pi^-$  radii are somewhat smaller than expected from the trend of our other data and are not easily explained by either hydrodynamic fits or the RQMD model. It may be that these fast pions left the source before the buildup of flow and so offer a glimpse of the hadronic system at an early point in its expansion.

We thank the staff of the CERN accelerators for

their excellent work. We are grateful for financial support from the Danish Natural Science Research Council; the Japanese Society for the Promotion of Science and the Ministry of Education, Science and Culture; the Swedish Science Research Council; the Austrian Fond für Förderung der Wissenschaftlichen Forschung; the US National Science Foundation and the Department of Energy.

- \* Deceased
- Proceedings of Quark Matter 2001, T.J. Hallman, D.E. Kharzeev, J.T. Mitchell and T. Ullrich , [Nucl. Phys. A698, 1c-676c (2002) and Quark Matter 2002 http://alice-france.in2p3.fr/qm2002/.
- [2] I. G. Bearden et al., Phys. Rev. Lett. 78, 2080 (1997).
- [3] I. G. Bearden et al., Eur. Phys. J. C 23, 237 (2002).
- [4] R. Hanbury-Brown, R. Q. Twiss, Nature **178** 1046 (1956) and G. Goldharber, S. Goldharber, W. Lee, A. Pais Phys. Rev. **120** 300 (1960).
- [5] Yu. M. Sinyukov, Nucl. Phys. A566 589c (1994).
- [6] T. Csörgő and B. Lörstad, Phys. Rev. C 54 1390 (1996).
- [7] U. A. Weidemann and U. Heinz, Phys. Rep **319** 145 (1999).
- [8] L. Ahle, et al., Phys. Rev. C66 054906 (2002).
- [9] A. Bamberger et al., Phys. Lett. B203, 320 (1988); T. Alber et al., Z. Phys. C66, 77 (1995).
- [10] H. Bøggild et al., Phys. Lett. **B302**, 510 (1993).
- [11] H. Appelshäuser et al., Eur. Phys. J. C2, 661 (1998).
- [12] M. M. Aggarwal et al., Eur. Phys. J. C16, 445 (2000).
- [13] I. G. Bearden *et al.*, Phys. Rev. C58, 1656 (1998).
- [14] I. G.!Bearden et al., Phys. Rev. Lett. 87, 112301 (2001).
- [15] C. Adler *et al.*, Phys. Rev. Lett. **87**, 082301 (2001).
- [16] K. Adcox et al., Phys. Rev. Lett. 88, 192302 (2002).
- [17] U. W. Heinz and B. V. Jacak, Ann. Rev Nucl. Part. Sci 49 529 (1999).
- [18] S. V. Akkelin and Yu. M. Sinyukov, Phys. Lett. B356, 525 (1995).
- [19] U. A. Wiedemann, P. Scotto, and U. Heinz, Phys. Rev. C53, 918 (1996).
- [20] T. Csörgő and B. Lörstad, Phys. Rev. C54, 1390 (1996).
- [21] C. Adler et al., Phys.Rev.Lett. 90 032301 (2003).
- [22] I. G. Bearden et al., Phys. Rev. C 66, 044907 (2002).
- [23] A. Braem *et al.*, Nucl. Inst. Meth. A **409** 426 (1998).
- [24] H. Bøggild *et al.*, Phys. Lett. **B458** 181 (1999).
- [25] S. V. Afanasiev et al. J. Phys. G 28, 1761 (2002)
- [26] F. Antinori et al., Eur. Phys. J. C14 633 (2000).
- [27] S. E. Koonin, Phys. Lett. **70B** 43 (1977).
- [28] R. Lednicky, V.L. Lyuboshitz, Yad. Fiz. **35** 1316 (1982) (Sov. J. Nucl. Phys. **35** 770(1982)).
- [29] S. Pratt, M.B. Tsang, Phys. Rev. C36 2390 (1987).
- [30] C. Ghisalberti et al., Nucl. Phys. A583 401c (1995).
- [31] S. Pratt, Phys. Rev. **D33**, 72 (1986).
- [32] W. A. Zajc et al., Phys. Rev. C29, 2173 (1984).
- [33] H. Appelshauser et al. Phys. Lett. B 467, 21 (1999).
- [34] H. Sorge, Phys Rev Cbf 52 3291 (1995).
- [35] J. P. Sullivan et al., Nucl. Phys. A 566, 531C (1994).
- [36] D. E. Fields *et al.*, Phys. Rev. C **52**, 986 (1995).
- [37] H.J.Drescher, M.Hladik, S.Ostapchenko, T.Pierog and K.Werner, Phys.Rev. C 65 054902 (2002). Phys. Lett. B 97, 363 (1980).

- [38] M. Murray and B. Holzer, Phys. Rev. C63 054901 (2001).
  [39] R. Scheibl and U. Heinz, Phys. Rev. C 59, 1585 (1999).