



LUND UNIVERSITY

Immune Tolerance Mechanisms in Human Breast Cancer

Mehmeti, Meliha

2019

Document Version:

Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (APA):

Mehmeti, M. (2019). *Immune Tolerance Mechanisms in Human Breast Cancer*. [Doctoral Thesis (compilation), Department of Translational Medicine]. Lund University: Faculty of Medicine.

Total number of authors:

1

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00



Immune Tolerance Mechanisms in Human Breast Cancer

MELIHA MEHMETI

TRANSLATIONAL MEDICINE, MALMÖ | FACULTY OF MEDICINE | LUND UNIVERSITY





**FACULTY OF
MEDICINE**

Department of
Translational Medicine, Malmö

Lund University, Faculty of Medicine
Doctoral Dissertation Series 2019:45
ISBN 978-91-7619-774-5
ISSN 1652-8220



Immune Tolerance Mechanisms in Human Breast Cancer

Immune Tolerance Mechanisms in Human Breast Cancer

Meliha Mehmeti



LUND
UNIVERSITY

DOCTORAL DISSERTATION

by due permission of the Faculty of Medicine, Department of Translational
Medicine Malmö, Lund University, Sweden.

To be defended at the main lecture hall, Pathology building, Skåne University
Hospital, Malmö on Friday 10th of May at 9.15 a.m.


Faculty opponent

Professor Susanna Cardell

Department of Microbiology and Immunology,
Sahlgrenska Academy, Gothenburg University, Sweden

| | | |
|---|---------------------------|-------------------------|
| Organization LUND UNIVERSITY Faculty of Medicine Department of Translational Medicine, Malmö | Document name | |
| | DOCTORAL DISSERTATION | |
| | Date of issue: 2019-05-10 | |
| Author: Meliha Mehmeti | Sponsoring organization | |
| Title and subtitle: Immune Tolerance Mechanisms in Human Breast Cancer | | |
| Abstract One of the most critical events in cancer progression involves the acquired ability of the tumor to avoid the immune system. One way to achieve a successful evasion is establishment of an immunosuppressive and tumor-promoting microenvironment. In this thesis we explored various immunosuppressive tolerance mechanisms in the context of breast cancer. We report that the immune cell receptor TLR4 is expressed in human triple-negative breast cancer cells. TLR4 is there able to respond to both PAMPs and DAMPs, inducing inflammatory, tumor-promoting cytokines. Furthermore, TLR4 expression is correlated with a shorter recurrence-free survival in ER+PgR- breast cancer patients. We further report that S100A9, a TLR4 DAMP, is expressed both by malignant cells and anti-inflammatory CD163+ immune cells in breast cancer. S100A9 binding to TLR4 elicits secretion of inflammatory, tumor-promoting cytokines and S100A9 expression is correlated with worse survival in ER+PgR- breast cancer patients. Further on in this thesis we show that Wnt5a, a factor implicated both in inflammation and cancer, is a novel endogenous TLR2/4 ligand. Through its binding to TLR2/4, Wnt5a is able to act as a DAMP but provokes an anti-inflammatory and a pro-inflammatory cytokine profile in human and mouse immune cells respectively. Moreover, Wnt5a promotes the differentiation of monocytes into an anti-inflammatory Mo-MDSCs cell population. In the fourth project of this thesis, we studied the presence, origin and tumor-biological functions in breast cancer patients of another MDSC population, G-MDSCs. We report an enrichment of immunosuppressive G-MDSCs in metastatic breast cancer patients, while also showing that G-MDSCs comprise a morphologically heterogeneous population of alternatively-activated neutrophils promoting cell proliferation, angiogenesis as well as inhibiting myeloid immune cell infiltration. Finally, we report a potential novel immune tolerance mechanism in breast cancer involving the expression of the transcriptional regulator AIRE, a factor with highly important role in central tolerance in the thymus. We show that is it expressed in malignant cells, CAFs and TAMs in breast cancer and that expression of AIRE in infiltrating TAMs is correlated with worse outcome in breast cancer patients. Collectively, this thesis brings clarity on and introduces novel immune tolerance mechanisms exploited by breast cancer for the purpose of its progression | | |
| Key words: Breast cancer, immune tolerance, toll-like receptors, DAMP, Wnt5a, TAMP, myeloid-derived suppressor cells, autoimmune regulator | | |
| Classification system and/or index terms (if any) | | |
| Supplementary bibliographical information | | Language: English |
| ISSN and key title: 1652-8220 | | ISBN: 978-91-7619-774-5 |
| Recipient's notes | Number of pages: 87 | Price |
| | Security classification | |

I, the undersigned, being the copyright owner of the abstract of the above-mentioned dissertation, hereby grant to all reference sources permission to publish and disseminate the abstract of the above-mentioned dissertation.

Signature 

Date 2019-04-02

Immune Tolerance Mechanisms in Human Breast Cancer

Meliha Mehmeti



LUND
UNIVERSITY

© Meliha Mehmeti, 2019

Cover: “Akaleyi I” by Emma Lindström. The artwork is also available without text on page 87.

All illustrations (Fig. 1-8) are created with BioRender.

Faculty of Medicine
Department of Translational Medicine

ISBN 978-91-7619-774-5
ISSN 1652-8220

Printed in Sweden by Media-Tryck, Lund University
Lund 2019



MADE IN SWEDEN 

Media-Tryck is an environmentally certified and ISO 14001 certified provider of printed material. Read more about our environmental work at www.mediatryck.lu.se

Till mamma

“Bez muke nema nauke”

Table of Contents

| | |
|---|----|
| List of Papers | 10 |
| Paper not included in this thesis | 11 |
| List of Abbreviations | 12 |
| Abstract | 15 |
| Populärvetenskaplig sammanfattning | 17 |
| Introduction | 21 |
| Cancer | 21 |
| Breast Cancer | 23 |
| Epidemiology and Etiology | 23 |
| Breast Cancer Progression | 23 |
| Breast Cancer Classification | 25 |
| Histological classification | 25 |
| The Nottingham histological grade and TNM staging | 25 |
| Immunohistochemical markers | 26 |
| Molecular classification | 26 |
| Breast Cancer Treatment | 27 |
| The Tumor Microenvironment | 29 |
| The Innate Immune System | 33 |
| Monocytes | 33 |
| Macrophages | 34 |
| Neutrophils | 35 |
| Toll-Like Receptors, PAMPs and DAMPs | 36 |
| The Adaptive Immune System | 39 |

| | |
|--|----|
| Immune Tolerance | 41 |
| Autoimmune Regulator | 42 |
| Endotoxin Tolerance | 43 |
| Myeloid-Derived Suppressor Cells | 44 |
| The Immune System and Cancer | 47 |
| Cancer Immunotherapy | 49 |
| Wnt..... | 51 |
| Canonical Wnt signaling | 51 |
| Non-canonical Wnt signaling | 52 |
| Wnt5a | 52 |
| The Present Investigation..... | 55 |
| Aims | 55 |
| Paper I..... | 56 |
| Paper II | 58 |
| Paper III..... | 60 |
| Paper IV..... | 61 |
| Paper V | 64 |
| Conclusions..... | 67 |
| Acknowledgements..... | 69 |
| References..... | 73 |
| The Cover..... | 87 |

List of Papers

- I. Expression of functional toll like receptor 4 in estrogen receptor/progesterone receptor-negative breast cancer.
Meliha Mehmeti, Roni Allaoui, Caroline Bergenfelz, Lao H. Saal, Stephen P. Ethier, Martin E. Johansson, Karin Jirström and Karin Leandersson.
Breast Cancer Research. 17, 1, 130
- II. S100A9 expressed in ER(-)PgR(-) breast cancers induces inflammatory cytokines and is associated with an impaired overall survival.
Caroline Bergenfelz, Alexander Gaber, Roni Allaoui[#], **Meliha Mehmeti**[#], Karin Jirström, Tomas Leanderson and Karin Leandersson.
Br J Cancer. 2015 Oct 20;113(8):1234-43. [#]Equal contribution.
- III. Wnt5a is a TLR2/4-ligand that induces tolerance in human myeloid cells.
Meliha Mehmeti, Caroline Bergenfelz, Eva Källberg, Camilla Rydberg Millrud, Per Björk, Fredrik Ivars, Bengt Johansson-Lindbom, Sven Kjellström, Ingemar André and Karin Leandersson.
Accepted.
- IV. Human granulocytic myeloid-derived suppressor cells (G-MDSCs) in metastatic breast cancer patients is a heterogeneous population with tumor promoting capacity *in vivo*
Meliha Mehmeti, Caroline Bergenfelz, Anna-Maria Larsson Daniel Bexell, Robert Carlsson, Rebecka Hellsten, Niklas Loman, Kristian Riesbeck, Jonas Ahl, Camilla Rydberg Millrud, Gesine Paul-Visse, Lisa Rydén, Fredrika Killander and Karin Leandersson
Manuscript
- V. Autoimmune regulator (AIRE) expressed in tumor associated macrophages is associated with worse prognosis in breast cancer patients.
Meliha Mehmeti, Frida Björk Gunnarsdottir, Caroline Bergenfelz, Catharina Hagerling, Roni Allaoui, Martin Johansson, Karin Jirström and Karin Leandersson.
Manuscript

Paper not included in this thesis

- I. Docetaxel promotes the generation of anti-tumorigenic human macrophages.

Camilla Rydberg Millrud, **Meliha Mehmeti** and Karin Leandersson.

Experimental Cell Research. 2018

List of Abbreviations

| | | | |
|---------------|--|----------------|---|
| <i>AIRE</i> | autoimmune regulator | <i>JNK</i> | c-Jun N-terminal kinase |
| <i>AP-1</i> | activator protein 1 | <i>LPS</i> | lipopolysaccharide |
| <i>APC</i> | antigen presenting cell | <i>LRP</i> | lipoprotein receptor-related protein |
| <i>Arg1</i> | arginase-1 | <i>M1</i> | pro-inflammatory macrophage |
| <i>BMM</i> | bone marrow macrophage | <i>M2</i> | anti-inflammatory macrophage |
| <i>BRCA</i> | breast cancer gene | <i>MAPK</i> | mitogen activated protein kinase |
| <i>CAF</i> | cancer associated fibroblast | <i>MBC</i> | metastatic breast cancer |
| <i>CaMKII</i> | calmodulin-dependent kinase II | <i>MCP-1</i> | monocyte chemotactic protein-1 |
| <i>CAR</i> | chimeric antigen receptor | <i>MDSC</i> | myeloid-derived suppressor cell |
| <i>CARS</i> | compensatory anti-inflammatory response syndrome | <i>MHC</i> | major histocompatibility complex |
| <i>CD</i> | cluster of differentiation | <i>MMP</i> | matrix metalloproteinase |
| <i>CIS</i> | carcinoma <i>in situ</i> | <i>Mo-MDSC</i> | monocytic MDSC |
| <i>CTL</i> | cytotoxic T-lymphocyte | <i>mTEC</i> | medullary thymic epithelial cell |
| <i>CTLA-4</i> | cytotoxic T-lymphocyte-associated protein 4 | <i>MyD88</i> | myeloid differentiation primary response 88 |
| <i>DAMP</i> | damage-associated molecular pattern | <i>NET</i> | neutrophil extracellular trap |
| <i>DC</i> | dendritic cell | <i>NFκB</i> | nuclear factor kappa B |
| <i>DVL</i> | dishevelled | <i>NHG</i> | nottingham histological grade |
| <i>ECM</i> | extracellular matrix | <i>NK</i> | natural killer cell |
| <i>ER</i> | estrogen receptor | <i>NKT</i> | natural killer T cell |
| <i>FZD</i> | frizzled | <i>NO</i> | nitric oxide |
| <i>G-MDSC</i> | granulocytic-MDSCs | <i>PAMP</i> | pathogen-associated molecular pattern |
| <i>GSK3β</i> | glycogen synthase kinase-3 beta | <i>PBMC</i> | peripheral blood mononuclear cell |
| <i>HER2</i> | human epidermal growth factor receptor 2 | <i>PCP</i> | planar cell polarity |
| <i>HMGB1</i> | high mobility group box 1 | <i>PD-1</i> | programmed death 1 |
| <i>ICI</i> | immune checkpoint inhibitor | <i>PD-L1</i> | programmed death-ligand 1 |
| <i>IDO</i> | indoleamine 2,3-dioxygenase | <i>PgR</i> | progesterone receptor |
| <i>IFN</i> | interferon | <i>PMN</i> | polymorphonuclear |
| <i>IL</i> | interleukin | <i>PRR</i> | pattern-recognition receptor |
| <i>iNOS</i> | inducible nitric oxide synthase | <i>ROS</i> | reactive oxygen species |
| <i>IRF</i> | interferon regulatory factor | <i>SERM</i> | selective estrogen receptor |
| | | <i>SIRS</i> | systemic inflammatory response syndrome |

| | | | |
|-------------|--|-------------|--|
| <i>TAA</i> | tumor associated antigen | <i>TLR</i> | toll-like receptor |
| <i>TAMP</i> | tolerance-associated molecular pattern | <i>TNF</i> | tumor necrosis factor |
| <i>TAM</i> | tumor associated macrophage | <i>Treg</i> | regulatory T cell |
| <i>TCR</i> | T cell receptor | <i>TSA</i> | tissue specific antigen |
| <i>TDLU</i> | terminal duct lobular unit | <i>VEGF</i> | vascular endothelial growth factor |
| <i>TGF</i> | transforming growth factor | <i>Wnt</i> | wingless-related MMTV integration site |
| <i>Th</i> | T helper cell | | |
| <i>TNBC</i> | triple negative breast cancer | | |

Abstract

One of the most critical events in cancer progression involves the acquired ability of the tumor to avoid the immune system. One way to achieve a successful evasion is establishment of an immunosuppressive and tumor-promoting microenvironment.

In this thesis we explored various immunosuppressive tolerance mechanisms in the context of breast cancer. We report that the immune cell receptor TLR4 is expressed in human triple-negative breast cancer cells. TLR4 is there able to respond to both PAMPs and DAMPs, inducing inflammatory, tumor-promoting cytokines. Furthermore, TLR4 expression is correlated with a shorter recurrence-free survival in ER⁻PgR⁻ breast cancer patients. We further report that S100A9, a TLR4 DAMP, is expressed both by malignant cells and anti-inflammatory CD163⁺ immune cells in breast cancer. S100A9 binding to TLR4 elicits secretion of inflammatory, tumor-promoting cytokines and S100A9 expression is correlated with worse survival in ER⁻PgR⁻ breast cancer patients. Further on in this thesis we show that Wnt5a, a factor implicated both in inflammation and cancer, is a novel endogenous TLR2/4 ligand. Through its binding to TLR2/4, Wnt5a is able to act as a DAMP but provokes an anti-inflammatory and a pro-inflammatory cytokine profile in human and mouse immune cells respectively. Moreover, Wnt5a promotes the differentiation of monocytes into an anti-inflammatory Mo-MDSCs cell population. In the fourth project of this thesis, we studied the presence, origin and tumor-biological functions in breast cancer patients of another MDSC population, G-MDSCs. We report an enrichment of immunosuppressive G-MDSCs in metastatic breast cancer patients, while also showing that G-MDSCs comprise a morphologically heterogeneous population of alternatively-activated neutrophils promoting cell proliferation, angiogenesis as well as inhibiting myeloid immune cell infiltration. Finally, we report a potential novel immune tolerance mechanism in breast cancer involving the expression of the transcriptional regulator AIRE, a factor with highly important role in central tolerance in the thymus. We show that it is expressed in malignant cells, CAFs and TAMs in breast cancer and that expression of AIRE in infiltrating TAMs is correlated with worse outcome in breast cancer patients.

Collectively, this thesis brings clarity on and introduces novel immune tolerance mechanisms exploited by breast cancer for the purpose of its progression.

Populärvetenskaplig sammanfattning

Det är av största vikt att vårt immunförsvar kan skydda vår kropp från fara. För att kunna göra det använder immunsystemet olika receptorer som kan känna igen farosignaler från smittämnen som bakterier och virus (pathogen-associated molecular patterns, PAMPs). Dessa receptorer benämns som Toll-likareceptorer (TLRs) och fungerar som första försvarslinjen genom att känna igen olika faromolekyler. Vid bindning till dessa molekyler framkallar de ett inflammatoriskt svar, vilket i bästa fall eliminerar faran. Inflammatoriska svaret involverar bl.a. produktion av olika pro-inflammatoriska signalsubstanser. Det är samtidigt lika viktigt att immunsvaret är strikt kontrollerat eftersom en för stark inflammation kan leda till förödande skador på kroppen. För att dämpa inflammationen, aktiverar immunsystemet olika tolerans-mekanismer såsom anti-inflammatoriska celler (t.ex. myeloid-derived suppressor cells; MDSCs) som motvikt för att skapa balans.

Cancer är ett samlingsnamn för en grupp komplexa sjukdomar som uppstår som följd av skador i vårt DNA. Under normala förhållanden är celldelning en kontrollerad process som styrs av signaler från omgivningen, men på grund genetiska förändringar (mutationer) kan en cell emellertid uppnå förmågan av okontrollerad och oändlig celldelning och så småningom därmed utvecklas till en tumör. En cancercell skiljer sig från en normal cell; den beter sig annorlunda och ser annorlunda ut, något som immunsystemet reagerar mot. Efter upptäckandet av cancerceller initierar immunsystemet processen för avläsning av tumören där flera olika immunceller rekryteras till platsen för att delta i kampen. I de flesta fall avlägsnas tumören, men i vissa fall kommer tumören undan och med tiden lyckas den manipulera immunsystemet till sin fördel. En viktig del i cancerens framgång hänger på att undkomma immunsystemet, och det gör den bland annat genom att skapa en anti-inflammatorisk mikromiljö som främjar dess tillväxt.

Den här avhandlingen fokuserar på olika mekanismer i förvärvandet av immuntolerans i bröstcancer.

I avhandlingens första del var målet att studera uttrycket och funktionen av olika TLRs i bröstcancer. TLRs utgör en familj av tio olika mönster-igenkännande receptorer (PRRs) som känner igen olika PAMPs och som i vanliga fall uttrycks av olika immunceller. På senare år har dock flera studier rapporterat att TLRs även uttrycks av olika cancerformer där de bidrar till cancerprogression. I *Artikel 1* visar vi att flera TLRs (TLR2, 3, 4, 9) uttrycks av bröstcancerceller. Uttrycket var mest

uppenbart i cancerceller som tillhör den trippel-negativa bröstcancer-undergruppen, en mycket aggressiv typ av bröstcancer som är förknippad med dålig prognos. Vidare i arbetet upptäckte vi att TLR4 var fullt funktionell med förmågan att framkalla utsöndring av olika cancer-främjande signalsubstanser. Vi fann även att förekomsten av TLR4 är korrelerad med snabbare återfall hos ER⁺PgR⁻ bröstcancerpatienter.

Fara uppenbaras inte enbart i form av smittämnen. Förutom att aktiveras av PAMPs, kan TLRs även känna igen och aktiveras av kropps-egna molekyler (danger-associated molecular patterns; DAMPs). Vid trauma, stress eller radio/kemoterapi kan vävnader skadas och detta kan i sin tur leda till att celler börjar läcka ut sitt innehåll, bland dem DAMPs. Å andra sidan kan DAMPs aktivt utsöndras av cancerceller och därmed framkalla en steril inflammation. I avhandlingens andra delarbete fokuserade vi på S100A9, en DAMP som binder till TLR4. Tidigare studier har rapporterat att S100A9 är involverad i olika inflammatoriska processer och att den har förmågan att locka till sig anti-inflammatoriska celler (MDSCs). I *Artikel II* fann vi att S100A9 uttrycks av bröstcancer celler och genom att binda till TLR4 inducerar S100A9 utsöndring av olika cancer-främjande signalsubstanser. Dessutom kunde vi också visa att förekomsten av S100A9 är förknippad med sämre överlevnad hos ER⁺PgR⁻ bröstcancerpatienter. Sammanfattningsvis visar vi att bröstcancerceller har förvärvat en komplett uppsättning av verktyg för att kunna inducera inflammation som främjar dess tillväxt.

I *Artikel III* fokuserade vi på Wnt5a, ett ämne som har kopplats både med inflammation och cancer. Vi visar för första gången att Wnt5a binder till TLR2 och TLR4, och att genom denna bindning är den kapabel att framkalla ett pro-inflammatoriskt (DAMP) och ett anti-inflammatoriskt (tolerance-associated molecular pattern; TAMP) svar i mus respektive humana immunceller. Vi visar även att Wnt5a är förmögen att omvandla pro-inflammatoriska monocytter (vit blodkropp) till anti-inflammatoriska monocyt-MDSCs (Mo-MDSCs). Förekomsten av MDSCs kan vara livsviktigt i sjukdomar som kännetecknas av överdriven inflammation, såsom blodförgiftning, genom att dämpa inflammationen. Men i samband med cancer kan MDSCs förlama anti-cancerimmunsvar och därigenom bidra till cancerprogression. MDSCs är en heterogen cell population och förutom Mo-MDSCs innefattar den även granulocyt-MDSCs (G-MDSCs). Båda cellerna är kraftigt immundämpande, men i motsats till Mo-MDSCs är G-MDSCs ursprung samt tumör-främjande funktioner i människa mindre studerade. I *Artikel IV* visar vi att G-MDSCs utgör en heterogen cellpopulation av alternativt-aktiverade neutrofiler (granulär vit blodkropp) som är berikad hos metastaserande bröstcancerpatienter. Genom att skapa en bröstcancer-musmodell upptäckte vi att G-MDSCs bidrar till ökad cancertillväxt och bildandet av nya blodkärl. Dessutom hindrar G-MDSCs andra immunceller att infiltrera tumören. Således bidrar G-MDSCs till cancertillväxt.

I sin sökning av fara är det absolut nödvändigt att immunförsvaret kan diskriminera mellan farosignaler och normala kropps-specifika molekyler (själv-antigen) och därmed visa tolerans mot sig själv. För att undvika förödande autoimmuna sjukdomar, måste T celler (antigen-specifik vit blodkropp) gå igenom en mognadsprocess (central tolerans) i thymus där självtoleranta T celler klarar sig och får lämna thymus och självreaktiva T celler avlägsnas. En viktig komponent i mognadsprocessen är Autoimmune regulator (AIRE), ett ämne som uttrycks i thymus. Mutationer i AIRE är kopplad till olika autoimmuna sjukdomar. Ibland kan det hända att självreaktiva T celler undkommer eliminering och tar sig ut i perifera organ. Som backup, har kroppen installerat en perifer tolerans där självreaktiva T celler upptäcks och oskadliggörs ute i kroppen. Man har bland annat upptäckt uttrycket av AIRE i periferin. I sista delarbetet, *Artikel V*, har vi upptäckt att AIRE uttrycks i cancerceller men även i en anti-inflammatorisk, tumör-främjande immuncellpopulation (tumor-associated macrophage, TAM) hos bröstcancerpatienter där uttrycket av AIRE_{TAM} och bröstcancer är korrelerad med sämre prognos. Vi undersöker nu om AIRE i bröstcancer bidrar till immuntolerans genom att avvärja T celler som i normala fall skulle angripa canceren.

Introduction

Cancer

During thousands of years, cancer has proven enigmatic, challenging both physicians and patients. Already in 3000 BC, cancer was documented by Egyptians as an incurable illness. Considered as the second leading cause of death, cancer is estimated to deprive over 9 million lives worldwide ¹.

Cancer is a collective term that includes a group of diseases arising as a result of DNA mutations which subsequently leads to an uncontrolled division of abnormal cells. During the course of cell division, errors can occur in the replicating DNA. Additionally, various environmental factors, to which our body is exposed to including smoking and UV-light, can damage the DNA. Under normal circumstances, the multiple DNA repairing mechanisms employed by the cell are perfectly able to repair the damage, however in some cases, the cell fails to repair the damage and normal cells acquire mutations and eventually become cancerous. Cell division without restraint is generated in various ways, including mutations in protooncogenes involved in cell division and mutations in tumor suppressor genes involved in e.g. apoptosis and DNA repair. Loss of function in the latter further potentiates accumulation of a plethora of mutations aiding tumor progression ². Additionally, mutations can also be inherited e.g. BRCA1 and BRCA2 in breast cancer.

Besides an aberrant cell division, cancer development and progression involve a multiple-step process. In 2000, Weinberg and Hanahan summarized the main processes in cancer progression into six different events referred to as hallmarks of cancer ³. Since then, the list has expanded and four additional hallmarks have emerged including the tumor-promoting inflammation and the immune destruction avoidance (**Fig. 1**) ⁴. The hallmarks illustrate the complexity and heterogeneity of cancer. For a long time, cancer was believed to solely consist of tumor cells, however, today we know that this is not the case. In fact, cancer constitutes a microenvironment consisting of multiple cell-types, both cancerous and non-malignant, all believed to be involved in a reciprocal communication aiding tumor progression ⁵. As part of the tumor microenvironment, immune cells have been shown to play a crucial role in cancer progression and patient survival. In the last two decades, a major effort has been invested in studying the role the immune

system plays in cancer, and this thesis has focused on how the immunosuppressive cells are generated.




Figure 1. The hallmarks of cancer. Adapted from Hannahan and Weinberg, 2011 ⁴.

Breast Cancer

Epidemiology and Etiology

Breast cancer is the most prevalent malignant disease in women worldwide. In Sweden alone, over 8900 breast cancer incidences are reported annually, accounting for approximately 30 % of all cancer cases in women. Today, one of ten women are diagnosed with breast cancer and the incidences have constantly been increasing in the last 20 years, most probably due to an increasing aging population and the introduction of screening ^{6,7}. Nevertheless, the prognosis is generally good with a relative 5- and 10- year survival of approximately 90 % and 80 % respectively ⁸.

Risk factors associated with breast cancer development are plentiful. As with most cancers, breast cancer is a disease of aging, where around 70 % of all breast cancer incidences are reported in women above 55 years of age. The length of exposure to estrogen hormones has strongly been attributed to breast cancer development, comprising an early menarche, late menopause, late first birth as well as exposure to hormone replacement therapy. Other risk factors include diet, smoking, and mammographic density ⁹.

Among all risk factors associated with breast cancer development, after age, the genetic predisposition is the most important. It is estimated that around 5 %- 10 % of all breast cancer incidences arise as a result of somatic genetic mutations ¹⁰. *BRCA1* and *BRCA2* are tumor suppressors which play important roles in the maintenance and repair of DNA double strand breaks and somatic mutations in these genes account for around 40 % of all familial breast cancers ¹¹. This implies that other genes ought to be responsible for the remaining hereditary breast cancers, and although rare, mutations in *PTEN* and *TP53* have also associated with hereditary breast cancer ¹².

Breast Cancer Progression

A normal breast comprises two different compartments, a glandular tissue which represent the functional part of the breast and a surrounding stromal tissue. A mature glandular tissue has a tree-like appearance and consists of a terminal duct

lobular unit (TDLU), the functional unit of the breast, responsible for milk production and transportation. Both the lobules (alveoli) and the ducts are constructed of a lumen lined by an inner luminal epithelial and an outer myoepithelial cell layer and a basement membrane. The surrounding stroma is fat-rich, where fibroblasts, immune cells and blood vessels are also present (**Fig. 2**)^{13,14}. Majority of breast cancers originate from the TDLU and the development of breast cancer involves a series of pathological events which start with an appearance of lesions or flat epithelial atypia. At this stage, the cells in questions might display an abnormal response to both growth factors and apoptosis signals. This is followed by atypical hyperplasia and subsequently into the development of carcinoma *in situ* (CIS). The transition between atypical aplasia and CIS is characterised by increased genetic changes and genetic instability. Approximately one fifth of all detected breast cancers upon a mammography screening are accounted for by CIS cases^{15,16}. The shift from a carcinoma *in situ* to an invasive cancer involves further recruitment of stromal cells and extracellular matrix (ECM) proteins which aid the invasive process by producing factors that breaks down the basement membrane and thus allowing the spread of malignant cells into the surrounding stroma (**Fig. 3**)¹⁷.




Figure 2. Schematic illustration of the normal breast.

The functional unit of the normal breast comprise the terminal duct lobular unit (TDLU), a branch-like complex extending from the nipple. The TDLU is composed by an inner lumen lined by lumen epithelial cells, myoepithelial cells and basement membrane (BM).




Figure 3. A simplified overview of the breast cancer progression.

The breast cancer progression summarized in four main steps represented by the normal intact duct, the appearance of lesions leading to atypical hyperplasia, followed by carcinoma *in situ* characterized by the gain of genetic instability and recruitment of various immune and stromal cells resulting in invasive cancer featured by the disruption of basement membrane.

Breast Cancer Classification

Breast cancer is a heterogeneous disease. The morphological and immunohistopathological characteristics divides the tumor into several subgroups, providing a useful tool when it comes to the choice of treatment and prognosis.

Histological classification

Based on the growth pattern, tumors can be classified into various histological subgroups, where the two major groups constitute the invasive breast carcinoma and non-invasive carcinoma *in situ*. Around 50-80 % of breast cancers reported are invasive breast carcinomas¹⁸, representing the most common histological type. The invasive breast carcinomas are further sub-classified in smaller groups as reviewed in¹⁹. In addition, both the invasive and non-invasive carcinomas can origin from the lobular and the ductal unit of the breast¹⁹.

The Nottingham histological grade and TNM staging

The Nottingham histological grading (NHG) is recognized as a valuable prognostic tool when assessing the aggressiveness of breast tumors. The degree of

differentiation of a tumor is evaluated based on morphological characteristics including the mitotic count, tubule formation and nuclear polymorphism upon which the tumor is assigned a grade. The classification system includes grades I-III, where grade I tumors are well differentiated, grade II tumors are moderately differentiated, and grade III tumors are poorly differentiated. The NHG provides predictive information regarding the overall survival and recurrence free survival ²⁰.

Breast cancer progression can further be assessed by the use of TNM staging, which takes tumor size (T), lymph node involvement (N) and distant metastases (M) into account. By combining the values obtained from the three parameters, the clinical stage (I-IV) of the tumor is assessed ²¹.

Immunohistochemical markers

One of the most important steps in the process of determining the prognosis and the treatment of breast cancer is the assessment of estrogen receptor (ER) and progesterone receptor (PgR) expression as well as the HER2 amplification. ER and PgR expression, which is present among 80% of all breast cancer patients, is evaluated with immunohistochemistry. Patients exhibiting ER/PgR-positive tumors are candidates for endocrine therapy. Furthermore, the amplification status of HER2 gene (*ERBB2*) is assessed, and this helps to determine whether the patient is a candidate for HER2-targeted therapy. Approximately 15 % of all breast cancer patients display HER2 overamplification. Additionally, the expression of the proliferation marker Ki67 is also analyzed ²¹. Tumors that lack all three receptors (ER⁻PgR⁻HER2⁻) are referred to as triple negative breast cancer (TNBC).

Molecular classification

Gene expression profiling has proven groundbreaking for defining molecular subtypes of breast cancer. As the molecular background of breast cancer provides useful information when predicting clinical features such as the overall survival and relapse-free survival, assessing the differences in molecular characteristics of breast cancer is of high clinical relevance ²². Today, six different breast cancer subtypes are identified including; Luminal A, Luminal B, HER2 enriched, Basal-like, Normal-like and Claudin-low (summarized in **Table 1**) ^{19,22,23}.

Majority of Luminal A and Luminal B tumors are characterized by positive expression of ER and PgR and the absence of HER2 amplification ²¹. While most Luminal A tumors exhibit a low histological grade, a low proliferation index and good outcome, Luminal B tumors on the other hand display a higher histological grade, a high proliferation index and intermediate/poor prognosis ¹⁸. HER2-enriched tumors exhibit a varied ER and PgR expression and, as the name implies, they

display HER2 amplification. Additionally, HER2-enriched tumors display high histological grade and proliferation index and although HER2-enriched breast cancer patients benefit from targeted therapy, the outcome still remains poor ²². Most Basal-like tumors and Claudin-low tumors lack the expression of ER and PgR and lack over-amplification of HER2 and are commonly referred to as TNBC. Moreover, they are also characterized by a low expression of cell adhesion molecules and elevated mesenchymal features ²³. The scarcity of treatment options due to the lack of drug targetable receptors as well as the tumors display of high histological grade and proliferation, results in Basal-like breast cancer patients having a very poor outcome ^{18,22,24}.

Table 1. Molecular subgroups of breast cancer and their characteristics ^{18-20,22-24}.

| Subtype | Molecular profile | Frequency | Histological grade | Proliferation, Ki67 | Outcome |
|-------------|--|-----------|----------------------|---------------------|----------------------|
| Luminal A | ER ⁺ PgR ⁺ HER2 ⁻ | 50-60% | Low | Low | Good |
| Luminal B | ER ^{+/+} PgR ^{+/+} HER2 ^{+/+} | 10-20% | Intermediate or high | High | Intermediate or poor |
| HER2 | ER ⁻ PgR ⁻ HER2 ⁺ | 10-15% | High | High | Poor |
| Basal-like | ER ⁻ PgR ⁻ HER2 ⁻ | 10-20% | High | High | Poor |
| Claudin-low | ER ⁻ PgR ⁻ HER2 ⁻ | 12-14% | High | High | Poor |
| Normal-like | ER ^{+/+} HER2 ⁻ | 5-10% | Low | Low or intermediate | Intermediate |

Breast Cancer Treatment

Upon the discovery of breast cancer, the most prominent and common treatment is surgery. Depending on the stage of the tumor, its spread to adjacent lymph nodes and existence of metastases, adjuvant radiation- and/or chemotherapy is introduced to the treatment ²⁵. Further treatment options include endocrine therapy such as selective estrogen receptor modulators (SERMs) with Tamoxifen and aromatase inhibitors as the most usually employed for ER⁺ tumors, and targeted therapy as trastuzumab or lapatinib for HER2-amplified tumors ^{26,27}. As patients with TNBC do not benefit from the targeted or endocrine therapy mentioned above, the current treatment strategy remains surgery and/or chemotherapy ²⁸. Nevertheless, Lynparza, a PAPR (poly-ADP ribose polymerase) inhibitor, has been approved earlier this year by the FDA for treatment of breast cancers displaying BRCA1 and/or BRCA2 germline mutations ²⁹. As around 35 % of TNBCs display mutations in BRCA1 and around 8 % are positive for BRCA2 mutations ³⁰, it is comforting to state that a good portion of TNBC patients might benefit from this treatment in the future. In a phase III clinical trial, metastatic TNBC patients assigned to a combination therapy of atezolizumab (programmed death ligand 1 (PD-L1) inhibitor) and nab-paclitaxel

showed encouraging results of significantly prolonged progression-free survival (PFS) and overall survival (OS) in comparison to patients receiving nab-paclitaxel only. Moreover, the PFS and OS was longer in patients with positive PD-L1 tumors in comparison to patients lacking PD-L1 expression ³¹.

The Tumor Microenvironment

A tumor aspires to satisfy the requirements needed to tick all the boxes on the list of hallmarks of cancer. However, a tumor cannot sustain a successful growth by itself and therefore relies on help from non-transformed cells; either already present in the tumor microenvironment or acquired to the tumor site. The tumor microenvironment is a dynamic organ comprised of a mixture of malignant and non-malignant cells including cancer-associated fibroblasts (CAFs), various immune cells such as tumor-associated macrophages (TAMs), and cells composing the vasculature, including endothelial cells, pericytes and the ECM (**Fig. 4**)^{4,5,32}.

Fibroblasts are spindle-shaped cells of the connective tissue with important functions in the ECM homeostasis by producing ECM-regulating components such as collagen, fibronectin and matrix metalloproteinases (MMPs). Furthermore, they play an important role in wound repair. During a non-cancerous wound healing process, fibroblasts are transiently activated and are then referred to as myofibroblasts. However, in the context of cancer, the transient activation remains persistent. Consequently, CAFs encompass one of the most prominent non-malignant cells in the tumor microenvironment^{32,33}. They are involved in several tumor-promoting processes including chemoattraction, proliferation and invasion by secreting various factors such as CXCL16, TGF- β and MMPs^{5,33-35}.

A vascular network within the tumor microenvironment is crucial for the prosperity of a tumor as it is a vital source of oxygen and nutrients³⁶. This is highlighted by the fact that tumor cells residing too far away from a vessel, or a tumor growing at a faster speed than the vessel formation, often suffers from necrosis³⁷. Endothelial cells comprise a fundamental component of the vascular architecture playing an important role in tumor angiogenesis. Tumors secrete various angiogenesis-promoting factors, including the prototypic vascular endothelial growth factor (VEGF). Upon release, VEGF activates endothelial cells to produce MMPs which subsequently breaks down the ECM, facilitating the endothelial cell migration and *de novo* vessel-formation³⁶.

Another cell type present in the tumor microenvironment, aiding tumor progression are the so called pericytes. In healthy tissues, pericytes can be found lining around the basement membrane of blood vessels, and foremost, exert important functions in vessel development and maintenance³⁸. In tumors, the functional role of pericytes is rather unknown, however, they seem to exert a less firm grip and display an

abnormal coverage around blood vessels ³⁹. For instance, an increased pericyte coverage stabilizes the vessels, thus supporting tumor growth ⁴⁰. On the other hand, a low pericytes coverage results in increased vessel permeability and leakage favoring tumor invasion and metastasis. Additionally, a low pericyte coverage is correlated with worse outcome in breast cancer patients ⁴¹.




Figure 4. The tumor microenvironment.

The tumor microenvironment constitutes multiple cell-types, all involved in a reciprocal communication aiding the tumor progression.

The ECM represents another important factor of the tumor microenvironment driving tumor progression. It comprises numerous non-cellular components including collagen, fibronectin, proteoglycans and laminin, which in healthy tissue does not only provide structural support but also plays an important role in cell adhesion, cell proliferation and in the communication between cells ⁴². A tumor ECM differs largely from normal ECM, displaying remodeling and a stiffer structure. It has been demonstrated that tumor stiffness characterized by increased collagen crosslinking can stimulate integrin signaling, which furthermore, promoted tumor invasion in murine mouse model ⁴³. This is further supported by the notion that women with extensive mammographic breast density have an increased risk of developing breast cancer ⁴⁴. CAFs represent a major source of ECM components which also are subjected to degradation by different MMPs, an event which

subsequently triggers cell motility and invasion^{33,45}. MMPs on the other hand are involved in several key regulatory processes important in tumor progression including angiogenesis, invasion and metastasis, and have been associated with worse prognosis in various cancer types⁴⁵⁻⁴⁷.

The tumor microenvironment also consists of infiltrating immune cells. It has lately become apparent that the immune response acts in favor of the tumor and therefore is part of the tumors' natural microenvironment. How this is all orchestrated, and which immune cells are responsible, will be described in detail below.

The Innate Immune System

The human body is under constant threat from pathogens such as bacteria, viruses and fungi that when infecting the host, interfere with its normal functions. Over hundreds of millions of years, the immune system has evolved to act as a shield against exogenic intruders while at the same time maintaining tolerance to self. The immune system consists of two branches, the innate and the adaptive immune system, whose complementary functions provide the host with full protection.

The innate immune system comprises a general first-line defense that involves recognition of foreign patterns as being non-self, thus enabling an immune response. To fulfil its tasks, the innate immune system employs several defense mechanisms including physical barriers such as the skin and the secreted mucosal layer lining the respiratory- and gastrointestinal tract⁴⁸. In case the pathogen manages to breach the physical barriers, the innate immune system engages a variety of pattern recognition receptors (PRRs), including the most studied Toll-like receptors (TLRs). These sense the intruder and subsequently initiate a proper immune response by e.g. the upregulation of MHC and co-receptor molecules responsible for antigen presentation and the induction of different cytokines and chemokines which act as messengers playing important roles in the further activation and recruitment of different immune cells^{49,50,51}. The cellular defense of the innate immune system includes natural killer (NK) cells, monocytes, neutrophils, macrophages, dendritic cells (DCs), eosinophils and basophils, which exert various functions such as direct killing of the pathogen through the release of antimicrobial compound, wound healing, pathogen phagocytosis and antigen presentation to lymphocytes^{48,52}. Below, some important mediators of the innate immune response related to this thesis will be described.

Monocytes

Monocytes are bone-marrow derived blood cells from the myeloid lineage that comprise around 10 % of leukocytes in human blood. They are circulating cells playing important roles in the early inflammatory responses, host defense during infections, homeostasis and tissue repair as well as in the replenishment of both recruited and tissue resident macrophages^{53,54}. Morphologically, monocytes are

characterized by their diverse shape and size, kidney-shaped nucleus and a high cytoplasm-to nucleus ratio. The challenging task in identifying monocytes based on their morphology has been facilitated by the use of flow cytometry where monocytes are identified by the expression of the lipopolysaccharide (LPS) receptor CD14 and their light-scatter characteristics⁵⁴. Nonetheless, monocytes comprise a heterogenous cell population which is further divided into several subtypes based on the presence and the density of cell-surface markers CD14 and CD16 (Fcγ receptor III). They are thus grouped into the classical (CD14⁺⁺CD16⁻), the intermediate (CD14⁺⁺CD16⁺) and the non-classical (CD14⁺CD16⁺⁺) subtypes^{55,56,57}.

Besides playing a crucial role in host defense, monocytes have also been associated with cancer progression. One example is that monocytes, after being recruited by CCL2, secreted from metastatic tumor cells or the tissue stroma in the target organ, have been shown to generate metastatic associated macrophages which promote breast cancer metastasis⁵⁸. Monocytes isolated from renal cell carcinoma (RCC) patients displayed a protumor gene- and protein profile and promoted an invasive behavior and angiogenic activity of RCC cells when co-cultured *in vitro*. Additionally, the monocytes subsequently infiltrated the tumor giving rise to TAMs⁵⁹. Furthermore, metastatic behavior was also observed *in vitro* in interactions between various monocytic- and breast cancer cell lines, where monocytes promote motility and invasiveness of breast cancer cells^{60,61}. In addition, reprogramming of monocytes towards a myeloid immunosuppressive cell population by the tumor microenvironment, has been shown to be more prevalent in advanced cancer^{35,62}. The heterogeneity of monocytes as well as their ability to differentiate to different types of macrophages including M1, M2, TAMs and MDSCs (which will be reviewed later on in this thesis) demonstrates an incredible versatility and plastic nature of monocytes.

Macrophages

Over a century ago, macrophages were discovered by Ilya Metchnikoff⁶³, and ever since, we have learned about their versatile functions including their central role in the immune responses upon an infection, their tumoricidal ability and contribution to tissue remodeling and cancer progression. During an infection, macrophages can be recruited from circulating monocytes. Additionally, various tissues comprise a tissue-resident macrophage population, including microglia in the central nervous system and Kupffer cells in the liver and similarly to the monocyte-derived macrophages, they are involved in processes such as immune surveillance⁶⁴.

Macrophages comprise a versatile cell population, differing in both phenotype and function. Depending on the signals received from the microenvironment,

macrophages can either undergo a classical or alternative activation and thus differentiate towards the M1 or the M2 macrophages respectively. The polarization towards classically activated M1 macrophages is induced by TLR-ligands such as LPS and pro-inflammatory cytokines including IFN- γ and TNF- α ^{65,66}. M1 macrophages are characterized by their role in driving T_h1 immune responses, upregulation of pro-inflammatory mediators such as TNF- α and IL-12, elevated antigen presentation as well as antimicrobial and tumoricidal activities ⁶⁷. On the other side, the polarization towards alternatively activated M2 macrophages is induced by IL-4, IL-10 and IL-13. Functions employed by M2 macrophages include tissue-remodeling, wound healing and anti-inflammatory attributes with an IL-10^{high}IL-12^{low} cytokine profile as well as a high expression of scavenger receptors ^{65,66,68}. However, it should be taken into consideration that M1 and M2 polarization merely represents a simplification and that in-between these extremes, a wide spectrum of different polarization states exists.

Tumor associated macrophages (TAMs) have been described as one of the most prominent players in tumor progression. They represent a major leukocyte population within a tumor, sharing common attributes with M2 macrophages including the anti-inflammatory IL-10^{high}IL-12^{low} cytokine profile and tissue remodeling properties ⁶⁸. Additionally, TAMs display impaired phagocytosis ⁶⁹. The presence of TAMs has been described in various cancer types including breast cancer, classic Hodgkin's lymphoma and bladder cancer, where they have been correlated with higher tumor grade and worse overall survival ^{68,70-72}. Moreover, TAMs have also been described to promote proliferation, invasiveness and angiogenesis ⁶⁸.

Neutrophils

Neutrophils represent the predominant circulating human leukocyte population with an essential role in the innate immune responses ⁷³. In the host defense, neutrophils are among the first cells at site of infection and employ a variety of defense mechanisms including phagocytosis of microbes, release of antimicrobial peptides and induction of neutrophil extracellular traps (NETs) ^{74,75}. Neutrophils are short-lived cells with a half-life of approximately 8 h in circulation in humans. Nonetheless, their lifespan can extend during inflammatory conditions enabling them to also participate in inflammation resolution ⁷³. Neutrophils were originally believed to comprise a rather homogenous population, however, in recent years, it has been shown that neutrophils indeed are phenotypically and functionally heterogeneous with several subtypes, including the anti-tumor N1 and the pro-tumor N2 neutrophils, being described ^{76,77}.

In the context of cancer, neutrophils play a paradoxical role. Increased levels of neutrophils have been associated with poor outcome in various cancers⁷⁸⁻⁸⁰, while in others, neutrophils employ tumoricidal effects and are correlated with a better outcome⁸¹⁻⁸². Indeed, controversy seems to surround neutrophils. Sharing similar phenotype and function it has been proposed that neutrophils might give rise to, or even are equal to an immunosuppressive subset of granulocytic-myeloid derived suppressor cells (G-MDSCs).(reviewed in^{83,84}). This hypothesis will be discussed in more detail further below in this thesis.

Toll-Like Receptors, PAMPs and DAMPs

To be able to recognize invading pathogens, TLRs act as first line of defense by recognizing molecular patterns conserved among microbial species, and thus alarm the body of the danger⁸⁵. TLRs are members of the pattern recognition receptor (PRR) family and today there are ten (1-10) TLRs identified in humans. They are located both on the cellular membrane (TLR1, TLR2, TLR4, TLR5, TLR5, TLR6) and on intracellular endosomes (TLR3, TLR4, TLR7, TLR8 and TLR9) of various immune cells including monocytes, DCs, neutrophils, macrophages as well as epithelial cells⁸⁶⁻⁹¹. Besides being expressed on immune cells, TLR expression has also been reported in various types of cancers, including melanoma, neuroblastoma, esophageal cancer, breast cancer and renal cell carcinoma^{92,93,94,95,96,97}. An increasing amount of evidence reports that TLRs employ a variety of pro-tumor activities including proliferation, metastasis and are correlated with high relapse rate^{94,95,98}. Consequently, TLRs have been described as an important link between inflammation and tumor progression. On the other hand, TLR expression has also been demonstrated to exhibit reduced proliferation, a protecting role against tumor development and better outcome^{96,97,99}. TLRs are type I integral membrane glycoproteins belonging to the interleukin-1 receptor superfamily. They consist of an extracellular domain characterized by a leucine-rich repeats region and a conserved intracellular TIR (Toll/ IL-1R) domain, displaying homologies with the intracellular domain of the IL-1 receptor. TLRs are able to sense a broad range of molecular patterns conserved among microbial species. These recognizable molecules are referred to as pathogen-associated molecular patterns (PAMPs) and include e.g. LPS from Gram- bacteria, viral double stranded RNA, flagellin and CpG motifs recognized by TLR3, TLR4, TLR5 and TLR9 respectively¹⁰⁰. Upon ligation, the intracellular TIR domains dimerize resulting in a TLR conformational change necessary for the recruitment of intracellular adapter proteins essential for the downstream signaling¹⁰¹. All TLRs, except TLR3, signal through the MyD88-dependent pathway, which consequently activates NFκB and MAPK (AP-1, p38 and JNK) signaling pathways. Following their activation, NFκB, AP-1, p38 and

JNK induce the transcription factors of various pro-inflammatory mediators. TLR3 signals through the MyD88-independent (TRIF-TRAM) pathway which ultimately results in the activation of the IRF3 pathway resulting in the transcription of type I IFNs^{101,102}. In addition to the cell membrane location, TLR4 can also be internalized into the cytosolic endosomes and signal through TRIF-TRAM signaling pathway (Fig. 5)¹⁰³.

TLRs have been studied extensively as adjuvants in anti-cancer vaccine therapy. As vaccines based on antigens and peptides are poorly immunogenic, delivery of TLR agonists has been shown to enhance inflammatory responses^{104,105}. The idea of boosting the immune system to eradicate tumors is not new. Already a hundred years ago, William Coley discovered that administration of dead bacteria (*S. pyogenes* and *Serratia marcescens*) in cancer patients induced a fever which in many cases caused tumor regression¹⁰⁶.

TLR-induced inflammation can also be achieved in the absence of PAMPs. In situations of trauma and tissue injury, endogenous molecules, referred to as damage-associated molecular patterns (DAMPs), are released from necrotic or stressed cells¹⁰⁷. Additionally, in response to hypoxia as well as to radio- and chemotherapy-induced cell damage, tumors release DAMPs which consequently bind to TLR, thus provoking a sterile inflammation¹⁰⁸. An increasing list of intracellular DAMPs have been recognized, including high-mobility group box protein-1 (HMGB1), heat shock proteins, RNA and uric acid, just to mention a few^{107,109,110}. Generally, DAMPs act as alarmins to report danger and induce tissue repair¹¹⁰, however they have also been implicated in different malignancies¹¹¹.

S100A9 is a TLR4-binding DAMP that has been strongly associated with tumor progression^{112,113}. It belongs to the Ca²⁺-binding S100 protein family consisting of 21 members to date and exists as a S100A9 homodimer or a S100A8/S100A9 heterodimer¹¹⁴. Produced primarily by myeloid cells, in particular neutrophils and monocytes, the presence of S100A9 has been associated with various inflammatory processes. S100A9 proteins are for instance involved in the early macrophage differentiation, migration of phagocytes through modulation of the cytoskeleton and recruitment of inflammatory cells¹¹⁵⁻¹¹⁸. As mentioned above, S100A9 has been implicated with various malignancies. Expression of S100A9 was reported in numerous cancer types including hepatocellular carcinoma¹¹⁹, breast¹²⁰, colorectal¹²¹ and gastric cancer¹²². Increased expression of S100A9 has been correlated with worse overall survival in breast- and prostate cancer patients^{120,123}. In addition, S100A9 expression was correlated with high grade prostate cancers, whilst in a prostate cancer mouse model, S100A9^{-/-} mice display reduced tumor growth^{112,122}. Additionally, S100A9 promotes anti-inflammatory responses by accumulating immunosuppressive MDSCs¹²⁴. On the contrary, anti-tumor behavior employed by S100A9 has also been reported. In a study of acute myeloid leukemia

(AML), the S100A9 has been shown to arrest the proliferation and induce differentiation in AML cells as well as improve the survival in an AML mouse model ¹²⁵. In colorectal cancer, high density of S100A9 was detected in tumor infiltrating lymphocytes, suggesting that S100A9 might play a role in anti-tumor responses ¹²¹.




Figure 5. Overview of toll-like receptors and their ligands.

Ten different toll-like receptors are identified in humans, and they can be located both on the cellular membrane (TLR1, 2, 4, 5, 6 and 10) and in intracellular endosomes (TLR3, 7, 8 and 9). TLRs recognize a broad range of microbial pathogenic molecules (in black) and endogenous danger molecules from stressed and necrotic cells (in italic red). Upon binding to putative ligand, TLRs elicit inflammatory responses by activating various intracellular signaling pathways e.g. NFκB and IRF3.

The Adaptive Immune System

In case the innate immunity fails to quickly neutralize a pathogen or if an infection becomes too challenging for the innate immunity to eliminate, the adaptive immune system is activated. In contrast to the less specific innate responses, the adaptive immunity mediates a finely tuned antigen-specific response mediated by the antigen-specific T-cell receptors (TCRs) and B-cell receptors expressed by T cells and B cells respectively.

The T cells are divided into two major types: the CD4⁺ T-helper (T_h) cells which can differentiate into one of several T-helper subtypes (T_{H1}, T_{H2}, T_{H3}, T_{H9}, T_{H17}, T_{regs}, T_{FH}) each and one of them characterized by the type of cytokines they secrete or in case of T_{FH} their localization, and the effector CD8⁺ cytotoxic T cells (CTLs)¹²⁶. The contact between T cells and an antigenic peptide is mediated through TCR binding to either MHC class I or MHC class II molecules. MHC class I molecules, which are expressed on virtually all nucleated cells in the body, present cytosolic antigens recognized by CD8⁺ T cells, while MHC class II molecules, expressed by specialized antigen presenting cells (APCs), present foreign antigens recognized by CD4⁺ T cells. Depending on the nature of the encountered antigen, the T_h cells orchestrate a proper immune response by secreting a specific range of cytokines, thus provoking various immune activities such as enhancing T- and B cells responses, boosting antibody production and isotype switching as well as activating CTLs killing of infected cells¹²⁶. The killing mechanisms employed by the CD8⁺ CTLs include contact-dependent release of granules containing perforin, granzyme and apoptosis-inducing proteases as well as the upregulation of CTL- membrane bound Fas ligand which upon interaction of Fas receptor on the target cell, induces apoptosis¹²⁶.

Together with T cells, the adaptive immune system also comprises the B cells which when activated differentiate into antibody producing plasma cells that aid the combat through e.g. antibody dependent cellular cytotoxicity (ADCC), antibody-mediated neutralization of toxins, facilitating phagocytosis of bacteria as well as inhibiting their entry into host cells^{127,128}. Besides the highly antigen-specific responses, the other cornerstone of the adaptive immunity is its ability to create an immunological memory, meaning that upon a future encounter with same antigen, the adaptive immune responses are developed more promptly and effectively⁴⁸.

Besides the highly antigen-specific lymphocytes, the lymphoid cell family also comprises $\gamma\delta$ T cells and Natural Killer T (NKT) cells. Owing to their invariant TCRs and an innate immune receptor profile, they are referred to as “innate lymphocytes”. Contrary to the conventional T cells, $\gamma\delta$ T- and NKT cells are not dependent on the classical MHC:antigen presentation to become activated. Instead they utilize receptors such as NKD2 and CD1d to recognize their target. They also employ similar cytotoxic mechanisms as CTLs including the production of IFN γ and the release of granzyme and perforin, thus playing important role in anti-tumor immunity, although a pro-tumor role has been reported as well ¹²⁹⁻¹³³.

Immune Tolerance

Immune tolerance is a crucial aspect of the well-being and integrity of the body. It is imperative that the immune system is able to discriminate between self and non-self and with respect to this, central tolerance has been developed. It is a sophisticated system that takes place in the thymus, whose prime function is clonal selection: meaning, survival of self-MHC reactive T-cell progenitors, and death of self-antigen reactive T-cell progenitors¹³⁴. Central tolerance is an umbrella term comprising two main processes; the positive selection, taking place in the thymus cortex and the negative selection taking place in the thymus cortico-medullary junction and medulla. The T-cells that develop in the thymus derive from a small group of T cell progenitors presented as CD4⁻CD8⁻ (double negative, DN) cells which subsequently proliferate and give rise to a CD4⁺CD8⁺ (double positive, DP) TCR⁺ cell population. During the positive selection, majority of DP T-cell progenitors die by neglect as they lack or present very low affinity to either MHC class I and- MHC class II molecules. Conversely, DP T progenitor cells showing affinity to MHC class I differentiate to single positive (SP) CD8⁺ T cells, and DP T progenitor cells showing affinity to MHC class II on the other hand differentiate to single positive (SP) CD4⁺ T cells¹³⁵. In the thymic cortico-medullary junction and medulla, T cell progenitors displaying a too strong affinity to MHC:antigen molecules die by negative selection. The antigens are a broad range of tissue-specific antigens (TSAs) presented by MHC on the surface of medullary thymic epithelial cells (mTECs) and DCs, and T cell progenitors which have a too strong affinity to these, are hence deleted¹³⁶.

However, the central tolerance can be leaky and some T cells, despite being autoreactive, manage to escape the clonal deletion and enter the periphery where they potentially might give rise to autoinflammatory diseases¹³⁷. To retaliate towards these cells, self-tolerance is also maintained in the periphery through a process referred to as peripheral tolerance, which employs a variety of regulatory processes including the pivotal immunosuppressive regulatory T cells (T_{reg}), peripheral deletion, an anti-inflammatory cytokine environment, as well as upregulation of inhibitory co-receptors regulators programmed death 1 (PD-1) and cytotoxic T lymphocyte-associated antigen 4 (CTLA-4) (**Fig. 6**)^{134,138-142}. Immune tolerance is however not exclusively apparent during the process of self and- non-self-discrimination but can also become subject of manipulation by a cancer for its own advantage.

Autoimmune Regulator

Autoimmune regulator (AIRE) is a transcription factor primarily expressed in the thymus by mTECs. AIRE plays a critical role in the negative selection of T cells and mutations in the *AIRE* gene results in loss of self-tolerance and autoimmune diseases¹⁴³. AIRE induces a “promiscuous” gene expression of a broad range of TSAs^{144,145}, which subsequently are presented by MHC class II on the cell-membrane where they come in contact with T cells undergoing negative selection. Only self-tolerant T cells leave the thymus and collateral damage is avoided¹⁴⁶. Thymic DCs are also involved in the tolerization of T cells (**Fig. 6**)¹⁴⁷.

As some autoreactive T cells manage to escape the negative selection in the thymus, it is of highest importance to maintain a peripheral tolerance¹³⁷. Although AIRE is also expressed in non-thymic tissues, its functional significance in the peripheral tolerance remains poorly understood. Pollani *et al.* performed extensive immunohistochemistry of various human tissues and reported that, besides in the thymus, AIRE expression was observed in peripheral lymphoid organs, including lymph nodes, tonsils and gut-associated lymphoid tissues in colon and appendix.




Figure 6. Simplified overview of the mechanisms of central and peripheral tolerance.

Autoimmune regulator (AIRE) induces transcription of numerous tissue-specific antigens (TSAs), which subsequently are presented by MHC molecules on the cell surface of medullary thymic epithelial cells (mTECs) and dendritic cells (DCs). T cells displaying a too strong affinity for TSAs are subjected to clonal deletion. In case some auto-reactive T cells escape clonal deletion in the thymus, multiple mechanisms are employed in order to maintain a peripheral tolerance. Adapted from Mathis *et al.*, 2009¹⁴³

Additionally, they reported that AIRE was expressed by mature DCs, which displayed a tolerogenic phenotype by increased transcripts of various TSAs and the production of anti-inflammatory cytokine IL-10 as well as indoleamine 2,3-dioxygenase (IDO) ¹⁴⁸. AIRE expression in peripheral DCs has also been shown to reduce autoreactive CD4⁺ T cell viability, decrease the expression of PD-1 and increase the expression of TSAs in a diabetes type I mouse model, all indicating a tolerogenic role of AIRE ¹⁴⁹. Furthermore, it has been shown that T_{regs} deviate into an autoreactive CD4⁺ T convective cell population in *Aire*^{-/-} mice, demonstrating T_{reg} involvement in the peripheral tolerance ¹⁵⁰. On the contrary, another study showed that AIRE-induced genes in peripheral DCs did not include any TSAs but rather genes relevant in DCs maturation, suggesting functional divergence between peripheral and thymic AIRE-expressing cells ¹⁵¹. In cancer, AIRE expression has been assessed in a small cohort consisting of 39 breast cancer patients. However, AIRE expression was only evaluated in ER⁺ breast cancer patients and no conclusion regarding its localization or the functional role was drawn. However, when transfecting ER⁺ breast cancer cell lines with an AIRE vector, they exhibited increased G1-phase and reduced S-phase as well as increased apoptosis ¹⁵².

Endotoxin Tolerance

During an infection, it is of prime importance that a proper immune response is induced in order to clear the pathogen. It is likewise important that the immune response is tightly regulated as an overt inflammation might result in tissue damage and harm the host. One clinically relevant example demonstrating this is the course of sepsis, which usually is simplified as a two-step process. Following a bacterial infection, the systemic inflammatory response syndrome (SIRS) is induced, a state characterized by the release of various pro-inflammatory cytokines such as TNF- α and IL-6 which, if not controlled, might result in multi-organ failure and death ¹⁵³⁻¹⁵⁵. Subsequently, the second state is characterized by the compensatory anti-inflammatory response syndrome (CARS), as an effort to dampen the excessive inflammation. However, this state might pose the same danger as an overt inflammation, as immunocompromised patients are highly susceptible to secondary infections which might be life-threatening ^{155,156}.

Functional parallels can be drawn between CARS and endotoxin tolerance, a mechanism described as a state of reduced responsive ability of monocytes and macrophages upon a repetitive LPS stimulation ¹⁵⁷. Endotoxin tolerance was described over 70 years ago by Paul Beeson which discovered that rabbits repeatedly injected with typhoid vaccine displayed reduced fever ¹⁵⁸. Today, multiple molecular mechanisms of endotoxin tolerance have been reported. As the principal receptor for recognizing LPS ¹⁵⁹, it is not surprising that TLR4 and its

downstream signaling pathway is involved in endotoxin tolerance. Repetitive LPS stimulation decreases the expression of TLR4 in neutrophils¹⁶⁰ and macrophages¹⁶¹ impairing the TLR4-MyD88 complex formation in monocytes¹⁶². Furthermore, endotoxin tolerant monocytes fail to generate activation of IRAK-1¹⁶². Expression of an alternatively spliced form of MyD88 (MyD88s), lacking the intermediary domain crucial for IRAK-1 and IRAK-4 binding, diminishes the subsequent downstream activation of NFκB¹⁶³. MyD88s expression in monocytes has been shown to increase upon LPS treatment¹⁶⁴. Besides MyD88s, several other negative regulators of TLR4 downstream signaling pathway have been reported, including factors such as IRAK-M, MKP-1 and A20, which inhibit IRAK1/4, MAPK and TRAF6 respectively^{165,166}. Endotoxin tolerance can also be induced by p50p50 homodimer formation, which not only induces IL-10 and inhibits TNF-α transcription¹⁶¹, but also binds to various DNA promoters and by acting as a steric hinder, p50p50 inhibits the p65p50 activity¹⁶⁷. Endotoxin tolerant cells are characterized by a diminished ability to present antigens¹⁶⁸⁻¹⁷⁰ and to produce inflammatory cytokine such as IL-6, TNF-α and IL-1¹⁷¹, whilst producing anti-inflammatory cytokines such as IL-10¹⁷⁰.

Myeloid-Derived Suppressor Cells

First reports describing cells with immunosuppressive properties emerged in the late 70's^{172,173}, and in the mid 90's, the first evidence of suppressor cells in tumors emerged^{174,175}. Originally described with an ambiguous terminology, in 2007, the term myeloid-derived suppressor cells (MDSCs) was coined by Gabrilovich *et al.* to describe the origin as well as the distinctive immunosuppressive function of these cells¹⁷⁶.

MDSCs comprise a heterogenous immature myeloid cell population usually divided in two different subgroups; the polymorphonuclear or granulocytic MDSCs (G-MDSCs) and the monocytic MDSCs (Mo-MDSCs). The phenotype of human MDSCs still remains poorly defined, and characterizing them solely by their cell-surface marker expression is challenging due to overlap in marker expression with other myeloid cells⁸³. Generally, G-MDSCs are described as immature CD11b⁺CD33⁺HLA-DR^{-/low}Lin⁻, the latter as they lack lineage markers. However, the expression of CD15 and CD66b as well as a low density in Ficoll gradients has also been recorded. G-MDSCs have been described to resemble neutrophils both morphologically and phenotypically. Mo-MDSCs, which phenotypically and morphologically are similar to monocytes, are defined as CD14⁺CD11b⁺HLA-DR^{-/low}CD15⁻^{177,178}. In contrast to human MDSCs, characterizing mouse MDSCs phenotype is less challenging. Mouse MDSCs are described as CD11b⁺Gr1⁺ cells,

where G-MDSCs and Mo-MDSCs are defined as Ly6G⁺/Ly6C⁻ and Ly6G⁻/Ly6C⁺ respectively ¹⁴².

MDSCs have been described in different pathological conditions such as sepsis ¹⁷⁹⁻¹⁸¹, chronic infection¹⁸²⁻¹⁸⁴ and autoimmune disease ^{185,186}. However, the most extensive research on MDSCs has been focusing on cancer ^{178,187}, where MDSCs have been associated with tumor angiogenesis, metastasis and tumor burden both in cancer patients and in mouse models ¹⁸⁸⁻¹⁹⁰.

The expansion and activation of MDSCs has been described as a two-signal process which includes multiple factors produced by activated T cells, tumor stromal cells and tumor cells themselves ⁸⁴. The first signal involves expansion of MDSCs which is mediated by a broad range of factors including GM-CSF, G-CSF, M-CSF, IL-6, VEGF, and COX2, with STAT3 as the main signaling pathway through which they operate. The second signal involves activation of MDSCs, which on the other hand is mediated by NFκB through various factors such as TLR ligands and different pro-inflammatory proteins. ¹⁴². On the contrary, it has also been reported that the expansion signal is instead mediated through NFκB, while STAT3 is responsible for the MDSCs activation ¹⁹¹. Furthermore, S100A8 and S100A9 proteins have also been shown to play a role in the expansion of MDSCs ¹²⁴. However, the generation and activation models do not necessarily cover both G- and Mo-MDSC.

The most distinctive feature of MDSCs is their incredible immunosuppressive ability, which until today remains the golden standard for MDSCs characterization (**Fig. 7**). One potent mechanism through which MDSCs induce immunosuppression is by inhibition of T cell function. Production of arginase-1 (arg-1) and iNOS by MDSCs results in depletion of L-arginine, which is crucial for T cell proliferation. T cells present in an L-arginine- deficient environment remain in their G0-G1 cell-cycle phase ¹⁴². Another way MDSCs can impair T cell function is through the production of reactive oxygen species (ROS) which exert several tumor-promoting activities such as inducing downregulation of T cell receptor ζ chain and thus inhibiting T cell activity, enhancing the malignant cell resistance against CTL cytotoxicity and impairing the T cell infiltration into the tumor site ^{178,192}. MDSCs further induce immunosuppression by producing immunosuppressive cytokines including IL-10 and TGF-β, and are also able to promote angiogenesis through the production of e.g MMP9 ^{178,190}. MDSCs have also been reported to induce T_{regs}, which subsequently inactivate T cells, as well as to impair the cytotoxic function of natural killer cells. Generally, immunosuppressive function between Mo-MDSCs and G-MDSCs differ being characterized with following profiles; Mo-MDSCs: Arg/IL-10/IDO and G-MDSCs: ROS ^{84,178,193,194}.

While the function of MDSCs has been extensively studied, the origin of MDSCs still remains an unexplored field. There are various theories explaining their potential origin as reviewed in ^{83,84}. In *Paper IV*, we have put effort in exploring the

origin of G-MDSCs where we show that G-MDSCs from metastatic breast cancer (MBC) patients cluster together with healthy donor neutrophils in a gene expression profile array, sharing genes relevant in angiogenesis and immunosuppression among others leading us to the conclusion that G-MDSCs might actually be a subset of neutrophils. This controversial theory has been contemplated by others owing to the phenotypical and morphological resemblance between G-MDSCs and neutrophils, the fact that activated neutrophils have been discovered in the low-density mononuclear cell fraction of Ficoll density gradient in cancer patients, and most importantly, the ability of neutrophils to induce ROS-dependent immunosuppression^{83,195}. Indeed, MDSCs and neutrophils share many similarities and one recent study has shown that MDSCs are able to produce NETs, a mechanism traditionally performed by neutrophils¹⁹⁶.




Figure 7. Myeloid-derived suppressor cells and their multiple immunosuppressive functions.
 Illustration adapted from Millrud *et al.*, 2017⁸⁴.

The Immune System and Cancer

Over a century ago, Paul Ehrlich predicted that our immune system plays a protective role against cancer. Fifty years later, Burnet and Thomas continued on that path and proposed that our immune system, involving T cells as principal effectors, is able to detect and eliminate cancerous cells. Consequently, the term “immunosurveillance” was coined. The concept of immunosurveillance was however reconsidered and the theory that our immune system not only protects us from cancer but is also involved in processes of sculpting the cancer emerged. The dynamic process involving the immune system and its host-protective, tumoricidal actions against tumors, together with its involvement in shaping the tumor characteristics, gave rise to a new term referred to as “cancer immunoediting” including three phases; elimination, equilibrium and escape (**Fig. 8**)¹⁹⁷⁻¹⁹⁹.

During the elimination phase, the immune system is activated by tumor associated antigens (TAAs) and when primed, it is able to provoke anti-tumor responses¹⁹⁸. Innate lymphocytes, including NK, NKT and $\gamma\delta$ T cells are able to recognize and kill transformed cells, and lymphocyte tumor infiltration has been correlated with better prognosis in several cancer types²⁰⁰⁻²⁰⁴. Phagocytosis of TAAs by DCs consequently results in T cell activation through antigen presentation in the lymph nodes. Ultimately, tumor specific adaptive T cells are recruited to the tumor site and participate in the process of tumor elimination¹⁹⁸. IFN- γ and IL-12 represent some of the principal cytokines in tumor rejection^{139,198,205,206}, and other tumoricidal molecules employed by the immune system include perforin and TRAIL²⁰⁷⁻²⁰⁹. The impact of the adaptive immune system in anti-tumor immunity has been demonstrated by the RAG2^{-/-} mice characterized by their lack of mature T, B, $\gamma\delta$ T and NKT cells. These mice display an increased frequency of carcinogen-induced and spontaneous tumor development^{139,210}. The importance is further demonstrated in immunosuppressed patients which may develop various cancer forms following e.g. organ transplantation²¹¹.

In situations where a tumor survives the immunosurveillance, it subsequently enters the equilibrium phase. During this phase, there is still an ongoing anti-tumor response, however, not potent enough. Instead of completely eradicating the tumor, the surviving tumor cells are subjected to a “survival of the fittest” selection which ultimately results in the generation of immune-resistant tumors¹⁹⁷.




Figure 8. The three E's of of cancer immunoeediting.

During *elimination* phase, the highly immunogenic cancer cells are detected by the immune system and are successfully eradicated. Over time, the tumor enters the *equilibrium* phase, characterized by a balance between tumor death and tumor growth. Ultimately, cancer cells become non-immunogenic and the tumor creates an immunosuppressive environment in which it thrives and *escapes* the immune destruction. Adapted from Dunn *et al.* 2002, 2004 ^{197,198}.

During this phase, there seems to be a balance between anti-tumor and immunosuppressive cells and mediators ¹⁹⁸.

During the escape phase, the tumor fully engages in mechanisms of avoiding the immune system. One such mechanism includes camouflaging by e.g. reducing the expression of MHC class I molecules and TAAs ¹⁹⁸. One major obstacle of successful anti-tumor responses is T-cell exhaustion. In the context of an infection, a persistent activation of T cells may result in the development of exhausted T cells as a regulatory mechanism of bringing a balance between inflammation and host-protection. Several mechanisms lie behind T cell exhaustion including a reduced ability to produce IL-2 and IFN- γ cytokines as well as upregulation of PD-1 ^{212,213}. This is also seen in tumors. To further avoid anti-tumor immunity, tumors promote accumulation of immunosuppressive cells, generating an immunosuppressive environment. The most prominent immunosuppressive cells in a tumor microenvironment are the TAMs and MDSCs. Both cell types suppress T cell activity through the secretion of immunosuppressive mediators as described above

^{68,142,178}. Another potent immunosuppressing cell population include the CD4⁺CD25^{high}FOXP3⁺T_{regs}. The prevalence of T_{regs} is increased in various malignancies and they apply their immunosuppressive functions through e.g. expressing CTLA-4, affecting DC maturation, sequestering IL-2 and the release of anti-inflammatory IL-10 and TGF-β²¹⁴⁻²¹⁹. Essentially, the immunoediting process develops tumor cells resistant to an immuno-competent environment.

Cancer Immunotherapy

Cancer immunotherapy is a collective term for a broad range of therapeutic strategies which utilize the immune system to fight cancer. These therapeutics have expanded immensely in the last two decades and revolutionized the way various cancers are treated. It is therefore not surprising that James P. Allison and Tasuku Honjo were awarded with Nobel Prize in Physiology and Medicine, 2018 for their contribution to the field of cancer immunotherapy. Their discoveries are perhaps among the most notable success stories within the field which include the practice of inhibiting negative T cell regulators; CTLA-4 and PD-1, in order to boost the anti-tumor responses^{220,221}. A complete T cell activation requires binding of a TCR to an antigen-presenting MHC-molecule together with the binding of costimulatory molecules CD28 and B7 present on T cells and APCs respectively. Also present on T cells are CTLA-4 and PD-1, which are CD28 homologues that impair T cell activation by distinct ways; CTLA-4 by inhibiting the CD28:B7 binding by binding to B7 itself in a competitive manner, and PD-1 by binding to its own ligand PD-L1, which often is present on tumor cells²²². Currently, monoclonal antibody-mediated CTLA4 and PD-1 inhibition (also referred to as immune checkpoint inhibition (ICI)) is used to treat various malignancies including melanoma, kidney cancer, bladder cancer and non-small cell lung cancer. Several ongoing phase II and III clinical trials are exploring the effects of the ICI on multiple other malignancies^{222,223}. Apart from ICIs, another therapeutic strategy explored include adoptive therapy where autologous T cells are modified *ex vivo*. One example are the chimeric antigen receptor (CAR) T cells which are not dependent on the classical TCR:MHC binding for their activity, but are already loaded with an antigen. Non-Hodgkin lymphoma and acute lymphoblastic leukemia patients are currently benefiting from this immunotherapy as CD19-directed CAR T cells have been approved by the FDA last year^{224,225}. The use of monoclonal antibodies (targeted therapy) has successfully been used in some cancer types by means of targeting the TAAs. In breast cancer, the function of HER2/neu receptor is inhibited by Trastazumab²⁶.

Wnt

Wnt proteins constitute a family of secreted cysteine-rich glycoproteins highly conserved among species, holding important roles in tissue homeostasis, embryonic development, and cancer²²⁶⁻²²⁸. The significance of Wnt's in oncogenesis was first evident upon the discovery of *Integration 1 (Int1)* in mice, a gene to which the oncogenic mouse mammary tumor virus (MMTV) integrates²²⁹. Later, it was found that *Int1* and *Drosophila wingless (wg)* were orthologs²³⁰, and consequently by combining *wg* and *Int*, the term *Wnt1* was coined²³¹. To date, 19 Wnt proteins in mammals have been identified and they operate through multiple receptors and co-receptors including the seven-transmembrane G-coupled Frizzled protein family (FZD), receptor tyrosine kinase (RTK), ROR1, ROR2 and the low-density lipoprotein receptor-related proteins 5 and 6 (LRP 5/6)^{227,232}. To become fit for action, Wnt proteins are subjected to a series of post-translational modifications giving Wnt's their hydrophobic properties crucial for their secretion and function^{233,234}. Wnt proteins are generally divided into two different subgroups; the transforming vs. the non-transforming subgroup based on their ability to transform C57MG mammary epithelial cells or the canonical and the non-canonical subgroups based on the signaling pathway they activate²³⁵.

Canonical Wnt signaling

Wnt/ β -catenin signaling pathway is one of the main signaling pathways activated by Wnt proteins. It involves the transcription factor β -catenin whose transcriptional activity relies on a canonical Wnt protein (e.g. Wnt1, Wnt3a and Wnt8) binding to a FZD receptor and the LRP5/6 co-receptor^{236,237}. In the absence of a Wnt protein, β -catenin is bound and regulated by a destruction complex consisting of several intracellular proteins including Axin, casein kinase 1 (CKI), adenomatous polyposis coli (APC) and glycogen synthase kinase-3 beta (GSK3 β). β -catenin is subsequently phosphorylated and ubiquitinated and ultimately subjected to proteasome degradation, keeping intracellular β -catenin levels low. The engagement of a Wnt protein with its putative FZD receptor along with the co-receptor LRP5/6 ultimately results in the dissemblance of the destruction complex and the release of β -catenin. The disruption of the destruction complex is mediated through the translocation of

Dishevelled (Dvl) to the cellular membrane and its recruitment of GSK3 β and Axin complex²³⁷. Conclusively, the unbound and stable β -catenin is able to translocate to the nucleus and activate the transcription of various TCF/LEF inducible genes e.g. *c-myc*^{236,238}.

Non-canonical Wnt signaling

The best described non-canonical Wnt signaling pathways, which also are referred to as β -catenin independent, include the Wnt/Ca²⁺, the Wnt/planar cell polarity (PCP) and the Wnt/ROR2 signaling pathways. These pathways are activated by non-canonical Wnt proteins (typified by Wnt5a and Wnt11) through their engagement with various FZD receptors alone or in combination with their co-receptor ROR2^{227,237,239}.

Cell migration and cell polarity is regulated by the Wnt/Ca²⁺ signaling pathway. Activation of the signaling pathway is initiated upon the ligation of a non-canonical Wnt protein to its putative FZD receptor, resulting in the increase of intracellular calcium influx. The intracellular accumulation of calcium in turn activates various effectors such as protein kinase C (PKC), calcineurin and calmodulin- dependent protein kinase II (CaMKII), where the latter has been shown to play an inhibitory role of the β -catenin signaling pathway through its activation of e.g. nemo-like kinase (NLK)²⁴⁰⁻²⁴².

Wnt/PCP signaling pathway is another non-canonical pathway which controls the cytoskeletal reorganization. The cellular orientation and movement is achieved through the binding of a non-canonical Wnt protein binding to a FZD receptor which ultimately results in the activation of JNK and ROCK. The activation of the JNK and ROCK is mediated via Dvl which induces the activation of the small GTPases Cdc42, Rac and Rho^{237,241}. Wnt/PCP signaling pathway is crucial during normal development and is activated during cancer²⁴³.

Wnt5a

The most extensively investigated non-canonical Wnt protein is Wnt5a, a non-transforming secreted glycoprotein that activates the β -catenin-independent non-canonical Wnt signaling pathway. Since its discovery in 1990²⁴⁴, the biological role of Wnt5a has been studied in various fields including developmental biology, inflammatory diseases and cancer^{245,246}. The importance of Wnt5a in development is illustrated by the knock-out of Wnt5a (*Wnt5a*^{-/-}) in mice displaying a phenotype of facial abnormalities, dwarfism and shortened limbs and tails, dysmorphic ribs,

heart defects, respiratory dysfunction and death shortly upon birth^{227,246}. The role of Wnt5a in cancer has proven paradoxical as being recognized both as a tumor-suppressor and a tumor-promotor depending on the cancer type. Down-regulation of Wnt5a has been observed in several cancer types including breast cancer²⁴⁷, acute lymphoblastic leukemias^{248,249}, and colon cancer²⁵⁰. Reduced Wnt5a levels have been correlated with worse overall clinical outcome in breast- and colon cancer patients^{250,251}. A lower Wnt5a expression is also evident in the more aggressive ER⁻MDA-MB-231 cell-line as compared with the ER⁺ non-metastatic MCF-7 cell-line²⁴⁷. Furthermore, Wnt5a plays a tumor-suppressing role in thyroid carcinoma by inhibiting the proliferation and migration in a Wnt/Ca²⁺/CaMKII signaling-manner leading to β -catenin degradation²⁵², and Wnt5a hemizygous mice display increased B-cell proliferation and develop myeloid leukemia²⁴⁸. In contrast, the tumor-promoting ability of Wnt5a has been reported in gastric cancer where Wnt5a promotes migration and invasion by activating the GTP-binding protein Rac and the focal adhesion kinase (FAK) protein. In addition, Wnt5a positivity in gastric cancer patients is correlated with higher tumor stage and worse overall survival²⁵³. Similarly, Wnt5a promotes invasiveness and motility by activating PKC in metastatic melanoma²⁵⁴, induces increased proliferation and is associated with poor prognosis in non-small cell lung carcinoma patients²⁵⁵ and induces EMT in pancreatic cancer²⁵⁶. Additionally, in several melanoma cell lines, Wnt5a has also been shown to induce exosome-mediated secretion of several tumor promoting mediators such as IL-8, VEGF, MMP2 and IL-6²⁵⁷.

Besides its crucial role in embryonic development and cancer, several studies have also addressed the involvement of Wnt5a in immune cell regulation. Bergenfelz *et al.* have shown that Wnt5a inhibits the monocyte differentiation towards M1 macrophages and instead induces an alternative monocyte activation producing a tolerogenic phenotype of macrophages¹⁶¹. Similarly, the Wnt5a-induced tolerogenic cell phenotype has also been observed in monocytes differentiated towards immature dendritic cells^{258,259}. The tolerogenic DCs obtained displayed a reduced antigen-presenting ability as well as a reduced production of pro-inflammatory cytokines and increased producing anti-inflammatory cytokines²⁵⁹. Moreover, Wnt5a has been shown to reduce the generation of mature DCs²⁵⁸ and to inhibit the B-cell proliferation²⁴⁸ and T-cell development²⁶⁰. In contrast, Wnt5a has also been implicated in inflammatory responses. Increased expression of Wnt5a has been observed in several inflammatory diseases such as sepsis, atherosclerosis and rheumatoid arthritis²⁶¹⁻²⁶³. Blumenthal *et al.* have shown that Wnt5a plays a role in Th1 responses by regulating the expression of IL-12 and IFN- γ in human macrophages and PBMCs²⁶⁴. Treatment of human macrophages with recombinant Wnt5a significantly induced the release of several pro-inflammatory cytokines including IL-1 β , IL-6 and IL-8²⁶¹. Moreover, Wnt5a treatment enhanced the ability of human bone marrow stromal cells (BMSCs) to attract T-cells and monocytes

through several distinct pathways, including NF κ B and MAPK ²⁶³. It has been reported that recombinant Wnt5a induced pro-inflammatory cytokines in mouse macrophages in a TLR4-dependent pathway. The authors speculated that this might be an artifact caused by contaminated Wnt5a preparations ²⁶⁵. Indeed, this is a logical explanation, however, in this context, quite far from the truth. In *Paper III*, we show for the first time that Wnt5a in fact is a novel TLR2 and TLR4 ligand eliciting pro-and anti-inflammatory cytokine profiles in human and mouse monocytes respectively.

The Present Investigation

Aims

The general aim of this thesis is to investigate various molecular mechanisms of immune tolerance in human breast cancer, including the generation of myeloid-derived suppressor cells (MDSCs) through a novel Toll-like receptor 4 (TLR4) ligand (Wnt5a), and the expression of Autoimmune regulator (AIRE).

The specific aims were:

- I. To investigate the expression of TLRs in breast cancer, and to analyze the functional role of TLR4 and its clinical relevance in breast cancer patients.
- II. To investigate the expression and the localization pattern of the DAMP S100A9, its functional role as well as its clinical relevance in breast cancer patients.
- III. To investigate Wnt5a as a novel ligand for TLR2 and TLR4.
- IV. To investigate the prevalence and generation of G-MDSCs in human breast cancer patients.
- V. To investigate the expression of Autoimmune regulator (AIRE) and its functional characteristics in human breast cancer.

Paper I

Expression and function of toll-like receptor 4 correlates with worse overall survival in ER-/PR- breast cancer.

Background

Our immune system has a crucial task in protecting our body from infectious pathogens. In order to recognize intruders, the immune system has employed a group of pattern-recognition receptors known as TLRs. Being expressed on various immune cells, they are able to sense danger in form of exogenous PAMPs and endogenous DAMPs, thus inducing proper inflammatory responses^{85,87,88,107}. An increasing amount of evidence reports that TLRs are expressed on various cancer cell types, contributing to cancer progression^{94,95,98}. In *Paper I*, we aimed to investigate the expression of TLRs in multiple breast cancer cell lines with the focus on TLR4 expression and function. Additionally, we aimed to investigate the clinical relevance of TLR4 expression in breast cancer patients.

Results and Discussion

The expression of TLRs has previously been reported in several cancer types^{92,94,95}. We started off by investigating the mRNA expression pattern of TLR2, TLR3, TLR4, and TLR9 as well as TLR4 co-receptors MD2 and CD14 in seven different breast cancer cell lines; four TN (MDA-MB-231, MDA-MB-468, SUM-149 and SUM-159) and in three ER⁺PgR⁺ (MCF-7, T47D and CAMA). The results showed that *TLR2* and *TLR3*, *TLR4* and the co-receptor *CD14* are favorably expressed in TNBC cell lines, whilst *TLR9* was generally expressed in all cell lines. Moreover, the co-receptor *MD2* was expressed in all TNBC cell lines, except in MDA-MB-468, meaning that three out of four TN breast cancer cell lines contain all necessary components for a functional TLR4 receptor. Furthermore, immunohistochemical analysis revealed a cytoplasmic TLR4 expression in MDA-MB-231, SUM-149 and SUM-159 cell lines, while being completely absent in MDA-MB-468 and in the ER⁺PgR⁺ cell lines. Although TLR4 is generally located on the cellular membrane, several studies have reported intracellular TLR4^{91,93,266,267}. To investigate the TLR4 functionality, we stimulated all seven cell lines with LPS. Our results show a marked increase of IL-6 and IL-8 and a low increase in TNF- α in the supernatants of MDA-MB-231, SUM-149 and SUM-159 cell lines. Moreover, NF κ B activity was significantly increased when stimulating MDA-MB-231 cells with LPS. Constitutively active NF κ B has been observed to be more prevalent in Basal-like compared to Luminal breast cancers. It has furthermore been correlated with

multiple tumor promoting activities such as proliferation, angiogenesis and metastasis^{268,269}. The fact that TLRs studied in this project are preferentially expressed in TNBC tumors might be a causality to the constitutive activity of NFκB observed in ER⁻ breast cancers. In order to mimic a sterile inflammation, we treated TNBC cell lines with two DAMPs; HMGB1 and S100A9, and while HMGB1 had no effect, S100A9 induced a significant IL-6 and IL-8 increase in MDA-MB-231, SUM-149 and SUM-159 cell lines. These results are supported by previous findings showing similar S100A9 effects, where it induces IL-6 and IL-8 in a human monocytic cell line²⁷⁰. Furthermore, MDA-MB-231 cells transfected with siRNA against TLR4 showed a low, yet a significant decrease in both IL-6 and IL-8 release, suggesting other mechanisms involved in the IL-6/IL-8 induction. It has previously been shown that ER⁻ breast cancers display a rather inflammatory milieu characterized by cytokine production which is associated with tumor aggressiveness²⁷¹. Moreover, the lack of ER expression has also been ascribed as one of the prognostic factors correlated with worse outcome in breast cancer patients²⁷². In order to investigate the chemoattractant ability employed by different breast cancer cell lines, we co-cultured human primary CD11b⁺ PBMCs together with conditioned media from two ER⁺PgR⁺ (MCF-7 and T47D) and two TNBC (MDA-MB-231 and MDA-MB-468) cell lines. Our results demonstrated that myeloid cells migrated towards MDA-MB-468 and MDA-MB-231, however, only statistically significant towards the latter. Additionally, MDA-MB-231 cells stimulated with LPS displayed a significantly higher invasive behavior in comparison to their untreated counterparts. LPS-induced TLR4 activation has previously been reported to promote metastasis²⁷³. A report by Volk-Draper *et al.* has also shown that TLR4 stimulation by paclitaxel induced breast cancer invasion and metastasis *in vivo*²⁷⁴. Lastly, in a TMA of 144 breast cancer patients, we investigated the expression of TLR4 and analyzed its expression with different histological and clinical parameters. From our analysis, we could draw the conclusion that TLR4 correlates significantly with the ER⁻PgR⁻ breast cancer patient group and that high TLR4 expression is significantly correlated with worse recurrence-free survival. These results are in line with previous published evidence shown the TLRs involvement in carcinogenesis^{94,98}.

In conclusion, functional TLR4 is expressed and is correlated with worse recurrence-free survival in ER⁻PgR⁻ breast cancer patients. The ability of TLR4 to induce IL-6 and IL-8, two pro-tumorigenic cytokines, should be taken in serious consideration in terms of therapeutic targeting in order to reduce the tumor inflammation.

Paper II

S100A9 expressed in ER(-) PgR(-) breast cancers induces inflammatory cytokines and is associated with an impaired overall survival.

Background

S100A9 is a Ca²⁺-binding protein that has been implicated in various inflammatory processes and cancer. S100A9 acts as a DAMP by binding to TLR4 and subsequently eliciting NFκB activity. Several immune cells, most notably neutrophils and myeloid cells, secrete S100A9, and although S100A9 protein expression in breast cancer has been investigated previously, the reports were rather inadequate¹¹⁴⁻¹¹⁸. In this paper, we wished to further elaborate on the expression pattern and function of S100A9 in breast cancer. We were also interested in investigating whether S100A9 expression correlated with any clinicopathological features that would be of relevance in breast cancer patients.

Results and Discussion

The expression of S100A9 has predominantly been attributed to immune cells and knowledge regarding its expression pattern in breast cancer is rather undefined^{115,117}. Several attempts to explore the expression of S100A9 in breast cancer have been employed, both *in vitro* and in cohorts of patients. However, the cohorts used were either small or the assessments were incomprehensive²⁷⁵⁻²⁷⁸. To shed some light on this area, we started off by assessing the S100A9 expression in six different breast cancer cell lines: three ER⁺PgR⁺ (MCF-7, T47D and CAMA), two TNBC (MDA-MB-231 and MDA-MB-468) and one HER2⁺ (SKBR3). Gene expression analysis revealed that *S100A9* was only expressed in the TNBC MDA-MB-468 and HER2⁺ SKBR3 cell lines. These results were confirmed by the immunohistochemical (IHC) analysis on paraffin-embedded cell pellets from all mentioned cell lines, which demonstrated S100A9 protein expression in MDA-MB-468 and SKBR3 solely. We further investigated S100A9 protein expression pattern on a TMA consisting of 144 breast cancers. Our results demonstrated cytoplasmic, nuclear, membrane and stromal S100A9 expression in all compartments significantly correlated with ER⁻PgR⁻ hormone receptor status. Reports regarding S100A9 expression in different breast cancers have indicated a tendency towards ER⁻ breast cancers²⁷⁸. However, one study showed that S100A9 is primarily expressed by non-invasive cell lines and primary breast cancers compared to the invasive counterparts²⁷⁹. In order to study the inflammatory responses S100A9 stimulation exerts on breast cancers cell lines, we performed a cytokine bead array.

Our results show that S100A9 induced a significant release of IL-8 and IL-1 β in MDA-MB-231 cell line only. Also, S100A9 induced IL-6 release in T47D, CAMA and MDA-MB-231 cell lines, however, only significantly in the latter. In *Paper I*, we have demonstrated the ability of S100A9 to induce cytokine secretion, however, only in TNBC cell lines²⁸⁰. This experiment on the other hand includes a broader array of cell lines, providing a more comprehensive view of S100A9 effects in breast cancer. Similar effects exerted by S100A9 have been shown in myeloid cells²⁷⁰. Despite expressing the highest level of S100A9, MDA-MB-468 and SKBR3 cell lines did not respond to S100A9 treatment. This can be explained by our next experiment where we assessed the expression of TLR4. Neither MDA-MB-468 nor SKBR3 expressed *TLR4*, while on the other hand, *TLR4* was significantly expressed in the MDA-MB-231 cell line. This is further confirmed by our results from *Paper I* where we report the expression and the lack of expression of TLR4 in MDA-MB-231 and MDA-MB-468 cell lines respectively²⁸⁰. As a ligand for TLR4, S100A9 elicits a NF κ B activity²⁷⁰. To confirm this notion in breast cancer cells, we performed a Dual luciferase reporter assay with a NF κ B reporter in MDA-MB-231, MDA-MB-468 and SKBR3. As expected, NF κ B activity upon S100A9 stimulation was only observed in the MDA-MB-231 cell line. In order to investigate whether S100A9 expression was associated with any clinicopathological features or myeloid cell markers, we performed IHC on a TMA consisting of 144 breast cancers. Besides the correlation with ER⁻PgR⁻ breast cancers as mentioned above, stromal S100A9 expression significantly correlated with HER2 and proliferation marker Ki67 positivity, larger tumor size and nodal stage. The stromal S100A9 expression in HER2 positive cancers is in line with our qPCR results, where the HER2⁺ SKBR3 cell line was one of the cell lines that expressed S100A9. Similar observations of the correlation between S100A9 expression and HER2 positivity was previously reported by Arai *et al.*²⁷⁵. We further observed a significant correlation between stromal S100A9 expression and the anti-inflammatory CD163⁺ myeloid cell, but not with CD68⁺ macrophages. Interestingly, we have reported previously that CD163⁺ myeloid cells are located in the stroma and not in the nest of breast cancer patients and in addition, S100A9 expression has been described in MDSCs^{70,124}. The expression of S100A9, both by tumor and immune cells in different tumors has been correlated with worse outcome^{123,275,281}, therefore, it would be logical to speculate that the presence of S100A9 expression might have a negative impact on the outcome in the patients from our cohort. Indeed, the expression of S100A9 both in the stroma and the cytoplasm was significantly associated with a worse survival in ER⁻PgR⁻ breast cancer patients.

Paper III

Wnt5a is a TLR2/4-ligand that induces tolerance in human myeloid cells

Background

In *Paper I* and *II*, we have shown that TLR4 is expressed in ER⁻ breast cancers is functional and more importantly, that it responds to DAMPs. In *Paper III*, we continued on this path and aimed to investigate Wnt5a as a novel endogenous TLR2/TLR4 ligand. Wnt5a is a non-canonical protein with important functions in development, cancer and inflammation²⁴⁵. Previous studies have reported elevated levels of Wnt5a in sepsis and as a result of LPS stimulation in myeloid immune cells.

Results and Discussion

Several independent studies have reported the ability of Wnt5a to induce various inflammatory mediators^{161,263,265,282} and many of them in MAPK and NFκB-dependent manner, strikingly similar to those induced through TLR signaling pathways. A study from 2014 speculated that this might be a result of endotoxin-contaminated recombinant Wnt5a preparations²⁶⁵, however, we thought that it might be the other way around and hypothesized that Wnt5a is a ligand for TLR4. By means of several methodologies, including co-localization of transfected Wnt5a-HA and hTLR4-GFP in NIH3T3 cells, SRP binding Biacore analysis and *in vitro* binding assays, we confirm that Wnt5a is a novel ligand for TLR2 and TLR4. Confirming previous reports, we show that Wnt5a treatment of primary monocytes and a monocyte cell line; THP1, induces an anti-inflammatory cytokine profile characterized by the increased secretion of IL-10, IL-8 and IL-6 and the downregulation of TNF-α. On contrary, in primary mouse bone marrow macrophages (BMM) and the macrophage cell line RAW264.7, Wnt5a induced a pro-inflammatory cytokine profile characterized with an increased secretion of TNF-α and MCP-1 (CCL2). The contradictory Wnt5a-induced response in human and murine immune cells (anti-inflammatory vs pro-inflammatory respectively) was rather confusing in the beginning, however, when diving into the literature, we slowly realized that this same pattern has been observed by other research groups^{161,259,283,284}. When performing NFκB reporter assays, Wnt5a did not activate NFκB in THP1-Blue cells. Hence, we propose that Wnt5a operates through alternative signaling pathways, including non-classical NFκB p50-homodimer formation as we previously showed¹⁶¹, TLR4-PI3K pathway and cytokine secretion through

exocytosis^{161,257,285}. When stimulating BMM from *MyD88*^{-/-} and *TLR4*^{-/-} mice with Wnt5a, we could see a completely abrogated TNF- α release in *TLR4*^{-/-} BMM and partially in *MyD88*^{-/-} BMM. These results further confirm hypothesis that Wnt5a is a novel TLR4 ligand. On the other hand, an increased MCP-1 secretion was observed in both *TLR4*^{-/-} and *MyD88*^{-/-} BMM. Furthermore, when treating human primary monocytes with Wnt5a in combination with TLR4 ligands (PAMP; LPS or DAMPs; HMGB1 and S100A9), we could observe increased prevalence of CD14⁺HLA-DR^{low/-} Mo-MDSCs cells *in vitro*. A Wnt5a-induced immunosuppressive phenotype has been observed previously. Valencia *et al.* reported that Wnt5a promoted an immunosuppressive phenotype of mDCs characterized by the reduction of HLA-DR and CD86 and upregulation of PD-L1 and a rendered LPS-induced activation²⁵⁹. Similar findings were reported in a study which demonstrates that Wnt5a blocks the M1 differentiation and instead induces a tolerogenic CD14⁺HLA-DR^{low/-}co-receptor^{low/-} cell population¹⁶¹.

In conclusion, we report that Wnt5a is a novel TLR2/4 ligand and based on its opposite effects; anti-inflammatory in human vs pro-inflammatory in murine immune cells, we propose that it should be viewed as a tolerance-associated molecular pattern (TAMP) and a danger-associated molecular pattern (DAMP) in human and mouse myeloid immune cells respectively.

Paper IV

Human granulocytic myeloid-derived suppressor cells (G-MDSCs) in metastatic breast cancer patients is a heterogeneous population with tumor promoting capacity *in vivo*.

Background

Myeloid-derived suppressor cells (MDSCs) comprise a heterogeneous cell population with potent immunosuppressive properties that has been implicated in tumorigenesis. MDSCs are commonly divided into two subgroups; Mo-MDSCs and G-MDSCs. In *Paper III*, we described one potential mechanism involved in the generation of Mo-MDSCs. The origin of G-MDSCs, on the other hand, is less clear. One controversial hypothesis states that G-MDSCs are an immunosuppressive subset of neutrophils^{83,195}. In *Paper IV*, we aimed to shed new light on the origin of G-MDSCs in humans and gain better understanding on the effects G-MDSCs might exert on tumor biological function *in vivo*.

Results and Discussion

The enrichment of G-MDSCs has been observed in various types of cancers^{188,286-288}. In order to investigate the prevalence of G-MDSCs in metastatic breast cancer (MBC) patients, we performed flow cytometric analysis of cells from the Ficoll low-density mononuclear cell fraction by using antibodies specific for G-MDSCs cell surface markers. Our results showed that the levels of CD33⁺CD11b⁺CD15^{+/low}CD14^{low} G-MDSCs was significantly enriched in patients with MBC as compared to healthy blood donors. We next assessed whether the enrichment of G-MDSCs was correlated with clinicopathological features, however, we did not observe any. This is in contrast to our previously published results where we report a correlation between enrichment of Mo-MDSCs and disease progression in MBC patients⁶².

Further analysis revealed that G-MDSCs comprise a morphologically heterogeneous cell population, consisting of both blasts and polymorphonuclear (PMN) cells, where the PMN cells predominate in amount. The morphological heterogeneity has been reported previously^{179,289}, however, there is still no information regarding the functional differences or similarities between these cells. It is rather intriguing to speculate whether one type of the cells actually comprise the true G-MDSC cell population. We have tried to answer this question by performing single-cell RNA sequencing analysis on sorted G-MDSCs from MBC patients. However, we could unfortunately not harvest enough RNA to perform the analysis. Currently we are in the process of collecting samples in order to perform CyTOF mass spectrometry analysis, which hopefully will provide us with some enlightening answers.

We performed gene expression profiling analysis on G-MDSCs sorted from MBC and Gram⁺ sepsis patients as well as neutrophils and monocytes from healthy donors. Our results demonstrated that the gene expression from MBC G-MDSCs clustered with neutrophils from healthy donors, sharing similarities in gene expression relevant in angiogenesis, lymphangiogenesis and chemotaxis. Several overlaps between G-MDSCs and neutrophils have been reported. Alike G-MDSCs, neutrophils isolated on a density gradient have been found in the low-density mononuclear cell fraction^{289,290}. Moreover, an immature ring-shaped and banded morphology of neutrophils has been described as well²⁸⁹. Concerning the function, immunosuppressive features in neutrophils have been also reported^{195,291}. This supports the notion that G-MDSCs might be a subset of neutrophils. Surprisingly, gene expression profile from MBC G-MDSCs differed from G-MDSCs from Gram⁺ sepsis patients. One explanation to this discrepancy might be due to different generation mechanisms. While in the context of sepsis, G-MDSCs might be generated through stimulation by PAMPs, G-MDSCs generation in MBC patients might, on the other hand, be steered by DAMPs⁸⁴.

To broaden our knowledge about the effects that G-MDSCs exert on tumor-biological functions, we constructed a breast cancer xenograft model where we co-transplanted G-MDSCs sorted from MBC patients together with a TNBC cell line (MDA-MB-231) in immunocompromised NSG mice. Upon excising the tumors on day 21, we were struck by the apparent significant increase in tumor size in G-MDSCs/MDA-MB-231 xenografts compared to xenografts containing MDA-MB-231 cells only. As no difference in the Ki67 expression was observed, we concluded that the size difference is not a result of an increased proliferation. When staining for CD11b, CD163 and S100A9 (pan-myeloid cell, anti-inflammatory myeloid cell and MDSCs marker respectively ³⁵), we did not observe the presence of G-MDSCs, indicating that they did not survive until day 21. Despite their short lifespan, they seem to have an effect on the tumor growth early on in tumor development. This is in contrast to the transplantation survival of monocytes which survived in tumors until day 90 ³⁵. Immunohistochemical staining of the endothelial marker; CD31 and lymph vessel marker; Lyve-1, revealed an increase, although not significant, of both markers in G-MDSCs/MDA-MB-231 xenografts. These results are supported by the gene profile analysis where upregulation of genes relevant in angiogenesis and lymphangiogenesis was observed. As angiogenesis and lymphangiogenesis play essential role in tumor progression and act as source of nutrients ²⁹², the observed increase in blood vessels in G-MDSCs/MDA-MB-231 xenografts might be an explanation to the increased tumor size. Interestingly, we also observed an infiltration of murine myeloid cells (Ly6C⁺) in MDA-MB-231 xenografts while being completely absent in G-MDSCs/MDA-MB-231 xenografts, suggesting a novel G-MDSCs immunosuppressive mechanism. At this point we are however oblivious to the mechanism behind this process, and encourage further investigation in this particular field.

In conclusion, we report that G-MDSCs are enriched in MBC patients and comprise a heterogeneous cell population displaying both blast-like and PMN cell morphology. Furthermore, G-MDSCs contribute to tumorigenesis by inducing tumor growth and angiogenesis. We also report a novel G-MDSCs immunosuppression mechanism characterized by inhibition of immune cells infiltration within the tumor. Additionally, we hypothesize that G-MDSCs is an alternative immunosuppressive subset of neutrophils.

Paper V

Autoimmune regulator (AIRE) expressed in tumor associated macrophages is associated with worse prognosis in breast cancer patients.

Background

One highly critical aspect of the immune system is self-tolerance. Ideally, our immune system should recognize and eliminate pathogens and transformed cells while being tolerant to self. Autoimmune regulator (AIRE), is a transcription factor with an important role immune tolerance. Primarily, expressed by mTECs in thymus, AIRE induces transcription of a vast array of TSAs which subsequently are involved in the negative selection of T cells. The fate of T cells displaying a too high affinity for a TSA is death by apoptosis¹⁴³. AIRE expression has also been reported in peripheral organs, most notably in DCs but also in cancer. However, the role of AIRE in peripheral tolerance is to date not clearly understood, and in *Paper V*, we aimed to elaborate on the expression and prognostic value of AIRE in breast cancer patients.

Results and Discussion

In order to evaluate the expression of AIRE, we stained a TMA of 144 breast cancer patients by means of immunohistochemistry. We observed a positive AIRE expression both in the cytoplasm (AIRE_{cyt}) and the nucleus (AIRE_{nuc}) of breast cancer cells as well as in CAFs (AIRE_{CAF}) and TAMs (AIRE_{TAM}). Nuclear AIRE expression in BC cells inversely correlated with nodal stage and tumor size, while AIRE_{cyt} expression displayed an inverse correlation only with tumor size. However, we did observe a significantly positive correlation between AIRE_{cyt} and HER2⁺ BCs. It is worth mentioning that the HER2⁺ BC population in present cohort was rather small (n=12), and that the AIRE expression in HER2⁺ BCs deserves further investigation in a larger patient population. When assessing the AIRE_{TAM} expression, we observed that AIRE_{TAM} correlated with larger tumor size and NHG status. Furthermore, AIRE_{TAM} expression displayed a significant correlation with infiltrating TAMs, both in the stroma and with overall presence. AIRE_{TAM} also displayed an inverse correlation with AIRE_{nuc} and AIRE_{CAF}, shorter recurrence free survival and a borderline correlation with worse overall survival in breast cancer patients. These results indicate that AIRE expressing TAMs are most relevant regarding function in human BC. Additionally, AIRE_{TAM} expression also correlated with TAM (CD68⁺CD163⁺) infiltration in the stroma but not in the tumor nest. The

correlation was particularly apparent in the presence of TAM rich tumors. When looking into other immune cells, including CD3⁺ T cells, T_{regs}, T_h17⁺ cells, mDCs and S100A9⁺ myeloid cells, we could not observe any correlation between the above-mentioned parameters and AIRE_{TAM}. However, we noticed a borderline correlation between FoxP3⁺ cells and AIRE_{CAF}, suggesting that CAFs might play a role in the T_{regs} accumulation in BC. Next, we performed gene expression analysis of AIRE in several BCs cell lines, *in vitro* differentiated monocyte-derived mDCs, M0, M1 and M2 macrophages and human breast cancer stromal cells. *AIRE* expression in peripheral DCs, has been reported previously^{148,293} and this is further supported by our results. As expected, we observed *AIRE* expression in monocyte derived DCs (mDCs) on mRNA level. *AIRE* mRNA expression was also observed in breast cancer stromal cells and was particularly apparent in M1 and M2 macrophages. Interestingly, Kogawi *et al.* showed that CD14⁺ monocytes as well as *in vitro* differentiated mDCs expressed AIRE both on mRNA and protein level²⁹³. These results suggest that *AIRE* expression in mDCs as well as in monocyte-derived M1 and M2 might be acquired early on in their development and preserved after their differentiation. Surprisingly, we did not observe *AIRE* mRNA expression in any of the seven BC cell-lines and as such this is in conflict with our IHC results where we observed a clear AIRE protein expression. The prime function of AIRE in the thymus is to regulate the maturation of T cells by inducing expression of TSAs¹⁴³. Although the reports are quite few and inconstant, the expression of TSAs has also been observed in the periphery with the main hypothesis speculating that they are involved in the maintenance of peripheral tolerance²⁹⁴. Thus, it would be of interest to evaluate whether the AIRE expression in human BCs is functional by assessing the expression of TSAs.

In conclusion, when assessing the expression of AIRE in a cohort comprising 144 breast cancer patients, we report that AIRE is expressed in the cytoplasm and the nucleus of breast cancer cells (AIRE_{cyt}/ AIRE_{nuc}) as well as in CAFs (AIRE_{CAF}) and TAMs (AIRE_{TAM}). We also report a correlation between AIRE_{TAM} and larger tumor size and NHG status, infiltrating TAMs and shorter recurrence free survival. Furthermore, we show *AIRE* mRNA expression in monocyte-derived DCs, M1 and M2 macrophages. In the future, we are planning various functional assays to investigate potential immunosuppressive function in BC.

Conclusions

- I. TLR4 is predominantly expressed in TNBC cells and responds both to PAMPs and DAMPs by secreting inflammatory mediators. TLR4 expression is correlated with worse recurrence-free survival in ER⁻PgR⁻ breast cancer patients.
- II. S100A9 is expressed both in malignant cells and in anti-inflammatory CD163⁺ myeloid cells in human breast cancer. It induces secretion of pro-inflammatory mediators in a TLR4-dependent manner and its expression correlates with poor outcome in ER⁻PgR⁻ breast cancer patients.
- III. Wnt5a is a novel TLR2/4 ligand that acts as a tolerance-associated molecular pattern (TAMP) in human myeloid cells and damage-associated molecular pattern (DAMP) in murine myeloid cells.
- IV. G-MDSCs comprise a heterogeneous cell population enriched in metastatic breast cancer patients. G-MDSCs represent an alternative immunosuppressive subset of neutrophils that contribute to tumor progression.
- V. AIRE is expressed in malignant cells, in CAFs and in TAMs in human breast cancer, and its expression in TAMs correlates with worse outcome.

Acknowledgements

The work in this thesis was carried out at Clinical Research Center and Center for Molecular Pathology at the Department of Translational Medicine, Lund University, Malmö Sweden. The financial support was provided by Cancerfonden, Vetenskapsrådet, Allmänna sjukhuset i Malmö stiftelse för bekämpande av cancer, Gyllenstiernska Krapperupps foundation, Gunnar Nillson's Cancer Foundation, Åke Wibergs foundation, Alfred Österlunds foundation and Percy Falks foundation.

First and foremost, I would like to thank all funders for their financial support and all patients who, despite their difficult circumstances, chose to participate in our studies. Without you, the work in this thesis would not have been possible.

Next, I would like to take this opportunity and express my utmost gratitude to my supervisor, **Karin Leandersson**. Thank you for this journey, and oh, what a journey this was. As supervisor, you were a rock. As scientist, an inspiration, and as a person, truly admirable. I look up to you, and among my friends and colleagues, you will always be known as the coolest supervisor ever.

I also would like to thank my co-supervisors; Camilla Rydberg Millrud, Caroline Bergenfelz and Martin Johansson. **Camilla**, we started off as colleagues and you exemplified the beauty of a good one. With time, we reached beyond that and now I am happy to call you my friend. Thank you for all your help, all great talks and for always peppering me with the words "you know more than you think". **Carro**, you introduced me to the methods and lab techniques, and your dedicated way of working has always been a source of inspiration. Thank you for all the scientific input and for always having an open door when there was something I needed to ask, even if it meant calling you late evenings when the Calibur just refused to work. **Martin Johansson**, thank you for the invaluable histology sessions. With your help, I managed to navigate through the jungle of TMA scoring.

To past and present members of the Cancer Immunology group. **Roni**, my first office mate. For all the discussions about work, training and life in general and for being the best travel buddy during conferences, thank you! **Frida**, you came, and your personality knocked us all out. Thank you for all your help (especially with my thesis), the hugs in moments of despair, your knitting-lessons, for organizing the lab so nicely (!) and for enlightening me about the danger with cave diving! **Eva Källberg**, I preferred having a quick lunch in front of my computer, but since you

started here, that has changed. Now, I very much enjoy our conversations in the lunch room. All students along the way, **Eva, Marie, Robert** and especially **Marcus**.

To all the past and present members of CMP: **Lisa, Nick, Agi, Helén, Tamae, Ben, Totte, Johan, Susan, Anna, Rebecka, Giacomo, Kiki, Rebecca, Maite, Anders, Darina, Mathieu, Macarena, Greta, Giuseppe, Aseem, Elise** and **Elisabeth**. Thank you for all your help and for always being willing to help, even after all these years after our move. Thank you for all the fun moments we shared together, all the “fikas”, the spring excursions and the retreats. You are a fantastic group of people, and I feel so lucky to have had such wonderful colleagues to share my PhD journey with. I would especially like to thank my friend **Krzysiek**. For all your help and support, for all talks and laughter and for creating such a positive atmosphere wherever you are, Thank You!

To all the past and present members of CRC: **Mansi, Farnaz, Lena, Lisa, Elin, Diana, Puru, Zdenka, Syrina, Janina, Geriolda, Anita, Shakti, Qing, Puja, Gunilla** and **Tommy**. Thank you for all the lovely “fikas” and for all your help and support upon our move to CRC. I also would like to thank **Njainday**, for being a good friend and for making me feel closer to home.

To my dear friends, my constant support and source of joy, without whom my life would have been so boring! **Anna**, for being such a beautiful soul and for inspiring me about minimalism and what’s important in life. **Una**, there are few out there as fierce and hard-working as you, and I aspire to be more like you! **Negin**, my bohemian friend, with your special spice in life that is love, you remind us every day to live outside the box. My oldest friend **Iva**, I don’t see you often, but whenever I do, it feels like we were never apart. You will always be my “Doog”! To my dearest friend **Noémie**, I feel so blessed having you as my friend and one of the perks of my move will be moving closer to you!

To my childhood friends **Jasna, Goga, Azze** and **Hila**. What incredible women you’ve all become! I often think of you.

To my dear sister, **Minka**, for teaching me that no matter how hard life can get, there’s always a way to overcome it. My nieces, **Ena** and **Emma**, you have brought so much joy into my life. My baby brother, **Melko**, who’s not so little anymore. You were the greatest gift my parents gave me. You are so wise and with such an admirable approach in life. I love you immensely!

Mojoj proširenoj porodici u Njemačkoj, hvala vam na ljubaznim rečima i podršci.

Nana, babo i **majka** hvala vam za svu vašu ljubav i što ste uvek bili ponosni na mene.

Želim da se zahvalim mojoj **majci** i mom **ocu** na beskrajnoj ljubavi i neizmernoj podršci. Hvala vam sto ste se uvek trudili da mi pružite sve i uvek me ohrabrivali da ostvarim svoje snove. Volim vas! Posebno se zahvaljujem mojoj majci, čija me snaga uvek inspirirala i ojačala. Ovu tezu posvećujem tebi.

Last but not least, I want to thank my special person in this world, my husband **Nudzeim**. Being so far away from each other and lately with my heavily preoccupied mind, it has not been easy for you. Nonetheless, your positivity and patience deserve all praise. Thank you for all your love and support and for showing me how beautifully easy life can be. You are my best friend, my confidant and I love you beyond measure ♥.

References

- 1 Bray, F. *et al.* Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*, doi:10.3322/caac.21492 (2018).
- 2 Sudhakar, A. History of Cancer, Ancient and Modern Treatment Methods. *J Cancer Sci Ther* **1**, 1-4, doi:10.4172/1948-5956.100000e2 (2009).
- 3 Hanahan, D. & Weinberg, R. A. The hallmarks of cancer. *Cell* **100**, 57-70 (2000).
- 4 Hanahan, D. & Weinberg, R. A. Hallmarks of cancer: the next generation. *Cell* **144**, 646-674, doi:10.1016/j.cell.2011.02.013 (2011).
- 5 Wang, M. *et al.* Role of tumor microenvironment in tumorigenesis. *J Cancer* **8**, 761-773, doi:10.7150/jca.17648 (2017).
- 6 Cancerfonden. Cancerfondsrapporten 2018. (2018).
- 7 Bray, F. *et al.* Global Cancer Statistics 2018: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J Clin*, 1-31, doi:10.3322/caac.21492 (2018).
- 8 Socialstyrelsen. Cancer i siffror 2018- Populärvetenskapliga fakta om cancer. (2018).
- 9 Hulkkaa, B. S. & Moormanb, P. G. Reprint of breast cancer: hormones and other risk factors. *Maturitas* **61**, 203-213, doi:10.1016/j.maturitas.2008.11.016 (2008).
- 10 Sheikh, A. *et al.* The spectrum of genetic mutations in breast cancer. *Asian Pac J Cancer Prev* **16**, 2177-2185 (2015).
- 11 Shuen, A. Y. & Foulkes, W. D. Inherited mutations in breast cancer genes--risk and response. *J Mammary Gland Biol Neoplasia* **16**, 3-15, doi:10.1007/s10911-011-9213-5 (2011).
- 12 Turnbull, C. & Rahman, N. Genetic predisposition to breast cancer: past, present, and future. *Annu Rev Genomics Hum Genet* **9**, 321-345, doi:10.1146/annurev.genom.9.081307.164339 (2008).
- 13 Lanigan, F., O'Connor, D., Martin, F. & Gallagher, W. M. Molecular links between mammary gland development and breast cancer. *Cell Mol Life Sci* **64**, 3159-3184, doi:10.1007/s00018-007-7386-2 (2007).
- 14 Javed, A. & Lteif, A. Development of the human breast. *Semin Plast Surg* **27**, 5-12, doi:10.1055/s-0033-1343989 (2013).
- 15 Burstein, H. J., Polyak, K., Wong, J. S., Lester, S. C. & Kaelin, C. M. Ductal carcinoma in situ of the breast. *N Engl J Med* **350**, 1430-1441, doi:10.1056/NEJMra031301 (2004).
- 16 Bombonati, A. & Sgroi, D. C. The molecular pathology of breast cancer progression. *J Pathol* **223**, 307-317, doi:10.1002/path.2808 (2011).
- 17 McGee, S. F., Lanigan, F., Gilligan, E. & Groner, B. Mammary gland biology and breast cancer. Conference on Common Molecular Mechanisms of Mammary Gland Development and Breast Cancer Progression. *EMBO Rep* **7**, 1084-1088, doi:10.1038/sj.embor.7400839 (2006).
- 18 Weigelt, B., Geyerb, F. C. & Reis-Filhob, J. S. Histological types of breast cancer: How special are they? *Mol Oncol* **4**, 192-208, doi:10.1016/j.molonc.2010.04.004 (2010).
- 19 Malhotra, G. K., Zhao, X., Band, H. & Band, V. Histological, molecular and functional subtypes of breast cancers. *Cancer Biol Ther* **10**, 955-960 (2010).

- 20 Elston, C. W. & Ellis, I. O. Pathological prognostic factors in breast cancer. I. The value of
histological grade in breast cancer: experience from a large study with long-term follow-up.
Histopathology **19**, 403-410 (1991).
- 21 SweBCG. Bröstcancer- Nationellt vårdprogram v. 2.0. (2018).
- 22 Sorlie, T. *et al.* Gene expression patterns of breast carcinomas distinguish tumor subclasses
with clinical implications. *Proc Natl Acad Sci U S A* **98**, 10869-10874,
doi:10.1073/pnas.191367098 (2001).
- 23 Prat, A. & Perou, C. M. Deconstructing the molecular portraits of breast cancer. *Mol Oncol*
5, 5-23, doi:10.1016/j.molonc.2010.11.003 (2011).
- 24 Bertucci, F. *et al.* How basal are triple-negative breast cancers? *Int J Cancer* **123**, 236-240,
doi:10.1002/ijc.23518 (2008).
- 25 Socialstyrelsen. Screening för bröstcancer- Rekommendation och bedömningsunderlag.
(2014).
- 26 Anjum, F., Razvi, N. & Masood, M. A. Breast cancer therapy: a mini review. *MOJ Drug
Des Develop Ther* **1**, 35-38, doi: 10.15406/mojddt.2017.01.00006 (2017).
- 27 Chumsri, S., Howes, T., Bao, T., Sabnis, G. & Brodie, A. Aromatase, aromatase inhibitors,
and breast cancer. *J Steroid Biochem Mol Biol* **125**, 13-22, doi:10.1016/j.jsbmb.2011.02.001
(2011).
- 28 Wahba, H. A. & El-Hadaad, H. A. Current approaches in treatment of triple-negative breast
cancer. *Cancer Biol Med* **12**, 106-116, doi:10.7497/j.issn.2095-3941.2015.0030 (2015).
- 29 Administration., U. S. F. D. (2018).
- 30 Peshkin, B. N., Alabek, M. L. & Isaacs, C. BRCA1/2 mutations and triple negative breast
cancers. *Breast Dis* **32**, 25-33, doi:10.3233/BD-2010-0306 (2010).
- 31 Schmid, P. *et al.* Atezolizumab and Nab-Paclitaxel in Advanced Triple-Negative Breast
Cancer. *N Engl J Med* **379**, 2108-2121, doi:10.1056/NEJMoa1809615 (2018).
- 32 Pietras, K. & Ostman, A. Hallmarks of cancer: interactions with the tumor stroma. *Exp Cell
Res* **316**, 1324-1331, doi:10.1016/j.yexcr.2010.02.045 (2010).
- 33 Kalluri, R. & Zeisberg, M. Fibroblasts in cancer. *Nat Rev Cancer* **6**, 392-401,
doi:10.1038/nrc1877 (2006).
- 34 Bhowmick, N. A., Neilson, E. G. & Moses, H. L. Stromal fibroblasts in cancer initiation and
progression. *Nature* **432**, 332-337, doi:10.1038/nature03096 (2004).
- 35 Allaoui, R. *et al.* Cancer-associated fibroblast-secreted CXCL16 attracts monocytes to
promote stroma activation in triple-negative breast cancers. *Nat Commun* **7**, 13050,
doi:10.1038/ncomms13050 (2016).
- 36 Nishida N, Y. H., Nishida T, Kamura T, Kojiro M. . Angiogenesis in cancer. *Vascular Health
and Risk Management* **2**, 213-219. (2006).
- 37 Forster, J. C., Harriss-Phillips, W. M., Douglass, M. J. & Bezak, E. A review of the
development of tumor vasculature and its effects on the tumor microenvironment. *Hypoxia
(Auckl)* **5**, 21-32, doi:10.2147/HP.S133231 (2017).
- 38 Bergers, G. & Song, S. The role of pericytes in blood-vessel formation and maintenance.
Neuro Oncol **7**, 452-464, doi:10.1215/S1152851705000232 (2005).
- 39 Ribeiro, A. L. & Okamoto, O. K. Combined effects of pericytes in the tumor
microenvironment. *Stem Cells Int* **2015**, 868475, doi:10.1155/2015/868475 (2015).
- 40 Furuhashi, M. *et al.* Platelet-derived growth factor production by B16 melanoma cells leads
to increased pericyte abundance in tumors and an associated increase in tumor growth rate.
Cancer Res **64**, 2725-2733 (2004).
- 41 Cooke, V. G. *et al.* Pericyte depletion results in hypoxia-associated epithelial-to-
mesenchymal transition and metastasis mediated by met signaling pathway. *Cancer Cell* **21**,
66-81, doi:10.1016/j.ccr.2011.11.024 (2012).
- 42 Walker, C., Mojares, E. & Del Rio Hernandez, A. Role of Extracellular Matrix in
Development and Cancer Progression. *Int J Mol Sci* **19**, doi:10.3390/ijms19103028 (2018).
- 43 Levental, K. R. *et al.* Matrix crosslinking forces tumor progression by enhancing integrin
signaling. *Cell* **139**, 891-906, doi:10.1016/j.cell.2009.10.027 (2009).

- 44 Martin, L. J. & Boyd, N. F. Mammographic density. Potential mechanisms of breast cancer risk associated with mammographic density: hypotheses based on epidemiological evidence. *Breast Cancer Res* **10**, 201, doi:10.1186/bcr1831 (2008).
- 45 Egeblad, M. & Werb, Z. New functions for the matrix metalloproteinases in cancer progression. *Nat Rev Cancer* **2**, 161-174, doi:10.1038/nrc745 (2002).
- 46 Kasurinen, A. *et al.* High serum MMP-14 predicts worse survival in gastric cancer. *PLoS One* **13**, e0208800, doi:10.1371/journal.pone.0208800 (2018).
- 47 Bockelman, C. *et al.* Serum MMP-8 and TIMP-1 predict prognosis in colorectal cancer. *BMC Cancer* **18**, 679, doi:10.1186/s12885-018-4589-x (2018).
- 48 Chaplin, D. D. Overview of the immune response. *J Allergy Clin Immunol* **125**, S3-23, doi:10.1016/j.jaci.2009.12.980 (2010).
- 49 Suresh, R. & Mosser, D. M. Pattern recognition receptors in innate immunity, host defense, and immunopathology. *Adv Physiol Educ* **37**, 284-291, doi:10.1152/advan.00058.2013 (2013).
- 50 Luster, A. D. The role of chemokines in linking innate and adaptive immunity. *Curr Opin Immunol* **14**, 129-135 (2002).
- 51 Dempsey, P. W., Vaidya, S. A. & Cheng, G. The art of war: Innate and adaptive immune responses. *Cell Mol Life Sci* **60**, 2604-2621, doi:10.1007/s00018-003-3180-y (2003).
- 52 Koenderman, L., Buurman, W. & Daha, M. R. The innate immune response. *Immunol Lett* **162**, 95-102, doi:10.1016/j.imlet.2014.10.010 (2014).
- 53 Gordon, S. & Taylor, P. R. Monocyte and macrophage heterogeneity. *Nat Rev Immunol* **5**, 953-964, doi:10.1038/nri1733 (2005).
- 54 Auffray, C., Sieweke, M. H. & Geissmann, F. Blood monocytes: development, heterogeneity, and relationship with dendritic cells. *Annu Rev Immunol* **27**, 669-692, doi:10.1146/annurev.immunol.021908.132557 (2009).
- 55 Ziegler-Heitbrock, L. *et al.* Nomenclature of monocytes and dendritic cells in blood. *Blood* **116**, e74-80, doi:10.1182/blood-2010-02-258558 (2010).
- 56 Fingerle, G. *et al.* The novel subset of CD14⁺/CD16⁺ blood monocytes is expanded in sepsis patients. *Blood* **82**, 3170-3176 (1993).
- 57 Nockher, W. A. & Scherberich, J. E. Expanded CD14⁺ CD16⁺ monocyte subpopulation in patients with acute and chronic infections undergoing hemodialysis. *Infect Immun* **66**, 2782-2790 (1998).
- 58 Qian, B. Z. *et al.* CCL2 recruits inflammatory monocytes to facilitate breast-tumour metastasis. *Nature* **475**, 222-225, doi:10.1038/nature10138 (2011).
- 59 Chittezhath, M. *et al.* Molecular profiling reveals a tumor-promoting phenotype of monocytes and macrophages in human cancer progression. *Immunity* **41**, 815-829, doi:10.1016/j.immuni.2014.09.014 (2014).
- 60 Szabo, K. A. & Singh, G. Modulation of monocyte matrix metalloproteinase-2 by breast adenocarcinoma cells. *Breast Cancer Res* **7**, R661-668, doi:10.1186/bcr1261 (2005).
- 61 Mohamed, M. M., Cavallo-Medved, D. & Sloane, B. F. Human monocytes augment invasiveness and proteolytic activity of inflammatory breast cancer. *Biol Chem* **389**, 1117-1121, doi:10.1515/BC.2008.117 (2008).
- 62 Bergenfelz, C. *et al.* Systemic Monocytic-MDSCs Are Generated from Monocytes and Correlate with Disease Progression in Breast Cancer Patients. *PLoS One* **10**, e0127028, doi:10.1371/journal.pone.0127028 (2015).
- 63 Gordon, S. The macrophage: past, present and future. *Eur J Immunol* **37 Suppl 1**, S9-17, doi:10.1002/eji.200737638 (2007).
- 64 Halder, M. & Murphy, K. M. Origin, development, and homeostasis of tissue-resident macrophages. *Immunol Rev* **262**, 25-35, doi:10.1111/imr.12215 (2014).
- 65 Guha, I., Naskar, D. & Sen, M. Macrophage as a mediator of immune response: Sustenance of immune homeostasis. *Macrophage* **2**, doi:10.14800/Macrophage.709. (2015).
- 66 Sica, A. & Mantovani, A. Macrophage plasticity and polarization: in vivo veritas. *J Clin Invest* **122**, 787-795, doi:10.1172/JCI59643 (2012).
- 67 Qian, B. Z. & Pollard, J. W. Macrophage diversity enhances tumor progression and metastasis. *Cell* **141**, 39-51, doi:10.1016/j.cell.2010.03.014 (2010).

- 68 Mantovani, A., Sica, A., Allavena, P., Garlanda, C. & Locati, M. Tumor-associated macrophages and the related myeloid-derived suppressor cells as a paradigm of the diversity of macrophage activation. *Hum Immunol* **70**, 325-330, doi:10.1016/j.humimm.2009.02.008 (2009).
- 69 Baskic, D., Acimovic, L., Samardzic, G., Vujanovic, N. L. & Arsenijevic, N. N. Blood monocytes and tumor-associated macrophages in human cancer: differences in activation levels. *Neoplasma* **48**, 169-174 (2001).
- 70 Medrek, C., Ponten, F., Jirstrom, K. & Leanderson, K. The presence of tumor associated macrophages in tumor stroma as a prognostic marker for breast cancer patients. *BMC Cancer* **12**, 306, doi:10.1186/1471-2407-12-306 (2012).
- 71 Steidl, C. *et al.* Tumor-associated macrophages and survival in classic Hodgkin's lymphoma. *N Engl J Med* **362**, 875-885, doi:10.1056/NEJMoa0905680 (2010).
- 72 Hanada, T. *et al.* Prognostic value of tumor-associated macrophage count in human bladder cancer. *Int J Urol* **7**, 263-269 (2000).
- 73 Kolaczowska, E. & Kubes, P. Neutrophil recruitment and function in health and inflammation. *Nat Rev Immunol* **13**, 159-175, doi:10.1038/nri3399 (2013).
- 74 Kruger, P. *et al.* Neutrophils: Between host defence, immune modulation, and tissue injury. *PLoS Pathog* **11**, e1004651, doi:10.1371/journal.ppat.1004651 (2015).
- 75 Jaillon, S. *et al.* Neutrophils in innate and adaptive immunity. *Semin Immunopathol* **35**, 377-394, doi:10.1007/s00281-013-0374-8 (2013).
- 76 Rosales, C. Neutrophil: A Cell with Many Roles in Inflammation or Several Cell Types? *Front Physiol* **9**, 113, doi:10.3389/fphys.2018.00113 (2018).
- 77 Deniset, J. F. & Kubes, P. Recent advances in understanding neutrophils. *F1000Res* **5**, 2912, doi:10.12688/f1000research.9691.1 (2016).
- 78 Schmidt, H. *et al.* Elevated neutrophil and monocyte counts in peripheral blood are associated with poor survival in patients with metastatic melanoma: a prognostic model. *Br J Cancer* **93**, 273-278, doi:10.1038/sj.bjc.6602702 (2005).
- 79 Peng, B., Wang, Y. H., Liu, Y. M. & Ma, L. X. Prognostic significance of the neutrophil to lymphocyte ratio in patients with non-small cell lung cancer: a systemic review and meta-analysis. *Int J Clin Exp Med* **8**, 3098-3106 (2015).
- 80 Wislez, M. *et al.* Hepatocyte growth factor production by neutrophils infiltrating bronchioloalveolar subtype pulmonary adenocarcinoma: role in tumor progression and death. *Cancer Res* **63**, 1405-1412 (2003).
- 81 Yan, J. *et al.* Human polymorphonuclear neutrophils specifically recognize and kill cancerous cells. *Oncoimmunology* **3**, e950163, doi:10.4161/15384101.2014.950163 (2014).
- 82 Caruso, R. A. *et al.* Prognostic value of intratumoral neutrophils in advanced gastric carcinoma in a high-risk area in northern Italy. *Mod Pathol* **15**, 831-837, doi:10.1097/01.MP.0000020391.98998.6B (2002).
- 83 Pillay, J., Tak, T., Kamp, V. M. & Koenderman, L. Immune suppression by neutrophils and granulocytic myeloid-derived suppressor cells: similarities and differences. *Cell Mol Life Sci* **70**, 3813-3827, doi:10.1007/s00018-013-1286-4 (2013).
- 84 Millrud, C. R., Bergenfelz, C. & Leanderson, K. On the origin of myeloid-derived suppressor cells. *Oncotarget* **8**, 3649-3665, doi:10.18632/oncotarget.12278 (2017).
- 85 Kawai, T. & Akira, S. The role of pattern-recognition receptors in innate immunity: update on Toll-like receptors. *Nat Immunol* **11**, 373-384, doi:10.1038/ni.1863 (2010).
- 86 Takeuchi, O. & Akira, S. Pattern recognition receptors and inflammation. *Cell* **140**, 805-820, doi:10.1016/j.cell.2010.01.022 (2010).
- 87 Visintin, A. *et al.* Regulation of Toll-Like Receptors in Human Monocytes and Dendritic Cells. *The Journal of Immunology* **166**, 249-255, doi:<https://doi.org/10.4049/jimmunol.166.1.249> (2001).
- 88 Prince, L. R., Whyte, M. K., Sabroe, I. & Parker, L. C. The role of TLRs in neutrophil activation. *Current Opinion in Pharmacology* **11**, 397-403, doi:10.1016/j.coph.2011.06.007 (2011).
- 89 Greene, C. M. & McElvaney, N. G. Toll-like receptor expression and function in airway epithelial cells. *Arch Immunol Ther Exp* **53**, 418-427 (2005).

- 90 Abreu, M. T. Toll-like receptor signalling in the intestinal epithelium: how bacterial recognition shapes intestinal function. *Nat Rev Immunol* **10**, 131-144, doi:10.1038/nri2707. (2010).
- 91 Lim, K. H. & Staudt, L. M. Toll-like receptor signaling. *Cold Spring Harb Perspect Biol* **5**, a011247, doi:10.1101/cshperspect.a011247 (2013).
- 92 Sheyhidin, I. *et al.* Overexpression of TLR3, TLR4, TLR7 and TLR9 in esophageal squamous cell carcinoma. *World J Gastroenterol* **17**, 3745-3751, doi:10.3748/wjg.v17.i32.3745 (2011).
- 93 Hassan, F. *et al.* Intracellular expression of toll-like receptor 4 in neuroblastoma cells and their unresponsiveness to lipopolysaccharide. *BMC Cancer* **6**, 281, doi:10.1186/1471-2407-6-281 (2006).
- 94 Gonzalez-Reyes, S. *et al.* Study of TLR3, TLR4 and TLR9 in breast carcinomas and their association with metastasis. *BMC Cancer* **10**, 665, doi:10.1186/1471-2407-10-665 (2010).
- 95 Eiro, N. *et al.* Expression of TLR3, 4, 7 and 9 in cutaneous malignant melanoma: relationship with clinicopathological characteristics and prognosis. *Arch Dermatol Res* **305**, 59-67, doi:10.1007/s00403-012-1300-y (2013).
- 96 Ronkainen, H. *et al.* Absent Toll-like receptor-9 expression predicts poor prognosis in renal cell carcinoma. *J Exp Clin Cancer Res* **30**, 84, doi:10.1186/1756-9966-30-84 (2011).
- 97 Cai, Z. *et al.* Activation of Toll-like receptor 5 on breast cancer cells by flagellin suppresses cell proliferation and tumor growth. *Cancer Res* **71**, 2466-2475, doi:10.1158/0008-5472.CAN-10-1993 (2011).
- 98 Yang, H. *et al.* Reduced expression of Toll-like receptor 4 inhibits human breast cancer cells proliferation and inflammatory cytokines secretion. *J Exp Clin Cancer Res* **29**, 92, doi:10.1186/1756-9966-29-92 (2010).
- 99 Lowe, E. L. *et al.* Toll-like receptor 2 signaling protects mice from tumor development in a mouse model of colitis-induced cancer. *PLoS One* **5**, e13027, doi:10.1371/journal.pone.0013027 (2010).
- 100 Gomes, S. S., Decout, A. & Nigou, J. Pathogen-Associated Molecular Patterns (PAMPs). *Encyclopedia of Inflammatory Diseases*, ed. M. Parnham, Birkhäuser, Basel, 1055-1069, doi: 10.1007/978-3-0348-0620-6_35-1 (2014).
- 101 Akira, S. & Takeda, K. Toll-like receptor signalling. *Nat Rev Immunol* **4**, 499-511, doi:10.1038/nri1391 (2004).
- 102 Kawasaki, T. & Kawai, T. Toll-like receptor signaling pathways. *Front Immunol* **5**, 461, doi:10.3389/fimmu.2014.00461 (2014).
- 103 Kagan, J. C. *et al.* TRAM couples endocytosis of Toll-like receptor 4 to the induction of interferon-beta. *Nat Immunol* **9**, 361-368, doi:10.1038/ni1569 (2008).
- 104 Kuai, R. *et al.* Dual TLR agonist nanodiscs as a strong adjuvant system for vaccines and immunotherapy. *J Control Release* **282**, 131-139, doi:10.1016/j.jconrel.2018.04.041 (2018).
- 105 Steinhagen, F., Kinjo, T., Bode, C. & Klinman, D. M. TLR-based immune adjuvants. *Vaccine* **29**, 3341-3355, doi:10.1016/j.vaccine.2010.08.002 (2011).
- 106 Hopton Cann, S. A., van Netten, J. P. & van Netten, C. Dr William Coley and tumour regression: a place in history or in the future. *Postgrad Med J* **79**, 672-680 (2003).
- 107 Chen, G. Y. & Nunez, G. Sterile inflammation: sensing and reacting to damage. *Nat Rev Immunol* **10**, 826-837, doi:10.1038/nri2873 (2010).
- 108 Srikrishna, G. & Freeze, H. H. Endogenous damage-associated molecular pattern molecules at the crossroads of inflammation and cancer. *Neoplasia* **11**, 615-628 (2009).
- 109 Bianchi, M. E. DAMPs, PAMPs and alarmins: all we need to know about danger. *J Leukoc Biol* **81**, 1-5, doi:10.1189/jlb.0306164 (2007).
- 110 Piccinini, A. M. & Midwood, K. S. DAMPening inflammation by modulating TLR signalling. *Mediators Inflamm* **2010**, doi:10.1155/2010/672395 (2010).
- 111 Hernandez, C., Huebener, P. & Schwabe, R. F. Damage-associated molecular patterns in cancer: a double-edged sword. *Oncogene* **35**, 5931-5941, doi:10.1038/onc.2016.104 (2016).
- 112 Kallberg, E. *et al.* S100A9 interaction with TLR4 promotes tumor growth. *PLoS One* **7**, e34207, doi:10.1371/journal.pone.0034207 (2012).

- 113 Markowitz, J. & Carson, W. E., 3rd. Review of S100A9 biology and its role in cancer. *Biochim Biophys Acta* **1835**, 100-109, doi:10.1016/j.bbcan.2012.10.003 (2013).
- 114 Bresnick, A. R., Weber, D. J. & Zimmer, D. B. S100 proteins in cancer. *Nat Rev Cancer* **15**, 96-109, doi:10.1038/nrc3893 (2015).
- 115 Zwadlo, G., Bruggen, J., Gerhards, G., Schlegel, R. & Sorg, C. Two calcium-binding proteins associated with specific stages of myeloid cell differentiation are expressed by subsets of macrophages in inflammatory tissues. *Clin Exp Immunol* **72**, 510-515 (1988).
- 116 Roth, J., Vogl, T., Sorg, C. & Sunderkotter, C. Phagocyte-specific S100 proteins: a novel group of proinflammatory molecules. *Trends Immunol* **24**, 155-158 (2003).
- 117 Sunahori, K. *et al.* The S100A8/A9 heterodimer amplifies proinflammatory cytokine production by macrophages via activation of nuclear factor kappa B and p38 mitogen-activated protein kinase in rheumatoid arthritis. *Arthritis Res Ther* **8**, R69, doi:10.1186/ar1939 (2006).
- 118 Heizmann, C. W., Fritz, G. & Schafer, B. W. S100 proteins: structure, functions and pathology. *Front Biosci* **7**, d1356-1368 (2002).
- 119 Arai, K., Yamada, T. & Nozawa, R. Immunohistochemical investigation of migration inhibitory factor-related protein (MRP)-14 expression in hepatocellular carcinoma. *Med Oncol* **17**, 183-188 (2000).
- 120 Bergenfelz, C. *et al.* S100A9 expressed in ER(-)PgR(-) breast cancers induces inflammatory cytokines and is associated with an impaired overall survival. *Br J Cancer* **113**, 1234-1243, doi:10.1038/bjc.2015.346 (2015).
- 121 Kim, H. J. *et al.* Identification of S100A8 and S100A9 as serological markers for colorectal cancer. *J Proteome Res* **8**, 1368-1379, doi:10.1021/pr8007573 (2009).
- 122 Hermani, A. *et al.* Calcium-binding proteins S100A8 and S100A9 as novel diagnostic markers in human prostate cancer. *Clin Cancer Res* **11**, 5146-5152, doi:10.1158/1078-0432.CCR-05-0352 (2005).
- 123 Tidehag, V. *et al.* High density of S100A9 positive inflammatory cells in prostate cancer stroma is associated with poor outcome. *Eur J Cancer* **50**, 1829-1835, doi:10.1016/j.ejca.2014.03.278 (2014).
- 124 Sinha, P. *et al.* Proinflammatory S100 proteins regulate the accumulation of myeloid-derived suppressor cells. *J Immunol* **181**, 4666-4675 (2008).
- 125 Laouedj, M. *et al.* S100A9 induces differentiation of acute myeloid leukemia cells through TLR4. *Blood* **129**, 1980-1990, doi:10.1182/blood-2016-09-738005 (2017).
- 126 Bonilla, F. A. & Oettgen, H. C. Adaptive immunity. *The Journal of allergy and clinical immunology* **125**, 33-40, doi: <https://doi.org/10.1016/j.jaci.2009.09.017> (2010).
- 127 Ollila, J. & Vihinen, M. B cells. *Int J Biochem Cell Biol* **37**, 518-523, doi:10.1016/j.biocel.2004.09.007 (2005).
- 128 Teillaud, J. L. Antibody-dependent Cellular Cytotoxicity (ADCC). *eLS. John Wiley & Sons*, doi:10.1002/9780470015902.a0000498.pub2 (2012).
- 129 Silva-Santos, B., Serre, K. & Norell, H. gammadelta T cells in cancer. *Nat Rev Immunol* **15**, 683-691, doi:10.1038/nri3904 (2015).
- 130 Vivier, E., Ugolini, S., Blaise, D., Chabannon, C. & Brossay, L. Targeting natural killer cells and natural killer T cells in cancer. *Nat Rev Immunol* **12**, 239-252, doi:10.1038/nri3174 (2012).
- 131 Zou, C. *et al.* gammadelta T cells in cancer immunotherapy. *Oncotarget* **8**, 8900-8909, doi:10.18632/oncotarget.13051 (2017).
- 132 Ma, C. *et al.* Tumor-infiltrating gammadelta T lymphocytes predict clinical outcome in human breast cancer. *J Immunol* **189**, 5029-5036, doi:10.4049/jimmunol.1201892 (2012).
- 133 Terabe, M. & Berzofsky, J. A. The role of NKT cells in tumor immunity. *Adv Cancer Res* **101**, 277-348, doi:10.1016/S0065-230X(08)00408-9 (2008).
- 134 Bluestone, J. A. Mechanisms of tolerance. *Immunol Rev* **241**, 5-19, doi:10.1111/j.1600-065X.2011.01019.x (2011).
- 135 Sprent, J. & Kishimoto, H. The thymus and central tolerance. *Philos Trans R Soc Lond B Biol Sci* **356**, 609-616, doi:10.1098/rstb.2001.0846 (2001).

- 136 Oh, J. & Shin, J. S. The Role of Dendritic Cells in Central Tolerance. *Immune Netw* **15**, 111-120, doi:10.4110/in.2015.15.3.111 (2015).
- 137 Mueller, D. L. Mechanisms maintaining peripheral tolerance. *Nat Immunol* **11**, 21-27, doi:10.1038/ni.1817 (2010).
- 138 Nakanishi, J. *et al.* Overexpression of B7-H1 (PD-L1) significantly associates with tumor grade and postoperative prognosis in human urothelial cancers. *Cancer Immunol Immunother* **56**, 1173-1182, doi:10.1007/s00262-006-0266-z (2007).
- 139 Shankaran, V. *et al.* IFN γ and lymphocytes prevent primary tumour development and shape tumour immunogenicity. *Nature* **410**, 1107-1111, doi:10.1038/35074122 (2001).
- 140 Wu, J., Xie, A. & Chen, W. Cytokine regulation of immune tolerance. *Burns Trauma* **2**, 11-17, doi:10.4103/2321-3868.124771 (2014).
- 141 Wing, K. & Sakaguchi, S. Regulatory T cells exert checks and balances on self tolerance and autoimmunity. *Nat Immunol* **11**, 7-13, doi:10.1038/ni.1818 (2010).
- 142 Gabrilovich, D. I. & Nagaraj, S. Myeloid-derived suppressor cells as regulators of the immune system. *Nat Rev Immunol* **9**, 162-174, doi:10.1038/nri2506 (2009).
- 143 Mathis, D. & Benoist, C. Aire. *Annu Rev Immunol* **27**, 287-312, doi:10.1146/annurev.immunol.25.022106.141532 (2009).
- 144 Ruan, Q. G. *et al.* The autoimmune regulator directly controls the expression of genes critical for thymic epithelial function. *J Immunol* **178**, 7173-7180 (2007).
- 145 Kyewski, B., Derbinski, J., Gotter, J. & Klein, L. Promiscuous gene expression and central T-cell tolerance: more than meets the eye. *Trends Immunol* **23**, 364-371 (2002).
- 146 Palmer, E. Negative selection- clearing out the bad apples from the T-cell repertoire. *Nat Rev Immunol* **3**, 383-391, doi:10.1038/nri1085 (2003).
- 147 Klein, L., Kyewski, B., Allen, P. M. & Hogquist, K. A. Positive and negative selection of the T cell repertoire: what thymocytes see and don't see. *Nat Rev Immunol*. **14**, 377-391, doi:10.1038/nri3667. (2014).
- 148 Poliani, P. L. *et al.* Human peripheral lymphoid tissues contain autoimmune regulator-expressing dendritic cells. *Am J Pathol* **176**, 1104-1112, doi:10.2353/ajpath.2010.090956 (2010).
- 149 Li, D. *et al.* Aire-Overexpressing Dendritic Cells Induce Peripheral CD4+ T Cell Tolerance. *Int. J. Mol. Sci.* **17**, doi:doi:10.3390/ijms17010038 (2016).
- 150 Malchow S, L. D., Lee V, Nishi S, Socci ND, Savage PA. Aire enforces immune tolerance by directing autoreactive T cells into the regulatory T cell lineage. *Immunity*. **44**, 1102-1113, doi:0.1016/j.immuni.2016.02.009 (2016).
- 151 Sillanpaa, N. *et al.* Autoimmune regulator induced changes in the gene expression profile of human monocyte-dendritic cell-lineage. *Mol Immunol* **41**, 1185-1198, doi:10.1016/j.molimm.2004.06.004 (2004).
- 152 Bianchi, B. *et al.* Expression and prognostic significance of the autoimmune regulator gene in breast cancer cells. *Cell Cycle* **15**, 3220-3229., doi:10.1080/15384101.2016.1241918 (2016).
- 153 Asimakopoulos, G. Mechanisms of the systemic inflammatory response. *Perfusion* **14**, 269-277, doi:10.1177/026765919901400406 (1999).
- 154 Jean-Baptiste, E. Cellular mechanisms in sepsis. *J Intensive Care Med* **22**, 63-72, doi:10.1177/0885066606297123 (2007).
- 155 Bone, R. C., Grodzin, C. J. & Balk, R. A. Sepsis: a new hypothesis for pathogenesis of the disease process. *Chest* **112**, 235-243 (1997).
- 156 Hotchkiss, R. S., Monneret, G. & Payen, D. Sepsis-induced immunosuppression: from cellular dysfunctions to immunotherapy. *Nat Rev Immunol* **13**, 862-874, doi:10.1038/nri3552 (2013).
- 157 Fan, H. & Cook, J. A. Molecular mechanisms of endotoxin tolerance. *J Endotoxin Res.* **10**, 71-84, doi:10.1179/096805104225003997 (2004).
- 158 Beeson, P. B. Development of tolerance to typhoid bacterial pyrogen and its abolition by reticulo-endothelial blockade. *Proc Soc Exp Biol Med* **61**, 248-150 (1946).
- 159 Beutler, B. Tlr4: central component of the sole mammalian LPS sensor. *Curr Opin Immunol* **12**, 20-26 (2000).

- 160 Parker, L. C. *et al.* Endotoxin tolerance induces selective alterations in neutrophil function. *J Leukoc Biol* **78**, 1301-1305 (2005).
- 161 Bergenfelz, C. *et al.* Wnt5a induces a tolerogenic phenotype of macrophages in sepsis and breast cancer patients. *J Immunol* **188**, 5448-5458, doi:10.4049/jimmunol.1103378 (2012).
- 162 Medvedev, A. E., Lentschat, A., Wahl, L. M., Golenbock, D. T. & Vogel, S. N. Dysregulation of LPS-induced Toll-like receptor 4-MyD88 complex formation and IL-1 receptor-associated kinase 1 activation in endotoxin-tolerant cells. *J Immunol* **169**, 5209-5216 (2002).
- 163 Burns, K. *et al.* Inhibition of Interleukin 1 Receptor/Toll-like Receptor Signaling through the Alternatively Spliced, Short Form of MyD88 Is Due to Its Failure to Recruit IRAK-4. *J Exp Med* **197**, 263-268 (2003).
- 164 Janssens, S., Burns, K., Tschopp, J. & Beyaert, R. Regulation of interleukin-1- and lipopolysaccharide-induced NF-kappaB activation by alternative splicing of MyD88. *Curr Biol* **12**, 467-471 (2002).
- 165 Liu, D., Cao, S., Zhou, Y. & Xiong, Y. Recent advances in endotoxin tolerance. *J Cell Biochem*, 1-15 (2018).
- 166 Biswas, S. K. & Lopez-Collazo, E. Endotoxin tolerance: new mechanisms, molecules and clinical significance. *Trends Immunol* **30**, 475-487, doi:10.1016/j.it.2009.07.009 (2009).
- 167 Ziegler-Heitbrock, L. The p50-homodimer mechanism in tolerance to LPS. *J Endotoxin Res.* **7**, 219-222 (2001).
- 168 Newton, S. *et al.* Sepsis-induced changes in macrophage co-stimulatory molecule expression: CD86 as a regulator of anti-inflammatory IL-10 response. *Surg Infect (Larchmt)* **5**, 375-383, doi:10.1089/sur.2004.5.375 (2004).
- 169 Monneret, G. *et al.* The anti-inflammatory response dominates after septic shock: association of low monocyte HLA-DR expression and high interleukin-10 concentration. *Immunol Lett.* **95**, 193-198, doi: 10.1016/j.imlet.2004.07.009 (2004).
- 170 del Fresno, C. *et al.* Potent phagocytic activity with impaired antigen presentation identifying lipopolysaccharide-tolerant human monocytes: demonstration in isolated monocytes from cystic fibrosis patients. *J Immunol* **182**, 6494-6507, doi:10.4049/jimmunol.0803350. (2009).
- 171 Munoz, C. *et al.* Dysregulation of in vitro cytokine production by monocytes during sepsis. *J Clin Invest.* **88**, 1747-1754 (1991).
- 172 Bennett, J. A., Rao, V. S. & Mitchell, M. S. Systemic bacillus Calmette-Guerin (BCG) activates natural suppressor cells. *PNAS* **75**, 5142-5144 (1978).
- 173 Slavina, S. & Strober, S. Induction of Allograft Tolerance after Total Lymphoid Irradiation (TLI): Development of Suppressor Cells of the Mixed Leukocyte Reaction (MLR). *J Immunol* **123**, 942-946 (1979).
- 174 Pak, A. S. *et al.* Mechanisms of immune suppression in patients with head and neck cancer: presence of CD34(+) cells which suppress immune functions within cancers that secrete granulocyte-macrophage colony-stimulating factor. *Clinical Cancer Research* **1**, 95-103 (1995).
- 175 Young, M. R. *et al.* Mechanisms of immune suppression in patients with head and neck cancer: influence on the immune infiltrate of the cancer. *Int J Cancer.* **67**, 333-338. (1996).
- 176 Gabrilovich, D. I. *et al.* The terminology issue for myeloid-derived suppressor cells. *Cancer Res* **67**, 425; author reply 426, doi:10.1158/0008-5472.CAN-06-3037 (2007).
- 177 Gabrilovich, D. I. Myeloid-Derived Suppressor Cells. *Cancer Immunol Res.* **5**, 3-8 (2017).
- 178 Poschke, I. & Kiessling, R. On the armament and appearances of human myeloid-derived suppressor cells. *Clin Immunol* **144**, 250-268, doi:10.1016/j.clim.2012.06.003 (2012).
- 179 Janols, H. *et al.* A high frequency of MDSCs in sepsis patients, with the granulocytic subtype dominating in gram-positive cases. *J Leukoc Biol* **96**, 685-693, doi:10.1189/jlb.5HI0214-074R (2014).
- 180 Lai, D., Qin, C. & Shu, Q. Myeloid-derived suppressor cells in sepsis. *Biomed Res Int* **2014**, 598654, doi:10.1155/2014/598654 (2014).
- 181 Uhel, F. *et al.* Early Expansion of Circulating Granulocytic Myeloid-derived Suppressor Cells Predicts Development of Nosocomial Infections in Patients with Sepsis. *Am J Respir Crit Care Med* **196**, 315-327, doi:10.1164/rccm.201606-1143OC (2017).

- 182 Cai, W. *et al.* Clinical significance and functional studies of myeloid-derived suppressor cells in chronic hepatitis C patients. *J Clin Immunol* **33**, 798-808, doi:10.1007/s10875-012-9861-2 (2013).
- 183 Qin, A. *et al.* Expansion of monocytic myeloid-derived suppressor cells dampens T cell function in HIV-1-seropositive individuals. *J Virol* **87**, 1477-1490, doi:10.1128/JVI.01759-12 (2013).
- 184 Vollbrecht, T. *et al.* Chronic progressive HIV-1 infection is associated with elevated levels of myeloid-derived suppressor cells. *AIDS* **26**, F31-37, doi:10.1097/QAD.0b013e328354b43f (2012).
- 185 Ioannou, M. *et al.* Crucial role of granulocytic myeloid-derived suppressor cells in the regulation of central nervous system autoimmune disease. *J Immunol* **188**, 1136-1146, doi:10.4049/jimmunol.1101816 (2012).
- 186 Whitfield-Larry, F., Felton, J., Buse, J. & Su, M. A. Myeloid-derived suppressor cells are increased in frequency but not maximally suppressive in peripheral blood of Type 1 Diabetes Mellitus patients. *Clin Immunol* **153**, 156-164, doi:10.1016/j.clim.2014.04.006 (2014).
- 187 Talmadge, J. E. & Gabrilovich, D. I. History of myeloid-derived suppressor cells. *Nat Rev Cancer* **13**, 739-752, doi:10.1038/nrc3581 (2013).
- 188 Diaz-Montero, C. M. *et al.* Increased circulating myeloid-derived suppressor cells correlate with clinical cancer stage, metastatic tumor burden, and doxorubicin-cyclophosphamide chemotherapy. *Cancer Immunol Immunother* **58**, 49-59, doi:10.1007/s00262-008-0523-4 (2009).
- 189 Yang, L. *et al.* Abrogation of TGF beta signaling in mammary carcinomas recruits Gr-1+CD11b+ myeloid cells that promote metastasis. *Cancer Cell* **13**, 23-35, doi:10.1016/j.ccr.2007.12.004 (2008).
- 190 Yang, L. *et al.* Expansion of myeloid immune suppressor Gr+CD11b+ cells in tumor-bearing host directly promotes tumor angiogenesis. *Cancer Cell* **6**, 409-421, doi:10.1016/j.ccr.2004.08.031 (2004).
- 191 Chalmin, F. *et al.* Membrane-associated Hsp72 from tumor-derived exosomes mediates STAT3-dependent immunosuppressive function of mouse and human myeloid-derived suppressor cells. *J Clin Invest* **120**, 457-471, doi:10.1172/JCI40483 (2010).
- 192 Lu, T. & Gabrilovich, D. I. Molecular pathways: tumor-infiltrating myeloid cells and reactive oxygen species in regulation of tumor microenvironment. *Clin Cancer Res* **18**, 4877-4882, doi:10.1158/1078-0432.CCR-11-2939 (2012).
- 193 Meirow, Y., Kanterman, J. & Baniyash, M. Paving the Road to Tumor Development and Spreading: Myeloid-Derived Suppressor Cells are Ruling the Fate. *Front Immunol* **6**, 523, doi:10.3389/fimmu.2015.00523 (2015).
- 194 Stiff, A. *et al.* Nitric Oxide Production by Myeloid-Derived Suppressor Cells Plays a Role in Impairing Fc Receptor-Mediated Natural Killer Cell Function. *Clin Cancer Res* **24**, 1891-1904, doi:10.1158/1078-0432.CCR-17-0691 (2018).
- 195 Pillay, J. *et al.* A subset of neutrophils in human systemic inflammation inhibits T cell responses through Mac-1. *J Clin Invest* **122**, 327-336, doi:10.1172/JCI57990 (2012).
- 196 Alfaro, C. *et al.* Tumor-Produced Interleukin-8 Attracts Human Myeloid-Derived Suppressor Cells and Elicits Extrusion of Neutrophil Extracellular Traps (NETs). *Clin Cancer Res* **22**, 3924-3936, doi:10.1158/1078-0432.CCR-15-2463 (2016).
- 197 Dunn, G. P., Bruce, A. T., Ikeda, H., Old, L. J. & Schreiber, R. D. Cancer immunoeediting: from immunosurveillance to tumor escape. *Nat Immunol* **3**, 991-998, doi:10.1038/ni1102-991 (2002).
- 198 Dunn, G. P., Old, L. J. & Schreiber, R. D. The three Es of cancer immunoeediting. *Annu Rev Immunol* **22**, 329-360, doi:10.1146/annurev.immunol.22.012703.104803 (2004).
- 199 Ichim, C. V. Revisiting immunosurveillance and immunostimulation: Implications for cancer immunotherapy. *J Transl Med* **3**, 8, doi:10.1186/1479-5876-3-8 (2005).
- 200 Hung, K. *et al.* The central role of CD4(+) T cells in the antitumor immune response. *J Exp Med* **188**, 2357-2368 (1998).
- 201 Galon, J. *et al.* Type, density, and location of immune cells within human colorectal tumors predict clinical outcome. *Science* **313**, 1960-1964, doi:10.1126/science.1129139 (2006).

- 202 Clemente, C. G. *et al.* Prognostic value of tumor infiltrating lymphocytes in the vertical
growth phase of primary cutaneous melanoma. *Cancer* **77**, 1303-1310,
doi:10.1002/(SICI)1097-0142(19960401)77:7<1303::AID-CNCR12>3.0.CO;2-5 (1996).
- 203 Schumacher, K., Haensch, W., Roefzaad, C. & Schlag, P. M. Prognostic significance of
activated CD8(+) T cell infiltrations within esophageal carcinomas. *Cancer Res* **61**, 3932-
3936 (2001).
- 204 Crowe, N. Y., Smyth, M. J. & Godfrey, D. I. A critical role for natural killer T cells in
immunosurveillance of methylcholanthrene-induced sarcomas. *J Exp Med* **196**, 119-127
(2002).
- 205 Ni, L. & Lu, J. Interferon gamma in cancer immunotherapy. *Cancer Med* **7**, 4509-4516,
doi:10.1002/cam4.1700 (2018).
- 206 Tugues, S. *et al.* New insights into IL-12-mediated tumor suppression. *Cell Death Differ* **22**,
237-246, doi:10.1038/cdd.2014.134 (2015).
- 207 Street, S. E., Trapani, J. A., MacGregor, D. & Smyth, M. J. Suppression of lymphoma and
epithelial malignancies effected by interferon gamma. *J Exp Med* **196**, 129-134 (2002).
- 208 Smyth, M. J. *et al.* Nature's TRAIL--on a path to cancer immunotherapy. *Immunity* **18**, 1-6
(2003).
- 209 Smyth, M. J. *et al.* Perforin-mediated cytotoxicity is critical for surveillance of spontaneous
lymphoma. *J Exp Med* **192**, 755-760 (2000).
- 210 Shinkai, Y. *et al.* RAG-2-deficient mice lack mature lymphocytes owing to inability to
initiate V(D)J rearrangement. *Cell* **68**, 855-867 (1992).
- 211 Penn, I. Post-transplant malignancy: the role of immunosuppression. *Drug Saf* **23**, 101-113,
doi:10.2165/00002018-200023020-00002 (2000).
- 212 Wang, J. C., Xu, Y., Huang, Z. M. & Lu, X. J. T cell exhaustion in cancer: Mechanisms and
clinical implications. *J Cell Biochem* **119**, 4279-4286, doi:10.1002/jcb.26645 (2018).
- 213 Wherry, E. J. T cell exhaustion. *Nat Immunol* **12**, 492-499 (2011).
- 214 Shen, L. S. *et al.* CD4(+)/CD25(+)/CD127(low/-) regulatory T cells express Foxp3 and
suppress effector T cell proliferation and contribute to gastric cancers progression. *Clin*
Immunol **131**, 109-118, doi:10.1016/j.clim.2008.11.010 (2009).
- 215 Sasada, T., Kimura, M., Yoshida, Y., Kanai, M. & Takabayashi, A. CD4+CD25+ regulatory
T cells in patients with gastrointestinal malignancies: possible involvement of regulatory T
cells in disease progression. *Cancer* **98**, 1089-1099, doi:10.1002/encr.11618 (2003).
- 216 Liyanage, U. K. *et al.* Prevalence of regulatory T cells is increased in peripheral blood and
tumor microenvironment of patients with pancreas or breast adenocarcinoma. *J Immunol*
169, 2756-2761 (2002).
- 217 Woo, E. Y. *et al.* Cutting edge: Regulatory T cells from lung cancer patients directly inhibit
autologous T cell proliferation. *J Immunol* **168**, 4272-4276 (2002).
- 218 Ha, T. Y. The role of regulatory T cells in cancer. *Immune Netw* **9**, 209-235,
doi:10.4110/in.2009.9.6.209 (2009).
- 219 Janikashvili, N., Bonnotte, B., Katsanis, E. & Larmonier, N. The dendritic cell-regulatory T
lymphocyte crosstalk contributes to tumor-induced tolerance. *Clin Dev Immunol* **2011**,
430394, doi:10.1155/2011/430394 (2011).
- 220 Leach, D. R., Krummel, M. F. & Allison, J. P. Enhancement of antitumor immunity by
CTLA-4 blockade. *Science* **271**, 1734-1736 (1996).
- 221 Iwai, Y., Terawaki, S. & Honjo, T. PD-1 blockade inhibits hematogenous spread of poorly
immunogenic tumor cells by enhanced recruitment of effector T cells. *Int Immunol* **17**, 133-
144, doi:10.1093/intimm/dxh194 (2005).
- 222 Buchbinder, E. I. & Desai, A. CTLA-4 and PD-1 Pathways: Similarities, Differences, and
Implications of Their Inhibition. *Am J Clin Oncol* **39**, 98-106,
doi:10.1097/COC.000000000000239 (2016).
- 223 Yu, Y. & Cui, J. Present and future of cancer immunotherapy: A tumor microenvironmental
perspective. *Oncol Lett* **16**, 4105-4113, doi:10.3892/ol.2018.9219 (2018).
- 224 Gust, J., Taraseviciute, A. & Turtle, C. J. Neurotoxicity Associated with CD19-Targeted
CAR-T Cell Therapies. *CNS Drugs* **32**, 1091-1101, doi:10.1007/s40263-018-0582-9 (2018).


- 225 Page, D. P. *et al.* Tumor immunology and cancer immunotherapy: summary of the 2014
SITC prime. *Journal for Immuno Therapy of Cancer*, , doi:10.1186/s40425-015-0072-2
(2015).
- 226 Logan, C. Y. & Nusse, R. The Wnt signaling pathway in development and disease. *Annu
Rev Cell Dev Biol* **20**, 781-810, doi:10.1146/annurev.cellbio.20.010403.113126 (2004).
- 227 Oishi, I. *et al.* The receptor tyrosine kinase Ror2 is involved in non-canonical Wnt5a/JNK
signalling pathway. *Genes Cells* **8**, 645-654 (2003).
- 228 Steinhart, Z. & Angers, S. Wnt signaling in development and tissue homeostasis.
Development **145**, doi:10.1242/dev.146589 (2018).
- 229 Nusse, R. & Varmus, H. E. Many tumors induced by the mouse mammary tumor virus
contain a provirus integrated in the same region of the host genome. *Cell* **31**, 99-109 (1982).
- 230 Rijsewijk, F. *et al.* The Drosophila homolog of the mouse mammary oncogene int-1 is
identical to the segment polarity gene wingless. *Cell* **50**, 649-657 (1987).
- 231 Nusse, R. *et al.* A new nomenclature for int-1 and related genes: the Wnt gene family. *Cell*
64, 231 (1991).
- 232 Fukuda, T. *et al.* Antisera induced by infusions of autologous Ad-CD154-leukemia B cells
identify ROR1 as an oncofetal antigen and receptor for Wnt5a. *Proc Natl Acad Sci U S A*
105, 3047-3052, doi:10.1073/pnas.0712148105 (2008).
- 233 Willert, K. *et al.* Wnt proteins are lipid-modified and can act as stem cell growth factors.
Nature **423**, 448-452, doi:10.1038/nature01611 (2003).
- 234 Kurayoshi, M., Yamamoto, H., Izumi, S. & Kikuchi, A. Post-translational palmitoylation
and glycosylation of Wnt-5a are necessary for its signalling. *Biochem J* **402**, 515-523,
doi:10.1042/BJ20061476 (2007).
- 235 Wong, G. T., Gavin, B. J. & McMahon, A. P. Differential transformation of mammary
epithelial cells by Wnt genes. *Mol Cell Biol* **14**, 6278-6286 (1994).
- 236 MacDonald, B. T., Tamai, K. & He, X. Wnt/beta-catenin signaling: components,
mechanisms, and diseases. *Dev Cell* **17**, 9-26, doi:10.1016/j.devcel.2009.06.016 (2009).
- 237 Rao, T. P. & Kuhl, M. An updated overview on Wnt signaling pathways: a prelude for more.
Circ Res **106**, 1798-1806, doi:10.1161/CIRCRESAHA.110.219840 (2010).
- 238 He, T. C. *et al.* Identification of c-MYC as a target of the APC pathway. *Science* **281**, 1509-
1512 (1998).
- 239 Sugimura, R. & Li, L. Noncanonical Wnt signaling in vertebrate development, stem cells,
and diseases. *Birth Defects Res C Embryo Today* **90**, 243-256, doi:10.1002/bdrc.20195
(2010).
- 240 Ishitani, T. *et al.* The TAK1-NLK mitogen-activated protein kinase cascade functions in the
Wnt-5a/Ca(2+) pathway to antagonize Wnt/beta-catenin signaling. *Mol Cell Biol* **23**, 131-
139 (2003).
- 241 Komiya, Y. & Habas, R. Wnt signal transduction pathways. *Organogenesis* **4**, 68-75 (2008).
- 242 Veeman, M. T., Axelrod, J. D. & Moon, R. T. A second canon. Functions and mechanisms
of beta-catenin-independent Wnt signaling. *Dev Cell* **5**, 367-377 (2003).
- 243 Wang, Y. Wnt/Planar cell polarity signaling: a new paradigm for cancer therapy. *Mol Cancer
Ther* **8**, 2103-2109, doi:10.1158/1535-7163.MCT-09-0282 (2009).
- 244 Gavin, B. J., McMahon, J. A. & McMahon, A. P. Expression of multiple novel Wnt-1/int-1-
related genes during fetal and adult mouse development. *Genes Dev* **4**, 2319-2332 (1990).
- 245 Kikuchi, A., Yamamoto, H., Sato, A. & Matsumoto, S. Wnt5a: its signalling, functions and
implication in diseases. *Acta Physiol (Oxf)* **204**, 17-33, doi:10.1111/j.1748-
1716.2011.02294.x (2012).
- 246 Yamaguchi, T. P., Bradley, A., McMahon, A. P. & Jones, S. A Wnt5a pathway underlies
outgrowth of multiple structures in the vertebrate embryo. *Development* **126**, 1211-1223
(1999).
- 247 Leris, A. C., Roberts, T. R., Jiang, W. G., Newbold, R. F. & Mokbel, K. WNT5A expression
in human breast cancer. *Anticancer Res* **25**, 731-734 (2005).
- 248 Liang, H. *et al.* Wnt5a inhibits B cell proliferation and functions as a tumor suppressor in
hematopoietic tissue. *Cancer Cell* **4**, 349-360 (2003).

- 249 Roman-Gomez, J. *et al.* WNT5A, a putative tumour suppressor of lymphoid malignancies, is inactivated by aberrant methylation in acute lymphoblastic leukaemia. *Eur J Cancer* **43**, 2736-2746, doi:10.1016/j.ejca.2007.10.004 (2007).
- 250 Dejmek, J., Dejmek, A., Safholm, A., Sjolander, A. & Andersson, T. Wnt-5a protein expression in primary dukes B colon cancers identifies a subgroup of patients with good prognosis. *Cancer Research* **65**, 9142-9146, doi:10.1158/0008-5472.Can-05-1710 (2005).
- 251 Zhong, Z. *et al.* Decreased Wnt5a Expression is a Poor Prognostic Factor in Triple-Negative Breast Cancer. *Med Sci Monit*. **1**, 1-7, doi:10.12659/MSM.894821 (2016).
- 252 Kremenevskaja, N. *et al.* Wnt-5a has tumor suppressor activity in thyroid carcinoma. *Oncogene* **24**, 2144-2154, doi:10.1038/sj.onc.1208370 (2005).
- 253 Kurayoshi, M. *et al.* Expression of Wnt-5a is correlated with aggressiveness of gastric cancer by stimulating cell migration and invasion. *Cancer Res* **66**, 10439-10448, doi:10.1158/0008-5472.CAN-06-2359 (2006).
- 254 Weeraratna, A. T. *et al.* Wnt5a signaling directly affects cell motility and invasion of metastatic melanoma. *Cancer Cell* **1**, 279-288 (2002).
- 255 Huang, C. L. *et al.* Wnt5a expression is associated with the tumor proliferation and the stromal vascular endothelial growth factor--an expression in non-small-cell lung cancer. *J Clin Oncol* **23**, 8765-8773, doi:10.1200/JCO.2005.02.2871 (2005).
- 256 Bo, H. *et al.* Upregulation of Wnt5a promotes epithelial-to-mesenchymal transition and metastasis of pancreatic cancer cells. *BMC Cancer* **13**, 496, doi:10.1186/1471-2407-13-496 (2013).
- 257 Ekstrom, E. J. *et al.* WNT5A induces release of exosomes containing pro-angiogenic and immunosuppressive factors from malignant melanoma cells. *Mol Cancer* **13**, 88, doi:10.1186/1476-4598-13-88 (2014).
- 258 Bergenfelz, C. *et al.* Wnt5a inhibits human monocyte-derived myeloid dendritic cell generation. *Scand J Immunol* **78**, 194-204, doi:10.1111/sji.12075 (2013).
- 259 Valencia, J. *et al.* Wnt5a skews dendritic cell differentiation to an unconventional phenotype with tolerogenic features. *J Immunol* **187**, 4129-4139, doi:10.4049/jimmunol.1101243 (2011).
- 260 Liang, H. *et al.* Noncanonical Wnt signaling promotes apoptosis in thymocyte development. *J Exp Med* **204**, 3077-3084, doi:10.1084/jem.20062692 (2007).
- 261 Pereira, C., Schaer, D. J., Bachli, E. B., Kurrer, M. O. & Schoedon, G. Wnt5A/CaMKII signaling contributes to the inflammatory response of macrophages and is a target for the antiinflammatory action of activated protein C and interleukin-10. *Arterioscler Thromb Vasc Biol* **28**, 504-510, doi:10.1161/ATVBAHA.107.157438 (2008).
- 262 Christman, M. A., 2nd *et al.* Wnt5a is expressed in murine and human atherosclerotic lesions. *Am J Physiol Heart Circ Physiol* **294**, H2864-2870, doi:10.1152/ajpheart.00982.2007 (2008).
- 263 Rauner, M. *et al.* WNT5A is induced by inflammatory mediators in bone marrow stromal cells and regulates cytokine and chemokine production. *J Bone Miner Res* **27**, 575-585, doi:10.1002/jbmr.1488 (2012).
- 264 Blumenthal, A. *et al.* The Wingless homolog WNT5A and its receptor Frizzled-5 regulate inflammatory responses of human mononuclear cells induced by microbial stimulation. *Blood* **108**, 965-973, doi:10.1182/blood-2005-12-5046 (2006).
- 265 Yu, C. H. *et al.* Recombinant Wnt3a and Wnt5a elicit macrophage cytokine production and tolerization to microbial stimulation via Toll-like receptor 4. *Eur J Immunol* **44**, 1480-1490, doi:10.1002/eji.201343959 (2014).
- 266 Hornef, M. W., Normark, B. H., Vandewalle, A. & Normark, S. Intracellular recognition of lipopolysaccharide by toll-like receptor 4 in intestinal epithelial cells. *J Exp Med* **198**, 1225-1235, doi:10.1084/jem.20022194 (2003).
- 267 Uronen-Hansson, H. *et al.* Toll-like receptor 2 (TLR2) and TLR4 are present inside human dendritic cells, associated with microtubules and the Golgi apparatus but are not detectable on the cell surface: integrity of microtubules is required for interleukin-12 production in response to internalized bacteria. *Immunology* **111**, 173-178, doi:10.1111/j.0019-2805.2003.01803.x (2004).

- 268 Yamaguchi, N. *et al.* Constitutive activation of nuclear factor-kappaB is preferentially
involved in the proliferation of basal-like subtype breast cancer cell lines. *Cancer Sci* **100**,
1668-1674, doi:10.1111/j.1349-7006.2009.01228.x (2009).
- 269 Park, M. H. & Hong, J. T. Roles of NF-kappaB in Cancer and Inflammatory Diseases and
Their Therapeutic Approaches. *Cells* **5**, doi:10.3390/cells5020015 (2016).
- 270 Riva, M. *et al.* Induction of nuclear factor-kappaB responses by the S100A9 protein is Toll-
like receptor-4-dependent. *Immunology* **137**, 172-182, doi:10.1111/j.1365-
2567.2012.03619.x (2012).
- 271 Chavey, C. *et al.* Oestrogen receptor negative breast cancers exhibit high cytokine content.
Breast Cancer Res **9**, R15, doi:10.1186/bcr1648 (2007).
- 272 Skoog, L. *et al.* Estrogen receptor levels and survival of breast cancer patients. A study on
patients participating in randomized trials of adjuvant therapy. *Acta Oncol* **26**, 95-100
(1987).
- 273 Hsu, R. Y. *et al.* LPS-induced TLR4 signaling in human colorectal cancer cells increases
beta1 integrin-mediated cell adhesion and liver metastasis. *Cancer Res* **71**, 1989-1998,
doi:10.1158/0008-5472.CAN-10-2833 (2011).
- 274 Volk-Draper, L. *et al.* Paclitaxel therapy promotes breast cancer metastasis in a TLR4-
dependent manner. *Cancer Res* **74**, 5421-5434, doi:10.1158/0008-5472.CAN-14-0067
(2014).
- 275 Arai, K. *et al.* S100A8 and S100A9 overexpression is associated with poor pathological
parameters in invasive ductal carcinoma of the breast. *Curr Cancer Drug Targets* **8**, 243-
252 (2008).
- 276 Arai, K., Teratani, T., Kuruto-Niwa, R., Yamada, T. & Nozawa, R. S100A9 expression in
invasive ductal carcinoma of the breast: S100A9 expression in adenocarcinoma is closely
associated with poor tumour differentiation. *Eur J Cancer* **40**, 1179-1187,
doi:10.1016/j.ejca.2004.01.022 (2004).
- 277 Cormier, K. *et al.* Intracellular expression of inflammatory proteins S100A8 and S100A9
leads to epithelial-mesenchymal transition and attenuated aggressivity of breast cancer cells.
Anticancer Agents Med Chem **14**, 35-45 (2014).
- 278 McKiernan, E., McDermott, E. W., Evoy, D., Crown, J. & Duffy, M. J. The role of S100
genes in breast cancer progression. *Tumour Biol* **32**, 441-450, doi:10.1007/s13277-010-
0137-2 (2011).
- 279 Gumireddy, K. *et al.* ID1 promotes breast cancer metastasis by S100A9 regulation. *Mol*
Cancer Res **12**, 1334-1343, doi:10.1158/1541-7786.MCR-14-0049 (2014).
- 280 Mehmeti, M. *et al.* Expression of functional toll like receptor 4 in estrogen
receptor/progesterone receptor-negative breast cancer. *Breast Cancer Res* **17**, 130,
doi:10.1186/s13058-015-0640-x (2015).
- 281 Kawai, H., Minamiya, Y. & Takahashi, N. Prognostic impact of S100A9 overexpression in
non-small cell lung cancer. *Tumour Biol* **32**, 641-646, doi:10.1007/s13277-011-0163-8
(2011).
- 282 Zhao, Y. *et al.* Wnt5a promotes inflammatory responses via nuclear factor kappaB (NF-
kappaB) and mitogen-activated protein kinase (MAPK) pathways in human dental pulp
cells. *J Biol Chem* **289**, 21028-21039, doi:10.1074/jbc.M113.546523 (2014).
- 283 Halleskog, C. *et al.* Heterotrimeric G protein-dependent WNT-5A signaling to ERK1/2
mediates distinct aspects of microglia proinflammatory transformation. *J*
Neuroinflammation **9**, 111, doi:10.1186/1742-2094-9-111 (2012).
- 284 Halleskog, C. & Schulte, G. WNT-3A and WNT-5A counteract lipopolysaccharide-induced
pro-inflammatory changes in mouse primary microglia. *J Neurochem* **125**, 803-808,
doi:10.1111/jnc.12250 (2013).
- 285 Luo, L. *et al.* Rab8a interacts directly with PI3Kgamma to modulate TLR4-driven PI3K and
mTOR signalling. *Nat Commun* **5**, 4407, doi:10.1038/ncomms5407 (2014).
- 286 Wang, P. F. *et al.* Prognostic role of pretreatment circulating MDSCs in patients with solid
malignancies: A meta-analysis of 40 studies. *Oncoimmunology* **7**,
doi:10.1080/2162402X.2018.1494113 (2018).

- 287 Liu, Y. F. *et al.* Expansion and activation of granulocytic, myeloid-derived suppressor cells in childhood precursor B cell acute lymphoblastic leukemia. *J Leukoc Biol* **102**, 449-458, doi:10.1189/jlb.5MA1116-453RR (2017).
- 288 Marini, O. *et al.* Identification of granulocytic myeloid-derived suppressor cells (G-MDSCs) in the peripheral blood of Hodgkin and non-Hodgkin lymphoma patients. *Oncotarget* **7**, 27676-27688, doi:10.18632/oncotarget.8507 (2016).
- 289 Sagiv, J. Y. *et al.* Phenotypic diversity and plasticity in circulating neutrophil subpopulations in cancer. *Cell Rep* **10**, 562-573, doi:10.1016/j.celrep.2014.12.039 (2015).
- 290 Brandau, S. *et al.* Myeloid-derived suppressor cells in the peripheral blood of cancer patients contain a subset of immature neutrophils with impaired migratory properties. *J Leukoc Biol* **89**, 311-317, doi:10.1189/jlb.0310162 (2011).
- 291 Sippel, T. R. *et al.* Neutrophil degranulation and immunosuppression in patients with GBM: restoration of cellular immune function by targeting arginase I. *Clin Cancer Res* **17**, 6992-7002, doi:10.1158/1078-0432.CCR-11-1107 (2011).
- 292 Bacac, M. & Stamenkovic, I. Metastatic cancer cell. *Annu Rev Pathol* **3**, 221-247, doi:10.1146/annurev.pathmechdis.3.121806.151523 (2008).
- 293 Kogawa K, N. S., Katsuta H, Kudoh J, Tamiya S, Sakai Y, Shimizu N, Harada M. Expression of AIRE gene in peripheral monocyte/dendritic cell lineage. *Immunol Lett.* **80**, 195-198 (2002).
- 294 Eldershaw S A, S. D. M., and Narendran P Expression and function of the autoimmune regulator (Aire) gene in non-thymic tissue. *Clin Exp Immunol.* **163**, 296-308, doi:10.1111/j.1365-2249.2010.04316.x (2011).

The Cover



"Akaleyi I" by Emma Lindström.

