

LUND UNIVERSITY

Cortisol and Cardiovascular Reactivity and Habituation to a Virtual Reality Version of the Trier Social Stress Test

Jönsson, Peter H; Wallergård, Mattias; Österberg, Kai; Johansson, Gerd; Karlson, Björn

2009

Link to publication

Citation for published version (APA):

Jönsson, P. H., Wallergård, M., Österberg, K., Johansson, G., & Karlson, B. (2009). Cortisol and Cardiovascular Reactivity and Habituation to a Virtual Reality Version of the Trier Social Stress Test. Paper presented at The 8th International Conference on Occupational Stress and Health, San Juan, Puerto Rico.

Total number of authors: 5

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights. • Users may download and print one copy of any publication from the public portal for the purpose of private study

or research.

- You may not further distribute the material or use it for any profit-making activity or commercial gain
- · You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: https://creativecommons.org/licenses/

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117 221 00 Lund +46 46-222 00 00

Cortisol and Cardiovascular Reactivity and Habituation to a Virtual Reality Version of the Trier Social Stress Test

AIM

Previously we have shown that a Virtual Reality (VR) version of the Trier Social Stress Test (VR-TSST) elicits automomic responses comparable to the real world TSST. Briefley, the TSST involves a speech and performance of a math task in front of an audience. In this pilot study we examined if repeated stress provocations with VR-TSST would result in the characteristic habituation of the cortisol stress response.

METHOD

7 healthy men were confronted with the VR version of TSST twice (one week between sessions). Salivary cortisol, heart rate (HR), high frequency heart rate variability (HF-HRV, parasympathetic activity) and T-wave amplitude (TWA, inversly related to sympathetic activity) was assessed. Data were analyzed with repeated measures ANOVAs, with DAY (first and second) and CONDITION (baseline, preparation, speech, math-task, and following rest) as within group factors.



Peter Jönsson^{1*}, Mattias Wallergård², Kai Österberg¹, Gerd Johansson², & Björn Karlson¹ ¹ Occupational and Environmental Medicine, ² Ergonomics and Aerosol Technology, Lund University, Sweden * Corresponding Author: Peter_2.Jonsson@med.lu.se



Cortisol increased during stress at day 1, but not at day 2: DAY * CONDITION [F(6, 26) = 4.68, p < .05], quadratic trend [F(1, 6) = 6.96, p < .05]. HR and TWA did not differ between day 1 and 2. Main effects of CONDITION for HR [F(8,48) = 19.28, p < .0001] and for TWA [F(8, 48) = 12.81, p < .005]. HR increased [cubic trend F(1, 6) = 36.72, p < .005]; TWA decreased [cubic trend F(1, 6) = 14.36, p < .005]. HF-HRV n.s.

Cortisol secretion showed a marked increase during the first stress provocation, but habituated in the second session. HR and TWA reactivity were similar during both sessions, implicating a stable increase in sympathetic activity during VR-TSST. The results closely resemble those obtained in prior studies using the real life TSST. If these results can be replicated with larger samples, the VR-TSST may be used as a simple and standardized tool for social stress induction in experimental settings.



RESULTS

DISCUSSION