



LUND UNIVERSITY

Cucurbits as a model system for crop pollination management

Knapp, Jessica; Osborne, J.

Published in:
Journal of Pollination Ecology

2019

Document Version:
Peer reviewed version (aka post-print)

[Link to publication](#)

Citation for published version (APA):
Knapp, J., & Osborne, J. (2019). Cucurbits as a model system for crop pollination management. *Journal of Pollination Ecology*, 25(9), 89-102.

Total number of authors:
2

Creative Commons License:
CC BY

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

CUCURBITS AS A MODEL SYSTEM FOR CROP POLLINATION MANAGEMENT

Jessica L. Knapp*¹ and Juliet L. Osborne²

¹Lund University, Department of Biology, Lund, Sweden

²University of Exeter, Penryn Campus, Penryn, Cornwall, TR10 9FE, UK

Abstract—Cucurbit crops have steadily increased in production over the last 50 years, particularly in Asia where pioneering technological advancements and genetic improvements have created new hybrid varieties. Generally, cucurbits are dependent on insect-pollination for fruit set and are popular species for pollination studies. This review systematically summarises pollination research conducted in the major food genera of cucurbits: *Cucurbita*, *Cucumis*, and *Citrullus*, to ask: 1) what are cucurbits' requirement for pollination and their most effective pollinators? And 2) Does pollinator management increase pollinator visitation to, and yield of, cucurbit crops? These accounts of cucurbit pollination demonstrate that wild bee species such as *Bombus terrestris*, *B. impatiens* and *Eucera spp.* were frequently able to fulfil the pollination requirements of multiple cucurbit species. However, pollinator behaviour, pollen deposition on stigmas, and pollinators' contribution to yield vary between cucurbit species and study site. Nonetheless, the provision of additional floral resources at both field and farm scales may help to encourage pollination of cucurbit species whilst supporting pollinators' nutritional requirements beyond those already provided by the cucurbit crop. Synthesising studies on cucurbits' requirement for pollination and how pollinators vary spatially and temporally in the landscape can extend beyond cucurbit systems to inform growers and pollination ecologists of other pollinator-dependent crop species wishing to maximise pollination services, species conservation; or both.

Keywords: *Cucurbita*, *Cucumis*, *Citrullus*, pollination, pollinator populations

INTRODUCTION

Cucurbits (*Cucurbitaceae*) are a large plant family which include major food crops within the *Cucurbita* (squash, pumpkin, courgette), *Cucumis* (cucumber, melon) and *Citrullus* (watermelon) genera (Kumar 2016). Over centuries, cucurbits have been domesticated for their fleshy fruits, roots, leaves, shoots, seeds, and flowers for food and commodity goods and thus are economically important crops (Bates et al. 1990; Bisognin 2002). Cultivated cucurbits are grown in a cultiar of agricultural environments from widespread monocultures to small-scale traditional garden systems and many species are able to persist in environmental conditions usually considered marginal for agriculture (Bates et al. 1990).

From a biological viewpoint, cucurbits' co-evolution with insects has inspired much scientific intrigue. For example, their ability to produce bitter cucurbitacins as a defence against insect herbivory has led to research into whether these compounds can be used for biological control, particularly against beetles (Metcalf et al. 1982; Adler & Hazzard 2009; Cavanagh et al. 2010). Likewise, cucurbits' dependency on pollination (Free 1993) means that cucurbit flowers offer large quantities of nectar and pollen as floral rewards to visiting insects, such as solitary bees, bumblebees and honeybees (Tepedino 1981; Vidal et al. 2006). In particular, the North American squash and gourd bees belonging to the

genera *Eucera* (Fig. 2A) and *Xenoglossa* are thought to rely exclusively on *Cucurbita* pollen to rear their offspring (Hurd, Linsley & Michelbacher 1974; Tepedino 1981).

From an agricultural viewpoint, various mechanisms have been explored to improve cucurbit yield such as improving the sex expression of flowers (Rodriguez-Granados et al. 2017) and producing F₁ hybrid seed (Robinson 2000). Indeed, the yield (per hectare) of cucurbit crops has steadily increased over the last 50 years, particularly in Asia where pioneering technological advancements and genetic improvements, especially with seedless varieties not requiring pollination, have increased global production (Fig. 1) (McCreight et al. 2013). Likewise, and most relevant to this review, cucurbit yield can also be increased by improving pollination (Hoehn et al. 2008; Kouonon et al. 2009).

Cucurbits are a popular plant family for pollination studies, particularly within the major food genera of *Cucurbita* (Hurd et al. 1971; Willis & Kevan 1995; Artz et al. 2011; Petersen et al. 2014), *Cucumis* (Adamson et al. 2012; Ali et al. 2015; Motzke et al. 2015), and *Citrullus* (Kremen et al. 2004; Winfree et al. 2008; Pisanty et al. 2015). This is likely because many cucurbit species are monoecious (Box 1) with many ovules, so manipulating pollen deposition to examine its effect on yield is relatively straight forward. Because cucurbits are such a large and genetically diverse plant family, synthesising studies on their pollination can extend beyond these systems to inform pollination ecologists and growers of other pollinator-dependent species on how to maximise pollination services, species conservation; or both.

Received 23 February 2011, accepted 28 June 2011

*Corresponding author: jessica.knapp@biol.lu.se




FIGURE 1. Regional production (primary y axis) and average global yield (secondary y axis) of *Cucurbita* species from 1961 to 2013. Data source: FAOSTAT (Aggregate, may include official, semi-official, estimated or calculated data).



Figure 2(A) *Eucera pruinosus* visiting a staminate courgette flower for pollen, (B) *Bombus terrestris* visiting a staminate courgette flower for nectar, and (C) *B. terrestris* visiting a pistillate *Cucurbita pepo* flower for nectar. Fig. 2A was taken in California, United States, and Figs. 2B and 2C were taken in the United Kingdom. Fig. 2B and 2C © Daphne Wong.

This review aims to summarise pollination research conducted on the major food genera of cucurbits: *Cucurbita*, *Cucumis*, and *Citrullus* (Box 1). Specifically, we ask: 1) what are cucurbits' requirements for pollination and their most effective pollinators? And 2) Does pollinator management increase pollinator visitation or yield of cucurbit crops?

MATERIALS AND METHODS

We searched the ISI Web of Science, SCOPUS and Science Direct databases, and Google Scholar for studies that investigated pollination requirements, pollinator performance and pollination management in cucurbit crops. For all databases literature was searched between 1945 to September 2018 using the search terms: (Cucurbit*) OR (Cucumis) OR (Citrullus) OR (squash) OR (pumpkin) OR (courgette) OR (cucumber) OR (*melon), AND (Pollinat*). Results are summarised in text and in Tables. 1-4.

This review encompasses multiple ways in which pollination is directly or indirectly quantified (Box 2). Broadly, there are two approaches for quantifying pollination: the first directly estimates pollinator performance in terms of

pollinator behaviour and / or pollen deposition on stigmas, whilst the second indirectly estimates pollinators' contribution to yield, usually measured as seed set or fruit weight (Ne'eman et al. 2010). Therefore, in this review pollinator visitation to crop flowers, abundance at crop flowers, and pollen deposition (whilst stigmas are receptive) are all considered to be measures of pollinator performance. On the other hand, seed set, fruit weight, fruit weight per plant, fruit number, and percentage fruit set are all considered measures of yield.

Measures of yield should be interpreted carefully since they depend on multiple resources such as water, nutrients, and pollen availability. But broadly, fruit number and percentage fruit set reflect the number of flowers pollinated, whilst seed set, and average fruit weight reflect the quantity of pollen that a flower receives (affecting the number of seeds per fruit or the fruit size). Arguably, when other environmental factors that influence fruit production e.g. soil type, resource availability and cultivation practices cannot be standardised, single visit pollen deposition is the most direct measure of pollination success (Kremen et al., 2004). It should also be

BOX 1. Scientific and common names of different cucurbit species included within this review. For each species the most common sex expressions, breeding systems, as well as average flower ratios and flower longevity are listed based on Bomfim et al. (2016). G, gynoeocious, M, monoecious, A, andromonoecious, S, staminate; P, Pistillate.

Scientific name	Common name(s)	Origin	Sex expression(s)	Breeding system(s)	Average flower ratio (S:P)	Average anthesis
<i>Citrullus lanatus</i>	Watermelon	Tropical Africa	Monoecious is most common but can also be andromonoecious	Self-pollination and cross-pollination	5.1-13:1	≈ 8 hours in field
<i>Cucumis melo</i>	Melon	Tropical Africa	Andromonoecious is most common (especially American varieties), but can also be monoecious (especially European varieties)	Self-pollination and cross-pollination	12:1	≈ 12 hours in field
<i>Cucumis sativus</i>	Cucumber	India	Monoecious is most common but can also be gynoeocious, hermaphroditic, or andromonoecious.	Self-pollination and cross-pollination (M), or only cross-pollination (G), parthenocarpy.	10:1 (M), or all predominantly pistillate flowers (G)	≈ 7 hours in field, ≈ 9 hours in greenhouses
<i>Cucurbita maxima</i>	Buttercup squash etc.	South America	Monoecious	Self-pollination and cross-pollination	3.5-10:1	≈ 6 hours in field
<i>Cucurbita moschata</i>	Butternut squash, crookneck pumpkin etc.	Central America	Monoecious	Self-pollination and cross-pollination	3.5-10:1	≈ 6 hours in field
<i>Cucurbita pepo</i>	Courgette, zucchini, summer squash, pumpkin	Northern Mexico/Southern USA	Monoecious	Self-pollination and cross-pollination	3.5-10:1	≈ 6 hours in field

BOX 2. Measures of pollination included within this review.

Term	Definition
Fruit number	Number of fruits, a measure of yield.
Fruit weight	Weight per fruit or for all fruits per plant, a measure of yield.
Percentage fruit set	Initial number of female flowers/ Number of female flowers setting fruit X 100, a measure of yield.
Pollen deposition	Number of pollen grains deposited on stigmas following visitation by individual bee species or all bee species combined, for multiple visits. Single visit pollen deposition is included in Appendix I. Pollen deposition is a measure of pollinator behaviour.
Pollinator performance	A relative measure of pollinator effectiveness, based on pollinator behaviour (visitation (rate and/or timing), single visit pollen deposition) or contribution to yield (fruit weight, fruit number, seed set).
Seed set	Number of seeds per fruit, a measure of yield.
Visitation	Visitation rate to cucurbit flowers for individual bee species or all bee species combined, a measure of pollinator behaviour.

noted, in cucurbits that the actual timing of pollination can affect the final number of fruits: Cucurbit flowers open early in the morning, when both stigma receptivity and pollen viability are at their greatest (Bomfim et al. 2016) and diurnal pollen transfer is likely to be important in cucurbits since stigma receptivity rapidly declines within a day (Bomfim et al. 2016). Thus, flowers on which pollen is deposited early in the day are more likely to set fruit. Also, the presence of already pollinated fruits has been shown to significantly decrease the number of pistillate flowers and increase the likelihood of new fruit aborting in *Cucurbita pepo* (Stephenson et al. 1988).

RESULTS AND DISCUSSION

Pollinator dependency

Cucurbits have been described as having an ‘essential need’ for insect-mediated pollination (Free 1993; Klein et al. 2007) and research has shown that seed number (Roldán-Serrano & Guerra-Sanz 2005) and fruit set of *Cucurbita pepo* (Roldán-Serrano & Guerra-Sanz 2005; Vidal et al. 2010) and *Cucumis sativus* (Gingras, Gingras & DeOliveira 1999; Walters 2005) are positively correlated with the number of pollinator visits that each flower receives. Similarly, fruit has been shown to abort in the absence of pollination in *Cucumis sativus* (Motzke et al. 2015), *Cucumis melo* (Kouonon et al. 2009), *Cucurbita moschata* (Hoehn et al. 2008), and *Cucurbita pepo* (Martínez et al. 2014). Interestingly, the addition of honey bee colonies did not increase the yield of smaller-sized *Cucurbita pepo* varieties as much as larger ones; suggesting that alternative pollinator species may be more important in these varieties or that some smaller varieties may be able to set fruit with a smaller pollen load (Walters & Taylor 2006).

However, several varieties of *Cucurbita pepo* have been observed to set fruit in the absence of fertilisation, and therefore pollination, via parthenocarpy (Robinson & Reiners 1999; Kurtar 2003; Martínez et al. 2013; Knapp & Osborne

2017). This genetic trait is desirable for growers in crops that are usually pollinator-dependent because fruit is able to form in conditions that are adverse for pollinators, potentially extending geographic and climatic ranges of production (Knapp et al. 2016). Seedlessness in fruits, caused by the lack of fertilisation, can be an important measure of quality, for example in green-house grown *Cucumis sativus*, where seedlessness is generally appreciated by consumers (Knapp et al. 2016). However, evidence suggests that parthenocarpic varieties may still produce a greater quantity and quality of fruits, including a higher sugar content (Shin et al. 2007), when they are pollinated by insects (Martínez et al. 2013; Robinson & Reiners 1999; Nicodemo et al. 2013).

Whilst there have been extensive selective breeding programmes, use of growth hormones and even genetic modifications for parthenocarpy (Knapp et al. 2016), this review focuses on pollination of pollinator-dependent varieties. Nonetheless, evidence of parthenocarpy in cucurbit crops suggests that, to get a more complete picture of pollinator dependence in crops, varietal information is required – both in terms of pollinator dependence, but also in terms of choices that farmers are making (Klein et al. 2007; Melathopoulos et al. 2015). Realistically, the best way of obtaining this information is if the pollination requirements of each variety are tested by organisations or institutes conducting variety trials and that this information is made freely available alongside other trait details.

Relative importance of pollination to yield

Whilst pollination affects cucurbit yield, there are many other environmental factors which contribute and interact with each other to influence fruit set, such as nutrient and water availability, herbivore damage and weed competition (Tab. 1). If any of these factors are deficient then yield may decrease, widening the yield gap between actual and attainable yields (Bommarco et al. 2013). For example, Motzke et al. (2015) showed that weed control and fertilisation were able

TABLE I. Environmental factors affecting cucurbit yield. Effects were classed as 'positive', 'negative' or 'none' based on the direction of the relationship and statistical significance ($P < 0.05$) presented in the original studies.

Species	Sex expression, breeding system	Study location	Key findings	Citation
<i>Cucumis sativus</i>	Monoecious variety, some parthenocarpy observed in no pollination controls	Field study, Indonesia	When tested independently, pollination, weed control, and fertilisation had a positive effect on fruit weight, however, herbivore control had no effect on fruit weight. When tested in combination, pollination and weed control had a positive effect on fruit weight. 'Pollination' compared open pollination to no pollination.	(Motzke et al. 2015)
<i>Cucumis sativus</i>	Monoecious variety, assumed to be dependent on insect pollination.	Field study, USA	When tested independently, root herbivory had a negative effect on fruit weight, whilst leaf herbivory and pollination had no effect on fruit weight. When tested in combination, pollination and root herbivory, pollination and leaf herbivory, and pollination, root herbivory and leaf herbivory had no effect on fruit weight. When tested independently, root herbivory had a negative effect on seed set, leaf herbivory had positive effect on seed set and pollination had no effect on seed set. When tested in combination, pollination and root herbivory had a positive effect on seed set (pollination mitigated root herbivory), whilst pollination and leaf herbivory, and pollination, root herbivory and leaf herbivory had no effect on seed set. 'Pollination' compared hand pollination to open pollination.	(Barber et al. 2011)
<i>Cucurbita moschata</i>	Monoecious variety, assumed to be dependent on insect pollination.	Field study, USA	When tested independently, pollination and root herbivory had a positive effect on fruit weight, however, leaf herbivory had no effect on fruit weight. When tested in combination, pollination and root herbivory and pollination and leaf herbivory had no effect on fruit weight. Pollination, root herbivory and leaf herbivory, and their interactions had no effect on seed set. 'Pollination' compared hand pollination to open pollination.	(Hladun & Adler 2009)

to reduce the yield gap of *Cucumis sativus* by 45% and 18% respectively; however, these factors, even in combination, were unable to compensate for a total absence of pollination which increased the yield gap by 75% (the difference between open and non-pollinated flowers) (Tab. 1). In contrast, root herbivory had a negative effect on *Cucumis sativus* yield despite crop flowers receiving high levels of pollination, demonstrating that below-ground herbivores may have profound effects on plant performance (Barber et al. 2011) (Tab. 1). Soil nitrogen has also been shown to increase the number, weight and viability of pollen grains with flowers hand pollinated from pollen grown in higher nitrogen environments observed to produce *Cucurbita pepo* fruits with more seeds compared to hand pollinated flowers using pollen from lower nitrogen environments (Lau Tak-Cheung & Stephenson 1993).

Relatively few studies in cucurbits (Tab. 1) have examined how a plant's health interacts with the level of pollination it has received to influence fruit set. Drought stress, nutrient deficiencies and diseases such as cucumber mosaic virus and powdery mildew are relatively common in cucurbit production, often more so than direct pest damage (Agriculture and Horticulture Development Board 2013). Since biological control may reduce aphid populations, which

are common vectors of cucumber mosaic virus, and fungicides can reduce powdery mildew, fully-factorial experiments could be established to test the effect of disease control (biological and/or chemical) in relation to different levels of pollination (i.e. hand, open and no pollination), nutrient (e.g. fertiliser use), and water availability (e.g. irrigation and/or rain covers) on fruit set. Results from this type of experiment would help growers to realise the importance of pollination relative to other factors such as disease. A more complete understanding of the environmental factors affecting fruit set is vital to ensure that ecologists do not promote the conservation of one ecosystem service at the expense of another and that growers are able to prioritise key limiting services in their management for optimal crop yields.

Pollinator performance

Cucurbits are pollinated by multiple species of honeybees, bumblebees, and solitary bees (Tab. 2). Consequently, several studies have compared pollinator performance in cucurbit crops by exploring aspects of pollinator behaviour (visitation rate and diurnal activity patterns, single visit pollen deposition) and/ or their contribution to yield (fruit weight, fruit number, seed number) (Tab. 2).

TABLE 2. Key findings from studies looking at measures of pollinator performance such as, pollinator visitation to, and abundance at crop flowers, as well as pollen deposition in various cucurbit crops.

Species	Sex expression, breeding system	Key findings	Study location	Citation
<i>Citrullus lanatus</i>	Monoecious variety, no fruit set in no pollination controls.	<i>Lasioglossum</i> spp. were more effective than <i>A. mellifera</i> at increasing seed set.	Field study, Greece	(Garantonakis et al. 2016)
<i>Citrullus lanatus</i>	Monoecious variety, assumed to be dependent on insect pollination.	<i>Nomia oxybeloides</i> was more effective than <i>A. florea</i> at single-visit pollen deposition (no <i>Bombus</i> spp. present)	Field study, Pakistan	(Ali et al. 2015)
<i>Citrullus lanatus</i>	Monoecious variety with some perfect flowers, some fruit set in no insect pollination controls hypothesised to be from stray insects.	<i>Bombus impatiens</i> was more effective than <i>A. mellifera</i> at single-visit pollen deposition.	Field study, USA	(Campbell et al. 2018)
<i>Citrullus lanatus</i>	Monoecious variety, assumed to be dependent on insect pollination.	<i>Lasioglossum</i> spp. was more effective than <i>A. mellifera</i> at single-visit pollen deposition and visited earlier in the day (no <i>Bombus</i> spp. present)	Field study, Kenya	(Njoroge et al. 2010)
<i>Citrullus lanatus</i> and <i>Cucumis sativus</i>	Gynoecious and monoecious varieties, some parthenocarpy observed in no pollination controls but all aborted within 1 week.	<i>Bombus</i> spp. were more effective than <i>A. mellifera</i> for increasing seed set	Field study, USA	(Stanghellini et al. 1998)
<i>Cucumis melo</i>	Monoecious variety, assumed to be dependent on insect pollination.	<i>Lasioglossum</i> spp. was more effective than <i>A. florea</i> at single visit pollen deposition (no <i>Bombus</i> spp. present)	Field study, Pakistan	(Ali et al. 2015)
<i>Cucumis melo</i>	Andromonoecious variety, however, perfect flowers assumed to be dependent on pollination.	No difference between <i>A. mellifera</i> and <i>B. impatiens</i> at single-visit pollen deposition, however, <i>A. mellifera</i> had a lower pollen removal to pollen deposition ratio, suggesting it is a more effective pollinator.	Field study, USA	(Goodell & Thomson 2007)
<i>Cucurbita maxima</i>	Monoecious variety, assumed to be dependent on insect pollination.	<i>Bombus</i> spp. were more effective than <i>A. mellifera</i> at pollen deposition, however, <i>A. mellifera</i> had a higher visitation frequency than <i>Bombus</i> spp.	Field study, Germany	(Pfister et al. 2018)
<i>Cucurbita moschata</i>	Monoecious variety, no fruit set in no pollination controls.	High species richness of pollinators increased seed set.	Field study, Indonesia	(Hoehn et al. 2008)
<i>Cucurbita moschata</i>	Monoecious variety, assumed to be dependent on insect pollination.	<i>Eucera limitaris</i> was more abundant and more effective than <i>A. mellifera</i> at single-visit pollen deposition (no <i>Bombus</i> spp. present)	Field study, Mexico	(Canto-Aguilar & Veterinaria 2000)
<i>Cucurbita pepo</i>	Monoecious variety, assumed to be dependent on insect pollination.	<i>Bombus impatiens</i> was more effective than <i>A. mellifera</i> and <i>Eucera pruinosa</i> at single-visit pollen deposition, however, <i>A. mellifera</i> was a more abundant pollinator than <i>B. impatiens</i> and <i>P. pruinosa</i> .	Field study, USA	(Artz & Nault 2011)
<i>Cucurbita pepo</i>	Monoecious variety, assumed to be dependent on insect pollination.	<i>A. dorsata</i> was the most abundant pollinator, however, <i>Nomia</i> spp. were more effective than <i>A. dorsata</i> at single-visit pollen deposition and fruit set (no <i>Bombus</i> spp present)	Field study, Pakistan	(Ali et al. 2014)
<i>Cucurbita pepo</i>	Monoecious variety, assumed to be dependent on insect pollination.	<i>Bombus</i> spp. were more effective than <i>A. cerana</i> at single-visit pollen deposition, however, <i>A. cerana</i> was a more abundant pollinator than <i>Bombus</i> spp.	Field study, China	(Xie & An 2014)

When present, *Bombus* spp. are frequently observed to be the most effective pollinators of cucurbit crops, despite *Apis* spp. often being more abundant at crop flowers (Tab. 2). For example, Artz and Nault (2011) observed *Bombus impatiens* C., depositing more than three times the number of pollen grains per stigma and nearly always contacting the stigma compared to *A. mellifera* and *Eucera pruinosa* S, in *Cucurbita pepo*. On the other hand, *Eucera limitaris* was more abundant and more effective than *A. mellifera* at single-visit pollen deposition (no *Bombus* spp. present) in *Cucurbita moschata* (Canto-aguilar & Veterinaria 2000). Indeed, *P. pruinosa* has been shown to forage for *Cucurbita* pollen (Willis & Kevan 1995) and *Lassioglossum* spp. forage for *Citrullus* pollen (Njoroge et al. 2010) early in the morning and thus have a higher pollination potential. However, these foraging visits will only contribute to pollination if pistillate flowers are also visited. Outside of intensive production systems in North America and Europe, more diverse assemblages of pollinators have been observed to fulfil the pollination requirements of *Cucumis melo*, *Citrullus lanatus* (Ali et al. 2015), *Cucurbita pepo* (Ali et al. 2014) and *Cucurbita moschata* (Hoehn et al. 2008) (Tab. 2). Likewise, Pisanty et al. (2015) observed spatial and temporal variation in pollinator species' visitation to *Citrullus lanatus*, suggesting niche complementarity.

Since the majority of cucurbit species are monoecious (Box 1), consideration must also be given to whether some pollinator species show a preference for either staminate or pistillate flowers which could significantly influence pollen transfer. Indeed, bee species have been shown to preferentially choose and forage for longer in staminate flowers as their nectar has a higher sugar content (Knapp et al. 2018) and the nectaries are harder to access than pistillate flowers (Tepedino 1981; Artz & Nault 2011; Phillips & Gardiner 2015). In the United Kingdom, *Bombus terrestris* L. has been shown to have a more equal preference for staminate (Fig. 2B) and pistillate *Cucurbita pepo* flowers (Fig. 2C) and carried more loose pollen grains than *A. mellifera*, thought to be desirable for optimum pollen transfer and therefore, fruit set (Knapp et al. 2018). This study and others from outside North America (Tab. 2 and Tab. 4) have also shown that maximal *Cucurbita* yields can be achieved without *Eucera* and *Xenoglossa* species, North American bee species which are frequently cited as the most effective pollinators of *Cucurbita* crops (Hurd et al. 1974; Tepedino 1981).

Supplementing fields with managed pollinators

To mitigate potential fluctuations in wild bee abundance some cucurbit producers choose to supplement fields with managed bee species (Mader et al. 2010). Despite managed bee species being present in fields, visitation rates to crop flowers can be variable and benefits to crop yields sometimes not seen (Tab. 3). This could be because of a high abundance of wild pollinators and therefore little difference in pollination between control sites and those where managed pollinators have been added (Artz et al. 2011; Petersen et al. 2013, 2014, Tab. 3.). This suggests that in many cases pollination services by wild bees may be sufficient for optimal yields (Pfister et al. 2018). Therefore, before supplementing fields with managed pollinators, it would be advantageous to determine if a site is experiencing a pollination deficit. This can be done by

comparing yields from open and hand pollinated flowers (Petersen et al. 2014) or by combining pollinator visitation (Julier & Roulston 2009) or pollen deposition data at sites (Xie & An 2014; Pfister et al. 2018) with published data on the pollination requirements of the crop (Tab. 3). If farmers could use similar techniques (Fig. 3) then they could potentially reduce the amount that they spend on hiring in temporary pollination services.

Management for wild pollinators

Cucurbit yield can also increase if a crop is surrounded by more diverse habitats where increased species richness and abundance of wild pollinators can improve pollination services (Hoehn et al. 2008; Garibaldi et al. 2011) and provide insurance against any pollinator loss (Shuler et al. 2005). Improving the quantity and quality of pollen and nectar resources available for pollinators, and allowing areas to remain undisturbed for the insects to nest, mate, and hibernate could benefit pollinator populations and therefore reduce pollination deficits (Bommarco et al. 2013). This can be done at a field scale by providing additional areas for food and nesting and following principles of integrated pest management, as well as at a landscape scale by promoting larger areas of natural habitat (Tab. 4). However, the effectiveness of field scale pollinator-supporting practices are often variable (Tab. 4), with more simplistic landscapes generally showing greater benefits after interventions than complex landscapes (Batáry, Báldi, Kleijn, & Tschamtké, 2011; Scheper et al., 2013, Tab. 4).

Wild flowers co-flowering with crops have been shown to increase solitary bee abundance in *Cucumis melo* and *Citrullus lanatus* (Winfree et al. 2008) as well as *Bombus* abundance in *Cucurbita pepo* (Knapp et al. 2018) (Tab. 4). Naturally-occurring wild flowers such as agricultural weeds and hedgerow flowers are frequently overlooked floral resources for pollinators (Bretagnolle & Gaba 2015) despite being free and sustainable, and thus easily available to growers. These wild flowers are unlikely to be competing with cucurbit flowers for pollination services as both *B. terrestris* and *A. mellifera* have been observed to visit crop flowers more often than wild flowers in the morning when *Cucurbita pepo* flowers were open, before 'switching' to wild flowers after *C. pepo* senescence (Knapp et al. 2018). These findings provide the first evidence of bee fidelity (from non-specialists) to a *Cucurbita* crop.

Tillage has been shown to have variable results on bee abundance and visitation (Tab. 4.), despite *P. pruinosa* preferentially laying their eggs in crop areas at depths around 12 to 30 cm (Julier & Roulston 2009; Hurd et al. 1974). For example, no-tillage farms have been shown to have an almost three-fold increase in *P. pruinosa* density (Shuler et al. 2005), whilst other studies have observed no effect of tillage on *P. pruinosa* abundance (Julier & Roulston 2009, Minter & Bessin 2014). This may be a result of survey timings, as Julier & Roulston (2009) surveyed in mid-August, compared to Shuler et al. (2005) who surveyed in July when *P. pruinosa* is thought to emerge (Mathewson 1968). Given *P. pruinosa* requirement for cucurbit pollen, allowing areas to remain undisturbed for nesting and hibernation near to the crop may be particularly important (Mathewson 1968).

TABLE 3. Effect of supplemented commercial pollinators on cucurbit pollination. Effects are classified as 'positive', 'negative' or 'none' based on the direction of the relationship and statistical significance ($P < 0.05$) presented in the original studies.

Species	Sex expression, breeding system	Study location	Level of pollination	Key findings	Citation
<i>Cucumis sativus</i> , <i>Cucumis melo</i> , and <i>Cucurbita</i> (spp)	Several varieties assumed to be dependent on pollination. <i>C. sativus</i> described as self-fertile but benefitting from pollination	Field study, USA	Pollination not limiting- visitation rate from wild bees 83%	<i>A. mellifera</i> had no effect on <i>A. mellifera</i> visitation to crop flowers.	(Adamson et al. 2012)
<i>Cucurbita pepo</i> , <i>C. moschata</i> , and <i>C. maxima</i>	Monoecious variety assumed to be dependent on pollination.	Field study, USA	Pollination likely limiting yield - ~ 6 visits per hour in <i>A. mellifera</i> supplementation treatment compared to ~ 4 visits per hour in no supplementation treatment.	<i>A. mellifera</i> had a positive effect on fruit weight but no effect on the number of fruits per hectare.	(Walters & Taylor 2006)
<i>Cucurbita pepo</i>	Monoecious variety, no fruit set in no pollination controls.	Field study, USA	Pollination deficit not quantified. <i>B. impatiens</i> shown to be the most effective pollinator in the system.	<i>B. impatiens</i> had a positive effect on the number of fruits per plants but not fruit weight.	(Artz & Nault 2011)
<i>Cucurbita pepo</i>	Monoecious variety assumed to be dependent on pollination.	Field study, USA	Pollination likely limiting yield -visitation rate from wild bees 59%.	<i>A. mellifera</i> had a positive effect on <i>A. mellifera</i> visitation to crop flowers. <i>A. mellifera</i> had no effect on <i>B. impatiens</i> visitation to crop flowers. <i>A. mellifera</i> had a negative effect on <i>P. pruinosa</i> visitation to crop flowers. Field size had no effect on <i>B. impatiens</i> , <i>A. mellifera</i> , and <i>P. pruinosa</i> abundance.	(Artz et al. 2011)
<i>Cucurbita pepo</i>	Monoecious variety assumed to be dependent on pollination.	Field study, USA	Pollination deficit not quantified.	<i>A. mellifera</i> had no effect on <i>A. mellifera</i> visitation to crop flowers and fruit weight. <i>B. impatiens</i> had no effect <i>B. impatiens</i> visitation to crop flowers and fruit weight.	(Petersen et al. 2013)
<i>Cucurbita pepo</i>	Monoecious variety assumed to be dependent on pollination.	Field study, USA	Pollination not limiting to yield- no difference in yield between hand- and open-pollinated flowers.	<i>B. impatiens</i> had no effect on <i>B. impatiens</i> visitation, fruit weight, and seed set.	(Petersen et al. 2014)

Interestingly, in the studies we found, pesticide /insecticide use had no effect on bee visitation (Shuler et al. 2005; Motzke et al. 2015), pollen deposition (Kremen et al. 2004), or yield (Motzke et al. 2015) (Tab. 4). Whilst trap crops may not directly increase yields of focal cash crops (Tab. 4), they have proven successful at reducing the abundance of pests. For example, Adler & Hazzard (2009) observed

significantly fewer striped cucumber beetles (*Acalymma vittatum*) in *Cucurbita moschata*, the focal cash crop, with multiple cucurbit species as trap crops. Following a similar experimental design, Cavanagh et al. (2010) showed that trap crops could reduce insecticide use by 97% compared to control fields. Importantly, both studies found no effect of trap crops on pollinator visitation to *Cucurbita moschata*,

suggesting no competition between trap crops and focal crops for pollination services (Adler & Hazzard 2009; Cavanagh et al. 2010, Tab. 4).

At a landscape scale, pollination services to cucurbit crops by wild bees have been shown to relate to the amount of natural habitat surrounding a site (Tab. 4). For example, Kremen et al. (2004) found that pollination by native bees in *Citrullus lanatus* were strongly associated with the proportion of natural habitat within a 1 to 2.5 km radius of farm sites, meaning that based on the area of natural habitat, pollination services to a given site could be estimated (Kremen et al. 2004). The effect of natural habitat was also more important than organic versus conventional farming for predicting pollen deposition and pollinator abundance in *Citrullus lanatus* (Kremen et al. 2004, Tab. 4). As many agricultural systems are isolated from natural habitats, crop producers may need to provide floral resources and nesting sites suitable for pollinators. In the UK, farm stewardship schemes provide guidance on hedgerow and field margin management, particularly favoured by bumblebee species (Osborne et al. 2008; Carvell et al. 2015; Dicks et al. 2015; Wood et al. 2015). Costs can also be directly offset by increased profit from improved quality and quantity of yields. For example, considering an example of a non-cucurbit crop: the economic benefit of improved *Vaccinium corymbosum* L. yields following wild flower establishment has been shown to exceed the original cost of implementation (Blaauw & Isaacs 2014).

Large areas of mass-flowering crops, including the cucurbit crop itself, may dilute pollinator densities or, if the area is small, concentrate pollinator densities (Holzschuh et al. 2016). This will be especially pronounced if additional food and nesting sites are not provided, meaning that pollinators move transiently between available forage rather than increasing their population size (Holzschuh et al. 2016). In a UK study of *Cucurbita pepo*, whilst *B. terrestris* showed a strong fidelity to the crop flowers' bountiful nectar, no foragers were found returning to their colonies with *C. pepo* pollen loads (Knapp et al. 2018). Despite *C. pepo* pollen being relatively high in protein (Petersen et al. 2013), its large sticky grains may make it difficult for the bees to collect (Vaissière & Vinson 1994). Indeed, *B. terrestris* has been observed to remove excess pollen grains from their bodies early in the morning (Personal observations). Whilst *B. terrestris* has been observed to collect *Cucurbita* pollen in flight cages (Vaissière & Vinson 1994), no studies have observed *B. terrestris* collecting cucurbit pollen in open fields. Therefore, *B. terrestris* may avoid collecting *Cucurbita* pollen, since as a generalist species it can visit alternative, more easily obtainable (in open field settings) or more desirable pollen. Combining this empirical data on *Cucurbita pepo* nectar and pollen with model simulations using the novel bumblebee model *Bumble-BEEHAVE* (Becher and Twiston-Davies et al. 2018), showed that populations of *B. terrestris* can only benefit from the transient nectar source of *C. pepo* if alternative floral resources (particularly pollen) are also available to fulfil bees' nutritional requirements in space and time (Knapp et al. 2018).

The complexity of pollinator-supporting practices can become further complicated when species-level responses are

taken into consideration. For example, *Apis* spp. are less likely to increase their colony size in response to pollinator-supporting practices because their populations are artificially maintained by beekeepers (Tab. 4). Since previous research has shown that *Cucurbita* crops are primarily serviced by long range, generalist pollinators: *B. impatiens* and *A. mellifera* in the USA (Petersen et al. 2014) and *B. terrestris* and *A. mellifera* in the UK (Knapp and Osborne 2017) increasing forage and nesting may be less important to cucurbit crops, particularly in already heterogeneous environments (Tab. 4). This highlights the need to match pollinator-supportive management practices with the surrounding landscape and crops' individual requirements for pollination, since an increase in pollinator abundance and/or species richness may not necessarily be required for yield to be improved (Kleijn et al. 2015; Winfree et al. 2015). However, relying on a few generalist species for pollination services may be risky as any declines in their populations are also likely to affect specialist and/or less common bee species which may also be effective pollinators.

Value of pollination to growers

Quantifying the economic value of pollination can be useful for informing decision making at farm and policy levels (Hanley et al. 2014). Differentiating between the economic value of a crop's dependence on insect pollination and its pollination deficit will show the economic benefit of increasing pollinator populations as well as the economic cost that a decline in pollinator populations may have. Consequently, quantifying the economics of pollination is a fundamental way for growers to understand the implications that changes in pollinator populations may have on their yield and economic return. Economic valuations in the *Cucurbita pepo* variety 'Tosca' showed that whilst 56% of fruit set could be achieved via parthenocarp, the total economic value of insect pollination to production was estimated to be worth approximately £3,398/ha (Knapp & Osborne 2017). However, this economic valuation was based on the pollinator dependency and pollination deficit of just one *Cucurbita pepo* variety, thus inter-variety differences in pollinator dependence, or site-specific levels of pollination deficit may increase or decrease this economic value. An analogous example of this comes from apple crops: *Malus domestica* in the UK, where the variety 'Cox' is estimated to have a pollination deficit of £146/ha, compared to the variety 'Gala' which had a much higher pollination deficit of £6,459/ha (Garratt et al. 2013). This is because 'Gala' is more pollinator-dependent and has a larger pollination deficit due to high yields from hand-pollinated flowers, compared to 'Cox'. Whilst economic valuations are based on relatively simple estimates of pollinator dependence, pollination levels, and growing practices, which may not be representative of a larger spatial scale, they do clearly demonstrate the importance of pollinators to crop production (Gallai et al. 2009).

It would be useful to know how growers of different species of pollinator-dependent crops perceive pollination in relation to their crops' level of pollinator dependence, as well as how the different factors affecting fruit set are prioritised in farm management, relative to the empirical evidence. A social survey of growers would tell us how their attitudes




FIGURE 3. Experimental process for a grower to discover the extent to which their cucurbit variety depends on pollination, and to investigate the adequacy of the existing pollinator community (adapted from Corbet et al. 1991).

towards pollinators correspond to their management actions and if growers identify any key barriers or opportunities to integrating pollination in their management for optimal crop yields. To date, farmer surveys have mostly focused on ecosystem services in isolation, although see Andersson et al. (2015). For example, surveys identified that achieving consistent and reliable pollination is a priority for *V.*

corymbosum growers in Michigan and Florida in the USA (Integrated Crop Pollination Project, 2016). However, there is no way of knowing how much of a priority pollination is to these growers, relative to all the other factors which may affect yield such as pest control or soil quality. This information is critical to understand the likelihood of growers adapting

existing, or adopting new, sustainable pollination management.

Since growers are unlikely to know pollination rates for their crops and therefore their potential pollination deficit, it would be useful to develop a predictive model to determine if managed pollinators are required and/ or if longer-term pollinator habitat creation is warranted. Although other pollination service models exist (e.g. Olsson et al. 2015), BEE-STEWARD (www.beehave-model.net) has an interface which already enables users to simulate the effects that different management options, such as wild flower strips will have on bumblebee population dynamics and pollination services.

Conclusion

These studies highlight the importance of pollination for improving cucurbit yield, despite pollinator behaviour, pollen deposition on stigmas, and pollinators' contribution to yield varying across cucurbit species and study. From a biological view, this may be due to the relative attractiveness of flowers, and/or the pollinator dependency of the cucurbit species, whilst from a management view, the spatial and temporal context of study sites could affect the abundance and richness of wild pollinator species. Nonetheless, these accounts of cucurbit pollination demonstrate that wild bee species such as *B. terrestris*, *B. impatiens* and *Eucera* spp. are frequently able to fulfil the pollination requirements of multiple cucurbit species. Sufficient abundance of these wild bee species may be why in several cases, the addition of managed pollinator species had little or no effect on increasing yield. The provision of additional floral resources at both field and farm scales may help to encourage pollination of cucurbit species whilst supporting pollinators' nutritional requirements beyond those already provided by the cucurbit crop.

These findings extend beyond cucurbit systems (Fig. 3) to demonstrate how understanding a crop's requirement for pollination and how pollinators vary spatially and temporally in the landscape can aid growers in their decision making about what varieties and sites should be used. In doing so, growers may be able to increase their agricultural resilience and further their economic advantage. Nonetheless, further work is needed to understand how other environmental factors interact with pollination to influence fruit set so that growers can prioritise key regulating services in their management for optimal crop yields.

ACKNOWLEDGEMENTS

We would like to thank Benjamin Phillips and Lewis Bartlett for their valuable feedback on this manuscript. This work was funded as part of PhD studentship (CPI18) sponsored by the Agriculture and Horticulture Development Board, UK. JO was also supported by a grant from the Natural Environment Research Council UK [NE/J014893/1].

APPENDICES

Additional supporting information may be found in the online version of this article:

APPENDIX I. Pollen transfer in monoecious cucurbits.

REFERENCES

- Adamson NL, Roulston TH, Fell RD, Mullins DE (2012) From April to August—Wild Bees Pollinating Crops Through the Growing Season in Virginia, USA. *Environmental Entomology* 41:813–821.
- Adler LS, Hazzard R V (2009) Comparison of perimeter trap crop varieties: effects on herbivory, pollination, and yield in butternut squash. *Environmental entomology* 38:207–215.
- Agriculture and Horticulture Development Board (2013) *Crop Walkers' Guide - Outdoor Cucurbits*.
- Ali M, Saeed S, Sajjad A, Akbar A (2015) Linking Pollination Effectiveness and Interspecific Displacement Success in Bees. *Neotropical Entomology* 44:101–108. [online] URL: <http://link.springer.com/10.1007/s13744-014-0259-0>
- Ali M, Saeed S, Sajjad A, Bashir MA (2014) Exploring the Best Native Pollinators for Pumpkin (*Cucurbita pepo*) Production in Punjab, Pakistan. *Pakistan Journal of Zoology* 46:531–539.
- Andersson E, Nykvist B, Malinga R, Jaramillo F, Lindborg R (2015) A social – ecological analysis of ecosystem services in two different farming systems. *Ambio* 44:102–112.
- Artz DR, Hsu CL, Nault BA (2011) Influence of honey bee, *Apis mellifera*, hives and field size on foraging activity of native bee species in pumpkin fields. *Environmental Entomology* 40:1144–1158.
- Artz D, Nault B (2011) Performance of *Apis mellifera*, *Bombus impatiens*, and *Peponapis pruinosa* (Hymenoptera: Apidae) as pollinators of pumpkin. *Journal of Economic Entomology* 104:1153–1161.
- Barber NA, Adler LS, Bernardo HL (2011) Effects of above- and belowground herbivory on growth, pollination, and reproduction in cucumber. *Oecologia* 165:377–386.
- Batáry P, Báldi A, Kleijn D, Tschamtker T (2011) Landscape-moderated biodiversity effects of agri-environmental management: a meta-analysis. *Proceedings of the Royal Society B* 278:1894–1902.
- Bates DM, Robinson RW, Jeffrey C (1990) *Biology and Utilization of the Cucurbitaceae*. Comstock Publishing Associates, Ithaca and London.
- Becher MA, Twiston-Davies G, Penny TD, Goulson D, Rotheray EL, Osborne JL (2018) Bumble-BEEHAVE: A systems model for exploring multifactorial causes of bumblebee decline at individual, colony, population and community level. *Journal of Applied Ecology* 55:2790–2801.
- Bisognin DA (2002) Origin and evolution of cultivated cucurbits. *Ciência Rural* 32:715–723.
- Blaauw BR, Isaacs R (2014) Flower plantings increase wild bee abundance and the pollination services provided to a pollination-dependent crop. *Journal of Applied Ecology* 51:890–898.
- Bomfim I, Souza de Aragão F, Walters S (2016) Pollination in Cucurbit Crops. In: *Cucurbits: History, Nomenclature, Taxonomy, and Reproductive Growth*. pp 181–200.
- Bommarco R, Kleijn D, Potts SG (2013) Ecological intensification: Harnessing ecosystem services for food security. *Trends in Ecology and Evolution* 28:230–238.
- Bretagnolle V, Gaba S (2015) Weeds for bees? A review. *Agronomy for Sustainable Development* 35:891–909.
- Campbell JW, Kimmel CB, Bammer M, Stanley-Stahr C, Daniels JC, Ellis JD (2018) Managed and Wild Bee Flower Visitors and Their Potential Contribution to Pollination Services of Low-Chill Highbush Blueberry (*Vaccinium corymbosum* L.; Ericales: Ericaceae). *Journal of Economic Entomology*:1–6.

TABLE 4. Effect of pollinator-supporting practices on various metrics of pollination in cucurbits. Effects of these agricultural practices were classed as 'positive', 'negative' or 'none' based on the direction of the relationship and statistical significance ($P < 0.05$) presented in the original studies.

Species	Sex expression, breeding system	Study location	Level of pollination	Key findings	Citation
<i>Citrullus lanatus</i>	Monoecious varieties assumed to be dependent on insect pollination.	Field study, USA	Pollination deficit not quantified.	Insecticide use had no effect on pollen deposition. Amount of natural habitat surrounding a site had a positive effect on pollen deposition. Field size and organic farming had no effect on pollen deposition.	(Kremen et al. 2004)
<i>Cucumis melo</i> and <i>Citrullus lanatus</i>	Monoecious variety, assumed to be dependent on insect pollination.	Field study, USA	Pollination unlikely limiting yield – visitation from wild bees 62% and fields supplemented with <i>A. mellifera</i> .	Flowering weeds had no effect on bee visitation to crop flowers.	(Winfree et al. 2008)
<i>Cucumis sativus</i>	Monoecious variety, some parthenocarpy observed in no pollination controls	Field study, Indonesia	Pollination deficit not quantified. Only yield from none and open pollinated flowers.	Fertilisation had a positive effect on fruit weight and bee visitation to crop flowers. Insecticide use had no effect on fruit weight and bee visitation to crop flowers. Herbicide use and manual weeding had no effect on fruit weight and bee visitation to crop flowers.	(Motzke et al. 2015)
<i>Cucumis sativus</i>	Monoecious variety, assumed to be dependent on insect pollination.	Field study, USA	Pollination deficit not quantified. Fields were supplemented with <i>A. mellifera</i> .	Amount of natural habitat surrounding a site had a positive effect on wild bee abundance and richness.	(Lowenstein & Hueseth 2012)
<i>Cucurbita maxima</i>	Monoecious variety, assumed to be dependent on insect pollination.	Field study, Germany	Pollination not limiting to yield – more pollen grains deposited (~11,000) then required for fruit set (~2,500).	Amount of cropland surrounding a site had a negative effect on <i>Bombus</i> spp. visitation to crop flowers	(Pfister et al. 2018)
<i>Cucurbita moschata</i>	Monoecious variety, assumed to be dependent on insect pollination.	Field study, USA	Pollination not limiting to yield- visitation from wild bees was 87%.	Trap crops had no effect on fruit weight or bee visitation to crop flowers.	(Adler & Hazzard 2009)
<i>Cucurbita moschata</i>	Monoecious variety, assumed to be dependent on insect pollination.	Field study, USA	Pollination not limiting to yield- no difference in yield between hand- and open- pollinated flowers	Trap crops had no effect on fruit weight or bee visitation to crop flowers, but did reduce pest damage in the focal crop.	(Cavanagh et al. 2010)
<i>Cucurbita pepo</i>	Monoecious variety, some parthenocarpy observed in no pollination controls	Field study, UK	Pollination deficit not quantified.	Flowering weeds had a positive effect on <i>B. terrestris</i> abundance and no effect on <i>A. mellifera</i> abundance. Amount of natural habitat surrounding a site had a negative effect on <i>A. mellifera</i> abundance and no effect on <i>B. terrestris</i> abundance. Field size had no effect on <i>A. mellifera</i> or <i>B. terrestris</i> abundance.	(Knapp et al. 2019)

Tab. 4 continued

Species	Sex expression, breeding system	Study location	Level of pollination	Key findings	Citation
<i>Cucurbita pepo</i>	Monoecious variety, assumed to be dependent on insect pollination.	Field study, USA	Pollination unlikely limiting yield – visitation from wild bees 52% and fields supplemented with <i>A. mellifera</i> .	Floral strips had no effect on bee visitation to crop flowers or pollen deposition. Landscape diversity surrounding a site had a positive effect on bee visitation and pollen deposition.	(Phillips & Gardiner 2015)
<i>Cucurbita pepo</i>	Monoecious variety, assumed to be dependent on insect pollination.	Field study, USA	Pollination unlikely limiting yield as bee visitation was determined to be sufficiently high and some fields supplemented with <i>A. mellifera</i>	Soil clay content had a negative effect on <i>P. pruinosa</i> abundance. Tillage had no effect on <i>P. pruinosa</i> abundance. Amount of natural habitat surrounding a site had no effect on <i>P. pruinosa</i> abundance.	(Julier & Roulston 2009)
<i>Cucurbita pepo</i>	Monoecious variety, assumed to be dependent on insect pollination.	Field study, China.	Pollination unlikely limiting to yield – at most sites more pollen grains deposited then required for fruit set.	Amount of natural habitat surrounding a site had a positive effect on pollen deposition and <i>Bombus</i> spp. visitation but no effect on <i>A. cerana</i> visitation to crop flowers.	(Xie & An 2014)
<i>Cucurbita pepo</i>	Monoecious variety assumed to be dependent on pollination.	Field study, USA	Pollination deficit not quantified.	Landscape diversity surrounding a site had a positive effect <i>Bombus</i> spp. visitation and fruit weight.	(Petersen & Nault 2014)
<i>Cucurbita</i> spp.	Monoecious varieties assumed to be dependent on insect pollination.	Field study, USA	Pollination deficit not quantified.	Pesticide use had no effect on <i>Bombus</i> spp. or <i>P. pruinosa</i> visitation to crop flowers. Tillage had a negative effect on <i>P. pruinosa</i> visitation to crop flowers and no effect on <i>B. impatiens</i> visitation to crop flowers.	(Shuler et al. 2005)

Canto-Aguilar A, Veterinaria M (2000) Importance of conserving alternative pollinators: assessing the pollination efficiency of the squash bee, *Peponapis limitaris* in *Cucurbita moschata* (*Cucurbitaceae*). *Journal of Insect Conservation* 4:203–210.

Carvell C, Osborne JL, Bourke AFG, Freeman SN, Pywell RF, Heard MS (2015) Bumble bee species' responses to a targeted conservation measure depend on landscape context and habitat quality. *Ecological Applications* 21:1760–1771.

Cavanagh AF, Adler LS, Hazzard R V. (2010) Buttercup Squash Provides a Marketable Alternative to Blue Hubbard as a Trap Crop for Control of Striped Cucumber Beetles (Coleoptera: Chrysomelidae). *Environmental Entomology* 39:1953–1960. [online] URL: <https://academic.oup.com/ee/article-lookup/doi/10.1603/EN10056>

Corbet SA, Williams IH, Osborne JL (1991) Bees and the pollination of crops and wild flowers in the European community. *Bee World* 72:47–59.

Dicks L V., Baude M, Roberts SPM, Phillips J, Green M, Carvell C (2015) How much flower-rich habitat is enough for wild pollinators? Answering a key policy question with incomplete knowledge. *Ecological Entomology* 40:22–35. [online] URL: <http://doi.wiley.com/10.1111/een.12226>

Free JB (1993) *Insect pollination of crops*. Academic Press Inc, London.

Gallai N, Salles JM, Settele J, Vaissière BE (2009) Economic valuation of the vulnerability of world agriculture confronted with pollinator decline. *Ecological Economics* 68:810–821.

Garantonakis N, Varikou K, Birouraki A, Edwards M, Kalliakaki V, Andrinopoulos F (2016) Comparing the pollination services of honey bees and wild bees in a watermelon field. *Scientia Horticulturae* 204:138–144. [online] URL: <http://www.sciencedirect.com/science/article/pii/S0304423816301716>

Garibaldi LA, Steffan-Dewenter I, Kremen C, Morales JM, Bommarco R, Cunningham SA, Carvalheiro LG, Chacoff NP, Dudenhöffer JH, Greenleaf SS, Holzschuh A, Isaacs R, Krewenka K, Mandelik Y, Mayfield MM, Morandin LA, Potts SG, Ricketts TH, Szentgyörgyi H, Viana BF, Westphal C, Winfree R, Klein AM (2011) Stability of pollination services decreases with isolation from natural areas despite honey bee visits. *Ecology Letters* 14:1062–1072.

Garratt MPD, Breeze TD, Jenner N, Polce C, Biesmeijer JC, Potts SG (2013) Avoiding a bad apple: insect pollination enhances fruit quality and economic value. *Agriculture, Ecosystems and Environment* 184:34–40. [online] URL: <http://dx.doi.org/10.1016/j.agee.2013.10.032>

- Gingras D, Gingras J, DeOliveira D (1999) Visits of honeybees (Hymenoptera: Apidae) and their effects on cucumber yields in the field. *Journal of Economic Entomology* 92:435–438.
- Goodell K, Thomson JD (2007) Influence of bee species (Hymenoptera: Apiformes) with contrasting behaviors on pollen movement in a mustard, *Brassica rapa* (Brassicaceae) and the muskmelon *Cucumis melo* (Cucurbitaceae). *Entomologia Generalis* 29:237–252.
- Hanley N, Breeze TD, Ellis C, Goulson D (2014) Measuring the economic value of pollination services: Principles, evidence and knowledge gaps. *Ecosystem Services* 14:1–9. [online] URL: <http://linkinghub.elsevier.com/retrieve/pii/S2212041614001156>
- Heard MS, Carvell C, Carreck NL, Rothery P, Osborne JL, Bourke AFG (2007) Landscape context not patch size determines bumblebee density on flower mixtures sown for agri-environment schemes. *Biology Letters* 3:638–641.
- Hladun KR, Adler LS (2009) Influence of leaf herbivory, root herbivory, and pollination on plant performance in *Cucurbita moschata*. *Ecological Entomology* 34:144–152.
- Hoehn P, Tschamtker T, Tylianakis JM, Steffan-Dewenter I (2008) Functional group diversity of bee pollinators increases crop yield. *Proceedings. Biological sciences / The Royal Society* 275:2283–2291.
- Holzschuh A, Dainese M, Gonzalez-Varo J, Mudri-Stojnic S, Riedinger V, Rundlöf M, Scheper J, Wickens J, Wickens V, Bommarco R, Kleijn D, Potts S, Roberts S, Smith HG, Montserrat V, Vujic A, Steffan-Dewenter I (2016) Mass-flowering crops dilute pollinator abundance in agricultural landscapes across Europe. *Ecology Letters* 19:1228–1236.
- Hurd PD, Linsley EG, Michelbacher AD (1974) Ecology of the squash and gourd bee, *Peponapis pruinosa*, on cultivated cucurbits in California (Hymenoptera: Apoidea). *Smithsonian Contributions to Zoology*:1–17.
- Hurd P, Linsley G, Whitaker T (1971) Squash and Gourd Bees (*Peponapis*, *Xenoglossa*) and the Origin of the Cultivated Cucurbita. *Evolution* 25:218–234.
- Integrated Crop Pollination Project (2016) Florida Blueberry Grower Survey Report.
- Julier HE, Roulston TH (2009) Wild bee abundance and pollination service in cultivated pumpkins: farm management, nesting behavior and landscape effects. *Journal of Economic Entomology* 102:563–573.
- Kleijn D, Winfree R, Bartomeus I, Carvalheiro LG, Henry M, Isaacs R, Klein A-M, Kremen C, M'Gonigle LK, Rader R, Ricketts TH, Williams NM, Adamson NL, Ascher JS, Báldi A, Batáry P, Benjamin F, Biesmeijer JC, Blitzer EJ, Bommarco R, Brand MR, Bretagnolle V, Butten L, Cariveau DP, Chifflet R, Colville JF, Danforth BN, Elle E, Garratt MPD, Herzog F, Holzschuh A, Howlett BG, Jauker F, Jha S, Knop E, Krewenka KM, Le Féon V, Mandelik Y, May EA, Park MG, Pisanty G, Reemer M, Riedinger V, Rollin O, Rundlöf M, Sardiñas HS, Scheper J, Sciligo AR, Smith HG, Steffan-Dewenter I, Thorp R, Tschamtker T, Verhulst J, Viana BF, Vaissière BE, Veldtman R, Westphal C, Potts SG (2015) Delivery of crop pollination services is an insufficient argument for wild pollinator conservation. *Nature Communications* 6:7414.
- Klein A-MM, Vaissière BE, Cane JH, Steffan-Dewenter I, Cunningham SA, Kremen C, Tschamtker T (2007) Importance of pollinators in changing landscapes for world crops. *Proceedings of the Royal Society B: Biological Sciences* 274:303–313.
- Knapp JL, Bartlett LJ, Osborne JL (2016) Re-evaluating strategies for pollinator-dependent crops: How useful is parthenocarpy? *Journal of Applied Ecology* 54:1–9.
- Knapp JL, Becher MA, Rankin CC, Osborne JL (2018) *Bombus terrestris* in a mass-flowering pollinator-dependent crop: A mutualistic relationship? *Ecology and Evolution*
- Knapp JL, Osborne JL (2017) Courgette Production: Pollination Demand, Supply, and Value. *Journal of Economic Entomology* 110:1973–1979.
- Knapp JL, Shaw R., Osborne JL (2019) Pollinator visitation to mass-flowering courgette and co-flowering wild flowers: implications for pollination and pollinator conservation. *Basic and Applied Ecology*
- Kouonon LC, Jacquemart AL, Zoro Bi AI, Bertin P, Baudoin JP, Dje Y (2009) Reproductive biology of the andromonoecious *Cucumis melo* subsp. *agrestis* (Cucurbitaceae). *Annals of Botany* 104:1129–1139.
- Kremen C, Williams NM, Bugg RL, Fay JP, Thorp RW (2004) The area requirements of an ecosystem service: crop pollination by native bee communities in California. *Ecology Letters* 7:1109–1119.
- Kumar R (2016) Cucurbits: History, Nomenclature, Taxonomy, and Reproductive Growth. In: Pesarakli M (ed) *Handbook of Cucurbits: Growth, Cultural Practices, and Physiology*. pp 3–22.
- Kurtar E. (2003) An investigation of parthenocarpy in some summer squash (*Cucurbita pepo* L.) cultivars. *Journal of Agronomy* 2:209–213.
- Lau Tak-Cheung, Stephenson A (1993) Effects of soil nitrogen on pollen production, pollen grain size, and pollen performance in *Cucurbita pepo* (Cucurbitaceae). *American Journal of Botany* 80:763–768.
- Lowenstein DM, Huseth AS (2012) Response of Wild Bees (Hymenoptera: Apoidea: Anthophila) to Surrounding Land Cover in Wisconsin Pickling Cucumber. *Environmental Entomology* 41:532–540.
- Mader E, Spivak M, Evans E (2010) *Managing Alternative Pollinators: A Handbook for Beekeepers, Growers, and Conservationists*. NRAES, New York.
- Martínez C, Manzano S, Megías Z, Garrido D, Picó B, Jamilena M (2013) Involvement of ethylene biosynthesis and signalling in fruit set and early fruit development in zucchini squash (*Cucurbita pepo* L.). *BMC plant biology* 13:1–14.
- Martínez C, Manzano S, Megías Z, Garrido D, Picó B, Jamilena M (2014) Sources of parthenocarpy for zucchini breeding: relationship with ethylene production and sensitivity. *Euphytica* 200:349–362.
- Mathewson JA (1968) Nest Construction and Life History of the Eastern Cucurbit Bee, *Peponapis pruinosa* (Hymenoptera: Apoidea). *Journal of the Kansas Entomological Society* 41:255–261.
- McCreight J, Staub J, Wehner T, Dhillon N (2013) Gone Global: Familiar and Exotic Cucurbits Have Asian Origins. *HortScience* 48:1078–1089.
- Melathopoulos AP, Cutler GC, Tyedmers P (2015) Where is the value in valuing pollination ecosystem services to agriculture? *Ecological Economics* 109:59–70. [online] URL: <http://dx.doi.org/10.1016/j.ecolecon.2014.11.007>
- Metcalfe RL, Rhodes AM, Metcalfe RA, Ferguson J, Metcalfe ER, Lu P (1982) Cucurbitacin Contents and Diabroticite (*Coleoptera Chrysomelidae*) feeding upon *Cucurbita* spp. *Entomological Society of America* 11:931–937.
- Motzke I, Tschamtker T, T. W, Klein A (2015) Pollination mitigates cucumber yield gaps more than pesticide and fertilizer use in tropical smallholder gardens. *Journal of Applied Ecology* 52:261–269.

- Ne'eman G, Jürgens A, Newstrom-Lloyd L, Potts SG, Dafni A (2010) A framework for comparing pollinator performance: Effectiveness and efficiency. *Biological Reviews* 85:435–451.
- Nicodemo D, Malheiros EB, Jong D De, Couto RHN (2013) Enhanced production of parthenocarpic cucumbers pollinated with stingless bees and Africanized honey bees in greenhouses. *Semina: Ciências Agrárias* 34:3625–3634.
- Njoroge GN, Gemmill B, Bussmann R, Newton LE, Ngumi VM (2010) Diversity and efficiency of wild pollinators of watermelon (*Citrullus lanatus* (Thunb.) Mansf.) at Yatta (Kenya). *Journal of Applied Horticulture* 12:35–41.
- Olsson O, Bolin A, Smith HG, Lonsdorf E V. (2015) Modeling pollinating bee visitation rates in heterogeneous landscapes from foraging theory. *Ecological Modelling* 316:133–143.
- Osborne JL, Martin AP, Shortall CR, Todd AD, Goulson D, Knight ME, Hale RJ, Sanderson R. a. (2008) Quantifying and comparing bumblebee nest densities in gardens and countryside habitats. *Journal of Applied Ecology* 45:784–792.
- Petersen JD, Huseeth AS, Nault BA (2014) Evaluating pollination deficits in pumpkin production in New York. *Plant-Insect Interactions*:1247–1253.
- Petersen J, Nault B (2014) Landscape diversity moderates the effects of bee visitation frequency to flowers on crop production. *Journal of Applied Ecology*:1347–1356.
- Petersen JD, Reiners S, Nault B (2013) Pollination services provided by bees in pumpkin fields supplemented with either *Apis mellifera* or *Bombus impatiens* or not supplemented. *PLoS ONE* 8:1–8.
- Pfister SC, Eckert PW, Krebs J, Cresswell JE, Schirmel J, Entling MH (2018) Dominance of cropland reduces the pollen deposition from bumble bees. *Scientific Reports* 8:1–8.
- Phillips BW, Gardiner MM (2015) Use of video surveillance to measure the influences of habitat management and landscape composition on pollinator visitation and pollen deposition in pumpkin (*Cucurbita pepo*) agroecosystems. *PeerJ* 3:e1342. [online] URL: <https://peerj.com/articles/1342>
- Pisanty G, Afik O, Wajnberg E, Mandelik Y (2015) Watermelon pollinators exhibit complementarity in both visitation rate and single-visit pollination efficiency. *Journal of Applied Ecology* 53:360–370.
- Project ICP (2016) Michigan Blueberry Grower Survey Report.
- Robinson RW (2000) Rationale and Methods for Producing Hybrid Cucurbit Seed. *Journal of New Seeds* 1:49–67.
- Robinson RW, Reiners S (1999) Parthenocarpy in summer squash. *HortScience* 34:715–717.
- Rodríguez-Granados NY, Lemhemdi A, Choucha FA, Latrasse D, Benhamed M, Adnane B, Bendahmane A (2017) Sex Determination in *Cucumis*. In: Grumet R., Katzir N. G-MJ (ed) *Genetics and Genomics of Cucurbitaceae*, Plant Gene. Springer, Cham
- Roldán-Serrano S, Guerra-Sanz JM (2005) Reward attractions of zucchini flowers (*Cucurbita pepo* L.) to bumblebees (*Bombus terrestris* L.). *European Journal of Horticultural Science* 70:23–28.
- Scheper J, Holzschuh A, Kuussaari M, Potts SG, Rundlöf M, Smith HG, Kleijn D, Letters E (2013) Environmental factors driving the effectiveness of European agri-environmental measures in mitigating pollinator loss - a meta-analysis. *Ecology Letters* 16:912–920.
- Shin YS, Park SD, Kim JH (2007) Influence of pollination methods on fruit development and sugar contents of oriental melon (*Cucumis melo* L. cv. Sagyejeol-Ggul). *Scientia Horticulturae* 112:388–392.
- Shuler RE, Roulston TH, Farris GE (2005) Farming practices influence wild pollinator populations on squash and pumpkin. *Journal of Economic Entomology* 98:790–795.
- Stanghellini MS, Ambrose JT, Schultheis JR (1998) Using commercial bumble bee colonies as backup pollinators for honey bees to produce cucumbers and watermelons. *HortTechnology* 8:588–590.
- Stanghellini MS, Ambrose JT, Schultheis JR (2002) Diurnal activity, floral visitation and pollen deposition by honey bees and bumble bees on field-grown cucumber and watermelon. *Journal of Apicultural Research* 41:27–34.
- Stephenson AG, Devlin B, Horton JB (1988) The effects of seed number and prior fruit dominance on the pattern of fruit production in *Cucurbita pepo* (zucchini squash). *Annals of Botany* 62:653–661.
- Tepedino V (1981) The Pollination Efficiency of the Squash Bee (*Peponapis pruinosa*) and the Honey Bee (*Apis mellifera*) on Summer Squash (*Cucurbita pepo*). *Journal of the Kansas (Central States) Entomological Society* 54:359–377.
- Tscharntke T, Tylianakis JM, Rand TA, Didham RK, Fahrig L, Batáry P, Bengtsson J, Clough Y, Crist TO, Dormann CF, Ewers RM, Fründ J, Holt RD, Holzschuh A, Klein AM, Kleijn D, Kremen C, Landis DA, Laurance W, Lindenmayer D, Scherber C, Sodhi N, Steffan-Dewenter I, Thies C, van der Putten WH, Westphal C (2012) Landscape moderation of biodiversity patterns and processes - eight hypotheses. *Biological Reviews* 87:661–685.
- Vaissière BE, Vinson SB (1994) Pollen morphology and its effect on pollen collection by honey bees, *apis mellifera* L. (hymenoptera: Apidae), with special reference to upland cotton, *gossypium hirsutum* L. (malvaceae). *Grana* 33:128–138.
- Vidal MDG, Jong D De, Wien HC, Morse R a. (2006) Nectar and pollen production in pumpkin (*Cucurbita pepo* L.). *Revista Brasileira de Botânica* 29:267–273.
- Vidal MDG, Jong D De, Wien HC, Morse R a. (2010) Pollination and fruit set in pumpkin (*Cucurbita pepo*) by honey bees. *Revista Brasileira de Botânica* 33:106–113.
- Walters SA (2005) Honey bee pollination requirements for triploid watermelon. *HortScience* 40:1268–1270.
- Walters A, Taylor B (2006) Effects of honey bee pollination on pumpkin fruit and seed yield. *HortScience* 41:370–373.
- Willis DS, Kevan PG (1995) Foraging Dynamics of *Peponapis Pruinosa* (Hymenoptera: Anthophoridae) on Pumpkin (*Cucurbita Pepo*) in Southern Ontario. *The Canadian Entomologist* 127:167–175.
- Winfree R, Fox J, Williams NM, Reilly JR, Cariveau DP (2015) Abundance of common species, not species richness, drives delivery of a real-world ecosystem service. *Ecology Letters* 18:626–635. [online] URL: <http://doi.wiley.com/10.1111/ele.12424>
- Winfree R, Williams NM, Gaines H, Ascher JS, Kremen C (2008) Wild bee pollinators provide the majority of crop visitation across land-use gradients in New Jersey and Pennsylvania, USA. *Journal of Applied Ecology* 45:793–802.
- Wood TJ, Holland JM, Goulson D (2015) Pollinator-friendly management does not increase the diversity of farmland bees and wasps. *Biological Conservation* 187:120–126. [online] URL: <http://linkinghub.elsevier.com/retrieve/pii/S0006320715001755>
- Xie Z, An J (2014) The effects of landscape on bumblebees to ensure crop pollination in the highland agricultural ecosystems in China. *Journal of Applied Entomology* 138:555–565.

