LUND UNIVERSITY

The role of HOXB2 and HOXB3 in acute myeloid leukemia.

Lindblad, Oscar; Chougule, Rohit A.; Moharram, Sausan A.; Kabir, Nuzhat N; Sun, Jianmin; Kazi, Julhash U.; Rönnstrand, Lars

Published in:
Biochemical and Biophysical Research Communications

DOI:
[10.1016/j.bbrc.2015.10.071](https://doi.org/10.1016/j.bbrc.2015.10.071)

2015

Document Version:
Peer reviewed version (aka post-print)

[Link to publication](#)

Citation for published version (APA):
Lindblad, O., Chougule, R. A., Moharram, S. A., Kabir, N. N., Sun, J., Kazi, J. U., & Rönnstrand, L. (2015). The role of HOXB2 and HOXB3 in acute myeloid leukemia. *Biochemical and Biophysical Research Communications*, 467(4), 742-747. <https://doi.org/10.1016/j.bbrc.2015.10.071>

Total number of authors:
7

Creative Commons License:
CC BY-NC-ND

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

The role of HOXB2 and HOXB3 in acute myeloid leukemia

Oscar Lindblad^{1,2,3}, Rohit Chougule^{1,2}, Sausan A. Moharram^{1,2}, Nuzhat N. Kabir³, Jianmin Sun^{1,2},

Julhash U. Kazi^{1,2,4} and Lars Rönnstrand^{1,2*}

¹Division of Translational Cancer Research, Department of Laboratory Medicine, Lund University, Lund, Sweden.

²Lund Stem Cell Center, Department of Laboratory Medicine, Lund University, Lund, Sweden.

³Department of Hematology and Vascular Disorders, Skåne University Hospital, Lund, Sweden.

⁴Laboratory of Computational Biochemistry, KN Biomedical Research Institute, Barisal, Bangladesh.

* Corresponding author

Lars Rönnstrand

Division of Translational Cancer Research

Department of Laboratory Medicine

Lund University

Medicon Village 404:C3

223 63 Lund

Sweden

E-mail: lars.ronnstrand@med.lu.se Tel.: +46 46 222 6454

Abstract

Acute myeloid leukemia (AML) is a heterogeneous aggressive disease and the most common form of adult leukemia. Mutations in the type III receptor tyrosine kinase FLT3 are found in more than 30% of patients. Drugs against FLT3 have been developed for the treatment of AML, but they lack specificity, show poor response and lead to the development of a resistant phenotype upon treatment. Therefore, a deeper understanding of FLT3 signaling will facilitate identification of additional pharmacological targets in FLT3-driven AML. In this report, we identify HOXB2 and HOXB3 as novel regulators of oncogenic FLT3-ITD-driven AML. We show that HOXB2 and HOXB3 expression is upregulated in a group of AML patients carrying FLT3-ITD. Overexpression of HOXB2 or HOXB3 in mouse pro-B cells resulted in decreased FLT3-ITD-dependent cell proliferation as well as decreased colony formation and increased apoptosis. Expression of HOXB2 or HOXB3 resulted in a significant decrease in FLT3-ITD-induced AKT, ERK, p38 and STAT5 phosphorylation. Our data suggest that HOXB2 and HOXB3 act as tumor suppressors in FLT3-ITD driven AML.

Keywords Acute myeloid leukemia; HOXB2; HOXB3; FLT3; STAT5; colony formation

Introduction

Acute myeloid leukemia (AML) is a heterogeneous aggressive hematopoietic disorder which is the most common adult acute leukemia accounting for around 80% of cases. Genetic alterations lead to abrogated differentiation of hematopoietic cells. Therefore, the self-renewal ability of those cells is increased, and regulation of normal cell proliferation is disturbed. The major genetic changes include mutation in genes affecting cell proliferation (FLT3, KIT, NRAS/KRAS, JAK/STAT and PTPN11), myeloid differentiation (RUNX1/AML1 and CEBPA), cell cycle regulation or apoptosis (TP53, NPM1), and up-regulation of genes involved in stem-cell maintenance (HOXA, HOXB) [1-4].

The homeobox (HOX) genes are a family of homeodomain-containing transcription factors mainly involved in development. The human HOX genes are clustered in four different chromosomes, 7p15 (HOXA), 17q21 (HOXB), 12q13 (HOXC) and 2q31 (HOXD). HOXA family genes have been thoroughly studied in respect to AML [5-7]. While some studies suggest that HOXB family genes are up-regulated in certain types of AML, their roles in AML have not yet been defined [8-13]. A recent report suggests that HOXB4 expression is elevated in a group of AML patients and higher HOXB4 expression correlated with better prognosis [8]. Another report suggests that over-expression of HOXB6 in NB4 cells or in HL60 cells caused inhibition of the granulocytic and monocytic maturation, respectively [12].

The type III receptor tyrosine kinase FLT3 is expressed in almost all AML, and about 35% of AML patients carry an oncogenic *FLT3* mutation [14]. Among the several mutations that have been found, the internal tandem duplication (ITD) of the sequence that encodes the juxtamembrane domain is the most common mutation in FLT3. The presence of FLT3-ITD mutation portends a poor prognosis in AML. FLT3 mutations also occur in other types of leukemia to a lesser extent, including acute lymphoblastic leukemia [15]. While wild-type FLT3 requires its ligand FL for activation, oncogenic mutants are constitutively active. Activated FLT3 recruits SH2 domain-containing protein through phosphotyrosine residues resulting in activation of AKT, ERK, p38, and STAT5 [16-20].

In this report, we show that expression of HOXB2 and HOXB3 genes is upregulated in AML patients carrying the FLT3-ITD mutations. Loss of HOXB2 or HOXB3 expression in patients carrying FLT3-ITD mutations results in enrichment of oncogenic pathways. Overexpression of HOXB2 or HOXB3 significantly inhibits FLT3-ITD-induced cell proliferation and colony formation and further increases apoptosis.

Materials and methods

Plasmids and Antibodies: Plasmids expressing human HOXB2 and HOXB3 were generated by ligating the open reading frame (ORF) of the corresponding gene into the retroviral vector pMSCVneo. FLT3-ITD plasmid was described previously [21]. Anti-FLT3 antibody were described previously [22]. Anti-phosphotyrosine antibody 4G10 was purchased from Millipore (Life Technologies, Carlsbad, CA) and Anti-phospho p38 and anti-p38 antibodies were from BD Biosciences (Franklin Lakes, New Jersey). Anti-phospho-ERK1/2, anti-ERK2, anti-STAT5 and anti-AKT antibodies were from Santa-Cruz Biotechnology (Dallas, Texas) and anti-phospho AKT was from Epitomics (Abcam, Cambridge, UK). Anti- β -actin antibody was from Sigma-Aldrich (St. Louis, MO).

Cell culture and transfection: The murine pro-B cell line Ba/F3 was cultured in RPMI-1640 medium (Hyclone, Thermo Scientific, Waltham, MA) supplemented with 10% heat-inactivated fetal bovine serum (Life Technologies, Carlsbad, CA), 10 ng/ml murine interleukin 3 (IL3) and 1% penicillin and streptomycin. Generation of Ba/F3-FLT3-ITD cells was described previously [23]. FLT3-ITD-transfected Ba/F3 cells were then further transfected with the pMSCV-neo-HOXB2 or pMSCV-neo-HOXB3 construct or empty pMSCV-neo vector. Cells were selected with 0.8 mg/ml G-418 for 2 weeks. Transfected cells were maintained in the same medium like Ba/F3 medium as recommended before [24]. Cells were grown at 37°C in a humidified atmosphere containing 5% CO₂.

Immunoprecipitation and western blotting: Cells were starved of serum and cytokines for 4 hours and were washed once with cold PBS before lysis with Triton X-100 based lysis buffer. Cell lysates were

mixed with SDS and DTT containing loading buffer in a 1:1 ratio and boiled before separation by SDS-PAGE. For immunoprecipitation 1 µg anti-STAT5 antibody was added in cell lysates and was kept for 1 hour on ice followed by purification on protein G Dynabeads and SDS-PAGE analysis.

Apoptosis, cell proliferation, and colony formation assay: Cells were washed three times to remove cytokine before all experiments. Annexin V and 7-aminoactinomycin D (7-AAD) apoptosis kit (BD Biosciences) was used to measure apoptosis in cytokine depleted cells. Cells positive for annexin V and both annexin V/7-AAD were counted as apoptotic cells. To measure cell proliferation, 10,000 cells were seeded in each well of a 96-well plate and incubated for 48 hours. AlamarBlue (Molecular Probe) was used to measure cell viability. Semisolid methylcellulose medium (Stem Cell Technologies) was used for colony formation assay. Around 500 cells were seeded and cultured for seven days before counting colonies.

Microarray data analysis: The data set GSE14468 was used which was generated from a cohort of 598 newly diagnosed AML patients [25]. Gene expression was compared using significance analysis of microarrays (SAM) tools [26] and gene set enrichment analysis (GSEA) [27]. One-way ANOVA was used for statistical analysis. In statistical significance tests, "ns" represents not significant, "*" represents $p < 0.05$, "***" represents $p < 0.01$, and "****" represents $p < 0.001$.

Results

HOXB family proteins are upregulated in FLT3-ITD positive AML:

To understand the molecular difference between oncogenic FLT3-ITD positive and negative AML, we analyzed gene expression data of AML patients. We used expression data from bone marrow aspirates or peripheral blood samples of 598 cases of de novo AML. Using SAM tool we checked the differential gene expression. We observed that several HOXB family genes were upregulated in FLT3-ITD positive patients (Table S1). HOXB2 displayed 2.3-fold upregulation, HOXB3 4.4-fold, HOXB5 1.4-

fold and HOXB6 2.2 fold upregulation (Fig. S1A). Therefore, our data suggest that expression of several HOXB genes is deregulated in FLT3-ITD-driven AML.

HOXB2 and HOXB3 are independent prognostic markers in AML:

The HOXB family includes 10 genes, HOXB1-9 and 13. Since several HOXB-family genes were deregulated in FLT3-ITD driven AML, we checked whether expression of HOXB-family genes has any prognostic significance in AML. We transformed relative expression values to the Z-score and divided patients into two groups depending on higher and lower HOXB genes expression. We observed that higher expression of either HOXB2 (Fig. 1A) or HOXB3 (Fig. 1B) but not HOXB5 (Fig. S1B) correlated with poor prognosis compared to lower HOXB2 ($P=0.0053$) and HOXB3 ($P=0.0147$) expression. This also holds true for higher expression of HOXB6 (Fig. S1C), HOXB7 (Fig. S1D), HOXB8 (Fig. S1E) and HOXB9 (Fig. S1F). Lower HOXB2 (Fig. 1C) and HOXB3 (Fig. 1D) further correlated with better event-free survival ($P=0.0234$ and $P=0.0432$ respectively). Although HOXB2 and HOXB3 displayed prognostic significance in the total patient group independent of FLT3 mutations, we were unable to show any prognostic significance in only FLT3-ITD-dependent AML (data not shown) probably due to limited number of patient samples in each group. Since both HOXB2 and HOXB3 gene expression were upregulated in FLT3-ITD-dependent AML and since both genes expression profiles displayed independent prognostic significance, we checked whether HOXB2 and HOXB3 expression correlate with each other. We observed a strong correlation in between expression of the two genes ($r^2=0.8633$) suggesting that patients having higher HOXB2 expression will also have a higher HOXB3 expression and *vice versa* (Fig. 1E). In addition to FLT3-ITD positive AML patients (Fig. 1F), HOXB2 and HOXB3 expression was upregulated in patients carrying the NPM1 mutation (Fig. 1G). Although HOXB2 and HOXB3 expression correlated with FLT3-ITD and NPM1 mutations, expression neither correlated with patients age (Fig. S1G and S1H) nor with the patients sex (Fig. S1I). However, expression of both HOXB2 (Fig. S1J) and HOXB3 (Fig. S1K) was significantly downregulated in the FAB M3 group patients.

Loss of HOXB2 or HOXB3 expression correlates with enrichment of oncogenic pathways:

Since HOXB2 and HOXB3 expression were upregulated in FLT3-ITD positive patients, we wanted to analyze whether the loss of HOXB2 or HOXB3 expression results in the enrichment of any oncogenic pathways. Therefore, we analyzed enrichment of oncogenic pathways using gene set enrichment analysis (GSEA). We observed enrichment of several oncogenic pathways including loss of RB and p107 function, loss of SNF function and E2F3 pathways in FLT3-ITD positive AML patients with lower HOXB2 (Fig. 2A) or HOXB3 (Fig. 2B) expression. These results indicate a possible link between the loss of HOXB2 or HOXB3 expression and enhancement of oncogenic signaling in FLT3-ITD-driven AML.

HOXB2 or HOXB3 over-expression inhibits colony formation and cell proliferation and induces apoptosis:

Results from GSEA suggest that HOXB2 or HOXB3 plays a role in FLT3-ITD-dependent AML patients. To determine the role of HOXB2 and HOXB3 in FLT3-ITD signaling we generated two cell lines by stably transfecting the pro-B cell line Ba/F3 with FLT3-ITD together with HOXB2 or HOXB3, or empty control vector. Expression of FLT3-ITD and HOXB2 or HOXB3 was verified by western blotting (data not shown). Expression of HOXB2 or HOXB3 significantly reduced the size (Fig. 3A) and number (Fig. 3B) of colonies in semi-solid medium. Furthermore, cells expressing HOXB2 or HOXB3 displayed reduced FLT3-ITD-dependent cell proliferation (Fig. 3C) and significantly enhanced apoptosis (Fig. 4D) compared to the empty vector. These findings suggest that expression of HOXB2 and HOXB3 is essential for controlling FLT3-ITD-induced biological events.

HOXB2 or HOXB3 over-expression inhibits FLT3-ITD-induced phosphorylation of AKT, ERK1/2, p38, and STAT5:

Since we observed that HOXB2 and HOXB3 altered FLT3-ITD-dependent cell proliferation, apoptosis and colony formation, we analyzed FLT3 downstream signaling using phospho-specific antibodies. We observed that HOXB2 and HOXB3 overexpression blocked FLT3-ITD-dependent AKT (Fig. 4A),

ERK1/2 (Fig. 4B), p38 (Fig. 4C) and STAT5 (Fig. 4D) phosphorylation in Ba/F3 cells. Thus, we suggest that overexpression of HOXB2 or HOXB3 initiates a transcriptional program that inhibits FLT3-ITD downstream signaling.

Discussion

In this study, we aimed to address the role of HOXB2 and HOXB3 in FLT3-ITD-dependent AML. Expression of HOXB2 and HOXB3 was upregulated in FLT3-ITD-dependent patients, and both HOXB2 and HOXB3 can be used as an independent prognostic marker in AML. Patients carrying lower HOXB2 or HOXB3 have enriched oncogenic pathways enrichment and overexpression of HOXB2 or HOXB3 resulted in inhibition FLT3-ITD signaling as well as biological effects.

Although aberrant expression of homeobox genes is quite common in many cancers, the role of HOXB2 and HOXB3 in human cancer has not been thoroughly studied [28]. HOXB2 has been shown to be overexpressed in breast cancer, and overexpression of HOXB2 was correlated with better prognostic outcome [29]. HOXB2 acts as a tumor suppressor in breast cancer cells. Knockdown of HOXB2 using shRNA resulted in increased tumor growth [29]. We observed that overexpression of HOXB2 in Ba/F3 cells expressing FLT3-ITD resulted in negative regulation of FLT3-ITD-induced signaling as well as the corresponding biological outcomes such as cell proliferation and colony formation and enhancement of apoptosis. Although our observation in Ba/F3 cells in line with the observation in breast cancer cell lines [29], other studies in cervical cancer [30], lung adenocarcinomas [31] and pancreatic cancer [32] suggest overexpression of HOXB2 was associated with cancer progression. Therefore, we suggest that role of HOXB2 in cancer is context dependent.

HOXB3 expression correlated with HOXB2 expression in AML patients suggesting that HOXB3 display similar function to HOXB2 in AML. Although expression of HOXB3 increased cell proliferation in several tumors [33-35], we observed that overexpression of HOXB3 resulted in reduced cell proliferation and colony formation as well as increased apoptosis suggesting similar function to HOXB2 in AML.

Taken together, our study suggests that HOXB2 and HOXB3 have a tumor suppressor role in FLT3-driven AML. Since HOXB2 and HOXB3 are capable of limiting cell proliferation and colony formation as well as inducing apoptosis, overexpression of HOXB2 and HOXB3 keeps the oncogenic FLT3-ITD signaling in balance [36, 37]. It is well-known from a number of cell types that too strong oncogenic signaling can lead to induction of apoptosis[38, 39], so the tumors needs to keep the oncogenic signals at a moderate level in order to be transforming. Future studies are aiming at elucidating the downstream effectors of HOXB2 and HOXB3 and their role in oncogenic transformation.

Acknowledgements:

We thank Dr. Peter J. Valk at Erasmus University Medical Center for help with AML patient data. This research was funded by Kungliga Fysiografiska Sällskapet i Lund (JUK), Ollie and Elof Ericssons Stiftelse (JUK), Åke-Wiberg Stiftelse (JUK), Lars Hiertas Minne Stiftelse (JUK), Stiftelsen Olle Engkvist Byggmästare (JUK), Harald Jeansson's Stiftelse and Harald och Greta Jeansson's Stiftelse (JUK), Swedish Childhood Cancer Foundation (JUK and LR), Swedish Cancer Society (LR) and Swedish Research Council (LR). JUK is a recipient of Assistant Professorship (forskarassistenttjänst) grant from the Swedish Childhood Cancer Foundation.

References:

- [1] A. Renneville, C. Roumier, V. Biggio, O. Nibourel, N. Boissel, P. Fenaux, C. Preudhomme, Cooperating gene mutations in acute myeloid leukemia: a review of the literature, *Leukemia*, 22 (2008) 915-931.
- [2] K. Dohner, H. Dohner, Molecular characterization of acute myeloid leukemia, *Haematologica*, 93 (2008) 976-982.
- [3] R.G. Verhaak, C.S. Goudswaard, W. van Putten, M.A. Bijl, M.A. Sanders, W. Hagens, A.G. Uitterlinden, C.A. Erpelinck, R. Delwel, B. Lowenberg, P.J. Valk, Mutations in nucleophosmin (NPM1)

- in acute myeloid leukemia (AML): association with other gene abnormalities and previously established gene expression signatures and their favorable prognostic significance, *Blood*, 106 (2005) 3747-3754.
- [4] M. Alcalay, E. Tiacci, R. Bergomas, B. Bigerna, E. Venturini, S.P. Minardi, N. Meani, D. Diverio, L. Bernard, L. Tizzoni, S. Volorio, L. Luzi, E. Colombo, F. Lo Coco, C. Mecucci, B. Falini, P.G. Pelicci, Acute myeloid leukemia bearing cytoplasmic nucleophosmin (NPMc+ AML) shows a distinct gene expression profile characterized by up-regulation of genes involved in stem-cell maintenance, *Blood*, 106 (2005) 899-902.
- [5] K.L. Rice, J.D. Licht, HOX deregulation in acute myeloid leukemia, *The Journal of clinical investigation*, 117 (2007) 865-868.
- [6] T. Nakamura, D.A. Largaespada, M.P. Lee, L.A. Johnson, K. Ohyashiki, K. Toyama, S.J. Chen, C.L. Willman, I.M. Chen, A.P. Feinberg, N.A. Jenkins, N.G. Copeland, J.D. Shaughnessy, Jr., Fusion of the nucleoporin gene NUP98 to HOXA9 by the chromosome translocation t(7;11)(p15;p15) in human myeloid leukaemia, *Nature genetics*, 12 (1996) 154-158.
- [7] M. Camos, J. Esteve, P. Jares, D. Colomer, M. Rozman, N. Villamor, D. Costa, A. Carrio, J. Nomdedeu, E. Montserrat, E. Campo, Gene expression profiling of acute myeloid leukemia with translocation t(8;16)(p11;p13) and MYST3-CREBBP rearrangement reveals a distinctive signature with a specific pattern of HOX gene expression, *Cancer research*, 66 (2006) 6947-6954.
- [8] S. Umeda, K. Yamamoto, T. Murayama, M. Hidaka, M. Kurata, T. Ohshima, S. Suzuki, E. Sugawara, F. Kawano, M. Kitagawa, Prognostic significance of HOXB4 in de novo acute myeloid leukemia, *Hematology*, 17 (2012) 125-131.
- [9] H.C. Liu, L.Y. Shih, M.J. May Chen, C.C. Wang, T.C. Yeh, T.H. Lin, C.Y. Chen, C.J. Lin, D.C. Liang, Expression of HOXB genes is significantly different in acute myeloid leukemia with a partial tandem duplication of MLL vs. a MLL translocation: a cross-laboratory study, *Cancer genetics*, 204 (2011) 252-259.

- [10] M.S. Thakar, X.B. Zhang, B.C. Beard, G.E. Sale, E.B. Santos, L. Peterson, H.P. Kiem, B.M. Sandmaier, Transmission and expansion of HOXB4-induced leukemia in two immunosuppressed dogs: implications for a new canine leukemia model, *Experimental hematology*, 37 (2009) 1157-1166.
- [11] N.A. Fischbach, S. Rozenfeld, W. Shen, S. Fong, D. Chrobak, D. Ginzinger, S.C. Kogan, A. Radhakrishnan, M.M. Le Beau, C. Largman, H.J. Lawrence, HOXB6 overexpression in murine bone marrow immortalizes a myelomonocytic precursor in vitro and causes hematopoietic stem cell expansion and acute myeloid leukemia in vivo, *Blood*, 105 (2005) 1456-1466.
- [12] A. Giampaolo, N. Felli, D. Diverio, O. Morsilli, P. Samoggia, M. Breccia, F. Lo Coco, C. Peschle, U. Testa, Expression pattern of HOXB6 homeobox gene in myelomonocytic differentiation and acute myeloid leukemia, *Leukemia*, 16 (2002) 1293-1301.
- [13] D. Prevot, T. Voeltzel, A.M. Birot, A.P. Morel, M.C. Rostan, J.P. Magaud, L. Corbo, The leukemia-associated protein Btg1 and the p53-regulated protein Btg2 interact with the homeoprotein Hoxb9 and enhance its transcriptional activation, *The Journal of biological chemistry*, 275 (2000) 147-153.
- [14] D.G. Gilliland, J.D. Griffin, The roles of FLT3 in hematopoiesis and leukemia, *Blood*, 100 (2002) 1532-1542.
- [15] N.N. Kabir, L. Rönstrand, J.U. Kazi, FLT3 mutations in patients with childhood acute lymphoblastic leukemia (ALL), *Medical oncology*, 30 (2013) 462.
- [16] A. Puissant, N. Fenouille, G. Alexe, Y. Pikman, C.F. Bassil, S. Mehta, J. Du, J.U. Kazi, F. Luciano, L. Rönstrand, A.L. Kung, J.C. Aster, I. Galinsky, R.M. Stone, D.J. DeAngelo, M.T. Hemann, K. Stegmaier, SYK is a critical regulator of FLT3 in acute myeloid leukemia, *Cancer Cell*, 25 (2014) 226-242.
- [17] J.U. Kazi, M. Vaapil, S. Agarwal, E. Bracco, S. Pålman, L. Rönstrand, The tyrosine kinase CSK associates with FLT3 and c-Kit receptors and regulates downstream signaling, *Cell Signal*, 25 (2013) 1852-1860.

- [18] N.N. Kabir, J. Sun, L. Rönnstrand, J.U. Kazi, SOCS6 is a selective suppressor of receptor tyrosine kinase signaling, *Tumour biology : the journal of the International Society for Oncodevelopmental Biology and Medicine*, 35 (2014) 10581-10589.
- [19] N.N. Kabir, J.U. Kazi, Grb10 is a dual regulator of receptor tyrosine kinase signaling, *Molecular biology reports*, 41 (2014) 1985-1992.
- [20] J.U. Kazi, N.N. Kabir, A. Flores-Morales, L. Rönnstrand, SOCS proteins in regulation of receptor tyrosine kinase signaling, *Cell Mol Life Sci*, 71 (2014) 3297-3310.
- [21] J.U. Kazi, J. Sun, B. Phung, F. Zadjali, A. Flores-Morales, L. Rönnstrand, Suppressor of cytokine signaling 6 (SOCS6) negatively regulates Flt3 signal transduction through direct binding to phosphorylated tyrosines 591 and 919 of Flt3, *The Journal of biological chemistry*, 287 (2012) 36509-36517.
- [22] J.U. Kazi, L. Rönnstrand, FLT3 signals via the adapter protein Grb10 and overexpression of Grb10 leads to aberrant cell proliferation in acute myeloid leukemia, *Mol Oncol*, 7 (2013) 402-418.
- [23] O. Lindblad, T. Li, X. Su, J. Sun, N.N. Kabir, F. Levander, F. Zhao, G. Lu, L. Rönnstrand, J.U. Kazi, BEX1 acts as a tumor suppressor in acute myeloid leukemia, *Oncotarget*, 6 (2015) 21395-21405.
- [24] J.U. Kazi, J. Sun, L. Rönnstrand, The presence or absence of IL-3 during long-term culture of Flt3-ITD and c-Kit-D816V expressing Ba/F3 cells influences signaling outcome, *Experimental hematology*, 41 (2013) 585-587.
- [25] B.J. Wouters, B. Löwenberg, C.A. Erpelinck-Verschueren, W.L. van Putten, P.J. Valk, R. Delwel, Double CEBPA mutations, but not single CEBPA mutations, define a subgroup of acute myeloid leukemia with a distinctive gene expression profile that is uniquely associated with a favorable outcome, *Blood*, 113 (2009) 3088-3091.
- [26] V.G. Tusher, R. Tibshirani, G. Chu, Significance analysis of microarrays applied to the ionizing radiation response, *Proceedings of the National Academy of Sciences of the United States of America*, 98 (2001) 5116-5121.

- [27] A. Subramanian, P. Tamayo, V.K. Mootha, S. Mukherjee, B.L. Ebert, M.A. Gillette, A. Paulovich, S.L. Pomeroy, T.R. Golub, E.S. Lander, J.P. Mesirov, Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles, *Proceedings of the National Academy of Sciences of the United States of America*, 102 (2005) 15545-15550.
- [28] C. Abate-Shen, Deregulated homeobox gene expression in cancer: cause or consequence?, *Nat Rev Cancer*, 2 (2002) 777-785.
- [29] P.J. Boimel, C. Cruz, J.E. Segall, A functional in vivo screen for regulators of tumor progression identifies HOXB2 as a regulator of tumor growth in breast cancer, *Genomics*, 98 (2011) 164-172.
- [30] R. Lopez, E. Garrido, P. Pina, A. Hidalgo, M. Lazos, R. Ochoa, M. Salcedo, HOXB homeobox gene expression in cervical carcinoma, *International journal of gynecological cancer : official journal of the International Gynecological Cancer Society*, 16 (2006) 329-335.
- [31] K. Inamura, Y. Togashi, M. Okui, H. Ninomiya, M. Hiramatsu, Y. Satoh, S. Okumura, K. Nakagawa, T. Shimoji, T. Noda, Y. Ishikawa, HOXB2 as a novel prognostic indicator for stage I lung adenocarcinomas, *Journal of thoracic oncology : official publication of the International Association for the Study of Lung Cancer*, 2 (2007) 802-807.
- [32] D. Segara, A.V. Biankin, J.G. Kench, C.C. Langusch, A.C. Dawson, D.A. Skalicky, D.C. Gotley, M.J. Coleman, R.L. Sutherland, S.M. Henshall, Expression of HOXB2, a retinoic acid signaling target in pancreatic cancer and pancreatic intraepithelial neoplasia, *Clinical cancer research : an official journal of the American Association for Cancer Research*, 11 (2005) 3587-3596.
- [33] E. Eklund, The role of Hox proteins in leukemogenesis: insights into key regulatory events in hematopoiesis, *Crit Rev Oncog*, 16 (2011) 65-76.
- [34] Q. Li, F. Zhu, P. Chen, miR-7 and miR-218 epigenetically control tumor suppressor genes RASSF1A and Claudin-6 by targeting HoxB3 in breast cancer, *Biochemical and biophysical research communications*, 424 (2012) 28-33.

- [35] J.M. Bjornsson, N. Larsson, A.C. Brun, M. Magnusson, E. Andersson, P. Lundstrom, J. Larsson, E. Repetowska, M. Ehinger, R.K. Humphries, S. Karlsson, Reduced proliferative capacity of hematopoietic stem cells deficient in Hoxb3 and Hoxb4, *Molecular and cellular biology*, 23 (2003) 3872-3883.
- [36] J.U. Kazi, L. Rönstrand, Suppressor of cytokine signaling 2 (SOCS2) associates with FLT3 and negatively regulates downstream signaling, *Mol Oncol*, 7 (2013) 693-703.
- [37] J.U. Kazi, L. Rönstrand, Src-Like adaptor protein (SLAP) binds to the receptor tyrosine kinase Flt3 and modulates receptor stability and downstream signaling, *PloS one*, 7 (2012) e53509.
- [38] E. Oveland, L. Wergeland, R. Hovland, J.B. Lorens, B.T. Gjertsen, K.E. Fladmark, Ectopic expression of Flt3 kinase inhibits proliferation and promotes cell death in different human cancer cell lines, *Cell biology and toxicology*, 28 (2012) 201-212.
- [39] I. Alanazi, E. Ebrahimie, P. Hoffmann, D.L. Adelson, Combined gene expression and proteomic analysis of EGF induced apoptosis in A431 cells suggests multiple pathways trigger apoptosis, *Apoptosis : an international journal on programmed cell death*, 18 (2013) 1291-1305.

Figure legends

Figure 1: Overall survival of AML patients with higher and lower HOXB2 or HOXB3 expression: Z-score was used to divide higher and lower HOXB2 or HOXB3 expressing AML patients. (A) Overall survival of AML patients with higher or lower HOXB2 expression. . (B) Overall survival of AML patients with higher or lower HOXB3 expression. (C) The event-free survival of AML patients with higher or lower HOXB2 expression. (D) The event-free survival of AML patients with higher or lower HOXB3 expression. (E) Correlation in between HOXB2 and HOXB3 expression. (F) Expression of HOXB2 and HOXB3 in FLT3-ITD positive and negative AML patients. (G) Expression of HOXB2 and HOXB3 in NPM1 mutation-positive and negative AML patients.

Figure 2: GSEA shows enrichment of oncogenic pathways in lower HOXB2 or HOXB3 expressing AML patients: Z-score was used to divide higher and lower HOXB2 or HOXB3 expressing FLT3-ITD positive AML patients. (A) Loss of HOXB2 expression correlated with enrichment of several oncogenic pathways.

(B) AML patients with lower HOXB3 expression showed enrichment of several oncogenic pathways compared to patients with higher HOXB3 expression.

Figure 3: Expression of HOXB2 or HOXB3 inhibits cell proliferation, colony formation, and enhanced apoptosis: (A) About 500 stably transfected Ba/F3 cells were used to determine colony formation potential in the semi-solid medium. (B) The number of colonies in stably transfected Ba/F3 cells in each well of 24 well plate. (C) FLT3-ITD dependent cell viability in presence and absence of HOXB2 or HOXB3 expression was measured after 48 hours using AlamarBlue. (D) Apoptosis induced by overexpression of HOXB2 or HOXB3 was measured using Annexin-V and 7-AAD kit in cytokine depleted cells.

Figure 4: Expression of HOXB2 or HOXB3 decreases FLT3-ITD-induced downstream signaling: Total cell lysates were used for SDS-PAGE and Western blotting analysis with AKT (A), ERK (B) and p38 (C) antibodies. (D) Cell lysates were immunoprecipitated with an anti-STAT5 antibody followed by western blotting analysis.

Figure 1

Figure 2

Figure 3

Figure 4

Figure S1: Prognostic significance of HOXB-family proteins.

Table S1: Up-regulated genes in FLT3-ITD positive samples

Gene ID	Gene Name	Score (d)	Numerator (r)	Denominator (s+s0)	Fold Change	q-value (%)	Local fdr (%)
209392_at	ENPP2	9.03	1.54	0.17	2.91	0.00	0.00
228372_at	C10orf128	8.21	1.23	0.15	2.35	0.00	0.00
235391_at	FAM92A1	8.02	1.03	0.13	2.04	0.00	0.00
228904_at	HOXB3	8.02	2.13	0.27	4.37	0.00	0.00
227461_at	STON2	7.76	1.33	0.17	2.51	0.00	0.00
236892_s_at	HOXB-AS3	7.75	1.41	0.18	2.65	0.00	0.00
235852_at	STON2	7.70	1.23	0.16	2.34	0.00	0.00
228011_at	FAM92A1	7.68	0.95	0.12	1.93	0.00	0.00
200923_at	LGALS3BP	7.62	1.52	0.20	2.87	0.00	0.00
229437_at	MIR155 /// MIR155HG	7.52	1.44	0.19	2.72	0.00	0.00
204341_at	TRIM16	7.46	0.82	0.11	1.77	0.00	0.00
213110_s_at	COL4A5	7.20	1.76	0.24	3.39	0.00	0.01
211144_x_at	TARP /// TRGC2	7.17	1.34	0.19	2.53	0.00	0.01
215806_x_at	TARP /// TRGC2	7.17	1.36	0.19	2.56	0.00	0.01
216920_s_at	TARP /// TRGC2	7.14	1.34	0.19	2.52	0.00	0.01
209813_x_at	TARP	7.13	1.34	0.19	2.53	0.00	0.01
1553808_a_at	NKX2-3	7.12	1.58	0.22	3.00	0.00	0.01
201663_s_at	SMC4	7.01	0.92	0.13	1.89	0.00	0.02
201664_at	SMC4	6.92	0.92	0.13	1.89	0.00	0.02
215388_s_at	CFH /// CFHR1	6.91	0.91	0.13	1.88	0.00	0.02
219615_s_at	KCNK5	6.89	0.81	0.12	1.75	0.00	0.02
213800_at	CFH	6.87	0.71	0.10	1.64	0.00	0.03
210839_s_at	ENPP2	6.76	0.82	0.12	1.77	0.00	0.03
203372_s_at	SOCS2	6.73	1.57	0.23	2.97	0.00	0.03
203151_at	MAP1A	6.65	0.88	0.13	1.84	0.00	0.04
203373_at	SOCS2	6.65	1.43	0.21	2.69	0.00	0.04
1555600_s_at	APOL4	6.58	0.71	0.11	1.63	0.00	0.04
205453_at	HOXB2	6.55	1.20	0.18	2.29	0.00	0.05
232424_at	PRDM16	6.43	1.02	0.16	2.03	0.00	0.06
203897_at	LYRM1	6.39	0.74	0.12	1.67	0.00	0.06
206341_at	IL2RA	6.39	0.71	0.11	1.64	0.00	0.06
205366_s_at	HOXB6	6.30	1.14	0.18	2.20	0.00	0.07
210425_x_at	GOLGA8A /// GOLGA8B /// LOC100508892	6.29	1.06	0.17	2.08	0.00	0.07
236738_at	C3orf80	6.25	1.28	0.20	2.42	0.00	0.07
204438_at	MRC1	6.15	1.23	0.20	2.34	0.00	0.08
212070_at	GPR56	6.11	0.96	0.16	1.95	0.00	0.08

219304_s_at	PDGFD	6.10	0.92	0.15	1.89	0.00	0.08
204082_at	PBX3	6.08	1.30	0.21	2.46	0.00	0.08
210424_s_at	GOLGA8A /// GOLGA8B /// LOC100508892	6.00	0.83	0.14	1.78	0.00	0.09
204030_s_at	IQCJ-SCHIP1 /// SCHIP1	5.95	1.32	0.22	2.49	0.00	0.09
211269_s_at	IL2RA	5.92	0.70	0.12	1.62	0.00	0.10
208029_s_at	LAPTM4B	5.91	1.21	0.20	2.31	0.00	0.10
209014_at	MAGED1	5.90	0.72	0.12	1.65	0.00	0.10
214039_s_at	LAPTM4B	5.86	1.50	0.26	2.84	0.00	0.10
232979_at	MIR10A	5.82	0.65	0.11	1.57	0.00	0.10
242269_at	FLJ42875	5.80	0.69	0.12	1.61	0.00	0.10
213844_at	HOXA5	5.78	1.45	0.25	2.74	0.00	0.10
201427_s_at	SEPP1	5.74	1.72	0.30	3.29	0.00	0.10
241464_s_at	---	5.70	0.83	0.14	1.77	0.00	0.10
205190_at	PLS1	5.69	0.79	0.14	1.73	0.00	0.10
239237_at	LOC100506776	5.61	0.85	0.15	1.80	0.00	0.11
1554679_a_at	LAPTM4B	5.56	1.15	0.21	2.21	0.00	0.10
205227_at	IL1RAP	5.48	0.77	0.14	1.70	0.00	0.10
204485_s_at	TOM1L1	5.46	0.71	0.13	1.64	0.00	0.10
204224_s_at	GCH1	5.45	1.05	0.19	2.07	0.00	0.10
208798_x_at	GOLGA8A	5.44	1.04	0.19	2.05	0.00	0.10
1559597_at	---	5.42	0.94	0.17	1.93	0.00	0.10
243010_at	MSI2	5.40	0.79	0.15	1.72	0.00	0.09
236553_at	LOC100507520	5.38	0.68	0.13	1.60	0.00	0.09
208767_s_at	LAPTM4B	5.38	0.92	0.17	1.89	0.00	0.09
201328_at	ETS2	5.37	0.77	0.14	1.70	0.00	0.09
225237_s_at	MSI2	5.36	1.03	0.19	2.05	0.00	0.09
211031_s_at	CLIP2	5.36	0.80	0.15	1.75	0.00	0.09
239232_at	MSI2	5.36	0.76	0.14	1.69	0.00	0.09
208791_at	CLU	5.32	1.07	0.20	2.10	0.00	0.09
235521_at	HOXA3	5.30	1.22	0.23	2.33	0.00	0.08
226134_s_at	---	5.28	1.03	0.19	2.04	0.00	0.08
230670_at	IGSF10	5.28	0.71	0.13	1.63	0.00	0.08
226098_at	IFT80	5.24	0.64	0.12	1.55	0.00	0.07
235900_at	SPNS3	5.22	0.70	0.13	1.62	0.00	0.07
208792_s_at	CLU	5.22	0.99	0.19	1.98	0.00	0.07
1558956_s_at	IFT80	5.20	0.64	0.12	1.55	0.00	0.06
210365_at	LOC100506403 /// RUNX1	5.20	0.74	0.14	1.67	0.00	0.06
232088_x_at	LOC100271722	5.18	0.84	0.16	1.79	0.00	0.06
225240_s_at	MSI2	5.16	0.96	0.19	1.95	0.00	0.06

207223_s_at	PTBP3	5.16	0.57	0.11	1.48	0.00	0.05
214697_s_at	PTBP3	5.14	0.61	0.12	1.52	0.00	0.05
226670_s_at	PABPC1L	5.12	0.64	0.12	1.56	0.00	0.04
1559987_at	---	5.10	0.72	0.14	1.65	0.00	0.04
206310_at	SPINK2	5.03	1.23	0.24	2.35	0.00	0.02
209905_at	HOXA10-HOXA9 /// HOXA9	5.03	1.34	0.27	2.53	0.00	0.01
219218_at	BAHCC1	4.99	0.76	0.15	1.69	0.00	0.00
221942_s_at	GUCY1A3	4.99	0.87	0.18	1.83	0.00	0.00
202660_at	ITPR2	4.99	0.70	0.14	1.62	0.00	0.00
228365_at	CPNE8	4.98	1.06	0.21	2.09	0.00	0.00
206945_at	LCT	4.97	0.51	0.10	1.42	0.00	0.00
212820_at	DMXL2	4.94	0.97	0.20	1.96	0.00	0.00
227400_at	NFIX	4.93	0.58	0.12	1.49	0.00	0.00
1566764_at	MACC1	4.92	0.78	0.16	1.72	0.00	0.00
209409_at	GRB10	4.86	0.80	0.17	1.75	0.00	0.00
202862_at	FAH	4.85	0.60	0.12	1.51	0.00	0.00
227235_at	GUCY1A3	4.84	1.01	0.21	2.02	0.00	0.00
230743_at	HOXB-AS3	4.83	0.51	0.11	1.43	0.00	0.00
224901_at	SCD5	4.83	0.72	0.15	1.65	0.00	0.00
243579_at	MSI2	4.82	0.51	0.11	1.42	0.00	0.00
237246_at	---	4.80	0.62	0.13	1.53	0.00	0.00
214651_s_at	HOXA10-HOXA9 /// HOXA9 /// MIR196B	4.79	1.57	0.33	2.97	0.00	0.00
1553311_at	C20orf197	4.78	0.72	0.15	1.65	0.00	0.00
206298_at	ARHGAP22	4.76	0.67	0.14	1.59	0.00	0.00
238058_at	LOC150381	4.76	0.65	0.14	1.57	0.00	0.00
237594_at	---	4.75	0.68	0.14	1.60	0.00	0.00
212646_at	RFTN1	4.75	0.58	0.12	1.50	0.00	0.00
201069_at	MMP2	4.75	0.88	0.19	1.84	0.00	0.00
1557527_at	---	4.74	0.68	0.14	1.60	0.00	0.00
208797_s_at	GOLGA8A	4.72	0.56	0.12	1.48	0.00	0.00
200670_at	XBP1	4.72	0.51	0.11	1.43	0.00	0.00
202662_s_at	ITPR2	4.71	0.50	0.11	1.42	0.00	0.00
201329_s_at	ETS2	4.68	0.63	0.13	1.55	0.00	0.00
241342_at	TMEM65	4.68	0.71	0.15	1.63	0.00	0.00
206582_s_at	GPR56	4.67	0.56	0.12	1.47	0.00	0.00
207034_s_at	GLI2	4.66	0.48	0.10	1.40	0.00	0.00
206478_at	KIAA0125	4.65	1.01	0.22	2.01	0.00	0.00
201242_s_at	ATP1B1	4.64	0.97	0.21	1.96	0.00	0.00
206574_s_at	PTP4A3	4.63	0.61	0.13	1.52	0.00	0.00
206950_at	SCN9A	4.61	0.58	0.13	1.50	0.00	0.00

1552364_s_at	MSI2	4.57	0.62	0.13	1.53	0.00	0.00
209386_at	TM4SF1	4.56	0.75	0.16	1.68	0.00	0.00
210145_at	PLA2G4A	4.56	0.76	0.17	1.69	0.00	0.00
229530_at	GUCY1A3	4.56	0.82	0.18	1.77	0.00	0.00
1555037_a_at	IDH1	4.56	0.68	0.15	1.60	0.00	0.00
212364_at	MYO1B	4.55	0.54	0.12	1.46	0.00	0.00
213217_at	ADCY2	4.55	0.49	0.11	1.40	0.00	0.00
213395_at	MLC1	4.54	0.74	0.16	1.67	0.00	0.00
213413_at	STON1	4.54	0.54	0.12	1.45	0.00	0.00
212276_at	LPIN1	4.51	0.54	0.12	1.46	0.00	0.00
201487_at	CTSC	4.50	0.51	0.11	1.42	0.00	0.00
239580_at	GUCY1A3	4.50	0.63	0.14	1.55	0.00	0.00
205653_at	CTSG	4.50	1.08	0.24	2.11	0.00	0.00
205600_x_at	HOXB5	4.50	0.51	0.11	1.43	0.00	0.00
225647_s_at	CTSC	4.49	0.58	0.13	1.50	0.00	0.00
219412_at	RAB38	4.47	0.53	0.12	1.44	0.00	0.00
201193_at	IDH1	4.46	0.60	0.14	1.52	0.00	0.00
231838_at	PABPC1L	4.46	0.49	0.11	1.41	0.00	0.00
209360_s_at	LOC100506403 /// RUNX1	4.45	0.52	0.12	1.44	0.00	0.00
206067_s_at	WT1	4.44	0.74	0.17	1.67	0.00	0.00
217620_s_at	PIK3CB	4.39	0.48	0.11	1.40	0.00	0.00
226065_at	PRICKLE1	4.39	0.59	0.13	1.50	0.00	0.00
205899_at	CCNA1	4.38	1.05	0.24	2.07	0.00	0.00
213150_at	HOXA10	4.38	1.06	0.24	2.09	0.00	0.00
235753_at	HOXA7	4.37	0.76	0.17	1.69	0.00	0.00
202932_at	YES1	4.37	0.67	0.15	1.59	0.00	0.00
209193_at	PIM1	4.35	0.54	0.12	1.46	0.00	0.00
202948_at	IL1R1	4.35	0.61	0.14	1.53	0.00	0.00
225233_at	MSI2	4.34	0.60	0.14	1.51	0.00	0.00
212386_at	TCF4	4.34	0.85	0.20	1.80	0.00	0.00
205624_at	CPA3	4.34	1.13	0.26	2.19	0.00	0.00
217975_at	WBP5	4.34	0.91	0.21	1.88	0.00	0.00
223939_at	SUCNR1	4.33	0.90	0.21	1.87	0.00	0.00
229199_at	SCN9A	4.32	0.51	0.12	1.43	0.00	0.00
229971_at	GPR114	4.31	0.53	0.12	1.45	0.00	0.00
209537_at	EXTL2	4.30	0.49	0.11	1.40	0.00	0.00
202933_s_at	YES1	4.28	0.65	0.15	1.57	0.00	0.00
205518_s_at	CMAHP	4.28	0.66	0.15	1.58	0.00	0.00
244881_at	LMLN	4.28	0.46	0.11	1.38	0.00	0.00
1559214_at	---	4.26	0.74	0.17	1.68	0.00	0.00
244457_at	---	4.25	0.61	0.14	1.53	0.00	0.00

239963_at	---	4.25	0.75	0.18	1.69	0.00	0.00
202718_at	IGFBP2	4.22	0.86	0.20	1.82	0.00	0.00
1558770_a_at	PIK3R6	4.21	0.46	0.11	1.37	0.00	0.00
219602_s_at	PIEZO2	4.20	0.59	0.14	1.51	0.00	0.00
204069_at	MEIS1	4.20	0.95	0.23	1.93	0.00	0.00
205582_s_at	GGT5	4.19	0.42	0.10	1.34	0.00	0.00
1559067_a_at	LOC158402	4.19	0.54	0.13	1.45	0.00	0.00
222303_at	---	4.18	0.66	0.16	1.58	0.00	0.00
206847_s_at	HOXA7	4.18	0.49	0.12	1.40	0.00	0.00
238658_at	---	4.17	0.43	0.10	1.35	0.00	0.00
201243_s_at	ATP1B1	4.17	0.61	0.15	1.53	0.00	0.00
223075_s_at	AIF1L	4.16	0.67	0.16	1.59	0.00	0.00
221781_s_at	DNAJC10	4.15	0.43	0.10	1.35	0.00	0.00
212274_at	LPIN1	4.14	0.45	0.11	1.37	0.00	0.00
230064_at	---	4.11	0.56	0.14	1.47	0.00	0.00
219036_at	CEP70	4.11	0.62	0.15	1.54	0.00	0.00
214228_x_at	TNFRSF4	4.11	0.46	0.11	1.38	0.00	0.00
1560500_at	---	4.09	0.49	0.12	1.40	0.00	0.00
226784_at	TWISTNB	4.09	0.54	0.13	1.45	0.00	0.00
201952_at	ALCAM	4.09	0.63	0.15	1.55	0.00	0.00
232151_at	MACC1	4.08	0.51	0.13	1.42	0.00	0.00
202661_at	ITPR2	4.08	0.46	0.11	1.37	0.00	0.00
242172_at	---	4.06	0.64	0.16	1.55	0.00	0.00
242476_at	---	4.04	0.67	0.17	1.59	0.00	0.00
209695_at	PTP4A3	4.04	0.46	0.11	1.38	0.00	0.00
226069_at	PRICKLE1	4.03	0.57	0.14	1.48	0.00	0.00
1552623_at	HSH2D	4.03	0.65	0.16	1.57	0.00	0.00
222146_s_at	TCF4	4.03	0.64	0.16	1.55	0.00	0.00
236488_s_at	---	4.03	0.48	0.12	1.40	0.00	0.00
212387_at	TCF4	4.02	0.64	0.16	1.56	0.00	0.00
213891_s_at	TCF4	4.01	0.67	0.17	1.59	0.00	0.00