



LUND UNIVERSITY

Parmelia ernstiae - new to the Nordic lichen flora.

Thell, Arne

Published in:
Graphis Scripta

2003

Document Version:
Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (APA):
Thell, A. (2003). Parmelia ernstiae - new to the Nordic lichen flora. *Graphis Scripta*, 14(1), 10-10.

Total number of authors:
1

Creative Commons License:
Other

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Parmelia ernstiae – new to the Nordic lichen flora

ARNE THELL

Parmelia ernstiae – new to the Nordic lichen flora. *Graphis Scripta 14*: 10. Stockholm ISSN 0901-7593.

Parmelia ernstiae Feuerer & A. Thell, recently described from northwestern Germany, is reported from one locality in central Skåne, southernmost Sweden.

Arne Thell, Lund University, Botanical Museum, Östra Vallgatan 18, SE-223 61 Lund. E-mail: Arne.Thell@sysbot.lu.se

Parmelia ernstiae Feuerer & A. Thell is one of the few recently described macrolichens from Europe. It is distinguished from its closest relative, *Parmelia saxatilis* Ach., by a pruinose upper surface, including pruinose isidia, and an irregular, lacinate, strictly corticolous habit. The status as separate species was confirmed by unique DNA-sequences, most distinctly observed in the rDNA ITS region (ITS1-5.8S-ITS2), where the sequences differed at eight positions between the two species. ITS-sequences of *P. saxatilis* populations were identical, irrespective of their geographical origin (Feuerer & Thell 2002). However, a contemporary molecular study indicate further taxa within *Parmelia saxatilis* – in addition to *P. ernstiae* (Crespo et al. 2002).

P. ernstiae was until now known from northwestern Germany only, where it grows on bark of deciduous trees, preferably *Fraxinus excelsior*, in open or half-open agricultural areas. It is here reported from one locality in southernmost Sweden:

Sweden: Skåne: Eslöv par., 800 m SE Grässjåhus, on *Fraxinus excelsior* at the road, alt. 100 m, 17 February 2001, A. Thell 0101 (HBG), DNA865 (HBG, LD); 1 March 2003, A. Thell 0302 (LD, UPS).

Results

The rDNA ITS sequence of the Swedish sample (AY247007) was compared with the two German *P. ernstiae* sequences available in the GenBank, <http://www.ncbi.nlm.nih.gov>. It was identical with one of these (AF410834), collected in Schleswig-Holstein (T. Feuerer & A. Thell 64330 HBG), whereas the second sequence (AF410833), from the type material from Niedersachsen (G. Ernst HBG-4619), differed by absence of two nucleotides.

References

- Crespo, A., Molina, M. C., Blanco, O., Schroeter, B., Sancho, L. G. & Hawksworth, D. L. 2002. rDNA ITS and β -tubulin gene sequence analyses reveal two monophyletic groups within the cosmopolitan lichen *Parmelia saxatilis*. *Mycological Research* 106: 788-795.
- Feuerer, T. & Thell, A. 2002. *Parmelia ernstiae* – a new macrolichen from Germany. *Mitteilungen aus dem Institut für Allgemeine Botanik Hamburg* 30-32: 49-60.