



LUND UNIVERSITY

Liquid in accelerated motion

Pendrill, Ann-Marie; Fägerlind, Carl-Olof

Published in:
Physics Education

DOI:
[10.1088/0031-9120/50/6/648](https://doi.org/10.1088/0031-9120/50/6/648)

2015

[Link to publication](#)

Citation for published version (APA):
Pendriill, A.-M., & Fägerlind, C.-O. (2015). Liquid in accelerated motion. *Physics Education*, 50(6), 648-650.
<https://doi.org/10.1088/0031-9120/50/6/648>

Total number of authors:
2

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Liquid in accelerated motion

Carl-Olof Fägerlind¹ and Ann-Marie Pendrill²

1) Krusbärsvägen 4, SE 168 59 Bromma, Sweden

2) National Resource Centre for Physics Education, Lund University, Box 118, SE 221 00 Lund, Sweden

E-mail: C.Fagerlind@gmail.com and Ann-Marie.Pendrill@fysik.lu.se



Figure 1. A glass of cordial, resting on wooden triangle on a string

Abstract. What happens to a liquid in a swing or a loop? Accelerated motions lead to effects that may seem surprising. Consider a liquid in a glass placed on a wooden triangle, in turn attached to a string and moving fast in circle a vertical plane. The surface of the water remains parallel to the bottom of the glass - and orthogonal to the string - as long as gravity and the tension in the string are the only forces acting on the triangle. A key to the understanding is that the tangential acceleration of the liquid, the glass and the triangle, are all identical to the tangential component of the acceleration of gravity.

This is an author-created, un-copied version of an article in *Physics Education* **50** 648 (2015). IOP Publishing Ltd is not responsible for any errors or omissions in this version of the manuscript or any version derived from it. The Version of Record is available online at <http://iopscience.iop.org/article/10.1088/0031-9120/50/6/648/pdf>

1. Introduction

Have you ever swung a bucket full of water over your head, and marvelled about not getting wet. Most likely, you heard explanations from adults mentioning the "centrifugal force" as responsible for "pushing the water" away from the centre of the circle. Still, your physics textbook is more likely to insist that the centrifugal force does not exist. The water in the bucket is then instead described as accelerating towards the ground with an acceleration larger than the acceleration of gravity, g . The photo in Figure 1



Figure 2. During rotation - and other motions - the surface of the liquid remains parallel to the bottom of the glass, orthogonal to the string

shows a setup for a beautiful demonstration for the brave: A glass with red cordial, placed on a wooden triangle, suspended from a string in one of the corners. With a determined move of the arm, the triangle can be made to move a full circle, while the liquid remains parallel to the bottom of the glass, as in figure 2. For the less confident,

an inch or so of coloured liquid in a closed bottle with a string will also do [1].

2. Force and acceleration

The only forces acting on the triangle with the glass are the force, \mathbf{T} , from the tension in the string and the force of gravity, $m\mathbf{g}$, as shown in Figure 3. The force from the string is always in the direction of the string, in the radial direction towards the centre of rotation, whereas the force of gravity can be divided into a tangential component $F_t = -mg \cos \theta$ and a radial component $F_r = -mg \sin \theta$, denoted by dashed yellow lines in figure 4, which also defines the angle θ . Similarly, the resulting acceleration can be divided into tangential and radial components, as in figure 4, giving $a_t = -g \cos \theta$ and $a_r = -g \sin \theta - T/m$.

For the glass to remain in the triangle, the string must be extended with some tension. In the highest point, this implies that the downward acceleration must be at least g . The centripetal acceleration is given by v^2/L , where v is the speed and L is the distance between the centre of the circle and the centre of mass of the triangle, implying a minimum speed at the top, $v \geq \sqrt{gL}$.

For an angle close to 30° , as in the photo, the glass is lower by $\Delta H = L/2$. This implies an increased speed and the centripetal acceleration will be larger by $2g\Delta H/L$, giving, $a_c = g + 2g\Delta H/L = 2g$. The radial component of the the force of gravity can only provide an acceleration $g \sin \theta \approx g/2$. The tension in the string in the photo must thus be at least $3mg/2$, to provide the required centripetal force. A faster rotation implies a larger centripetal force, and a larger tension in the string, while leaving the tangential components of force and acceleration unaffected.



Figure 3. During a sufficiently fast rotation the surface of the liquid remains parallel to the bottom of the glass, orthogonal to the string

3. Conclusion

The key to understanding of the behaviour of the liquid in figure 2 is to note that the only force acting in the direction of the tangent is the force of gravity. This component, $F_t = mg \cos \theta$, leads directly to an acceleration $a_t = g \cos \theta$. From within the accelerating triangle, it seems like gravity has been turned off in the tangential direction. This is analogous to free fall causing an experience of weightlessness. An accelerometer measuring the vector $\mathbf{a} - \mathbf{g}$ will show zero in the tangential direction (see e.g. [2, 3]). It is a consequence of the equivalence between inertial and gravitational mass, which has been explored e.g. in [4]

The triangle, the glass and the liquid in the glass share the same acceleration, and the force to be added to gravity must point to the centre of rotation for all of them. The net force from liquid on a small droplet in the glass must also be orthogonal to the surface of the liquid - which thus remains orthogonal to the string and parallel to the bottom of the glass. The orthogonality between the liquid surface and the string holds also for other motion, e.g. in a glass taken along in a waveswinger amusement ride, or in a glass or bottle put on a swing [1]. That experiment has also been adapted for Science Days at the Bakken amusement park in Denmark [5].




Figure 4. Radial and tangential components of force and acceleration for the triangle on the string

Acknowledgements

We would like to express our appreciation to Astrid Fägerlind for performing the demonstration.

References

- [1] Pendrill A-M and Williams G 2005 *Swings and Slides* Phys. Educ. 40 527
- [2] Pendrill A-M and Rohlén J 2011 *Acceleration and rotation in a pendulum ride, measured using an iPhone 4* Phys. Educ. 46 676
- [3] Vernier, *Wireless Dynamic Sensor System*, <http://www.vernier.com/products/sensors/wdss/>
- [4] Pendrill A-M, Ekström P, Hansson L, Mars P, Ouattara L and Ryan U 2014 *The equivalence principle comes to school - falling objects and other middle school investigations* Phys. Educ. 49 425
- [5] Bakken, *Natur og Fag*, (bakken.dk), and Ulrik Lundby Hansen, private communication

Biographies



Carl-Olof Fägerlind is a recently retired upper-secondary school physics teacher and also a singer. He enjoys creating physics demonstrations and is part of the annual Faraday Christmas lecture at Stockholm university.



Ann-Marie Pendrill is professor in Science Communication and Physics Education at Lund University, where she is also the director of the Swedish National Resource Centre for Physics Education. She enjoys using outdoor environments for physics education, e.g. to illustrate relations between force and acceleration. (Foto Maja-Kristin Nylander)